1

Title: The Goldilocks Effect: Female geladas in mid-sized groups have higher fitness

Running head: The Goldilocks Effect

Authors: Elizabeth Tinsley Johnson[1*], Jacob A. Feder[2*], Thore J. Bergman[3,4], Amy Lu[2,5], Noah Snyder-

Mackler[6,7**], Jacinta C. Beehner[3,8**]

Affiliations:

[1] Department of Integrative Biology, Michigan State University, East Lansing, MI 48824, USA.

[2] Interdepartmental Program in Anthropological Sciences, Stony Brook University, Stony Brook, NY

11794-4364, USA.

[3] Department of Psychology, University of Michigan, Ann Arbor, MI 48109-1043, USA.

[4] Department of Ecology & Evolution, University of Michigan, Ann Arbor, MI 48019-1085, USA.

[5] Department of Anthropology, Stony Brook University, Stony Brook, NY 11794-4364, USA.

[6] School of Life Sciences, Arizona State University, Tempe, AZ 85287-4701, USA.

[7] Center for Evolution and Medicine, Arizona State University, Tempe, AZ 85287-1701, USA.

[8] Department of Anthropology, University of Michigan, Ann Arbor, MI 48109-1107, USA.

[*] Co-first authors

[**] Co-senior authors

Corresponding authors: etinsley@msu.edu; nsnyderm@asu.edu; jbeehner@umich.edu

Abstract:

The cost-benefit ratio of group-living is thought to vary with group size: individuals in "optimally-sized"

groups should have higher fitness than individuals in groups that are either too large or too small.

However, the relationship between group size and individual fitness has been difficult to establish for

long-lived species where the number of groups studied is typically quite low. Here we present evidence

for optimal group size that maximizes female fitness in a population of geladas (*Theropithecus gelada*).

Drawing on 14 years of demographic data, we found that females in small groups experienced the

highest death rates, while females in mid-sized units exhibited the highest reproductive performance.

This group-size effect on female reproductive performance was largely explained by variation in infant

mortality (and, in particular, by infanticide from immigrant males) but not by variation in reproductive

rates. Taken together, females in mid-sized units are projected to attain optimal fitness due to

conspecific infanticide and, potentially, predation. Our findings provide insight into how and why group

size shapes fitness in long-lived species.

Keywords: optimal group size, fitness, folivore paradox, infanticide, reproduction, survival

INTRODUCTION

Variability in group size within species reflects a balance between the costs and benefits of group living [1,2]. For example, individuals living in large groups may experience high within-group foraging competition (a cost: e.g., [3–5]) but low predation risk (a benefit: e.g., [3,6–9]); or, high parasite loads (a cost: e.g., [10]) but improved resource defense (a benefit: e.g., [11]). Individuals living in optimally-sized groups, where (by definition) the benefits outweigh the costs, should have the highest lifetime reproductive success (i.e., fitness [12]) compared to others in the population [13]. Although the link between group size and fitness has long been the focus of socioecological theory, it has been challenging to demonstrate this relationship with empirical data. This is, in large part, because group size fluctuates over an individual's lifespan, making it difficult to align a specific group size with a clear fitness metric (especially in long-lived species). As a result, testing the effects of group size requires long-term data across multiple groups of variable size.

Ecological costs and benefits at both extremes of group size support the idea that mid-sized groups are optimal. For example, the feeding costs of living in large groups are well-documented: variables that indicate greater within-group feeding competition, such as home range size, day range length, and feeding time are often positively correlated with group size (carnivores: [14], primates: [15,16]), and those that reflect the consequences of such competition, such as body condition, fertility, and survival are often negatively correlated with group size [17–22]. Despite the costs of living in large groups, individuals are often found in groups significantly larger than the theoretical species optimum [23,24], likely because splitting into two smaller groups is equally, if not more, costly. This may be particularly true if small groups are more susceptible to predation [25,26] or if between-group competition pushes competitively weaker groups into inferior home ranges (carnivores: [27], primates: [28,29]). Perhaps as a consequence of costs at both extremes, individuals in both large and small groups exhibit higher

glucocorticoid levels, greater ranging disadvantages, and experience more extreme energetic conditions

when compared to individuals in mid-sized groups (e.g., [16,30–32], reviewed in [5]).

However, many species do not exhibit group sizes that conform to predictions based on ecological

factors (e.g. [33]). For example, although folivores are thought to experience low within-group feeding

competition, females in numerous folivorous taxa live in groups that are smaller than those found in

frugivorous species (the "folivore paradox": [34], e.g. [35]). For some folivorous species, within-group

feeding competition may still be high, thus limiting group size (primates: [36], elephants: [37], deer:

[38]). Alternatively (or additionally), in many folivorous primates, females may form smaller groups

because these groups are less attractive to immigrant males and thus less vulnerable to infanticide

[39,40]. By contrast, in some species, infanticide risk may favor larger groups that are better able to

prevent male immigration or defend dependent offspring [41]. As a result, conspecific threat of

infanticide has the potential to shape group size such that mid-sized groups are optimal, but isolating the

effects of such social factors from environmental factors has proven challenging.

Geladas are folivorous monkeys that feed on widely-dispersed food resources consisting primarily of

graminoid (grass and sedge) leaves [42,43]. Due to this ubiquitous resource base, geladas face low

within-group feeding competition [44] and (as predicted: [34]) live in large groups for predator

protection [45,46]. However, gelada social organization is not so simple: individuals form small

reproductive groups (hereafter, "units", varying in size from 1-12 adult females: [47]) which, in turn,

unite to form large bands (100+ individuals) that travel and forage together in the same home range

(forming a "multi-level" society: [46]). The large foraging band likely represents the ecological unit

(i.e., by offering some degree of protection from predation, [48]) and conforms to the predictions of

standard socioecological theory [34]. In contrast, the small reproductive unit represents the social unit

and, as such, is likely shaped by social factors. Specifically, in this population, male immigrations

(called "takeovers" because the new male replaces the dominant group male) dramatically alter female

reproductive patterns via infanticide and the Bruce effect [49,50]. Infanticide is the leading cause of

infant mortality: immigrant males can kill up to half of the dependent infants in a unit [51]. Moreover,

larger units experience more takeovers than smaller units [52].

Ecological factors like habitat quality are unlikely to vary significantly among units in the same band;

therefore, we predict that females in larger units incur disproportionately higher fitness costs due to male

takeovers compared to females in smaller units. Alternatively (or additionally), when a takeover occurs,

infants in small groups may have a higher chance of being killed due to poor infant defense. If both large

and small units incur fitness costs due to infanticide, then we predict that females in mid-sized units will

exhibit the highest fitness. Finally, adult female survival may also be affected by unit size: females in

small units may be more vulnerable to predation (e.g. [25]), while females in larger units may be more

vulnerable to disease (e.g. [53]). If both factors affect female longevity in geladas, then we predict the

lowest adult female death rates in mid-sized units.

To test these predictions, we first examined the evidence for an optimal unit size for gelada females.

Specifically, we tested if unit size predicted female death rates and/or reproductive performance (the

production of surviving infants). Second, because reproductive performance could be a product of both

female fertility and/or infant mortality, we explored the effect of unit size on these two components.

Specifically, we assessed variation in both interbirth intervals, which should track both female energetic

condition (e.g., [54,55], reviewed in [7]) and takeover-related fetal loss [49,56], and in infant mortality,

which is predominantly driven by infanticide for this population [51]. Third, we examined how the

cause of infant deaths varied by unit size, specifically considering the extent to which maternal death

and/or infanticide explained these patterns. Taken together, our results show that females in mid-sized

units appear to display optimal fitness, resulting from the combined effects of lower adult female death

rates and infanticides in these units.

MATERIALS AND METHODS

Study site and subjects

The data for this study derive from 14 years of observation (Jan 2006-Jul 2020) on a population of wild

geladas living in the Simien Mountains National Park, in northern Ethiopia (13°13.5' N latitude). The

Simien Mountains Gelada Research Project (SMGRP, formerly the University of Michigan Gelada

Research Project) has collected near-daily behavioral, demographic, genetic, and hormonal data from

individuals since Jan 2006. All gelada subjects are habituated to human observers on foot and are

individually recognizable. We used longitudinal data from 200 adult females in 46 reproductive units

(20 original "founding" units that later became 26 "daughter" units due to fissions).

Unit size

The identities of all individuals in a unit were recorded each day the unit was seen. For each month of

the study period, we recorded the total number of adult females in each unit; where changes in unit size

occurred (i.e., due to adult female deaths or subadult female maturations), we used the maximum

number of adult females in a unit in a given month. We focus on adult females because we have

longitudinal records of the number of adults in each unit for the entire study period but only started

recording the total number of individuals (including juveniles and infants) in 2012. However, when we

compared the number of adult females in a unit to the total number of individuals in a unit for the subset

of data where we have both (2012-2020), we found that both values were highly correlated (Pearson

correlation coefficient = 0.76, *p*-value = 2.2 x 10^{-16}).

137 Gelada units contain one reproductively dominant "leader male" as well as 0-3 "follower males" that

138 receive few reproductive opportunities but can deter male takeovers [47]. Because the number of

139 females and the number of males could have contrasting impacts on female fitness components, and

140 because the number of males was weakly correlated with the number of females (Pearson correlation

141 coefficient = 0.21, p-value = 2.2 x 10^{-16}), we did not include males when calculating group size. In all

142 relevant analyses, we considered males separately as covariates.

144 Changes in the number of adult females within a unit were primarily due to female maturations and

145 deaths. Maturations were recorded as the first observation of a sex skin swelling (details are outlined in

146 [57]). Deaths were recorded as the first day an adult female was no longer observed with a unit for more

147 than three consecutive encounters with that unit (and not observed in a different unit, as in the case of

148 transfers or fissions).

150 Dates of fissions, fusions, and female transfers were recorded as the first day the unit females were no

151 longer observed together and subsequently observed either in a separate daughter unit with a new leader

152 male (for fissions), together with non-unit females and a new leader male (for fusions), or associating

153 individually with non-unit females and a new leader male (for the rare cases of female dispersal). In all

154 cases, we immediately identified known females in daughter units or in new units following their

155 disappearance from their natal unit.

157 All models included female unit size as a continuous predictor variable (see supplementary figure S1 for

158 the observed distribution of unit sizes). However, for visualization purposes in our figures, we also

159 categorized units into small, medium (i.e., mid-sized), and large units based on the observed range in

160 unit sizes. Specifically, cut-offs were determined by calculating the tertiles of the observed monthly

distribution of sizes: "small" indicates units of less than or equal to 4 adult females; "mid-sized"

indicates units from 5-7 adult females; and "large" indicates units of 8 or more adult females.

Adult female death rates

For this analysis, we included data on 200 adult female geladas. All adult females had known or

estimated birth dates from which we calculated age. Dates of birth were known for 56 adult females

born during the study, while the remaining females' birth dates were estimated using their size at the

start of observation (n=44), back-calculated using life history milestones (i.e., maturation or first birth,

n=40), or extrapolated from their total number of infants (n=60). The mean age at the mid-point of the

study (2013) for all females was 11.86 +/- 5.14 SD years (range = 3.54 - 28.90 years). There was no

mean difference in female age across unit sizes (estimated or known; females estimate = -2.01 +/- 2.35

SE, *p*-value = 0.394), but large units skewed towards having more young females than small

(Kolmogorov-Smirnov D = 0.094, *p*-value = 0.049) and mid-sized units (D = 0.104, *p*-value = 0.012;

supplementary figure S2).

To assess whether the rate at which adult females died varied by unit size, we constructed a binomial

generalized linear mixed model (GLMM) using the *lme4* package (version 1.1.20: [58]) in R (version

3.6.0: [59]). Here, the unit of observation was the female-year (Jan 1-Dec 31, unless truncated by

maturation, death, or the start or end of the study period). The dependent binary variable indicated

whether the female died during the study year, which was modeled as a function of the following

predictors: unit size (the average number of females in the unit, including both the linear and the

quadratic term), the female's age at the start of the year (linear and quadratic), and the number of males

(i.e., the total number of adult follower males plus one leader male). We controlled for the repeated

measures of female, year, and unit as crossed random effects (due to unit fissions, females can reside in

multiple units during their lifetimes). Additionally, we constructed mixed effects Cox proportional

hazards models on adult female survival using the R package *coxme [60]*. For these models, we

incorporated uncertainty in female ages by sampling randomly from females' minimum and maximum

birth dates. Model averaged results from 1000 survival model simulations were congruent with the

results from the GLMM (supplementary table S6).

Reproductive performance

For the majority of infants born during the study period (n = 352 out of 394 total), the date of birth was

known within a day or two. For those infants where the exact date of birth was unavailable (n = 42), we

were able to assign a birth date within one month of the actual birth date based on established

morphological and behavioral criteria (i.e., the size of the infant, infant motor skills, presence/absence of

the umbilical cord; for more details, see [57]). Conception dates were retrospectively assigned by

subtracting the mean gestation length (n = 183 days; see [57]) in the population from the observed or

estimated date of birth.

We used a binary scoring system to assign female reproductive performance on a monthly basis.

Females were assigned a 1 in a given month if they conceived an infant that survived to 1.5 years of age,

which is the mean age of weaning in this population (regardless of actual weaning date). Females were

assigned a 0 if they conceived an offspring that died before reaching 1.5 years of age or if they did not

conceive during the calendar month. We excluded all months and births after 2018, as the survival

outcomes of these infants were not resolved by the end of the study period.

To assess the effects of unit size on female reproductive performance, we constructed a binomial

GLMM. The dependent binary variable indicated whether a female conceived a surviving offspring

during each female-month (273 surviving offspring, 996.5 female-years total, 188 females). We

modeled this outcome variable as a function of the following predictors: female age (both the linear and

quadratic term, to control for the known effects of female age on reproductive output, reviewed in [61]),

number of females, and the number of males. We controlled for the repeated measures of individual

identity, unit, and year as crossed random effects. To capture the effect of unit size variation beyond the

month of conception, we additionally modeled reproductive performance using 6-month bins (i.e., Jan-

Jun, Jul-Dec) that were centered on the two primary conception seasons [50]. We used the same binary

outcome listed above and included the average number of females and males as fixed effects. The results

from this separate analysis (supplementary table S7) were congruent with those generated from the

model using monthly data.

Given the long lifespans of geladas, we do not have the data necessary to directly calculate lifetime

reproductive success. However, we estimated this value by pairing median ages at death with

reproductive rates across the three unit size categories (i.e., small, medium, large). Specifically, we first

calculated median ages at death for adult females at each unit size category from survival curves

including right-censored females (supplementary figure S3). Second, we calculated reproductive rates

by summing the total number of surviving offspring within each unit size category and then dividing by

the total number of female-years. To approximate lifetime reproductive success, we subtracted the

population mean age at first birth (6.06 years; range: 4.88-7.56: [57]) from the median ages at death and

multiplied these reproductive lifespans by their corresponding reproductive rates. Although these static

estimates may not reflect true differences in lifetime reproductive success (because unit size changes

across the lifespans of individual females, especially following fissions and fusions), we use these

estimated fitness differences to underscore the potential costs of remaining at suboptimal unit sizes and

to highlight the need for female strategies to avoid these costs.

Interbirth intervals

Using the birth dates of all infants in this population, we calculated interbirth intervals (IBIs) between

females' successive offspring. We constructed two linear mixed models (LMM) using the *lme4* package.

For both models, the dependent variable was the length of the IBI in days, which was modeled as a

function of unit size at the first infant's birth (both linear and quadratic), the mother's age, and the

infant's sex. In the first model, we utilized only IBIs following surviving offspring (n = 187), as infant

mortality substantially shortens IBIs in this population and elsewhere [62]. However, because IBIs

where takeovers occur may capture instances of takeover-related fetal loss (i.e., the "Bruce effect":

[49]), which can lengthen IBIs [56], in the second model we only included the subset of IBIs that did not

contain takeovers or infant deaths (n = 82). Rather than reflecting fetal loss, variation in these "takeover-

independent" IBIs should reflect the ecological advantages (e.g., higher food intake) that some females

have over others in their ability to wean offspring and conceive again.

Infant mortality

The disappearance of any infant prior to the average age at weaning in this population (1.5 years of age:

[57]) was assumed to be a case of infant mortality. To assess infant mortality and its causes across unit

sizes, we constructed a mixed-effects Cox proportional hazards model using the R package *coxme* [60].

For this model, we used all 394 infants born during the study period. Infants entered the dataset at birth

and were censored at death (n = 90) or if their social units stopped being observed (n = 17) and exited

the dataset when they survived to 1.5 years of age (n = 287). We used the number of females present

during the month of the infant's birth as a fixed effect, including both linear and quadratic terms. We

included birth-year, unit, and maternal ID as random effects. Schoenfeld residuals showed no significant

deviations from the assumption of proportional hazards.

Takeover rates

We recorded the dates of all observed male takeovers ($n = 80$) of known reproductive units (following

[63]) as well as the number of mature females in the unit at the time of takeover. We calculated the

length of time each unit was seen at a given size for each year the unit was observed (244 total unit-

years) and counted the number of takeovers that occurred during these periods. To calculate the

influence of unit size (i.e., number of adult females) on the likelihood of takeover, we modeled the

number of takeovers a unit experienced as the dependent variable in a Poisson GLMM. Since takeovers

are more likely to occur during longer observation windows, we included the number of continuous

months the unit was seen at a given size as an offset term. To control for repeated measures across units

and time, we included unit and year as random intercepts. Finally, we included two fixed effects

variables: the number of adult females in the unit and the average number of males.

Causes of infant mortality

To examine whether the probable causes of infant mortality varied with unit size, we assigned the cause

of mortality based on the following characteristics: if the infant's death occurred within 6 months of a

takeover, the cause of death was recorded as "infanticide" ($n = 31$) [50,51]. If the infant died within 3

months of their mother's death or disappearance, the cause of death was recorded as "maternal death" (n

$= 27$). If maternal and infant disappearances co-occurred within the 6 months following a male takeover,

however, the cause of death was categorized as "maternal death" ($n = 4$). All other causes of infant

deaths were recorded as "unknown" ($n = 32$). For each of these mortality outcomes, we constructed a

binomial GLMM using the *lme4* package. Here, we modeled whether the infant died of each respective

mortality cause (i.e., maternal death, infanticide, and unknown) as a binary variable, using unit size

(both linear and quadratic) as a fixed effect and unit as a random effect. For these three analyses, we

used only infants whose survival outcomes were known (n = 377; i.e., no right-censored offspring). All

figures were constructed using *ggplot2* [64], and prediction intervals were extracted from mixed-models

using the *effects* package [65].

RESULTS

Adult female death rates

Of the 200 adult females included, 90 died before the end of the study period (median age at death =

16.2 years, including right-censored females). Unsurprisingly, the odds of dying increased substantially

with age (age estimate = 30.07 +/- 4.06 SE, p-value = 2.1 x 10^{-5}). Females in small units had the highest

death rates (females estimate = -9.39 +/- 4.78 SE, p-value = 0.049, figure 1a, supplementary table S1).

Specifically, females in small units had an annual death rate of 10.4%, while females in medium and

large units had an annual death rate of 5.8% and 7.3% respectively. Including right-censored data, the

median age at death was 16.0 years in small units, 19.7 years in mid-sized units, and 16.0 years in large

units.

Reproductive performance

Although reproductive units in this population can range in size from 1-12 adult females, females in

units in the middle of this range (i.e., between 5-7 adult females, hereafter "mid-sized" units) had the

highest reproductive performance. Females in mid-sized units were most likely to conceive surviving

offspring (females^{2} estimate = -18.14 +/- 7.69 SE, p-value = 0.018, figure 1c, supplementary table S1)

and were 17.6% more likely than females in small units and 39.8% more likely than females in large

units to produce a surviving offspring. Given the median ages at death calculated above, females are

projected to produce approximately 4.35 surviving infants in mid-sized units, compared to 2.70 infants

in small units, and 2.27 infants in large units.

Figure 1. Female fitness is optimal in mid-sized units. (a) Females in mid-sized units had the lowest annual probability of death, calculated as the number of deaths observed at each unit size divided by the total number of female-years. Confidence bands show the lower and upper limits of the model predictions, with colors following the unit size categories described in *Methods*. The size of each point is proportional to the number of female-years observed at each unit size. **(b)** Females in mid-sized units showed the highest rates of conceiving surviving offspring. Points indicate yearly rates of conceiving surviving offspring at each unit size, generated from raw data. The size of each point is proportional to the number of female-years observed at each unit size. Confidence bands show the lower and upper limits of the model predictions. Although the model output specified monthly probabilities, we converted these to annual probabilities for the sake of visualization.

Interbirth intervals

Interbirth intervals tended to be slightly longer in larger units (females estimate = 492.6 +/- 242.6 SE, *p*-value = 0.044; females^2 estimate = 388.9 +/- 226.4 SE, *p*-value = 0.088; figure 2a). However, this effect was no longer significant when only IBIs that were uninterrupted by takeovers were included for analysis (females estimate = 39.2 +/- 230.3 SE, *p*-value = 0.867; females^2 estimate = 376.8 +/- 224.8 SE, *p*-value = 0.107; figure 2b, supplementary table S2).

Infant mortality

Out of 394 infants, 90 (22.8%) died before reaching 1.5 years. Infants born into small and large units were more likely to die before this age than infants born into medium-sized units (HR=1.06 +/- 0.017 SE, *p*-value = 0.0014; figure 2c, supplementary table S3). Specifically, 28.1% and 33.8% of infants born

328   into small and large units died before weaning, while only 10.4% of infants born into mid-sized units

329   died before reaching this age.

**Figure 2. Unit size does not influence takeover-independent interbirth intervals, but does impact infant
survival. (a)** Including intervals where takeovers occurred, females in large units show slightly longer interbirth
intervals. Colored bands show the 95% confidence interval from the linear mixed model. **(b)** Excluding cases
where takeovers occurred, there is no relationship between unit size and interbirth interval. **(c)** Infant survival was
much higher in mid-sized units than in small and large units. Unit size was modeled as a continuous variable, and
colored categories here were used for visualization purposes only.

Takeover rates

340   Takeover frequency increased with unit size (females estimate = 7.33 +/- 2.48 SE, *p*-value = 0.003,

341   figure 3a, supplementary table S4). On average, small units experienced male takeovers once every 4.15

342   years, mid-sized units once every 3.17 years, and large units once every 1.61 years. However, the

343   number of males did not influence the frequency of takeovers (males estimate = -0.08 +/- 0.11 SE, *p*-

344   value = 0.458, supplementary table S4).

Causes of infant mortality

Infanticides and maternal death accounted for 38.9% and 25.6% of all dependent infant deaths respectively. Despite the fact that small units experienced the lowest takeover rates, females in mid-sized units experienced the lowest infanticide rates (females^2 estimate = 10.81 +/- 3.72 SE, *p*-value = 0.004, figure 3b, supplementary table S5): 9.1% and 12.8% of infants born into small and large units experienced infanticide respectively, while only 3.0% of infants in mid-sized units experienced infanticide.

Second, infants in mid-sized units were about half as likely to die as a result of maternal death as females in small and large units (females^2 estimate = 5.25 +/- 3.38 SE, *p*-value = 0.12, figure 3b, supplementary table S5), though this effect was not significant. Specifically, 9.1% and 8.3% of infants born into small and large units died following their mothers' deaths, while only 4.5% of infants in mid-sized groups died under these circumstances. Lastly, there was no association between unit size and infant mortalities of unknown cause.

Fig. 3. Takeovers and the causes of infant mortality vary with unit size. (**a**) Annual takeover rates increased with unit size. The dotted line represents the population mean (0.341 takeovers per unit-year). (**b**) Infants in small and large units were more likely to die of infanticide, but not maternal death or unknown causes. For the plots above, unit size categories are used solely for visualization purposes.

DISCUSSION

DISCUSSION

Taken together, our results show that females in mid-sized units have the highest fitness. Females in

smaller units die at the highest rates and females in both smaller and larger units have lower

reproductive performance, primarily as a result of takeover-related infanticide. Specifically, male

takeovers are more frequent in larger units and are more likely to lead to infanticide when they do occur

in smaller units. We also show that infant death rates, but not takeover-independent female fertility (i.e.,

uninterrupted interbirth intervals), vary by unit size, with the lowest infant death rates occurring in mid-

sized units. As a result, females in mid-sized units demonstrate the highest reproductive performance

and, when factoring in variation in adult female death rates, are expected to produce more offspring over

the course of their reproductive lifespans than are females in small or large units.

Infants in small and large units were more likely to die from infanticide than infants in mid-sized units.

Conspecific threat of infanticide has been suggested to limit group sizes in folivores (i.e., the "folivore

paradox" [39]). However, larger groups have also been suggested to act as a deterrent to male takeover

and infanticide [41]. Here we show that both forces may work in conjunction to result in optimal mid-

sized groups. Infanticide in large units was driven by high rates of male takeovers. By contrast, we are

less certain why infanticide in small units was also high, despite infrequent takeovers. One possibility is

that small units have poor infant defense (e.g. as suggested in gray langurs: [41]). Another possibility is

that new males may be more likely to commit infanticide in smaller units because smaller units present

fewer immediate reproductive opportunities (i.e., receptive, cycling females) than larger units. A third

possibility is that infanticide risk may follow a "dilution effect" (similar to predation risk: [66]), where

infants in large units are buffered by an increased number of potential targets.

Our proxy for ecological factors, interbirth interval [7], did not explain variation in reproductive

performance across unit sizes (after removing cases where male-mediated fetal loss may have occurred).

This is not surprising given the relatively dispersed and non-monopolizable diet of geladas. This

analysis, however, did not consider whether unit size interacts with other metrics that might track

ecological advantages, such as reproductive aging, age at maturation, or age at first birth. Based on a

recent analysis on female maturations for this population [67], we expect that females in small units will

have later ages at maturation and first birth to avoid inbreeding (since their fathers have longer tenures),

and we expect that females in large units will have earlier ages at maturation and first birth due to the

"Vandenbergh effect" that accompanies male takeovers. However, even if these predictions are

supported in future analyses, we do not expect these unit-size differences to significantly alter our

reproductive success estimates for two reasons: first, the observed range in age at first birth in this

population is narrow [57]; and second, in similarly long-lived primates, age at first birth had a very

modest impact on variation in lifetime fitness [68,69]. Future analyses that address the potential for

different female reproductive strategies, which require more complete female lifespans than we have

currently, will allow us to determine the effects of unit size on realized lifetime reproductive success.

Adult female death, which contributes to low lifetime reproductive success, was highest in small units.

Although female deaths could be triggered by takeover-related injuries [56], smaller units experienced

less frequent takeovers than mid-sized or large units, and we did not observe an increase in adult female

deaths after takeovers (in contrast to infant deaths, which do increase following takeovers: [56]).

Females in small units may be more likely to die if females in larger units monopolize food resources.

Indeed, strong between-group competition has been shown in other folivorous species (e.g., [70,71]);

however, in geladas, feeding-related aggression between units is relatively rare and low-intensity [72].

Moreover, if between-group competition were strong, we would expect females in the smallest units to

show the longest interbirth intervals. We think the most likely explanation is that females in small units

are more vulnerable to predation. Although the large foraging bands characteristic of gelada social

structure likely act as a deterrent to predators [48], large units may monopolize more central locations

within the band during the day or sleeping positions on the cliffs at night, thus leaving small units

vulnerable at the periphery. This question will require simultaneous targeted data collection on relative

group position - a focus of our future research.

What, if anything, can females in suboptimal groups do? In broad strokes, they have three theoretical

choices: they can disperse to a new unit entirely, they can add females to their too-small units, or they

can subtract females from their too-large units. In this population, female dispersal is rare (n = 2 across

the entire study period; 0.008 per unit-year); reproductive units consist of close maternal kin, and

females tend to remain in the same unit they were born into [73]. Adding females can be accomplished

through demographic processes (maturation); however, small units, given their lower reproductive

performance and high adult female death rates, require more time to grow than mid-sized or large units.

Some small units may never reach an optimal size, or, worse, could become "sinks", at risk of dying out

entirely (and we have seen at least one of our units go extinct). This raises the possibility that unit size

and female fitness are bidirectionally related: units grow or shrink according to female reproductive

performance and survival, while group size itself can also impose fitness costs, creating a feedback loop.

Therefore, for some small units, a more effective way to add females may involve joining with another

unit ("fusions"). Fusions are rare (n = 3 across the entire study period; 0.012 per unit-year) and may

involve a number of hurdles (e.g., resistance of leader males and/or unrelated females), but these

observations suggest that this is nevertheless a viable option for gelada females.

Although some small units may become sinks, overall demographic analyses from this population

indicate that the stochastic growth rate is 3.5% (Sloan et al., in revision). Thus, the more common

problem (for mid-sized and large units) is that they continue to grow in size. Subtracting females from

units can occur when one group splits permanently into two daughter groups ("fissions"), which we

routinely observe for large units (n = 11; 0.045 per unit-year). Compared to other primate taxa, gelada

social organization may offer more opportunities for fissions to occur, as the band may buffer the

ecological costs of fissions. In sum, unit size changes through frequent, slow, and small demographic

processes (i.e., maturations and deaths) in addition to rare, rapid, and large group dynamics (i.e., fusions

and fissions). Future analyses will focus on the relative importance that these events have in structuring

groups and on identifying the social network metrics that anticipate changes in group dynamics.

Takeover frequency was positively associated with the number of females in a unit (as previously

reported in this population [52]), but was not associated with the number of males in a unit. This stands

in contrast to our previous research showing that multimale units (consisting of a leader male and 1+

follower males) experienced fewer takeovers per unit-year than units with only one adult leader male

across all unit sizes [47]. In geladas, the consequences of males for female fitness may depend on the

reproductive strategies of individual follower males and on the ability of leader males to regulate group

membership (rather than solely on quantity of males), which require further data and analysis. Indeed, in

a number of primates, group composition (e.g., a higher ratio of males to females) has been shown to

reduce the risk of takeovers and/or infanticide (e.g. [74], reviewed in [75]). For example, reproductive

success of female white-faced capuchins (*Cebus capucinus*) was both negatively correlated with group

size and positively correlated with the ratio of males to females in the group [76]. In contrast, multimale

groups of ursine colobus monkeys (*Colobus vellerosus*) had higher rates of male immigration and

infanticide, perhaps because these dominant males were ineffective at deterring immigrants [40].

Whether female geladas in small or large *multi-male* units demonstrate higher reproductive performance

than females in small or large *single-male* units remains to be seen.

The link between group size and individual fitness has long been the focus of socioecological theory, yet

it has been difficult to demonstrate this relationship for long-lived species with empirical data. Here, we

leveraged long-term data from a population of wild geladas to provide insight into how and why group

size shapes female fitness – specifically, adult female deaths and conspecific infanticide.

ACKNOWLEDGEMENTS

We would like to thank the Ethiopian Wildlife Conservation Authority (EWCA) and the wardens and

staff of the Simien Mountain National Park for their permission and ongoing support for our long-term

research project. Additionally, we are grateful to our excellent field team across the years, most

especially E. Jejaw, A. Fanta, S. Girmay, J. Jarvey, and M. Gomery for their assistance with field data

collection. We also owe thanks to A. Marshall, B. Dantzer, and J. Mitani for providing valuable

feedback on data analyses. **Funding:** This work was supported by: National Science Foundation (grant

numbers BCS-0715179, IOS-1255974, BCS-1340911, IOS-1854359; BCS-1723237; BCS-17123228;

BCS-2010309); Leakey Foundation (multiple grants); National Institutes of Health (grant number R00-

AG051764); National Geographic Society (grant numbers 8100-06, 8989-11, 50409R-18; the Fulbright

Scholar Program; Sigma Xi; American Society of Primatologists; Wildlife Conservation Society; and

the University of Michigan. **Author contributions:** ETJ, TJB, NSM, and JCB designed the study. ETJ,

AL, TJB, NSM, and JCB collected the data. ETJ, JAF, AL, TJB, NSM, and JCB analyzed the data. ETJ,

JAF, and NSM wrote the paper, with contributions and edits from all authors. ETJ, JAF, AL, TJB, NSM,

and JCB provided funding support. **Competing interests:** Authors declare no competing interests. **Data**

and materials availability: All data and code used in this analysis are available at:

https://github.com/GeladaResearchProject/TinsleyJohnson_Feder_et_al_Goldilocks_2021.

REFERENCES:

1.   Alexander RD. 1974 The evolution of social behavior. *Annu. Rev. Ecol. Syst.* **5**, 325–383.

2.   Terborgh J, Janson CH. 1986 The socioecology of primate groups. *Annu. Rev. Ecol. Syst.* **17**, 111–136.

3.   Janson CH, Goldsmith ML. 1995 Predicting group size in primates: foraging costs and predation risks. *Behav. Ecol.* **6**, 326–336.

4.   Grove M. 2012 Space, time, and group size: a model of constraints on primate social foraging. *Anim. Behav.* **83**, 411–419.

5.   Markham AC, Gesquiere LR. 2017 Costs and benefits of group living in primates: an energetic perspective. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372**, 20160239.

6.   Pulliam HR. 1973 On the advantages of flocking. *J. Theor. Biol.* **38**, 419–422.

7.   van Schaik CP. 1983 Why are diurnal primates living in groups? *Behaviour* **87**, 120–144.

8.   Whiteside MA, Langley EJG, Madden JR. 2016 Males and females differentially adjust vigilance levels as group size increases: effect on optimal group size. *Anim. Behav.* **118**, 11–18.

9.   Rasa OAE. 1989 The costs and effectiveness of vigilance behaviour in the Dwarf Mongoose: implications for fitness and optimal group size. *Ethol. Ecol. Evol.* **1**, 265–282.

10.   Rifkin JL, Nunn CL, Garamszegi LZ. 2012 Do animals living in larger groups experience greater parasitism? A meta-analysis. *Am. Nat.* **180**, 70–82.

11.   Wrangham RW. 1980 An ecological model of female-bonded primate groups. *Behaviour* **75**, 262–300.

12.   Clutton-Brock TH. 1988 *Reproductive success: studies of individual variation in contrasting breeding systems*. Chicago: University of Chicago Press.

13.   Giraldeau L-A. 1988 The stable group and the determinants of foraging group size. In *The ecology of social behavior* (ed CN Slobodchikoff), pp. 33–53. New York: Academic Press.

14.   Carbone C, Cowlishaw G, Isaac NJB, Rowcliffe JM. 2005 How far do animals go? Determinants of day range in mammals. *Am. Nat.* **165**, 290–297.

15.   Majolo B, de Bortoli Vizioli A, Schino G. 2008 Costs and benefits of group living in primates: group size effects on behaviour and demography. *Anim. Behav.* **76**, 1235–1247.

16.   Markham AC, Gesquiere LR, Alberts SC, Altmann J. 2015 Optimal group size in a highly social mammal. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 14882–14887.

17.   Gesquiere LR, Altmann J, Archie EA, Alberts SC. 2018 Interbirth intervals in wild baboons: environmental predictors and hormonal correlates. *Am. J. Phys. Anthropol.* **166**, 107–126.

18.   Clutton-Brock TH, Albon SD, Guinness FE. 1982 Competition between female relatives in a

matrilocal mammal. *Nature* **300**, 178–180.

19. Borries C, Larney E, Lu A, Ossi K, Koenig A. 2008 Costs of group size: lower developmental and
reproductive rates in larger groups of leaf monkeys. *Behav. Ecol.* **19**, 1186–1191.

20. McComb K, Moss C, Durant SM, Baker L, Sayialel S. 2001 Matriarchs as repositories of social
knowledge in African elephants. *Science* **292**, 491–494.

21. Waterman JM. 2006 Delayed maturity, group fission and the limits of group size in female Cape
ground squirrels (Sciuridae: *Xerus inauris*). *J. Zool.* **256**, 113–120.

22. Creel S, Creel NM. 2015 Opposing effects of group size on reproduction and survival in African
wild dogs. *Behav. Ecol.* **26**, 1414–1422.

23. Sibly RM. 1983 Optimal group size is unstable. *Anim. Behav.* **31**, 947–948.

24. Krause J, Ruxton GD. 2002 *Living in Groups*. Oxford: Oxford University Press.

25. Clutton-Brock TH, Gaynor D, McIlrath GM, Maccoll ADC, Kansky R, Chadwick P, Manser M,
Skinner JD, Brotherton PNM. 1999 Predation, group size and mortality in a cooperative mongoose,
Suricata suricatta. *J. Anim. Ecol.* **68**, 672–683.

26. Guindre-Parker S, Rubenstein DR. 2020 Survival benefits of group living in a fluctuating
environment. *Am. Nat.* **195**, 1027–1036.

27. Mosser A, Packer C. 2009 Group territoriality and the benefits of sociality in the African lion,
Panthera leo. *Anim. Behav.* **78**, 359–370.

28. Cheney DL, Seyfarth RM. 1987 The influence of intergroup competition on the survival and
reproduction of female vervet monkeys. *Behav. Ecol. Sociobiol.* **21**, 375–386.

29. Harris TR. 2006 Between-group contest competition for food in a highly folivorous population of
black and white colobus monkeys (*Colobus guereza*). *Behav. Ecol. Sociobiol.* **61**, 317–329.

30. Pride RE. 2005 Optimal group size and seasonal stress in ring-tailed lemurs (*Lemur catta*). *Behav.*
Ecol. **16**, 550–560.

31. Papageorgiou D, Farine DR. 2020 Group size and composition influence collective movement in a
highly social terrestrial bird. *Elife* **9**, e59902.

32. Williams CK, Lutz RS, Applegate RD. 2003 Optimal group size and northern bobwhite coveys.
Anim. Behav. **66**, 377–387.

33. Rudolph K, Fichtel C, Schneider D, Heistermann M, Koch F, Daniel R, Kappeler PM. 2019 One
size fits all? Relationships among group size, health, and ecology indicate a lack of an optimal
group size in a wild lemur population. *Behav. Ecol. Sociobiol.* **73**, 132.

34. Isbell LA. 1991 Contest and scramble competition: patterns of female aggression and ranging
behavior among primates. *Behav. Ecol.* **2**, 143–155.

35. Koenig A, Borries C. 2002 Feeding competition and infanticide constrain group size in wild

hanuman langurs. *Am. J. Primatol.* **57**, 33–34.

36. Snaith TV, Chapman CA. 2007 Primate group size and interpreting socioecological models: Do
folivores really play by different rules? *Evol. Anthropol.* **16**, 94–106.

37. Archie EA, Morrison TA, Foley CAH, Moss CJ, Alberts SC. 2006 Dominance rank relationships
among wild female African elephants, *Loxodonta africana*. *Anim. Behav.* **71**, 117–127.

38. Focardi S, Pecchioli E. 2005 Social cohesion and foraging decrease with group size in fallow deer
(*Dama dama*). *Behav. Ecol. Sociobiol.* **59**, 84–91.

39. Steenbeek R, van Schaik CP. 2001 Competition and group size in Thomass langurs (*Presbytis
thomasi*): the folivore paradox revisited. *Behav. Ecol. Sociobiol.* **49**, 100–110.

40. Teichroeb JA, Wikberg EC, Bădescu I, Macdonald LJ, Sicotte P. 2012 Infanticide risk and male
quality influence optimal group composition for *Colobus vellerosus*. *Behav. Ecol.* **23**, 1348–1359.

41. Treves A, Chapman CA. 1996 Conspecific threat, predation avoidance, and resource defense:
implications for grouping in langurs. *Behav. Ecol. Sociobiol.* **39**, 43–53.

42. Jarvey JC, Low BS, Pappano DJ, Bergman TJ, Beehner JC. 2018 Graminivory and fallback foods:
annual diet profile of geladas (*Theropithecus gelada*) living in the Simien Mountains National Park,
Ethiopia. *Int. J. Primatol.* **39**, 105–126.

43. Fashing PJ, Nguyen N, Venkataraman VV, Kerby JT. 2014 Gelada feeding ecology in an intact
ecosystem at Guassa, Ethiopia: variability over time and implications for theropith and hominin
dietary evolution. *Am. J. Phys. Anthropol.* **155**, 1–16.

44. Dunbar R, Dunbar P. 1975 *Social dynamics of gelada baboons*. Basel: S. Karger.

45. Pappano DJ, Snyder-Mackler N, Bergman TJ, Beehner JC. 2012 Social 'predators' within a
multilevel primate society. *Anim. Behav.* **84**, 653–658.

46. Snyder-Mackler N, Beehner JC, Bergman TJ. 2012 Defining higher levels in the multilevel
societies of geladas (Theropithecus gelada). *Int. J. Primatol.* **33**, 1054–1068.

47. Snyder-Mackler N, Alberts SC, Bergman TJ. 2012 Concessions of an alpha male? Cooperative
defence and shared reproduction in multi-male primate groups. *Proc. Biol. Sci.* **279**, 3788–3795.

48. Dunbar RIM. 1986 The social ecology of gelada baboons. In *Ecological aspects of social evolution*
(eds DI Rubenstein, RW Wrangham), pp. 332–351. Princeton: Princeton University Press.

49. Roberts EK, Lu A, Bergman TJ, Beehner JC. 2012 A Bruce effect in wild geladas. *Science* **335**,
1222–1225.

50. Tinsley Johnson E, Snyder-Mackler N, Lu A, Bergman TJ, Beehner JC. 2018 Social and ecological
drivers of reproductive seasonality in geladas. *Behav. Ecol.* **29**, 574–588.

51. Beehner JC, Bergman TJ. 2008 Infant mortality following male takeovers in wild geladas. *Am. J.
Primatol.* **70**, 1152–1159.

52. Bergman TJ, Ho L, Beehner JC. 2009 Chest color and social status in male geladas (*Theropithecus*
gelada). *Int. J. Primatol.* **30**, 791–806.

53. Ezenwa VO. 2004 Host social behavior and parasitic infection: a multifactorial approach.
Behavioral Ecology **15**, 446–454.

54. Koenig A, Borries C, Chalise MK, Winkler P. 1997 Ecology, nutrition, and timing of reproductive
events in an Asian primate, the Hanuman langur (*Presbytis entellus*). *J. Zool.* **243**, 215–235.

55. Emery Thompson M, Muller MN, Wrangham RW. 2012 The energetics of lactation and the return
to fecundity in wild chimpanzees. *Behav. Ecol.* **23**, 1234–1241.

56. Schneider-Crease I, Chiou KL, Snyder-Mackler N, Bergman TJ, Beehner JC, Lu A. 2020 Beyond
infant death: the hidden costs of male immigration in geladas. *Animal Behaviour* **159**, 89–95.

57. Roberts EK, Lu A, Bergman TJ, Beehner JC. 2017 Female reproductive parameters in wild geladas
(*Theropithecus gelada*). *Int. J. Primatol.* **38**, 1–20.

58. Bates D, Maechler M, Bolker B, Walker S, Christensen R, Singmann H, Others. 2019 Lme4:
Linear mixed-effects models using Eigen and S4 (Version 1.1-20).

59. R Core Team. 2020 R: A language and environment for statistical computing. R Core Team,
Vienna, Austria.

60. Therneau T. 2015 coxme: mixed effects Cox models. R package version 2.2-3. *URL: http://CRAN.*
R-project. org/package= coxme

61. Pusey A. 2012 Magnitude and sources of variation in female reproductive performance. In *The*
evolution of primate societies (eds JC Mitani, J Call, PM Kappeler, RM Palombit, JB Silk), pp.
343–366. Chicago: University of Chicago Press.

62. Beehner JC, Bergman TJ. 2008 Infant mortality following male takeovers in wild geladas. *Am. J.*
Primatol. **70**, 1152–1159.

63. Tinsley Johnson E, Snyder-Mackler N, Lu A, Bergman TJ, Beehner JC. 2018 Social and ecological
drivers of reproductive seasonality in geladas. *Behav. Ecol.* **29**, 574–588.

64. Wickham H. 2011 ggplot2. *WIREs Comp Stat* **3**, 180–185.

65. Fox J, Others. 2003 Effect displays in R for generalised linear models. *J. Stat. Softw.* **8**, 1–27.

66. Foster WA, Treherne JE. 1981 Evidence for the dilution effect in the selfish herd from fish
predation on a marine insect. *Nature* **293**, 466–467.

67. Lu A, Feder JA, Snyder-Mackler N, Bergman TJ, Beehner JC. 2021 Male-mediated maturation in
wild geladas. *Curr. Biol.* **31**, 214–219.e2.

68. Weibel CJ, Tung J, Alberts SC, Archie EA. 2020 Accelerated reproduction is not an adaptive
response to early-life adversity in wild baboons. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 24909–24919.

69. Bercovitch FB, Berard JD. 1993 Life history costs and consequences of rapid reproductive

maturation in female rhesus macaques. *Behav. Ecol. Sociobiol.* **32**, 103–109.

70. Koenig A. 2000 Competitive regimes in forest-dwelling Hanuman langur females (*Semnopithecus*
entellus). *Behav. Ecol. Sociobiol.* **48**, 93–109.

71. Harris TR, Chapman CA, Monfort SL. 2009 Small folivorous primate groups exhibit behavioral
and physiological effects of food scarcity. *Behav. Ecol.* **21**, 46–56.

72. Jarvey JC. 2016 The importance of underground foods in female gelada (*Theropithecus gelada*)
socioecology. University of Michigan. See https://deepblue.lib.umich.edu/handle/2027.42/134702.

73. Snyder-Mackler N, Alberts SC, Bergman TJ. 2014 The socio-genetics of a complex society: female
gelada relatedness patterns mirror association patterns in a multilevel society. *Mol. Ecol.* **23**, 6179–
6191.

74. Ostner J, Kappeler PM. 2004 Male life history and the unusual adult sex ratios of redfronted lemur,
Eulemur fulvus rufus, groups. *Anim. Behav.* **67**, 249–259.

75. Teichroeb JA, Jack KM. 2017 Alpha male replacements in nonhuman primates: Variability in
processes, outcomes, and terminology. *Am. J. Primatol.* **79**, e22674.

76. Fedigan LM, Jack KM. 2011 Two girls for every boy: The effects of group size and composition on
the reproductive success of male and female white-faced capuchins. *Am. J. Phys. Anthropol.* **144**,
317–326.