Microbial adaptation to venom is common in snakes and spiders.

E. Esmaeilishirazifard[1,2†], L. Usher[1,2†], C. Trim[3], H. Denise[4], V. Sangal[5], G.H. Tyson[6], A. Barlow[7], K.F. Redway[1], J.D. Taylor[1,2,8], M. Kremyda-Vlachou[1], T. D. Loftus[9], M.M.G. Lock[9], K. Wright[1], A. Dalby[1], L.A.S. Snyder[10], W. Wuster[11], S. Trim[9], and S.A. Moschos[1,2,5.*]

Affiliations:

[1]Department of Biomedical Sciences, Faculty of Science and Technology, University of Westminster, London, U.K., E.U.

[2]Westminster Genomic Services, Faculty of Science and Technology, University of Westminster, London, U.K., E.U.

[3]School of Human and Life Sciences, Faculty of Social and Applied Sciences, Canterbury Christ Church University, Canterbury, Kent, U.K.

[4]EMBL-EBI European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK.

[5]Department of Applied Sciences, Faculty of Health and Life Sciences, Northumbria University, Newcastle, Tyne and Wear, UK.

[6]Food and Drug Administration, Center for Veterinary Medicine, Office of Research. Laurel, MD, USA.

[7]Institute for Biochemistry and Biology, University of Potsdam, Potsdam, Germany.

[8]School of Environment and Life Sciences, University of Salford, Salford, Greater Manchester, U.K.

[9]Venomtech Limited, Discovery Park, Sandwich, Kent, UK.

[10]School of Life Sciences, Pharmacy, and Chemistry, Kingston University, Kingston Upon Thames, London, UK.

[11]School of Biological Sciences, College of Natural Sciences, Bangor University, Bangor, Wales, U.K.

*Correspondence to: sterghios.moschos@northumbria.ac.uk.

†These authors contributed equally to this manuscript.

Abstract: Animal venoms are considered sterile sources of antimicrobial compounds with strong membrane disrupting activity against multi-drug resistant bacteria. However, bite wound infections are common in developing nations. Investigating the oral and venom microbiome of five snake and two spider species, we evidence viable microorganisms potentially unique to venom for black-necked spitting cobras (*Naja nigricollis*). Among these are two novel sequence types of *Enterococcus faecalis* misidentified by commonly used clinical biochemistry procedures as *Staphylococcus*; the genome sequence data of venom-specific isolates feature an additional 45 genes, at least 11 of which improve membrane integrity. Our findings challenge the dogma of venom sterility and indicate an increased primary infection risk in the clinical management of venomous animal bite wounds.

One Sentence Summary: Independent bacterial colonization of cobra venom drives acquisition of genes antagonistic to venom antimicrobial peptides.

Keywords: venom, microbiome, adaptation, infection, snake, spider, scorpion.

Introduction

The rise of single and multi-drug resistant (MDR) bacterial infections suggests the end of the antibiotic golden era might be approaching fast. Discovery of novel antimicrobials is therefore an urgent priority of exceptional socioeconomic value. Crude preparations of animal venoms exhibit strong antibiotic potencies, including against clinical MDR bacterial isolates such as *Mycobacterium tuberculosis* (Xie et al. 2003). With antimicrobial properties described for crotalid (pit viper) venom as early as 1948 (Glaser 1948), relevant compounds have been isolated from most animal venoms, including those of spiders, scorpions, and insects, as well as aquatic species. Examples include phospholipase-A2 enzymes, L-amino acid oxidases, cathelicidins, C-type lectins, and hydrophobic/cationic peptides, as well as venom toxin domains (Samy et al. 2013), which may act by physically disrupting bacterial cell membranes through pore formation (Stocker and Traynor 1986; Perumal Samy et al. 2017). Accordingly, venomous animal bite or sting (envenomation) wound infections are considered rare (Talan et al. 1991) and are attributed to secondary infection (Powers 2005). Yet over three quarters of snake bite victims may develop mono- or polymicrobial envenomation wound infections, characterized by *Bacteroides*, *Morganella, Proteus*, *and Enterococcus* (Wagener et al. 2017; Mao et al. 2016) – bacterial taxa commonly found in the gut. Indeed, *Enterococcus faecalis and Morganella morganii* have been independently reported as the most common Gram positive and Gram negative infections across several countries (Mao et al. 2016; Lam et al. 2011; Wagener et al. 2017). Historically associated with the oral snake microbiome (Jorge et al.), these bacteria are thought to originate from prey faeces (Garg et al. 2009) persisting in the snake oral cavity (Lam et al. 2011) with a diversity similar to that of the snake gut (Iqbal et al. 2014). Yet no 'fixed' oral microbiome was observed in early systematic studies, beyond a seasonal variation of diversity (Blaylock 2001). Curiously,

non-venomous snake mouths were reportedly more sterile than those of venomous snakes (Blaylock 2001), a counterintuitive finding independently reproduced elsewhere (Lam et al. 2011). More recently, the oral microbiome of the non-venomous Burmese python (*Python bivittatus*) has also been reported to be native and not derived from prey guts (Costello et al. 2010).

As venom glands are connected to the tip of envenomation apparatus via a persistently open duct which is continuously exposed to the environment (Trim and Trim 2013), envenomation apparatus could be compared to clinical catheterisation assemblies: a transcutaneous needle resting on a non-sterile environment, connected to a continually open duct, leading to a liquid vessel. Such devices develop biofilms within a few days, making weekly catheter replacement necessary (Loveday et al. 2014). Unlike the high flow rates of catheters, however, envenomation apparatus normally ejects venom only sporadically. Captive snakes are often fed weekly and can fast for months whereas large arachnids are fed typically on a monthly basis. Wild animals may also undergo hibernation for several months when venom expulsion frequency can be assumed to be zero. We therefore postulated that the anatomy of envenomation apparatus allows it to be colonised by microbes and that their intermittent use may facilitate bacterial persistence, adaptation and establishment within antimicrobial venom.

Results

The snake venom microbiome varies on account of host species and not on account of the oral flora.

Applying established culture-free methods (Muyzer et al. 1993) on commercially available *Bothrops atrox* venom (*fer-de-lance*; viperidae) and a venom sample from a captive *Bitis arietans* (African puff adder), we first optimised microbial DNA extraction for this unusual

biological matrix **(Fig. S1)**. Given animal availability, behavioural, and sampling limitations, we next focused our efforts on five snake, two spider and two scorpion species (**Table 1**). We collected a swab (O) of the oral cavity (snakes) or fang/aculeus surface (spiders/scorpions) and two consecutive envenomation samples (E1 and E2), expecting the second venom sample to have fewer contaminants from bacterial plugs possibly forming on the envenomation apparatus. In agreement with previous reports (Jorge et al.; Blaylock 2001), principle coordinate analysis and unsupervised clustering (**Fig. S2**) failed to discriminate the swab microbiomes by host species, suggesting the common diets and water sources in captivity would have the biggest impact on the swab data.

This lead us to hypothesise that captive animals would feature more closely related microbiota in their venoms compared to commercial or wild samples. We therefore compared the venom microbiomes of all snakes using the same approach (**Fig. 1**). High Shannon-Weiner indices indicated considerably diversity in snake venom microbiomes, however, surprisingly closer relationships were observed between *B. arietans* and other *Viperidae*, despite samples spanning captive and wild animals; an exception was *B. atrox* venom, which was characterized principally by *Gammaproteobacteria.* Focusing on *B. arietans* also failed to cluster samples by origin (**Fig. S2**), despite the disparate locations across South Africa where wild *B. arietans* samples were collected (**Fig. 1D**). In contrast, *N. nigricollis* microbiomes largely formed a distinct cluster (**Fig. 1**) characterized *bacteroidia* (*bacteroidaceae*), a taxon less common among *Viperidae*. This could reflect anatomical differences in elapid (cobra) fang location at the front of the mouth compared to the sheathed nature of the longer, hinged viperid fangs, whose tips rest at the back of the oral cavity. In contrast, spider species did not seem to influence venom microbiome consistency and exhibited lower biodiversity (**Fig. S4)**. These results likely reflected

Table 1: Animals sampled for the presence of microbiomes in venom.

Common name	Scientific name	Short name	Origin	Preservation method	Number of animals.
Snakes					
Puff adder	*Bitis arietans*	*B.are*	Captivity	Flash-frozen	1
			Commercial	Lyophilised	1
			Wild	Air-dried	8
Black-necked cobra	*Naja nigricollis*	*N.nig*	Captivity	Flash-frozen	3
Fer-de-lance	*Bothrops atrox**	*B.atr*	Captivity	Flash-frozen	3
Western diamond rattlesnake	*Crotalus atrox**	*C.atr*	Captivity	Flash-frozen	2
Taipan	*Oxyuranus scutellatus**	*O.scu*	Captivity	Flash-frozen	2
Spiders					
Indian ornamental	*Poecilotheria regalis*	*P.reg*	Captivity	Flash-frozen	3
Salmon pink	*Lasiodora parahybana***	*L.par*	Captivity	Flash-frozen	5
Scorpions					
Desert hairy scorpion	*Hardurus arizonensis***	*H.ari*	Captivity	Flash-frozen	3
Asian forest scorpion	*Heterometrus spinifer***	*H.spi*	Captivity	Flash-frozen	3

 * Venom produced by one animal only

** Yields ranged <1 - 30 ul.

Figure 1: Snake venom microbiomes cluster on account of host species. Viperid venom microbiomes cluster separately from *N. nigricollis* with the exception of *B. atrox* as determined by A) PCoA, B) UPGMA tree and C) class-level taxonomic profiling following 16S rRNA phylogenetic analysis. Dots in (A) and (B) are coloured by species (red: *B. arietans*; green: *N. nigricollis*; light blue: *B. atrox*; orange: *C. atrox;* dark blue: *O. scutallatus*), represent data of captivity individuals, are labelled with short species name, enumerated for individual number and identified for the envenomation number (E1 or E2) of the sample. The 8 wild (red dots B1-B8) and the commercially sourced, lyophilised (red L dot) *B. arietans* sample are independently labelled. Relative taxonomic diversity profiles in (C) are aligned to the UPGMA tree sample

labels, with the Shannon-Weiner Index (SWI) of each sample indicated. The geographical origin of the wild *B. areitans* samples collected in South Africa are shown in D.

vertebrate/invertebrate anatomical differences and the limited venom yield from invertebrates (<1-30 μl) vs snakes (100-1,000 μl).

A fifth of the *N. nigricollis* venom microbiome is distinct to that of fangs.

Encouraged by the distinctive bacterial taxonomies in *N. nigricollis* venom, the availability of animals under controlled conditions, and the paired nature of the fang swab and envenomation samples, we delved deeper into this dataset. Thus, fang microbiomes appeared to form a distinct cluster to that of venom microbiomes (**Fig. 2A, B)** suggesting the venom gland might be a distinct ecological niche (**Fig. 2C**). We therefore asked if any bacterial taxa were unique to subsets of these samples. Operational taxonomic unit (OTU) incidence analysis within each animal (**Fig. 2D**) suggested some 60% of OTUs were shared between corresponding venoms and fangs. Yet, importantly, up to 20% of these appeared to be unique to venom, and some 15% were unique to the fang (**Fig. 2E**), indicating an OTU continuum between the two microenvironments, with unique taxa in each site. Common sample types also featured a majority of common taxa, and OTUs unique to each site in each animal (**Fig. 2F**). However, taxa unique to each sample type (O, E1 or E2) were rarely found across all animals. These results suggested that although the microbiome between each snake fang and venom was largely common, venom contained distinct organisms.

Figure 2: The intra- and inter-individual relationship of venom and oral microbiomes in *N. nigricollis.* Comparison of the oral and venom microbiomes in three *N. nigricollis* individuals by A) PCoA, B) UPGMA tree and C) class-level taxonomic profiling following 16S rRNA phylogenetic analysis indicates separate clustering of the microbiotae in the two microenvironments. D) Within animal incidence comparisons of operational taxonomic units (OTUs) suggest E) unique taxa exist within the oral but also the venom microenvironments. F) Between animal comparisons per niche (E1, E2, Oral) indicate most OTUs are shared but some OTUs are unique to each animal for each site. Dots in (A) and (B) represent individual *N. nigricollis* (N.nig) animal data and are coloured/labelled by sample type (black: oral; red: envenomation 1 (E1); blue: envenomation 2 (E2)). Relative taxonomic diversity profiles in (C)

are aligned to the UPGMA tree sample labels, with the Shannon-Weiner Index (SWI) of each sample indicated. The 'venom' histogram in E) represents the sum OTU fraction found in the two envenomation samples per individual (+/- standard deviation).

The venom flora in snakes and spiders is viable.

Testing microbiome viability on identification agar (**Supp. Table 1**) yielded less growth with swab samples. Where this was significant, it was not usually matched by similar growth from the corresponding venom samples, further suggesting that the venom bacteria were probably not mouth contaminants. Strikingly, substantial and consistent growth was encountered amongst *N. nigricollis* (**Fig. 3A)** and *P. regalis* (**Table S1**) samples on blood agar. Unexpectedly, neither the wild (air dried) nor the commercial (lyophilised) venom samples yielded any growth, although colonies were obtained in blood agar from the captive *B. arietans*, underscoring the impact of venom handling on microbiome viability.

Well-established clinical microbial biochemistry tests identified the multiple, punctate white colonies from *N. nigricollis* almost universally as *Staphylococcus spp.*, albeit with assay confidence intervals (CI; **TABLE S2**) below 50%. In contrast, *Stenotrophomonas maltophila* (80.4% CI) was present in five out of six *P. regalis* (all animals positive) and two *Lasiodora parahybana* (salmon pink tarantula) venom samples, but not on any fang swabs. Perplexed by the *N. nigricollis* results we sequenced the isolates on the Ion Torrent PGM.

Viable bacteria in *N. nigricollis* venom are two new *E. faecalis* sequence types.

Resequencing against putative reference genomes (**Table S2**) demonstrated less than 6% base

alignment across all isolates. Instead, BLASTn analysis of the largest contig per isolate after *de*

**Figure 3: Whole genome sequencing identifies viable bacteria in *N. nigricollis* venom as

two, animal-specific, *E. faecalis* strains.** (A) White punctate colonies were recovered in blood

agar (BA) and MacConkey agar (MAC) blinded cultures of individual oral swab (O) and two

consecutive envenomation samples (E1 and E2) obtained from three captivity *N. nigricollis*

snakes. N/D: none detected. (B) Blinded multiple sequence alignment (ClustalO followed by

ClustalW phylogeny) of homologous sequences across the *de novo* assembled genomes against

the *E. faecalis* V583 *KatA* gene (distance to V583 *KatA* indicated in V583 track) suggests two

sequence groups reflecting the history and housing of the host animals. (C) Blinded MST construction based on the MLST of the *N. nigricollis*-derived isolates against nine *E. faecalis* reference genomes again separate samples into two distinct clusters that reflect the history and housing of the host animals. Partially available allele data are included in this analysis and allelic difference instances between nearest neighbours are annotated in white boxes. (D) Blinded complete genome MLST against a custom schema generated using three closely related *E. faecalis* reference genomes cluster these isolates by animal of origin (animals 1 (light blue), animal 2 (dark blue) and animal 3 (orange)). The host animal colour scheme depicted in D) is also used in B) and C).

novo assembly identified *E. faecalis* V583 as the closest likely relative, resulting in >80% base alignment, at an average coverage of 51.2x. This was puzzling given the catalase positive isolate biochemistry vs. the generally accepted catalase negative nature of *E. faecalis* (Fisher and Phillips 2009). However, the *E. faecalis* V583 *katA* gene, previously reported to encode a haem-dependent cytoplasmic catalase (Frankenberg et al. 2002), was confirmed by BLASTn amongst all isolates at 99% identity, explaining the biochemical misclassification. Blinded multiple sequence alignment (MSA) further revealed two *katA* alleles: one shared between isolates from animals 1 and 2 (allele 1) vs. another found in animal 3 isolates (allele 2; **Fig. 3B)** varying by less than 20 single nucleotide polymorphisms to the V583 allele (**Fig. S5A**). Interestingly, these alleles grouped isolates according to the origin and joint housing histories of animals 1 and 2 vs animal 3.

To explore isolate relationships further we generated minimum spanning trees (MST; **Fig. 3C**) by multi-locus sequence typing (MLST; **Table 2**), including at core genome level (cgMLST;

Fig. 3D and **Fig. S5B-D**) (Mellmann et al. 2011; de Been et al. 2015). Comparisons to five

complete reference genomes of the closely related *E. faecium* succeeded only for the *gyd* (alelles

16, 19) and *adk* (allele 18) loci. In contrast, MLST succeeded for all *E. faecalis* loci (**Table 2**),

grouping isolates in line with the *katA* allele observations (**Fig. 3B**), and identifying two novel

sequence types featuring two new alleles for *pstS* and *yqiL* (**Fig. S6**) as confirmed by Sanger

Table 2: Novel sequence types of *E. faecalis* recovered from fangs and venoms of *N. nigricollis*.

Animal no.	Sample Isolate origin	Sample Blinding code	gdh	gyd	pstS	gki	aroE	xpt	yqil
1	O	S22	22	6	31	13	11	35	8
	E1	V36	22	6	31	13	11	35	8
	E2	V29	22	6	31	13	11	35	8
2	O	S17	22	6	31	13	11	35	8
	E1	V31	22	6	31	13	11	35	8*
	E2	V28	22	6	31	13	11	35	8**
3	O	S3	18	1	New	24	83	47	New
	E1	V33	18	1	New	24	83	47	New
	E2	V23	18	1	New	24	83	47	New

O: oral swab sample

E1: envenomation 1

E2: envenomation 2

*: single base pair deletion in NGS data not validated by Sanger sequencing
**: homopolymer single base extension not validated by Sanger sequencing

sequencing. MLST also indicated closer relationships to the *E. faecalis* strains OG1RF, D32, and DENG1, with 87.5% +/- 1.7 of OG1RF cgMLST targets accepted for distance calculations vs D32 (78.8% +/- 2.1) and DENG1 (77.4% +/- 1.5). Pairwise comparisons of the resulting custom cgMLST schema including 5041 loci found across all the *N. nigricollis*-derived isolates also grouped isolates by their host animal (**Fig. 3C**) in line with the *katA* and MLST locus observations. Collectively, these results indicated independent acquisition of two separate and novel *E. faecalis* strains across these three animals.

Genomic evidence of *E. faecalis* isolate adaptation to venom

Including in cgMLST comparisons an additional 3060 loci found in some, but not all of the isolates (**Fig. S5D**) identified between 290 and 831 allelic differences occurring within each animal. Furthermore, whilst 80.9% of alleles varied between the two nearest neighbour isolates from the two strains, venom isolates from animals 1 and 2 were divergent by 7.15 – 10.3% to their oral isolate counterparts. Given the well-described plasticity of the *E. faecalis* genome (OG1RF: 2.74 Mb vs V583: 3.36 Mb), we next examined mobile genetic element divergence.

Detecting the *repA-2* gene from plasmid pTEF2 (**Table S3**) amongst all isolates in accordance to MLST-derived isolate groupings suggested only plasmid fragments were found in these genomes. However, as pTEF2 is one of three *E. faecalis* V583 plasmids associated to vancomycin resistance (Paulsen et al. 2003), we confirmed fragments from all three pTEF plasmids (**Fig. 4A**), consistent to MLST profiles (**Table S4, Table 2**), and with some pTEF sequence elements not found in secondary envenomation isolates from animals 1 and 2. Thus, the average per base read pTEF1/pTEF2 ratios for isolates from animal 3 were 0.25 (+/- 0.01), as

Figure 4: Comparative genomics of mobile and core genomic chromosomal elements of venom-tolerant *E. faecalis* (A) Circos coverage plots of the vancomycin resistance-associated, V583 plasmids pTEF1, pTEF2 and pTEF3 in the *E. faecalis* isolates obtained from oral, envenomation 1 (E1) and envenomation 2 (E2) samples from three *N. nigricollis* individuals, reinforce the two sequence type groupings and highlight within animal variation (green arcs) indicative of sample-specific variation (lack of reads) across E2 samples in animals 1 and 2. The central plot for each plasmid and animal reflects GC content. All data is represented in 50 nt blocks. (B) Blinded maximum likelihood tree of the core genomic alignments for the 6 *N. nigricollis*- derived *E. faecalis* isolates against the V583 and OG1RF reference strains.

elements with high (>95%) sequence identity to these plasmids were also known, highly mobile sequences common to other plasmids (e.g. the *E. faecalis* Bac41 bacteriocin locus) (Kurushima

et al. 2016). These results therefore indicated the presence of highly mobile sequences in these compared to 0.90 (+/-0.15) and 0.81 (+/-0.05) for isolates from animal 1 and 2, respectively; similarly, the pTEF2/pTEF3 average per base read ratios were 0.68 (+/-0.02) for isolates from animal 3 vs. 1.57 (+/-0.04) for isolates from animal 1 and 2. Most importantly, however, many of the genomic isolates, either on plasmids or integrated in the bacterial chromosome, that appeared to participate in the observed genomic divergence of *E. faecalis* within each animal.

Closer examination of the draft genome assemblies indicated that the three *E. faecalis* isolates from animal 3 shared close genomes of ~2.9 Mb with 2,772 to 2,836 genes. The rest of the genomes varied from 3.04 to 3.24 Mb in size, and encompassed 3,128 to 3,282 genes (**Table S3**). The *E. faecalis* pan-genome, including the strains OG1RF and V583, contained a total of 5,130 genes of which 1,977 belonged to the core genome. As with other analyses, the core genomic diversity separated the snake-derived strains into two major groups (**Fig. 4B**): three isolates with smaller genome sizes corresponded to animal 3 forming group A, and the remaining six isolates with larger genome sizes forming group B. However, in line with cgMLST data (**Fig. 3C**) OG1RF and V583 were quite distinct from both of these groups. Comparison of the annotated genomes indicated that 235 genes that were specific to group A and were absent from group B isolates (**Table S5**). The majority of these genes (144/235) encoded hypothetical proteins. Among the rest, two genes were involved in capsule biosynthesis, and one pesticidal crystal protein cry22Aa, which may contribute to the virulence or fitness of these isolates. In addition, three genes belonged to a CRISPR-Cas type II system, a prokaryotic immune system that protects bacteria from invading nucleic acids, including bacteriophages and plasmids; these were absent from other isolates. However, a small proportion of the genes were mobile genetic elements that encoded phage associated proteins, integrases, and transposes (**Table S5**), further

supporting the hypothesis that these group A-specific genes might have been acquired by recombination. Of the 321 genes specific to group B, again the largest fraction (203/321) encoded hypothetical proteins (**Table S6**). However, a number of genes encoded phage-associated proteins and transposases.

Most interestingly, examination of the nine genomes from the snake-derived isolates identified 45 genes unique to them and absent from the OG1RF and V583 reference genomes. In contrast to Group A- and Group B-specific genes, 21 (46.7%) of these 45 genes were of known function (**Table S7**). UniProt functional annotation indicated that 11 of these 21 (52.4%) genes were associated to cell wall/membrane integrity, with four additional genes (19.1%) associated to pathogen foreign protein and toxin defence. Pathway analysis via DAVID using *B. subtilis* orthologues identified significant enrichment (11.11 fold, p=0.00802) in the two component system pathway, specifically genes responsive to cationic antimicrobial peptides, cell wall active antimicrobials, bacitracin efflux, and salt stress.

Projecting primary infection clinical risk from venom-tolerant *E. faecalis* isolates.

Given these viable *E. faecalis* strains could potentially infect envenomation wounds, we next examined the genomic data for known resistance determinants that might facilitate opportunistic primary infection. None of these strains had any acquired resistance genes to any antimicrobial classes (**Table S8**), although all isolates featured *lsaA*, which confers intrinsic streptogramin resistance to *E. faecalis* (Singh et al. 2002): MSA further reinforced isolate grouping based on host animals (**Fig. S7**). Since horizontally acquired genes are largely responsible for resistance to vancomycin, aminoglycosides, macrolides, and tetracycline in *Enterococcus* (Hollenbeck and Rice 2012), these strains were considered likely to be susceptible to drugs in each of these drug

classes. In addition, the absence of known resistance-associated mutations in *gyrA*, *parC*, and the

23S rRNA genes also increased the likelihood that these isolates would be susceptible to

oxazolidinones and fluoroquinolones. Thus, these data indicated that several available

antimicrobials would likely be effective in treating infections caused by these strains of *E.*

faecalis. However, a gene related to macrolide export (macB5) was detected in the venom

microbiome strains (**Table S7**).

On the other hand, one additional concern was the potential misidentification of these isolates as

Staphylococcus using standard diagnostic methods: this could potentially impact treatment

decision-making. Although many of the same antibiotics, including vancomycin and linezolid,

would be considered for treatment of both staphylococci and enterococci, there are some

potential differences. For instance, oxacillin is often employed as a first-line agent to treat

Staphylococcus (Andrews et al. 2000). This drug is not effective against enterococci, as the use

of penicillins for *E. faecalis* infections would typically involve ampicillin, usually in

combination with an aminoglycoside (Arias et al. 2010). In addition, cephalosporins such as

cefotaxime are considered second-line therapies for coagulase-negative staphylococci such as *S.*

epidermidis (Aldridge). However, enterococci are intrinsically resistant to this class of drugs, and

their prevalence in the gut actually tends to increase in response to cephalosporin therapy (de

Vries-Hospers et al. 1991). Thus, while the *E. faecalis* strains in this study did not have any

known acquired resistance determinants, if they were to cause infections, ensuring their proper

identification would be critical to issuing correct treatment and achieving positive clinical

outcomes.

Discussion

In contrast to the generally held view that venoms are both antimicrobial (Xie et al. 2003; Glaser 1948; Samy et al. 2013; Stocker and Traynor 1986) and sterile (Talan et al. 1991; Palappallil 2015; Powers 2005), despite contrasting reports since the 1940's (Pasricha and Abedin 1941), we show that microorganisms are common and viable in the venoms of both vertebrates and invertebrates. As with previous work (Jorge et al.; Garg et al. 2009; Iqbal et al. 2014), our data support the precept that prey faeces might seed the oral and the venom microbiome. However, significant adaptation takes place in these bacteria to persist in venom, occurring in parallel across distinct bacterial isolates and host animals probably through horizontal gene transfer, to contribute genes unique to these isolates. Of the genes with known function, an unusually large fraction (71.5%) is involved in maintaining bacterial membrane integrity or toxin defence. Notably, disruption of membrane integrity appears to be the most common mechanism of action for known, venom-derived antimicrobial peptides and enzymes (Stocker and Traynor 1986; Perumal Samy et al. 2017) of long-standing (Glaser 1948) and significant biotherapeutic interest, including MDR (Xie et al. 2003) and nociception (Trim and Trim 2013). It is unclear at present to which extent this form of parallel convergent evolution extends beyond *Enterococcus* or other antimicrobial resistance mechanisms, such as antibiotic resistance genes against last resort antibiotics found on mobile genetic elements (Liu et al. 2016; McGann et al. 2016) found in multiple continents.

From a clinical perspective, identification of *E. faecalis* as the most prevalent culturable microbe across our European *N. nigricollis* venom samples strikingly reflects three independent clinical reports across Africa and Asia that this non-sporulating microbe is the most common Gram positive infection cultured from infected envenomation wounds (Mao et al. 2016; Lam et al.

2011; Wagener et al. 2017). Further epidemiological data from countries with high envenomation incidence (Mao et al. 2016; Lam et al. 2011; Wagener et al. 2017; Johnston et al. 2017; Monteiro et al. 2002) challenge the consensus view in developed nations that venom is sterile, opportunistic primary infection upon envenomation is uncommon, and that venom wound infection is a consequence of poor hygiene or poor debridement practice (Talan et al. 1991; Powers 2005). Accordingly, in an experimental rabbit model of dermonecrosis (Monteiro et al. 2002) caused by *Loxosceles intermedia* (recluse spider) venom, *Clostridium perfringens* recovered from the spider fang and venom enhanced disease symptoms. *Stenotrophomonas*-like bacteria were also found to dominate cone snail venom microbiomes (Torres et al. 2017), indicating that microbial venom adaptation may extend well beyond snakes, spiders, scorpions, and snails. The cone snail study also reported comparable microbiomes in samples collected across the Pacific basin as well as Atlantic specimens. Furthermore, building upon the few instances of polymicrobial infection reported clinically (Mao et al. 2016; Lam et al. 2011; Wagener et al. 2017), the reports on *L. intermedia* (Monteiro et al. 2002)*, Conus* (Torres et al. 2017), and herein suggest that diverse microbes effectively co-colonise venom glands in host species-specific manner, and thus envenomation wounds. Taken together, these studies support a review of the current standards of care for envenomation wound management (Palappallil 2015) beyond simply managing the severe tissue damage and necrosis that might be caused by venomous bites, to include clinical microbiology on envenomation wounds upon presentation. This would be particularly relevant to individuals immunocompromised through disease or malnutrition, e.g. in developing nations where envenomation incidence rates are high, or to children on a venom/colony forming unit dose per body weight basis.

Yet common diagnostic methods mistook *E. faecalis* for *Staphylococcus,* which could lead to unfavourable clinical decision making. It is unclear at present how common such misidentification events might be, contributing to MDR through inappropriate antibiotic use. At least one retrospective study reported higher incidence of *Staphylococcus spp.* in envenomation wounds (Garg et al. 2009), and Blaylock's seminal snake oral flora studies also reported *Proteus* and *Staphylococcus* (Blaylock 2001): both relied on the same methods we used that resulted in pathogen misidentification. Our results therefore further support use of PCR/sequencing methods as they become more relevant to resource limited settings (Quick et al. 2016), and suited to the point of need (Shah et al. 2017), in line with World Health Organisation ASSURED criteria. Understanding the sensitivity of these methods will be crucial in their reliable implementation in envenomation care. It is therefore noteworthy that despite the limited biomass levels in these samples, species-level OTU analysis on MG-RAST (Wilke et al. 2016) correctly identified *E. faecalis* as one of the principle aerobic isolates in *N. nigricollis* venom. Thus, a simple phylogenetic or metagenomic approach, combined with local herpetogeography knowledge, could quickly and accurately inform clinical action regarding antivenom administration. This is because, unlike incidents involving venomous pets, envenomating animal capture is both rare and contraindicated to minimise further envenomation injuries (Mao et al. 2016). Culprit animals are also commonly misidentified through inadequate description (Mao et al. 2016), and antibody-based venom identification kits have so far proven unreliable (Johnston et al. 2017), generally complicating antivenom selection.

To conclude, we evidence that vertebrate and invertebrate animal venoms host diverse, viable microbiomes, with isolates genetically adapted to venom antimicrobials of medical interest against MDR. These results challenge perceptions on the sterility of venom and absence of

primary infection risk upon envenomation, pointing to modern nucleic acid testing technologies

to better inform envenomation care and antibiotic use.

Materials and Methods

Animals, sampling and microbial culture

All samples analyzed in this study were provided by Venomtech Ltd., with the exception of freeze-

dried *B. arietans* venom (Latoxan, Portes les Valence, France) and field collected samples

collected in South Africa. Briefly, captive animals were housed in 2 m by 1m wooden, glass

fronted vivaria with large hide, thermal gradient and water *ad-libitum*. All procedures for venom

collection and swabbing were approved as unregulated under the Animals (Scientific Procedures)

Act 1976. Venom was collected by standard techniques; briefly snakes were restrained behind the

head and presented to a collection vessel. Snakes freely bit into the vessel until envenomation was

observed. Each snake was presented to two sterile collection vessels in succession, one for the first

envenomation with potential fang plug, and the other for the second flow (labelled E1 and E2,

respectively). While the snake was positioned over the second vessel, the oral cavity was swabbed

with a sterile swab with individual collection tubes (invasive sterile swab with transport media,

DeltaLab, VWR, Lutterworth, UK). The venom collection vessels were clear, sterile 125 ml

polypropylene containers (ThermoFisher Scientific Ltd., Paisley, UK) covered by 2 x 9 cm^2 pieces

of parafilm stretched to fit (ThermoFisher Scientific Ltd.). The collection vessel was secured to a

bench during collection. Samples collected in the field were from wild puff adders sampled as part

of a previous phylogeographic study (Barlow et al. 2013). Venom samples were collected using a

similar method to that described for captive animals, except that the entire venom sample was

collected in a single collection vessel. Samples were lyophilised by storing <100 μl venom aliquots

in a vacuum-sealed container that was half-filled with silica gel. Following drying, venom samples were stored in a refrigerator at 5°C.

Lasiodora parahybana and *Poecilotheria regalis* were housed in 5 and 8 litre polypropylene boxes (Really Useful Products Ltd, Normanton, UK), respectively, with moist vermiculite (Peregrine Livefoods Ltd., Ongar, UK), plastic hide and 5 cm water bowl, as previously described.[58] Arachnids were anaesthetised with a rising concentration of carbon dioxide, the fangs were swabbed with a sterile swab which was then placed in an individual 1 ml sterile, DNA free, polypropylene collection tube (FluidX Ltd, Nether Alderley, UK), and venom was subsequently collected from arachnids by electrical stimulation. All samples were stored at -80°C. The same transport swabs (VWR) as those used for snakes were also used for invertebrate oral / aculear swabbing. Samples were stored at 4°C and cultured within 24 hours of collection. Aerobic microbial viability was determined by plating swabs or aliquoting 10 µl volumes of venom samples onto oxalated whole horse blood agar, MacConkey agar, or mannitol salt agar (ThermoFisher Scientific) plates and incubating at 30°C for 72 hours. Biochemical isolate identification was undertaken using API® strips (20E, 20NE and Staph) interpreted via the APIWEB interface (BioMerieux, Basingstoke, UK). All isolates were stored on beads at -80°C at the University of Westminster microbial isolate library. *Naja nigricollis* sub-culture was performed by restoring cryogenically stored bacteria on lysogeny broth agar (ThermoFisher Scientific) grown for 48 hours at 30°C, and single colony overnight culture in lysogeny broth (ThermoFisher Scientific) using aerated culture (300 rpm).

DNA extraction

Neat venom samples or samples diluted in 18 megaohm water previously confirmed as bacterial DNA free by 16S PCR were subjected to DNA extraction using TRIzol[TM], PureLink Genomic DNA kits or MagMAX Cell-Free DNA kits (ThermoFisher Scientific) according to the manufacturer's instructions. For combined extraction of Gram positive and Gram negative bacteria from liquid samples, diluted samples were split in equal volumes and processed according to the manufacturer's Gram wall specific lysis protocols, with lysates combined prior to DNA binding onto columns by simple admixture. DNA content was then analyzed by Nanodrop (ThermoFisher Scientific) spectrophotometry and purified material was stored at -80°C until further analysis.

16S phylogenetic library preparation and sequencing.

For short amplicon library preparation, the hypervariable V3 region of the 16S rDNA gene was amplified from 20 ng of DNA using the primers 5′-CCTACGGGAGGCAGCAG-3′ and 5′-ATTACCGCGGCTGCTGG-3′ (Integrated DNA Technologies BVBA, Leuven, Belgium),(Muyzer et al. 1993) 1U Platinum® PCR SuperMix High Fidelity (ThermoFisher Scientific) and 10 μM of primer-mix. The reaction mixes were incubated at 94 °C for 5 min followed by 30 cycles of 30 seconds at 94 °C, 30 seconds at 55 °C and 1 min at 72 °C and then final elongation at 72 °C for 10 min using a Techne Prime Thermal cycler (ColePalmer, Staffordshire, UK). PCR products (193 bp) were confirmed by 2% w/v agarose gel electrophoresis in TAE buffer (ThermoFisher Scientific).

NGS library preparation was carried out using the Ion Plus Fragment Library Kit according to the manufacturer's instructions (Rev. 3, ThermoFisher Scientific), except that reactions were reduced

to 1/5th volumes. Pooled libraries were diluted to ~26 pM for templating on the Ion OneTouch 2 system (ThermoFisher Scientific) using the Ion PGM Template OT2 200 v2 kit according to the manufacturer's instructions (Rev. B, ThermoFisher Scientific). Templated samples were sequenced on the Ion Torrent Personal Genome Machine (PGM; ThermoFisher Scientific) system on a single 318 Ion Chip (ThermoFisher Scientific) using the Ion PGM^{TM} 200 Sequencing kit according to the manufacturer's instructions (Rev G., ThermoFisher Scientific).

Whole genome sequencing.

DNA extracted from cultured isolates was mechanically sheared using the Covaris S220 Focused-ultrasonicator (Covaris, Brighton, UK). NGS libraries were generated using the NEBNext Fast DNA Library Prep Set for Ion Torrent (New England Biolabs, Hitchin, UK). Pooled samples were size selected with the LabChip XT (LabChip XT DNA 300 Assay Kit; PerkinElmer, Seer Green, UK) and diluted to 26 pM for templating with the Ion OneTouch 2 system using the Ion PGM Template OT2 200 kit. Templated samples were sequenced on the Ion PGM using the Ion PGM™ Sequencing 200 v2 reagent Kit (ThermoFisher Scientific) and Ion 318™ v2 Ion Chip (ThermoFisher Scientific).

Bioinformatic analyses

Raw Ion Torrent sequencing data reads were quality controlled and demultiplexed using the standard Ion Server v. 4.0 pipeline (ThermoFisher Scientific). Referenced and *de novo* assemblies were carried out using TMAP v.4.0 and the SPAdes plugin in the Ion Server. Phylogenetic data analyses were carried out after independent data deposition and curation on the MG-RAST v.3.0 pipeline (Wilke et al. 2016) (project IDs MGP5177 and MGP5617) which uses a BLAST approach

and the EBI-METAGENOMICS v.1 (project ID ERP004004) pipeline (Hunter et al. 2014) which uses a Hidden Markov Model approach. Raw 16S sequencing reads were deposited in the European Nucleotide Archive (PRJEB4693). Quality control for both resources included length and quality filtering followed by a dereplication step where sequences with identical 50 nucleotides in 5' positions were clustered together. MG-RAST taxonomy annotation involved RNA identification using VSearch, and assignments using a custom database generated by 90% identity clustering of SILVA, GreenGenes and RDP prokaryotic databases. EBI-METAGENOMICS identified rRNA using Hidden Markov Models present in the RDP databases and assigned taxonomy using Qiime and the GreenGenes prokaryotic database.

For post-processing analyses, the EBI-curated dataset was analyzed using MEGAN v.5.5.3 (Huson et al. 2016). Classical multi-locus sequence typing (http://efaecalis.mlst.net/) and cgMLST (Mellmann et al. 2011; de Been et al. 2015) were carried out using Ridom SeqSphere+ v.4.0 running on a 2 core, 10 GB RAM, 500 GB hard disk Biolinux v.8.0 installation on a VirtualBox virtual machine instance on a 16GB RAM, 1TB hard disk Apple iMac. Extended cgMLST analysis to include partially detected loci, excluded loci annotated as 'failed' due to sequencing error suggesting genuine *E. faecalis* genomic divergence occurring within each animal. Plasmid detection was carried out using the PlasmidFinder v.1.3 server (Carattoli et al. 2014), followed by NCBI BLASTn analysis to detect shorter fragments, e.g. the same 398 nt fragment of *repA-2* in animal 3 isolates (<40% of the full-length gene) at 90.1% identity to the plasmid-borne reference sequence. Single gene comparisons and multiple sequence analyses were carried out using TCoffe and MView on the EMBL-EBI server, with base conservation visualized by BoxShade v.3.3.1 on

mobyle.pasteur.fr. Genome-level plasmid coverage analyses were carried out by NCBI BLASTn and comparisons were visualized using Circos v.0.69-4.

The sequencing reads were assembled using SPAdes v.3.9.0 (Bankevich et al. 2012), and the draft assemblies were annotated using Prokka (Seemann 2014) before NCBI deposition (BioProject No. PRJNA415175). The genome sequences of *E. faecalis* strains V583 and OG1RF (Accession numbers NC_004668.1 and NC_017316.1, respectively) were obtained from GenBank and were re-annotated using Prokka to have an equivalence of annotation for comparative analyses. The genomes were compared using the program Roary (Page et al. 2015; Tange). A maximum-likelihood tree was constructed from the core genomic alignment using IQ-Tree (Nguyen et al. 2015) with 100,000 ultra-fast bootstraps and 100,000 SH-aLRT tests. The tree was visualized using Interactive Tree Of Life (iTOL) (Letunic and Bork 2016).

To identify acquired resistance genes, nucleotide BLAST analysis was performed on the ResFinder (Zankari et al. 2012) and NCBI (https://www.ncbi.nlm.nih.gov/pathogens/) resistance gene databases using cutoffs of 50% length and 85% identity to known resistance determinants. Additional BLAST analysis was performed to identify single nucleotide polymorphisms in the quinolone resistance determining region (QRDR) of *gyrA* and *parC* (Kanematsu et al. 1998). Additional mutational analysis was performed on region V of the 23S rRNA-encoding genes (Marshall et al. 2002).

BLASTP was performed in Ensembl Bacteria (release 38), against the *E. faecalis* V583 and *E. faecalis* (GCA_000763645) to obtain further geneID's from significant matches. *Bacillus*

subtilis orthologue gene ID's were collated as this species is the closest relative to *E. faecalis* (VetBact.org) with the most comprehensive genome annotation required for gene ontolgy and KEGG pathway analysis. From the 45 genes unique to venom isolates, useable *B. subtilis* GeneID's were obtained for 22, of which 12 of these successfully converted to ENTREZ Gene ID's using the functional annotation tool (DAVID Bioinformatics resource 6.8)(Huang et al. 2009b, 2009a), selecting *B. subtilis* as the background species.

References and Notes:

Aldridge KE. Cefotaxime in the treatment of staphylococcal infections. Comparison of in vitro and in vivo studies. *Diagn Microbiol Infect Dis* **22**: 195–201. http://www.ncbi.nlm.nih.gov/pubmed/7587040 (Accessed December 19, 2017).

Andrews JM, Boswell FJ, Wise R. 2000. Establishing MIC breakpoints for coagulase-negative Staphylococci to oxacillin. *J Antimicrob Chemother* **45**: 259–61. http://www.ncbi.nlm.nih.gov/pubmed/10660515 (Accessed December 19, 2017).

Arias CA, Contreras GA, Murray BE. 2010. Management of Multi-Drug Resistant Enterococcal Infections. *Clin Microbiol Infect* **16**: 555–62. http://www.ncbi.nlm.nih.gov/pubmed/20569266 (Accessed December 19, 2017).

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, et al. 2012. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J Comput Biol* **19**: 455–477.

http://www.ncbi.nlm.nih.gov/pubmed/22506599 (Accessed December 19, 2017).

Barlow A, Baker K, Hendry CR, Peppin L, Phelps T, Tolley KA, Wüster CE, Wüster W. 2013.
Phylogeography of the widespread African puff adder (Bitis arietans) reveals multiple
Pleistocene refugia in southern Africa. *Mol Ecol* **22**: 1134–57.
http://doi.wiley.com/10.1111/mec.12157 (Accessed December 19, 2017).

Blaylock RS. 2001. Normal oral bacterial flora from some southern African snakes.
Onderstepoort J Vet Res **68**: 175–82. http://www.ncbi.nlm.nih.gov/pubmed/11769348
(Accessed December 19, 2017).

Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller
Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using
PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* **58**:
3895–903. http://www.ncbi.nlm.nih.gov/pubmed/24777092 (Accessed December 19, 2017).

Costello EK, Gordon JI, Secor SM, Knight R. 2010. Postprandial remodeling of the gut
microbiota in Burmese pythons. *ISME J* **4**: 1375–1385.
http://www.ncbi.nlm.nih.gov/pubmed/20520652 (Accessed December 19, 2017).

de Been M, Pinholt M, Top J, Bletz S, Mellmann A, van Schaik W, Brouwer E, Rogers M, Kraat
Y, Bonten M, et al. 2015. Core Genome Multilocus Sequence Typing Scheme for High-
Resolution Typing of Enterococcus faecium ed. K.C. Carroll. *J Clin Microbiol* **53**: 3788–
3797. http://www.ncbi.nlm.nih.gov/pubmed/26400782 (Accessed December 19, 2017).

de Vries-Hospers HG, Tonk RH, van der Waaij D. 1991. Effect of intramuscular ceftriaxone on
aerobic oral and faecal flora of 11 healthy volunteers. *Scand J Infect Dis* **23**: 625–33.

http://www.ncbi.nlm.nih.gov/pubmed/1767260 (Accessed December 19, 2017).

Fisher K, Phillips C. 2009. The ecology, epidemiology and virulence of Enterococcus. *Microbiology* **155**: 1749–1757. http://www.ncbi.nlm.nih.gov/pubmed/19383684 (Accessed December 19, 2017).

Frankenberg L, Brugna M, Hederstedt L. 2002. Enterococcus faecalis heme-dependent catalase. *J Bacteriol* **184**: 6351–6. http://www.ncbi.nlm.nih.gov/pubmed/12399505 (Accessed December 19, 2017).

Garg A, Sujatha S, Garg J, Acharya NS, Parija SC. 2009. Wound infections secondary to snakebite. *J Infect Dev Ctries* **3**: 221–223.

Glaser HSR. 1948. Bactericidal Activity of Crotalus Venom in Vitro. *Copeia* **1948**: 245. http://www.jstor.org/stable/1438710?origin=crossref (Accessed December 18, 2017).

Hollenbeck BL, Rice LB. 2012. Intrinsic and acquired resistance mechanisms in enterococcus. *Virulence* **3**: 421–569. http://www.ncbi.nlm.nih.gov/pubmed/23076243 (Accessed December 19, 2017).

Huang DW, Sherman BT, Lempicki RA. 2009a. Bioinformatics enrichment tools: Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**: 1–13.

Huang DW, Sherman BT, Lempicki RA. 2009b. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**: 44–57.

Hunter S, Corbett M, Denise H, Fraser M, Gonzalez-Beltran A, Hunter C, Jones P, Leinonen R, McAnulla C, Maguire E, et al. 2014. EBI metagenomics—a new resource for the analysis

and archiving of metagenomic data. *Nucleic Acids Res* **42**: D600–D606.

https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkt961 (Accessed December

19, 2017).

Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, Ruscheweyh H-J, Tappu R. 2016.

MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale

Microbiome Sequencing Data. ed. T. Poisot. *PLoS Comput Biol* **12**: e1004957.

http://dx.plos.org/10.1371/journal.pcbi.1004957 (Accessed December 19, 2017).

Iqbal J, Sagheer M, Tabassum N, Siddiqui R, Khan NA. 2014. Culturable Aerobic and

Facultative Anaerobic Intestinal Bacterial Flora of Black Cobra (Naja naja karachiensis) in

Southern Pakistan. *ISRN Vet Sci* **2014**: 878479.

http://www.ncbi.nlm.nih.gov/pubmed/25002979 (Accessed December 19, 2017).

Johnston CI, Ryan NM, Page CB, Buckley NA, Brown SG, O'Leary MA, Isbister GK. 2017.

The Australian Snakebite Project, 2005-2015 (ASP-20). *Med J Aust* **207**: 119–125.

http://www.ncbi.nlm.nih.gov/pubmed/28764620 (Accessed December 18, 2017).

Jorge MT, de Mendonça JS, Ribeiro LA, da Silva ML, Kusano EJ, Cordeiro CL. [Bacterial flora

of the oral cavity, fangs and venom of Bothrops jararaca: possible source of infection at the

site of bite]. *Rev Inst Med Trop Sao Paulo* **32**: 6–10.

http://www.ncbi.nlm.nih.gov/pubmed/2259834 (Accessed December 18, 2017).

Kanematsu E, Deguchi T, Yasuda M, Kawamura T, Nishino Y, Kawada Y. 1998. Alterations in

the GyrA subunit of DNA gyrase and the ParC subunit of DNA topoisomerase IV

associated with quinolone resistance in Enterococcus faecalis. *Antimicrob Agents*

Chemother **42**: 433–5. http://www.ncbi.nlm.nih.gov/pubmed/9527801 (Accessed December 19, 2017).

Kurushima J, Ike Y, Tomita H. 2016. Partial Diversity Generates Effector Immunity Specificity of the Bac41-Like Bacteriocins of Enterococcus faecalis Clinical Strains. *J Bacteriol* **198**: 2379–90. http://www.ncbi.nlm.nih.gov/pubmed/27353651 (Accessed December 19, 2017).

Lam KK, Crow P, Ng KHL, Shek KC, Fung HT, Ades G, Grioni A, Tan KS, Yip KT, Lung DC, et al. 2011. A cross-sectional survey of snake oral bacterial flora from Hong Kong, SAR, China. *Emerg Med J* **28**: 107–114. http://www.ncbi.nlm.nih.gov/pubmed/20511640 (Accessed December 19, 2017).

Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**: W242–W245. http://www.ncbi.nlm.nih.gov/pubmed/27095192 (Accessed December 19, 2017).

Liu Y-Y, Wang Y, Walsh TR, Yi L-X, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, et al. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* **16**: 161–168. http://linkinghub.elsevier.com/retrieve/pii/S1473309915004247 (Accessed December 15, 2017).

Loveday HP, Wilson JA, Pratt RJ, Golsorkhi M, Tingle A, Bak A, Browne J, Prieto J, Wilcox M. 2014. Epic3: National evidence-based guidelines for preventing healthcare-associated infections in nhs hospitals in england. *J Hosp Infect* **86**.

Mao YC, Liu PY, Hung DZ, Lai WC, Huang ST, Hung YM, Yang CC. 2016. Bacteriology of

Naja atra snakebite wound and its implications for antibiotic therapy. *Am J Trop Med Hyg* **94**: 1129–1135.

Marshall SH, Donskey CJ, Hutton-Thomas R, Salata RA, Rice LB. 2002. Gene dosage and linezolid resistance in Enterococcus faecium and Enterococcus faecalis. *Antimicrob Agents Chemother* **46**: 3334–3336.

McGann P, Snesrud E, Maybank R, Corey B, Ong AC, Clifford R, Hinkle M, Whitman T, Lesho E, Schaecher KE. 2016. Escherichia coli Harboring *mcr-1* and *bla* CTX-M on a Novel IncF Plasmid: First Report of *mcr-1* in the United States. *Antimicrob Agents Chemother* **60**: 4420–4421. http://www.ncbi.nlm.nih.gov/pubmed/27230792 (Accessed December 15, 2017).

Mellmann A, Harmsen D, Cummings CA, Zentz EB, Leopold SR, Rico A, Prior K, Szczepanowski R, Ji Y, Zhang W, et al. 2011. Prospective Genomic Characterization of the German Enterohemorrhagic Escherichia coli O104:H4 Outbreak by Rapid Next Generation Sequencing Technology ed. N. Ahmed. *PLoS One* **6**: e22751. http://www.ncbi.nlm.nih.gov/pubmed/21799941 (Accessed December 19, 2017).

Monteiro CLB, Rubel R, Cogo LL, Mangili OC, Gremski W, Veiga SS. 2002. Isolation and identification of Clostridium perfringens in the venom and fangs of Loxosceles intermedia (brown spider): Enhancement of the dermonecrotic lesion in loxoscelism. *Toxicon* **40**: 409–418.

Muyzer G, de Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified

genes coding for 16S rRNA. *Appl Env Microbiol* **59**: 695–700.

http://www.ncbi.nlm.nih.gov/pubmed/7683183 (Accessed December 19, 2017).

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **32**:

268–274. http://www.ncbi.nlm.nih.gov/pubmed/25371430 (Accessed December 19, 2017).

Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D,

Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis.

Bioinformatics **31**: 3691–3693. http://www.ncbi.nlm.nih.gov/pubmed/26198102 (Accessed

December 19, 2017).

Palappallil DS. 2015. Pattern of Use of Antibiotics Following Snake Bite in a Tertiary Care

Hospital. *J Clin Diagn Res* **9**: OC05-9. http://www.ncbi.nlm.nih.gov/pubmed/26435981

(Accessed December 18, 2017).

Pasricha CL, Abedin Z. 1941. The Sterility of Snake Venom Solutions. *Ind Med Gaz* **76**: 276–

277. http://www.ncbi.nlm.nih.gov/pubmed/29013551 (Accessed December 19, 2017).

Paulsen IT, Banerjei L, Myers GSA, Nelson KE, Seshadri R, Read TD, Fouts DE, Eisen JA, Gill

SR, Heidelberg JF, et al. 2003. Role of Mobile DNA in the Evolution of Vancomycin-

Resistant Enterococcus faecalis. *Science (80-)* **299**: 2071–2074.

http://www.ncbi.nlm.nih.gov/pubmed/12663927 (Accessed December 19, 2017).

Perumal Samy R, Stiles BG, Franco OL, Sethi G, Lim LHK. 2017. Animal venoms as

antimicrobial agents. *Biochem Pharmacol* **134**: 127–138.

http://www.ncbi.nlm.nih.gov/pubmed/28288817 (Accessed December 19, 2017).

Powers DW. 2005. Stings and bites: what to do about envenomation injuries. *Emerg Med Serv* **34**: 67, 69–75, quiz 99. http://www.ncbi.nlm.nih.gov/pubmed/15900872 (Accessed December 18, 2017).

Quick J, Loman NJ, Duraffour S, Simpson JT, Severi E, Cowley L, Bore JA, Koundouno R, Dudas G, Mikhail A, et al. 2016. Real-time, portable genome sequencing for Ebola surveillance. *Nature* **530**: 228–232. http://www.nature.com/doifinder/10.1038/nature16996.

Samy R, Gopalakrishnakone P, Satyanarayanajois S, Stiles B, Chow V. 2013. Snake Venom Proteins and Peptides as Novel Antibiotics Against Microbial Infections. *Curr Proteomics* **10**: 10–28. http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1570-1646&volume=10&issue=1&spage=10 (Accessed December 18, 2017).

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**: 2068–2069. http://www.ncbi.nlm.nih.gov/pubmed/24642063 (Accessed December 19, 2017).

Shah K, Bentley E, Tyler A, Richards KSR, Wright E, Easterbrook L, Lee D, Cleaver C, Usher L, Burton JE, et al. 2017. Field-deployable, quantitative, rapid identification of active Ebola virus infection in unprocessed blood. *Chem Sci* **8**: 7780–7797. http://www.ncbi.nlm.nih.gov/pubmed/29163915 (Accessed December 19, 2017).

Singh K V, Weinstock GM, Murray BE. 2002. An Enterococcus faecalis ABC homologue (Lsa) is required for the resistance of this species to clindamycin and quinupristin-dalfopristin. *Antimicrob Agents Chemother* **46**: 1845–50. http://www.ncbi.nlm.nih.gov/pubmed/12019099 (Accessed December 19, 2017).

Stocker JF, Traynor JR. 1986. The action of various venoms on Escherichia coli. *J Appl*

Bacteriol **61**: 383–388.

Talan DA, Citron DM, Overturf GD, Singer B, Froman P, Goldstein EJ. 1991. Antibacterial activity of crotalid venoms against oral snake flora and other clinical bacteria. *J Infect Dis* **164**: 195–8. http://www.ncbi.nlm.nih.gov/pubmed/2056205 (Accessed December 18, 2017).

Tange O. Gnu parallel-the command-line power tool. *usenix.org*. https://www.usenix.org/publications/login/february-2011-volume-36-number-1/gnu-parallel-command-line-power-tool (Accessed December 19, 2017).

Torres JP, Tianero MD, Robes JMD, Kwan JC, Biggs JS, Concepcion GP, Olivera BM, Haygood MG, Schmidt EW. 2017. Stenotrophomonas-Like Bacteria Are Widespread Symbionts in Cone Snail Venom Ducts. *Appl Environ Microbiol* **83**: AEM.01418-17. http://www.ncbi.nlm.nih.gov/pubmed/28986377 (Accessed December 19, 2017).

Trim SA, Trim CM. 2013. Venom: the sharp end of pain therapeutics. *Br J pain* **7**: 179–88. http://www.ncbi.nlm.nih.gov/pubmed/26516522 (Accessed December 19, 2017).

Wagener M, Naidoo M, Aldous C. 2017. Wound infection secondary to snakebite. *South African Med J* **107**: 315–319. http://www.ncbi.nlm.nih.gov/pubmed/28395682 (Accessed December 18, 2017).

Wilke A, Bischof J, Gerlach W, Glass E, Harrison T, Keegan KP, Paczian T, Trimble WL, Bagchi S, Grama A, et al. 2016. The MG-RAST metagenomics database and portal in 2015. *Nucleic Acids Res* **44**: D590–D594. http://www.ncbi.nlm.nih.gov/pubmed/26656948 (Accessed December 19, 2017).

Xie JP, Yue J, Xiong YL, Wang WY, Yu SQ, Wang HH. 2003. In vitro activities of small

    peptides from snake venom against clinical isolates of drug-resistant Mycobacterium

    tuberculosis. *Int J Antimicrob Agents* **22**: 172–174.

Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM,

    Larsen M V. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob

    Chemother* **67**: 2640–2644. http://www.ncbi.nlm.nih.gov/pubmed/22782487 (Accessed

    December 19, 2017).

Acknowledgments: We would like to thank: Drs. Pamela Greenwell and Caroline Smith for their invaluable input on non-standard DNA extraction methodology options suited to unusual samples; Dr. Patrick Kimmit for his input on microbial characterisation; Mr Peter Gibbens for housing and venom collection from captive *N. nigricollis* and *B. areitans*. **Funding:** This work was funded by the University of Westminster, University of Northumbria, and Venomtech Ltd. ;

Author contributions: a MMGL and TDL sampled, and CT and ST prepared the library of captive animal venoms. WW and AB collected and prepared the wild snake samples. EE, JT, PG and SAM optimized and performed the DNA extractions and 16S PCR. JT and EE performed the preliminary and main study library preps and next generation sequencing experiments, respectively. AD, HD, PK, LS, and SAM performed the phylogenetic data quality control, curation and analysis. KFR and SAM performed the microbial cultures and biochemical characterization. MKV and LU grew the *E. faecalis* isolates and performed the whole genome sequencing. MKV, KW and SAM performed the *E. faecalis* isolate genomic characterization and MLST+ analysis. GT performed *E. faecalis* resistome analysis. VS performed the *E. faecalis* isolate pangenome data reduction and ST identified the venom resistance gene ontology subset.

SAM conceived the study and designed experiments together with ST. All authors contributed equally to the overall interpretation of the dataset and manuscript preparation; **Competing interests:** Authors declare no competing interests; and **Data and materials availability:** Phylogenetic data are deposited on MG-RAST (project IDs MGP5177 and MGP5617) and the EBI-METAGENOMICS servers (project ID ERP004004). Raw 16S sequencing reads were deposited in the European Nucleotide Archive (PRJEB4693). Annotated draft *E. faecalis* genome assemblies are deposited on NCBI (BioProject No. PRJNA415175).

Supplementary Materials:

Figures S1-S7

Tables S1-S8