Article

Classification: BIOLOGICAL SCIENCES – Biochemistry

Protein nitration is a physiological regulator of cardiac lactate dehydrogenase active

site loop mobility and activity

Cristian Nogales[a,1,*], Jeroen Frijhoff[b,1], Detlef Thoenges[c], Richard B. Sessions[d], Albert

Sickmann[e], Tatiana Nedvetskaya[f], Cesar Ibarra-Alvarado[g], Vera Schmitz[h], Peter Rösen[h],

Axel Goedecke[i], Aleksandar Jovanović[j], David H. Perlman[k], Anna Klinke[l], Stephan Baldus[l],

Ana I. Casas[a], Péter Bai[m], Csaba Szabo[n], Martin Feelisch[o], Harald H.H.W. Schmidt[a,*].

[a]Department of Pharmacology and Personalised Medicine, Maastricht University,

Maastricht, The Netherlands; [b]Medtronic Bakken Research Center, Maastricht, The

Netherlands; [c]DiaSys Diagnostic Systems GmbH, Holzheim, Germany; [d]School of

Biochemistry, University of Bristol, Bristol, UK; [e]Leibiniz-Institut für Analytische

Wissenschaften – ISAS, Dortmund, Germany; [f]Rudolf-Buchheim-Institue for Pharmacology,

Justus-Liebig-University, Giessen, Germany; [g]School of Chemistry, Autonomous University

of Querétaro. Querétaro, México; [h]German Diabetes Research Institute, Düsseldorf,

Germany; [i]Heinrich-Heine-University, Department Cardiovascular Physiology, Düsseldorf,

Germany; [j]University of Nicosia Medical School, Nicosia, Cyprus; [k]Merck Exploratory

Science Center, Cambridge, Massachusetts; [l]Cologne Cardiovascular Research Center

CCRC, Cologne, Germany; [m]Department of Medical Chemistry, Faculty of Medicine,

University of Debrecen, Debrecen, Hungary and MTA-DE Lendület Laboratory of Cellular

Metabolism, Debrecen, Hungary; [n]Department of Oncology, Microbiology and Immunology;

University of Fribourg, Fribourg, Switzerland; [u]University of Southampton Medical School,

Southampton, UK.

[1]Both first authors contributed equally to this work.

[*]To whom correspondence may be addressed. Email: h.schmidt@maastrichtuniversity.nl or

c.nogales@maastrichtuniversity.nl

Key words: protein nitration, nitro-tryptophan, nitro-tyrosine, biomarker

Word count (main text including references): 24578 characters (including spaces)

Abstract

Protein tyrosine nitration, detected as anti-nitrotyrosine immunoreactivity, is considered one

of the most relevant disease biomarkers of oxidative stress. The mechanism of nitration,

target protein and functional consequences remain often unclear. Here we first extend

protein tyrosine nitration from pathology to physiology as additional mechanism of post-

translational regulation. We focus on a prominent protein band surprisingly nitrotyrosine

immunopositive under basal conditions in mouse, rat and pig heart and more so in diabetes

and myocardial stress. Upon purification, we identify it as lactate dehydrogenase (LDH) and

its basal nitration depending on NO synthase (NOS) and myeloperoxidase (MPO),

respectively. Surprisingly, we locate LDH's nitration by MALDI-TOF mass spectrometry not

to a tyrosine but the C-terminal tryptophan, Trp-324. Molecular dynamics simulations

suggested that Trp-324 nitration restricts the interaction of the active site loop with the C-

terminal α-helix essential for activity, which was corroborated by an apparent lower V_{max}. In

summary, protein nitration is not merely a disease marker but also a physiological event

involving both the eNOS/NO and the MPO/nitrite pathways. In the case of LDH, to our

knowledge the first protein identified to be basally regulated by nitration, this limits its activity,

which is aggravated under cardiac metabolic stress conditions.

Significance

Protein nitration is the most widely used disease marker indicating oxidative stress. How nitration occurs and its functional consequences have remained unclear in many cases. Here we show that protein nitration is not limited to disease but also occurs physiologically. We identify a key enzyme of energy metabolism in the heart as a target of nitration and two mechanisms equally responsible for its formation resulting in lower activity. Thus, protein nitration does not necessarily indicate a disease process but a physiological protein modification, e.g. for dynamic regulation of enzymatic activity.

\body

Introduction

Increased levels of reactive oxygen species (ROS), i.e. oxidative stress, is considered a

common mechanism of several cardiovascular and other disease states (1). In concert with

different nitrogen species (2–4), ROS can lead to protein nitration, e.g. tyrosine residues

(NO_2Tyr). Tyrosine nitration is the most frequently used and considered the most robust

disease marker of oxidative stress (5–7). Its detection relies mainly on NO_2Tyr specific

antibodies (8–10). With respect to the nitrogen source, anti-NO_2Tyr immunopositive signals

are generally interpreted as a hallmark of nitric oxide (NO) being scavenged by superoxide

and intermediate peroxynitrite (9, 11, 12), although other mechanisms exist (13). Regarding

the functional consequences of protein nitration few *in vitro* data exist suggesting mainly

loss of protein function correlating with disease severity (14–17). *In vivo* data are lacking.

Here, we address three major knowledge gaps in our understanding of protein nitration:

specificity, origin*,* and disease relevance.

Results

Development of an anti-nitrotyrosine antibody panel as biomarkers of disease-

relevant oxidative stress. We started out by examining a panel of anti-NO_2Tyr antibodies

(**Fig. 1**), including those directed against carrier-linked NO_2Tyr or against H_2O_2/NO_2^- treated

proteins (see Supplemental Information (**SI**) **Table S1**). We tested these both in rodent (rat)

and, for later purification, also in porcine tissues. Surprisingly we obtained signals already

under physiological conditions. The immunoreaction, however, varied greatly in protein band

size and intensity. Two polyclonal (pAb), pAb1 and pAb2, and one monoclonal antibody

(mAb), mAb1, displayed the most robust signals (**SI Fig. S1**). The specificity of anti-NO_2Tyr

antibodies (18–22) is rarely validated, e.g. by antibody pre-absorption with free NO_2Tyr (11).

In our case, pAb1 signals were partially blocked by pre-absorption with 3 mM NO_2Tyr. mAb1

showed very little non-specific background whilst pAb2 exhibited moderate sensitivity with

high anti-NO_2Tyr specificity.

A prominent nitrotyrosine immunoreactive protein band was detected in mouse, rat

and pig heart under physiological conditions. To conduct a tissue/species screen, we

selected three antibodies, pAb1, pAb2, and mAb1, from our original panel (see **SI Table**

S1). Three proteins were consistently detected under basal conditions and with apparent

molecular weights of 70, 45 and 38 kDa, of which the latter gave the by far the strongest

signal in heart (**Fig. 2A**). Pre-incubation with free NO_2Tyr or NO_2BSA (3 mM) or nitro-group

reduction (21), at least partially, blocked the pAb1 dependent signal. Therefore, subsequent

efforts were directed at identifying the nature of the NO_2Tyr positive 38 kDa protein and its

site of nitration.

Upon purification, the 38 kDa nitrated protein was identified as LDH with lower V_{max}.

Using porcine heart, we isolated the NO_2Tyr immunoreactive 38 kDa protein by ion

exchange (Q Sepharose, QS) and affinity (Blue Sepharose, BS) chromatography (**Fig. 2B**). The partially purified 38 kDa protein was stained by colloidal Coomassie Blue and subjected to peptide mass fingerprinting using MALDI-TOF MS. Based on the NCBIprot database most ions present in the spectrum of tryptic peptides matched LDH as the primary protein component of the sample (**Fig. 2C**).

Next, we tested the effect of nitration on the enzyme activity by nitrating cardiac LDH *in vitro* with increasing amounts of H_2O_2/NO_2^-. Indeed, higher nitration levels of LDH (NO_2-LDH) (**Fig. 2E**) correlated with decreased V_{max} (**Fig. 2F**) and unchanged K_M. Our data suggest therefore that protein nitration is not only a physiological event for cardiac LDH, but also modulates enzyme activity by lowering V_{max}.

Nitrated LDH is increased in diabetes and myocardial stress. To investigate whether NO_2-LDH is further increased in response to metabolic stress, we investigated heart extracts from rat models of streptozotocin-induced short-term (ST) and long-term (LT) diabetes mellitus (23) as well as a doxorubicin (DoxoR) induced mouse model of cardiac stress (24) (**Fig. 3A**). Diabetic rats were sacrificed 3 (ST) and 16 (LT) weeks after inducing diabetes at week 10. Using the pAb1, NO_2-LDH signals were significantly increased in all three models when compared against healthy controls (**Fig. 3B/C**). Immunosignals of pAb1 pre-absorbed with NO_2Tyr or anti-LDH wildtype antibody did not show any significant change. In both diabetes mellitus models and doxorubicin-treatment animals, levels of cardiac LDH were unchanged versus controls, indicating that indeed the NO_2-LDH/LDH ratio was increased.

Cardiac LDH nitration is lowered in eNOS-KO and MPO-KO mice. We next wanted to address our second goal, to understand the mechanistic origin of the nitrogen in NO_2-LDH. Usually, it is assumed that protein nitration derives from peroxynitrite from the reaction between $^\bullet$NO and $O_2^{\bullet-}$ (25). However, another major source, not involving de novo NO

synthesis is frequently overlooked, i.e. MPO-catalyzed nitration using nitrite in the presence

of hydrogen peroxide (26) (**Fig. 4A**). To analyze this *in vivo*, we had to switch to mouse

models, i.e. eNOS-KO and MPO-KO lines. Indeed, we observed that NO_2-LDH signals in

heart tissue were not only reduced in eNOS-KO but also in MPO-KO mice when compared

to their respective WT litter mates (**Fig. 4B**). These results therefore indicated that both

pathways contribute to basal nitration of cardiac LDH.

Further MALDI-TOF mass spectrometry analysis of the nitration site identifies the

nitrated residue as tryptophan and not tyrosine. A similarly frequent assumption about

NO_2Tyr immunosignals is specificity with respect to the nitrated amino acid. Other nitration

target amino acids, however, have been reported, most prominently tryptophan (27–29). To

identify the physiologically nitrated LDH residue(s), heart tissue was homogenized and

subjected to 2D-PAGE in-gel digestion prior to MALDI-TOF MS analysis. Surprisingly,

signals consistent with a nitrated tryptophan at position 324 of peptide 319-328

(SADTLWGIQK) of LDH (NO_2Trp-LDH) were detected (**Fig. 5A**). We cannot exclude

nitration of other peptides of LDH that were below our level of detection. Nevertheless, our

results strongly point to NO_2Trp being detected by the supposedly anti-NO_2Tyr pAb1.

NO_2Trp-LDH molecular dynamics simulations suggested higher restriction of the

activity-essential interaction of the active site loop with the C-terminal α-helix. Since

we did observe a V_{max} effect on NO_2-LDH, we wanted to investigate whether there is a likely

structural correlate for this via Trp-324. We therefore modeled the nitration into the structure

of tetrameric and binary human LDH, and then performed molecular dynamics simulations.

For distance metrics of human LDH and key structural components refer to **Fig. 5B**. The

root-mean squared fluctuations (RMSF) of all atoms in each residue were averaged across

the four subunits in each combined 3.5 µs trajectory (**SI Fig. S2**). Subtracting the wildtype

tetramer (WT) RMSF from the T4N (tetramer with NO_2Trp-324 residue on each subunit) highlighted differences in mobility between the dynamics of the WT compared with NO_2Trp-LDH (**Fig. 5C**). Positive values in the resulting graphs indicated greater mobility of residues in T4N compared with WT and *vice versa*. Moreover, distance histograms (**Fig 5D/E**) show an increased mobility in the binary complexes compared with the ternary, expressed as a broadening of the histograms.

We found that NO_2Trp-LDH weakened the interaction between α-H and the active site loop. This decoupling led to a greater population of closed and "over-closed" states of the T4N active-site loop. By corollary, in WT LDH, the open active-site loop states are partly stabilized by interaction with the α-H helix. Therefore, nitration of LDH is expected to decrease LDH activity, consistent with our enzyme-kinetic findings (see **Fig. 2D-E**).

Discussion

We here close three major knowledge gaps in our understanding of protein nitration: specificity, origin, and disease relevance. With respect to the specificity of protein nitration, a vast heterogeneity of proteins are detected by supposedly uniform anti-NO_2Tyr antibodies. Moreover, tryptophan nitration (NO_2Trp) was a cofounding factor of our apparent NO_2Tyr-immunosignals. Protein NO_2Trp formation has been described in isolated proteins treated with H_2O_2/NO_2^- *in vitro* (30–33), peroxynitrite-exposed cells (34), and as physiological modification during aging in rat heart mitochondria (28). Trp is an alternative target of nitration that deserves more attention (35, 36) and immunosignals by anti-NO_2Tyr antibodies should rather be interpreted as NO_2Tyr/ NO_2Trp signals unless purified and chemically validated. Rebrin et al. (28) reported Trp nitration of succinyl-CoA:3-ketoacid CoA transferase (SCOT), which could not be reproduced by Wang et al. (37). Whether Trp nitration has a functional consequence on SCOT remained unclear because both activation and inactivation were observed (28). In hindsight, SCOT nitration is also the first documented example where Trp nitration has been mistaken as Tyr nitration based on cross-reactivity of a supposedly NO_2Tyr specific antibody (38), and many more of these examples may exist in the literature.

With respect to the origin of the nitrating nitrogen in NO_2Tyr/ NO_2Trp positive proteins, we show that both eNOS/NO and MPO/nitrite pathways, and possibly others, need to be equally considered and that the *a priori* assumption that NO_2Tyr immunosignals indicate NO scavenging by superoxide is not valid.

With respect to disease relevance, probably our most relevant finding is that protein nitration is a physiological event. This represents a similar dogma shift as for ROS formation, which also should not be *a priori* considered as oxidative stress and a disease marker. Thus, both ROS and reactive nitrogen and protein nitration represent physiological events, even though they may be upregulated in disease as we show for diabetes and cardiac stress.

Not many specific functional consequences have been described for protein nitration.

Assuming Trp nitration and activity modulation of both LDH (decreasing activity) and SCOT

(increasing activity) would occur, this may be a protective mechanism, allowing the heart to

better utilize ketone metabolism for energy production at a time when other metabolic

processes may be diminished (27, 28). Molecular dynamics simulations were performed on

NO_2-LDH, aiming to understand the impairment of the enzyme activity (see **Fig. 5**). While

differences in activity between WT and T4N of a factor of two is significant in physiological

terms, it corresponds to minor changes in the free energy states in the catalytic process.

The catalytic cycle of LDH is a well understood ordered bi-bi reaction mechanism where

substrate and product enter and leave the active-site, with concomitant closing and opening

of the active-site loop, after cofactor binding. Detailed kinetic characterization of LDH from

Bacillus stearothermophilus (39) and *Plasmodium falciparum* (40) has revealed this loop

movement to be a rate limiting step in the reaction mechanism. Any perturbation of this

delicately balanced conformational change is expected to compromise the catalytic activity

of the enzyme. We suggest that conformational movement of the active-site loop in cardiac

LDH is also a rate limiting step and the weakening of the interaction between this loop and

the α-H helix in T4N (as described in our results) is responsible for the impaired catalytic

activity.

In conclusion, cardiac LDH appears to be the first protein basally regulated by

nitration and not only a loss-of-function biomarker of metabolic stress. NO_2-LDH has a lower

V_{max} and the nitration is further increased under cardiac metabolic stress. Furthermore, *in*

vivo NO_2Tyr immunoreactivity not necessarily reflects Tyr nitration or solely NO scavenging.

Instead, it may indicate nitrite/MPO-dependent Trp nitration and physiological signaling. This

has wide implications on re-interpreting previous data using NO_2Tyr immunoreactivity as a

biomarker and on our understanding of nitration as a physiological protein modification and

signaling mechanism.

Material and Methods

Detailed experimental procedures are provided in *SI Materials and Methods*.

Footnotes

Footnote 1. Several publications on protein NO_2Tyr immunoreactions report the use of

NO_2Tyr concentrations of up to 10 mM to block the signal and demonstrate specificity; this

exceeds the solubility of NO_2Tyr. Moreover, these saturated solutions are highly acidic and

require buffering to ensure that any blockade of an immunosignal is due to competition with

free NO_2Tyr and not due to pH-induced antibody denaturation.

Author contributions: H.H.H.W.S. designed research; J.F., D.T., T.N., C.I.A., A.S., P.B.,

C.S., V.S., P.R., A.J., A.G., D.H.P., R.B.S. and M.F. performed research; A.S., A.G., A.K.,

S.B., D.H.P., M.F. and H.H.H.W.S. contributed new reagents/analytic tools; J.F., D.T.,

D.H.P., and H.H.H.W.S. analyzed data; and C.N., J.F., C.S., D.H.P., M.F., A.I.C. and

H.H.H.W.S. wrote the paper.

The authors declare no conflict of interests.

This article is a PNAS regular research article.

This article contains supplemental information (SI Appendix).

Acknowledgements

Work in A.J. laboratory was supported by BBSRC, BHF and MRC; in P.B. laboratory by NKFIH (K123975); in H.H.H.W.S. laboratory by an ERC Advanced Investigator Grant (294683 - RadMed), an ERC proof-of-concept grant (139-101052 - SAVEBRAIN) and the EU Horizon 2020 programme, REPO-TRIAL. The authors would like to thank G. Jansen for technical help, and Dr. P.M.H. Schiffers for carefully reading earlier stages of the manuscript. We also thank the Advanced Computing Research Centre at Bristol and BrisSynBio for access to the HPC machines BlueCrystal and BlueGem. We thank the HECBiosim Consortium for access to the UK National HPC facility Archer.

References

1. Sies H (2015) Oxidative stress: a concept in redox biology and medicine. *Redox Biol* 4:180–183.

2. Beckman JS, Koppenol WH (1996) Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol Physiol* 271(5 Pt 1):C1424-37.

3. Radi R (2004) Nitric oxide, oxidants, and protein tyrosine nitration. *Proc Natl Acad Sci* 101(12):4003–4008.

4. Armitage ME, Wingler K, Schmidt HHHW, La M (2009) Translating the oxidative stress hypothesis into the clinic: NOX versus NOS. *J Mol Med* 87(11):1071–1076.

5. Viappiani S, Schulz R (2006) Detection of specific nitrotyrosine-modified proteins as a marker of oxidative stress in cardiovascular disease. *AJP Hear Circ Physiol* 290(6):H2167–H2168.

6. Shishehbor MH, Hazen SL (2004) Inflammatory and oxidative markers in atherosclerosis: Relationship to outcome. *Curr Atheroscler Rep* 6(3):243–50.

7. Ho E, Karimi Galougahi K, Liu C-C, Bhindi R, Figtree GA (2013) Biological markers of oxidative stress: Applications to cardiovascular research and practice. *Redox Biol* 1(1):483–491.

8. Fernández E, García-Moreno J-M, Martín de Pablos A, Chacón J (2013) May the Evaluation of Nitrosative Stress Through Selective Increase of 3-Nitrotyrosine Proteins Other Than Nitroalbumin and Dominant Tyrosine-125/136 Nitrosylation of Serum α-Synuclein Serve for Diagnosis of Sporadic Parkinson's Disease? *Antioxid Redox Signal* 19(9):912–918.

9. Halliwell B (1997) What nitrates tyrosine? Is nitrotyrosine specific as a biomarker of peroxynitrite formation in vivo? *FEBS Lett* 411(2–3):157–160.

10. Ye YZ, Strong M, Huang Z-QQ, Beckman JS (1996) Antibodies that recognize nitrotyrosine. *Methods Enzymol* 269:201–209.

11. Reiter CD, Teng R-J, Beckman JS (2000) Superoxide Reacts with Nitric Oxide to Nitrate Tyrosine at Physiological pH via Peroxynitrite. *J Biol Chem* 275(42):32460–32466.

12. Breusing N, Grune T (2010) Biomarkers of protein oxidation from a chemical, biological and medical point of view. *Exp Gerontol*. doi:10.1016/j.exger.2010.04.004.

13. Sampson JB, Ye Y, Rosen H, Beckman JS (1998) Myeloperoxidase and Horseradish Peroxidase Catalyze Tyrosine Nitration in Proteins from Nitrite and Hydrogen Peroxide. *Arch Biochem Biophys* 356(2):207–213.

14. Bigelow DJ (2009) Nitrotyrosine-modified SERCA2: a cellular sensor of reactive nitrogen species. *Pflügers Arch - Eur J Physiol* 457(3):701–710.

15. Knyushko T V., Sharov VS, Williams TD, Schöneich C, Bigelow DJ (2005) 3-Nitrotyrosine Modification of SERCA2a in the Aging Heart: A Distinct Signature of the Cellular Redox Environment †. *Biochemistry* 44(39):13071–13081.

16. Li H, Gutterman DD, Rusch NJ, Bubolz A, Liu Y (2004) Nitration and Functional Loss of Voltage-Gated K+ Channels in Rat Coronary Microvessels Exposed to High Glucose. *Diabetes* 53(9):2436–2442.

17. Haddad IY, Zhu S, Ischiropoulos H, Matalon S (1996) Nitration of surfactant protein A results in decreased ability to aggregate lipids. *Am J Physiol Cell Mol Physiol* 270(2):L281-8.

18. Akaike T, et al. (1996) Pathogenesis of influenza virus-induced pneumonia: involvement of both nitric oxide and oxygen radicals. *Proc Natl Acad Sci* 93(6):2448–2453.

19. Akaike T, et al. (2003) 8-Nitroguanosine formation in viral pneumonia and its implication for pathogenesis. *Proc Natl Acad Sci* 100(2):685–690.

20. Evans TJ, et al. (1996) Cytokine-treated human neutrophils contain inducible nitric oxide synthase that produces nitration of ingested bacteria. *Proc Natl Acad Sci*

1   93(18):9553–9558.

21.   Hayashi T, et al. (2006) Modulating role of estradiol on arginase II expression in hyperlipidemic rabbits as an atheroprotective mechanism. *Proc Natl Acad Sci* 103(27):10485–10490.

22.   Szabo C, et al. (1998) Protection against peroxynitrite-induced fibroblast injury and arthritis development by inhibition of poly(ADP-ribose) synthase. *Proc Natl Acad Sci* 95(7):3867–3872.

23.   Graham ML, Janecek JL, Kittredge JA, Hering BJ, Schuurman HJ (2011) The streptozotocin-induced diabetic nude mouse model: Differences between animals from different sources. *Comp Med*. doi:10.1152/ajprenal.00292.2015.

24.   Berthiaume JM, Wallace KB (2007) Adriamycin-induced oxidative mitochondrial cardiotoxicity. *Cell Biology and Toxicology* doi:10.1007/s10565-006-0140-y.

25.   Peluffo G, Radi R (2007) Biochemistry of protein tyrosine nitration in cardiovascular pathology. *Cardiovasc Res* 75(2):291–302.

26.   van der Vliet A, Eiserich JP, Halliwell B, Cross CE (1997) Formation of Reactive Nitrogen Species during Peroxidase-catalyzed Oxidation of Nitrite. *J Biol Chem* 272(12):7617–7625.

27.   Nuriel T, Hansler A, Gross SS (2011) Protein nitrotryptophan: Formation, significance and identification. *J Proteomics* 74(11):2300–2312.

28.   Rebrin I, Brégère C, Kamzalov S, Gallaher TK, Sohal RS (2007) Nitration of Tryptophan 372 in Succinyl-CoA:3-Ketoacid CoA Transferase during Aging in Rat Heart Mitochondria †. *Biochemistry* 46(35):10130–10144.

29.   Souza JM, Peluffo G, Radi R (2008) Protein tyrosine nitration—Functional alteration or just a biomarker? *Free Radic Biol Med* 45(4):357–366.

30.   Suzuki T, et al. (2004) Nitration and nitrosation of N-acetyl-l-tryptophan and tryptophan residues in proteins by various reactive nitrogen species. *Free Radic Biol*

1   *Med* 37(5):671–681.

31.   Yamakura F, Ikeda K (2006) Modification of tryptophan and tryptophan residues in proteins by reactive nitrogen species. *Nitric Oxide* 14(2):152–161.

32.   Yamakura F, et al. (2005) Nitrated and Oxidized Products of a Single Tryptophan Residue in Human Cu,Zn-Superoxide Dismutase Treated with Either Peroxynitrite-Carbon Dioxide or Myeloperoxidase-Hydrogen Peroxide-Nitrite. *J Biochem* 138(1):57–69.

33.   Yamakura F, Matsumoto T, Taka H, Fujimura T, Murayama K (2003) 6-Nitrotryptophan: a specific reaction product of tryptophan residue in human Cu, Zn-SOD treated with peroxynitrite. *Adv Exp Med Biol* 527:745–9.

34.   Ikeda K, et al. (2007) Detection of 6-nitrotryptophan in proteins by Western blot analysis and its application for peroxynitrite-treated PC12 cells. *Nitric Oxide* 16(1):18–28.

35.   Alvarez B, et al. (1996) Peroxynitrite-Dependent Tryptophan Nitration. *Chem Res Toxicol* 9(2):390–396.

36.   Herold S (2004) Nitrotyrosine, dityrosine, and nitrotryptophan formation from metmyoglobin, hydrogen peroxide, and nitrite. *Free Radic Biol Med* 36(5):565–579.

37.   Wang Y, et al. (2010) The Nitrated Proteome in Heart Mitochondria of the db/db Mouse Model: Characterization of Nitrated Tyrosine Residues in SCOT. *J Proteome Res* 9(8):4254–4263.

38.   Marcondes S, Turko I V, Murad F (2001) Nitration of succinyl-CoA:3-oxoacid CoA-transferase in rats after endotoxin administration. *Proc Natl Acad Sci* 98(13):7146–7151.

39.   Wilks HM, et al. (1990) Designs for a broad substrate specificity keto acid dehydrogenase. *Biochemistry* 29(37):8587–8591.

40.   Shoemark DK, Cliff MJ, Sessions RB, Clarke AR (2007) Enzymatic properties of the

lactate dehydrogenase enzyme from Plasmodium falciparum. *FEBS J* 274(11):2738–2748.

41. Viera L, Ye YZ, Estévez AG, Beckman JS (1999) [39] Immunohistochemical methods to detect nitrotyrosine. *Methods in Enzymology*, pp 373–381.

42. Görg A, Weiss W, Dunn MJ (2004) Current two-dimensional electrophoresis technology for proteomics. *Proteomics* 4(12):3665–3685.

43. Perlman DH, Berg EA, O'Connor PB, Costello CE, Hu J (2005) Reverse transcription-associated dephosphorylation of hepadnavirus nucleocapsids. *Proc Natl Acad Sci* 102(25):9020–9025.

44. Karas M, Hillenkamp F (1988) Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *Anal Chem* 60(20):2299–2301.

45. Sousa da Silva AW, Vranken WF (2012) ACPYPE - AnteChamber PYthon Parser interfacE. *BMC Res Notes* 5(1):367.

46. Lindorff-Larsen K, et al. (2010) Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins Struct Funct Bioinforma* 78(8):NA-NA.

47. Abraham MJ, et al. (2015) GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1–2:19–25.

48. Humphrey W, Dalke A, Schulten K (1996) VMD: Visual molecular dynamics. *J Mol Graph* 14(1):33–38.

49. Pettersen EF, et al. (2004) UCSF Chimera?A visualization system for exploratory research and analysis. *J Comput Chem* 25(13):1605–1612.

50. Pacher P (2003) Potent Metalloporphyrin Peroxynitrite Decomposition Catalyst Protects Against the Development of Doxorubicin-Induced Cardiac Dysfunction. *Circulation* 107(6):896–904.

51. Perlman DH, et al. (2009) Mechanistic Insights Into Nitrite-Induced Cardioprotection

1  Using an Integrated Metabolomic/Proteomic Approach. *Circ Res* 104(6):796–804.

Figures and figure legends

anti-NO$_2$Tyr antibody based biomarker detection

Knock-out	anti-NO$_2$Tyr Ab panel screening	Mechanism

Enzymatic sources of RNS

RNS → NO$_2$

Disease specificity

NO$_2$ ↔ NO$_2$

Mechanism

p38^{nt} antibody-guided purification
MALDI-MS protein identification
NO$_2$-LDH activity

NO$_2$

NO$_2$-LDH MALDI-TOF MS-based residue identification

Function

Protein structure

Mouse

Rat

Pig

Human

Figure 1. Experimental outline for an anti-NO$_2$Tyr antibody-based biomarker detection approach. Central panel: Antibody panel screening, an anti-NO$_2$Tyr antibody panel was tested in porcine and rat tissues under physiological conditions. Different antibodies might recognize different nitrated proteins (green and orange antibodies) or one antibody might detect more than one nitrated protein (blue antibody). Right panel: A 38 kDa band was especially prominent and, subsequently, purified in large mammals and identified via MALDI-TOF as LDH. NO$_2$-LDH activity was also assessed. We identified the nitrated residue and simulated the effect of the nitration in the protein structure and mechanism using human LDH. Left panel: Different rodent models, as well as knock-out mouse models were used to elucidate the effects of NO$_2$-LDH in disease, diabetes and myocardial stress, and the enzymatic sources of reactive nitrogen species (RNS) leading to LDH nitration, respectively.

Figure 2. Screening of different native tissue homogenates identifies a 38 kDa band in heart as major physiologically NO$_2$Tyr-immuno positive protein. Further purification and MALDI-MS analysis identified the 38 kDa band as LDH. *(A)* Immunopositive pig and rat tissues from Fig S1 were further analyzed by SDS-PAGE under reducing conditions (dithiothreitol) and subsequent western blot. The band pattern was different for all three polyclonal (pAb) or monoclonal (mAb) antibodies: pAb1, mAb1 and pAb2. The most intense NO$_2$Tyr immunopositive protein band at approximately 38 kDa (p38^{nt}) was detected using pAb1 in heart. *(B)* p38^{nt} was purified from porcine heart homogenate by affinity (Blue Sepharose) followed by ion-exchange (Q-Sepharose) chromatography. Purification steps were tracked via protein nitrotyrosine detection using pAb1. *(C)* Purified cardiac p38^{nt} was cut out from the gel, digested with trypsin and analyzed

by mass spectrometry and peptide mass fingerprint analysis. Most peptide ions detected

were matched to lactate dehydrogenase: 11 ions matched to LDH (■) out of a total of 20

detected, with 5 ions matching to the next best protein assignment, malate dehydrogenase

(MDH, ☐), identifying p38^{nt} as LDH with a $p < 0.05$. *(D)* LDH nitration was assessed by

quantitative western blot analysis using pAb1 (n=9). *(E)* Vmax[LDH] remained unchanged

up to an equimolar peroxynitrite:LDH ratio, and then decreased at a ratio of 10 (n=8). Data

represent means ±SEM.

Figure 3. Pathological LDH nitration levels in short-term (ST) and long-term (LT)

diabetic disease models and an oxidative stress myocardial dysfunction model. *(A)*

Heart homogenates of ST and LT streptozotocin-treated diabetic rats or doxorubicin-treated

mice (DoxoR) were analyzed by SDS-PAGE and immunoblot analysis for LDH nitration

using the pAb1 antibody. *(B)* In all three animal models, the relative NO_2-LDH signal intensity

was increased. In diabetes, the long-term model yielded higher values than the short-term

model, which may indicate either higher nitrosative stress, or, more likely, accumulation of

LDH nitration. The highest increase was observed in the doxorubicin model. Data are

expressed as % of untreated control animals (open bars) and represent means ± SEM of

n=6-18 experiments.

Figure 4. *In vivo* genetic validation of reactive nitrogen species (RNS) sources. *(A)*

Events leading to oxidized and nitrated LDH. RNS (represented as $NO_2^-/H_2O_2/ONOO^-$) can

nitrate LDH, resulting in decreased V_{max}. *(B)* Cardiac tissue was obtained from endothelial

nitric oxide synthase (eNOS) and myeloperoxidase (MPO) knock-out (KO), and the

respective wild-type (WT), mice. Basal nitration of LDH (p38^{nt}) was analyzed by immunoblot,

and immunosignals quantified densitometrically. Data are expressed as % of control (i.e., of

the respective WT animals) and represent means ± SEM of n=3-5 animals. NO_2-LDH

depended on both eNOS and MPO.

A

NO₂-LDH MALDI-TOF MS-based residue identification

Intensity [a.u.]

1000   2500 m/z

319-328
1176.61

319-328 + O-C
1180.60

319-328 + O-2H
1190.60

Nitro/Ox

³¹⁹SADTLWDIQK₃₂₈

319-328 + O
1192.61

319-328
1186.65

319-328 + NO₂-O
1205.65

319-328 + 2O
1208.61

319-328 -2H
1174.59

319-328 + NO₂
1221.66

Intensity [a.u.]

1170   1230 m/z

B

Protein structure
WT human LDH

6.2Å

7.6Å

- ▬ b-K--b-L loop
- ▬ a-H C-terminal helix
- ▬ Active-site loop
- ▬ a-2G
- ▬ Pyruvate
- ▬ NADH

C

WT and T4N backbone RMSF differences per residue

Binary complex (NADH)

RMSF (nm)

0.10
0.05
0.00
-0.05
-0.10

0   50   100   150   200   250   300   350
Residue number

Ternary complex (NADH + pyruvate)

RMSF (nm)

0.10
0.05
0.00
-0.05
-0.10

0   50   100   150   200   250   300   350
Residue number

D

Association of the active site with a-H (m:LH)

Binary complex

Count frequency (nm⁻¹)

WT
T4N

0.2   0.6   1.0   1.4
Distance (nm)

Ternary complex

Count frequency (nm⁻¹)

0.2   0.6   1.0   1.4
Distance (nm)

E

Openness of the active site loop (m:LO)

Binary complex

Count frequency (nm⁻¹)

0.2   0.6   1.0   1.4
Distance (nm)

Ternary complex

Count frequency (nm⁻¹)

0.2   0.6   1.0   1.4
Distance (nm)

Figure 5. Identification of LDH Trp-324 as physiologically nitrated and oxidized. The interaction of the active site loop with the C-terminal α-helix is essential and restricted in NO$_2$Trp-LDH. *(A)* Rat heart homogenate was subjected to 2D-PAGE and Coomasie staining; spots were subjected to in-gel digestion and MALDI-TOF MS analysis. Left panel: Mass spectrum derived from a spot corresponding to rat LDH over the range m/z 600 to 2700. Right panel: An expansion of the same spectrum over the range m/z 1170 to 1230, showing peptide ions assigned primarily to the peptide 319-328 (319SADTLWDIQK328), and its oxidized and nitrated derivatives. Major peptide ions are labeled with their observed m/z values and corresponding LDH mature protein amino acid intervals, as assigned by Mascot™, and additionally by manual comparison to predicted values calculated from the LDH sequence and oxidative and nitrative modifications. The ions correspond to the peptide 319-328 Trp (W)-nitrated species (319-328 + NO2) and its typical high energy decomposition product (319-328 + NO2-O), as well as ions corresponding to the characteristic family of Trp oxidation forms, hydroxytryptophan (319-328 + Ox), didehydro-hydroxytryptophan (319-328 + Ox-2H), dihydroxytryptophan (319-328 + 2Ox), and kynurenin (319-328 + Ox-C). Other peptide ions display single (+ Ox) and double (+ 2Ox) oxidation, or didehydrogenation (- 2H). *(B)* Distance metrics mapped onto the crystal structure of WT (1I0Z). Color scheme: β-K--β-K loop, dark blue; α-helix, pink; active-site loop, green; α-2G, light blue; pyruvate, grey; NADH, cyan. Residue Trp-324 is shown in space filling. *(C)* Plot of the differences between T4N and WT backbone RMSF per residue (T4N – WT) in the binary complex. Structures are highlighted using the same color scheme as previously described plus the juxtaposed loop to the C-terminal α-helix which is highlighted in light orange. (D – E) Histograms of the distribution of distances. Blue WT; Red T4N. Metrics m:LH (107:CB-326:CB) showing the juxtaposition of the active-site loop and α-C-terminal helix and m:LO (103:CA-243:CB) showing the open/closed state of the active-site loop.

anti-NO$_2$Tyr antibody based biomarker detection

Knock-out

Enzymatic sources of RNS

RNS → NO$_2$

Disease specificity

anti-NO$_2$Tyr Ab panel screening

NO$_2$ ↔ NO$_2$

Mechanism

p38^{nt} antibody-guided purification
MALDI-MS protein identification
NO$_2$-LDH activity

NO$_2$

NO$_2$-LDH MALDI-TOF MS-based residue identification

Function

Protein structure

Mouse

Rat

Pig

Human

A

pAb1

Pig / Rat

anti-NO₂Tyr antibody screening

pAb1

Pig / Rat

pAb2

Pig / Rat

B

p30⁺ antibody-guided purification

H   FT1   FT2   E

C

MALDI-MS protein identification

Lactata dehydrogenase

11 hits
9 hits

D

NO₂-LDH activity

Control   1   10

E

Pig

Control   1   10

A

Enzymatic sources of RNS

$eNOS$

Diet

$NO \xrightarrow{O_2} NO_2 \rightarrow NO_2^-$

O_2^-

H_2O_2

MPO

$ONOO^-$

RNS-dependent protein nitration

B

Mouse

NO_2-LDH Immunosignal [% of control]

A

NO$_2$-LDH MALDI-TOF MS-based residue identification

Intensity [a.u.]

1000   2500 m/z

319-328
1176.61

319-328 + O-C
1180.60

319-328 + O
1192.61

319-328 + O-2H
1190.60

Nitro/Ox
^{319}SADTLWDIQK$_{328}$

319-328
1186.65

**319-328
-2H**
1174.59

319-328 + NO$_2$-O
1205.65

319-328 + 2O
1208.61

319-328 + NO$_2$
1221.66

Intensity [a.u.]

1170   1230 m/z

B

Protein structure
WT human LDH

6.2Å

7.6Å

β-K--β-L loop
α-H C-terminal helix
Active-site loop
α-2G
Pyruvate
NADH

C

WT and T4N backbone RMSF differences
per residue

Binary complex (NADH)

RMSF (nm)

0.10
0.05
0.00
-0.05
-0.10

0   50   100 150 200 250 300 350
Residue number

Ternary complex (NADH + pyruvate)

RMSF (nm)

0.10
0.05
0.00
-0.05
-0.10

0   50   100 150 200 250 300 350
Residue number

D

Association of the
active site with α-H
(m:LH)

Binary complex

Count frequency (nm^{-1})

WT
T4N

0.2   0.6   1.0   1.4
Distance (nm)

Ternary complex

Count frequency (nm^{-1})

0.2   0.6   1.0   1.4
Distance (nm)

E

Openness of the
active site loop
(m:LO)

Binary complex

Count frequency (nm^{-1})

0.2   0.6   1.0   1.4
Distance (nm)

Ternary complex

Count frequency (nm^{-1})

0.2   0.6   1.0   1.4
Distance (nm)

Antigen \ Antibody	pAb1 1:3000		pAb2 1:3000		mAb1 1:3000	
NO$_2$Tyr-Block		+		+		+
NO$_2$BSA (Control)						
Pul. Artery (pig)						
Heart (pig)						
Lung (rat)						
Kidney (rat)						
Cerebellum (rat)						
Spleen (rat)						
Cortex (rat)						