Article

Tryptophan nitration physiologically regulates cardiac lactate dehydrogenase active

site loop mobility and activity

Cristian Nogales[1,*], Jeroen Frijhoff[1], Detlef Thoenges, Richard B. Sessions, Albert

Sickmann, Tatiana Nedvetskaya, Cesar Ibarra-Alvarado, Vera Schmitz, Peter Rösen, Axel

Goedecke, Aleksandar Jovanović, David H. Perlman, Anna Klinke, Stephan Baldus, Ana I.

Casas, Péter Bai, Csaba Szabo, Martin Feelisch & Harald H.H.W. Schmidt[*].

[1]Both first authors contributed equally to this work.

[*]To whom correspondence may be addressed. Email: h.schmidt@maastrichtuniversity.nl

or c.nogales@maastrichtuniversity.nl

Abstract (140 words)

Immunopositive protein tyrosine nitration is considered a disease marker for oxidative

stress. The mechanism of nitration, target proteins and functional consequences, however,

remain often unclear. We here focus on a prominent protein band in mouse, rat and pig

heart surprisingly nitrotyrosine immunopositive under basal conditions. Upon purification,

we identify it as lactate dehydrogenase (LDH) and its basal nitration mechanism

depending on NO synthase and myeloperoxidase. Surprisingly, we locate LDH nitration by

mass spectrometry not to tyrosine but to a C-terminal tryptophan, Trp-324. Molecular

dynamics simulations suggested that Trp-324 nitration restricts the interaction of the active

site loop with the C-terminal α-helix essential for activity explaining the observed lower

V_{max}. Protein nitration is thus not merely a disease marker but also a physiological event.

In cardiac LDH, this extends to tryptophan modification limiting enzyme activity under

basal and metabolic stress conditions.

Increased levels of reactive oxygen species (ROS), i.e. oxidative stress, is considered a

common mechanism of several cardiovascular and other disease states [1]. In concert with

different nitrogen species [2–4], ROS can lead to protein nitration, e.g. tyrosine residues

(NO_2Tyr). Tyrosine nitration is the most frequently used and considered the most robust

disease marker of oxidative stress [5–7]. Its detection relies mainly on NO_2Tyr specific

antibodies [8–10]. With respect to the nitrogen source, anti-NO_2Tyr immunopositive signals

are generally interpreted as a hallmark of nitric oxide (NO) being scavenged by superoxide

and intermediate peroxynitrite [9,11,12], although other mechanisms exist [13]. Regarding the

functional consequences of protein nitration few *in vitro* data exist suggesting mainly loss

of protein function correlating with disease severity [14–17]. *In vivo* data are lacking.

Moreover, little is known about the protein factors that govern other nitration sites, e.g.

tryptophan, and about the effects of the modification in protein structure and function [18].

Here, we address three major knowledge gaps in our understanding of protein nitration:

specificity, origin, and disease relevance.

Results

Development of an anti-nitrotyrosine antibody panel as biomarkers of disease-

relevant oxidative stress.

We started out by examining a panel of anti-NO_2Tyr antibodies (Fig. 1), including those

directed against carrier-linked NO_2Tyr or against H_2O_2/NO_2^- treated proteins (see

Supplementary Table 1). We tested these in rodent (rat) and, for later purification, also in

porcine tissues. Surprisingly we obtained signals already under physiological conditions.

The immunoreaction, however, varied greatly in protein band size and intensity. Two

polyclonal (pAb), pAb1 and pAb2, and one monoclonal antibody (mAb), mAb1, displayed

the most robust signals (Supplementary Fig. 1). The specificity of anti-NO_2Tyr antibodies

[19–23] is rarely validated, e.g. by antibody pre-absorption with free NO_2Tyr [11]. In our case,

pAb1 signals were partially blocked by pre-absorption with 3 mM NO_2Tyr. mAb1 showed

very little non-specific background whilst pAb2 exhibited moderate sensitivity with high

anti-NO_2Tyr specificity.

A prominent nitrotyrosine immunoreactive protein band was detected in mouse, rat

and pig heart under physiological conditions.

To conduct a tissue/species screen, we selected three antibodies, pAb1, pAb2, and mAb1,

from our original panel. Three proteins were consistently detected under basal conditions

and with apparent molecular weights of 70, 45 and 38 kDa, of which the latter gave the by

far the strongest signal in heart (Fig. 2a). Pre-incubation with free NO_2Tyr or NO_2BSA (3

mM) or nitro-group reduction [22], at least partially, blocked the pAb1 dependent signal.

Therefore, subsequent efforts were directed at identifying the nature of the NO_2Tyr positive

38 kDa protein and its site of nitration.

Upon purification, the 38 kDa nitrated protein was identified as LDH with lower V_{max}.

Using porcine heart, we isolated the NO_2Tyr immunoreactive 38 kDa protein by ion

exchange (Q Sepharose, QS) and affinity (Blue Sepharose, BS) chromatography (Fig. 2b).

The partially purified 38 kDa protein was stained by colloidal Coomassie Blue and

subjected to peptide mass fingerprinting using MALDI-TOF MS. Based on the NCBIprot

database most ions present in the spectrum of tryptic peptides matched LDH as the

primary protein component of the sample (Fig. 2c).

Next, we tested the effect of nitration on the enzyme activity by nitrating cardiac LDH *in*

vitro with increasing amounts of H_2O_2/NO_2^-. Indeed, higher nitration levels of LDH (NO_2-

LDH) (Fig. 2e) correlated with decreased V_{max} (Fig. 2f) and unchanged K_M. Our data

suggest therefore that protein nitration is not only a physiological event for cardiac LDH,

but also modulates enzyme activity by lowering V_{max}.

Nitrated LDH is increased in diabetes and myocardial stress.

To investigate whether NO_2-LDH is further increased in response to metabolic stress, we

investigated heart extracts from rat models of streptozotocin-induced short-term (ST) and

long-term (LT) diabetes mellitus [24] as well as a doxorubicin (DoxoR) induced mouse model

of cardiac stress [25] (Fig. 3a). Diabetic rats were sacrificed 3 (ST) and 16 (LT) weeks after

inducing diabetes at week 10. Using the pAb1, NO_2-LDH signals were significantly

increased in all three models when compared against healthy controls (Fig. 3b-c).

Immunosignals of pAb1 pre-absorbed with NO_2Tyr or anti-LDH wildtype antibody did not

show any significant change. In both diabetes mellitus models and doxorubicin-treatment

animals, levels of cardiac LDH were unchanged versus controls, indicating that indeed the

NO_2-LDH/LDH ratio was increased.

1 **Cardiac LDH nitration is lowered in NO synthase (eNOS) knockout (KO) and**

2 **myeloperoxidase (MPO) KO mice.**

3 We next wanted to address our second goal, to understand the mechanistic origin of the

4 nitrogen in NO_2-LDH. Usually, it is assumed that protein nitration derives from peroxynitrite

5 from the reaction between $^{\bullet}NO$ and $O_2^{\bullet -}$ [26]. However, another major source, not involving

6 de novo NO synthesis is frequently overlooked, i.e. MPO-catalyzed nitration using nitrite in

7 the presence of hydrogen peroxide [27] (Fig. 4a). To analyze this *in vivo*, we had to switch to

8 mouse models, i.e. eNOS-KO and MPO-KO lines. Indeed, we observed that NO_2-LDH

9 signals in heart tissue were not only reduced in eNOS-KO but also in MPO-KO mice when

10 compared to their respective WT litter mates (Fig. 4b). These results therefore indicated

11 that both pathways contribute to basal nitration of cardiac LDH. Moreover, we also

12 compared LDH nitration to oxidation and showed that oxidation occurred at lower

13 peroxynitrite concentrations and was half-maximal when LDH activity was still unchanged

14 (Supplementary Fig. 2).

15

16 **Further MALDI-TOF mass spectrometry analysis of the nitration site identifies the**

17 **nitrated residue as tryptophan and not tyrosine.**

18 A similarly frequent assumption about NO_2Tyr immunosignals is specificity with respect to

19 the nitrated amino acid. Other nitration target amino acids, however, have been reported,

20 most prominently tryptophan [28–30]. To identify the physiologically nitrated LDH residue(s),

21 heart tissue was homogenized and subjected to 2D-PAGE in-gel digestion prior to MALDI-

22 TOF MS analysis. Surprisingly, signals consistent with a nitrated tryptophan at position

23 324 of peptide 319-328 (SADTLWGIQK) of LDH (NO_2Trp-LDH) were detected (Fig. 5a).

24 We cannot exclude nitration of other peptides of LDH that were below our level of

25 detection. Nevertheless, our results strongly point to NO_2Trp being detected by the

26 supposedly anti-NO_2Tyr pAb1.

NO$_2$Trp-LDH molecular dynamics simulations suggested higher restriction of the activity-essential interaction of the active site loop with the C-terminal α-helix.

Since we did observe a V$_{max}$ effect on NO$_2$-LDH, we wanted to investigate whether there is a likely structural correlate for this via Trp-324. We therefore modeled the nitration into the structure of tetrameric and binary human LDH, and then performed molecular dynamics simulations. For distance metrics of human LDH and key structural components refer to Fig. 5b. The root-mean squared fluctuations (RMSF) of all atoms in each residue were averaged across the four subunits in each combined 3.5 µs trajectory (Supplementary Fig. 3). Subtracting the wildtype tetramer (WT) RMSF from the T4N (tetramer with NO$_2$Trp-324 residue on each subunit) highlighted differences in mobility between the dynamics of the WT compared with NO$_2$Trp-LDH (Fig. 5c). Positive values in the resulting graphs indicated greater mobility of residues in T4N compared with WT and *vice versa*. Moreover, distance histograms (Fig. 5d-e) show an increased mobility in the binary complexes compared with the ternary, expressed as a broadening of the histograms.

We found that NO$_2$Trp-LDH weakened the interaction between α-H and the active site loop. This decoupling led to a greater population of closed and "over-closed" states of the T4N active-site loop. By corollary, in WT LDH, the open active-site loop states are partly stabilized by interaction with the α-H helix. Therefore, nitration of LDH is expected to decrease LDH activity, consistent with our enzyme-kinetic findings (see Fig. 2d-e).

Discussion

We here close three major knowledge gaps in our understanding of protein nitration: specificity, origin, and disease relevance. With respect to the specificity of protein nitration, a vast heterogeneity of proteins are detected by supposedly uniform anti-NO_2Tyr antibodies. Moreover, tryptophan nitration (NO_2Trp) was a cofounding factor of our apparent NO_2Tyr-immunosignals. Protein NO_2Trp formation has been described in isolated proteins treated with H_2O_2/NO_2^- *in vitro* [31–34], peroxynitrite-exposed cells [35], and as physiological modification during aging in rat heart mitochondria [29]. Trp is an alternative target of nitration that deserves more attention [36,37] and immunosignals by anti-NO_2Tyr antibodies should rather be interpreted as NO_2Tyr/ NO_2Trp signals unless purified and chemically validated. Rebrin et al. [29] reported Trp nitration of succinyl-CoA:3-ketoacid CoA transferase (SCOT), which could not be reproduced by Wang et al. [38]. Whether Trp nitration has a functional consequence on SCOT remained unclear because both activation and inactivation were observed [29]. In hindsight, SCOT nitration is also the first documented example where Trp nitration has been mistaken as Tyr nitration based on cross-reactivity of a supposedly NO_2Tyr specific antibody [39], and many more of these examples may exist in the literature.

With respect to the origin of the nitrating nitrogen in NO_2Tyr/ NO_2Trp positive proteins, we show that both eNOS/NO and MPO/nitrite pathways, and possibly others, need to be equally considered and that the *a priori* assumption that NO_2Tyr immunosignals indicate NO scavenging by superoxide is not valid. Nevertheless, the effects of KO of eNOS and MPO are significant but not great. Previous data shows that NO modifications persist in the absence of active NO synthesis and dietary NOx intake (e.g., in eNOS$^{-/-}$ animals on a low-NOx diet and under the acute action of the NOS-inhibitor, L-NIO, or in Wistar rats on low-NOx diet with chronic administration of L-NMMA) [40]. Moreoever, NO levels remain essentially unchanged upon deletion of each NOS isoform. Accordingly, these findings

reflect the action of a compensatory mechanism that maintains basal levels of NO

metabolites in tissues e.g. constant upregulation of remaining NOSs in an attempt to

restore basal levels.

With respect to disease relevance, probably our most relevant finding is that protein

tryptophan nitration is a physiological event. This represents a similar dogma shift as for

ROS formation, which also should not be *a priori* considered as oxidative stress and a

disease marker. Thus, both ROS and reactive nitrogen and protein nitration represent

physiological events, even though they may be upregulated in disease as we show for

diabetes and cardiac stress (see Fig. 3).

Not many specific functional consequences have been described for protein nitration.

Assuming Trp nitration and activity modulation of both LDH (decreasing activity) and

SCOT (increasing activity) would occur, this may be a protective mechanism, allowing the

heart to better utilize ketone metabolism for energy production at a time when other

metabolic processes may be diminished [28,29]. Molecular dynamics simulations were

performed on NO_2-LDH, aiming to understand the impairment of the enzyme activity (see

Fig. 5). While differences in activity between WT and T4N of a factor of two is significant in

physiological terms, it corresponds to minor changes in the free energy states in the

catalytic process. The catalytic cycle of LDH is a well understood ordered bi-bi reaction

mechanism where substrate and product enter and leave the active-site, with concomitant

closing and opening of the active-site loop, after cofactor binding. Detailed kinetic

characterization of LDH from *Bacillus stearothermophilus* [41] and *Plasmodium falciparum* [42]

has revealed this loop movement to be a rate limiting step in the reaction mechanism. Any

perturbation of this delicately balanced conformational change is expected to compromise

the catalytic activity of the enzyme. We suggest that conformational movement of the

active-site loop in cardiac LDH is also a rate limiting step and the weakening of the

interaction between this loop and the α-H helix in T4N (as described in our results) is

responsible for the impaired catalytic activity. Whether tryptophan nitration effects the

conversion of either pyruvate or other 2-oxo/-2-hydroxy acids warrants further

investigation.

Like many enzymes, LDH post-transcriptional activity is regulated by phosphorylation and

acetylation of amino acid residues. The tyrosine kinase FGFR1 has been shown to directly

phosphorylate LDH at Y10 and Y83 [43]. Y10 phosphorylation of LDH promotes active,

tetrameric LDHA formation, whereas phosphorylation of Y83 promotes NADH substrate

binding. Our finding obviously promotes the opposite, suggesting an antagonistic effects of

tyrosine phosphorylation and tryptophan nitration. Inhibition of LDH reduces aerobic

glycolysis, gene transcription, channel complex regulation, molecular chaperones and cell

cycle regulation. Cells are then forced to use oxidative phosphorylation and pyruvate

enters the mitochondria. This leads to reactive oxygen species (ROS) generation [44]. Since

the nitration is already a ROS-induced event, this may be a mechanism of ROS-induced

ROS formation [45], which in heart is protective triggering for example capillary density and

angiogenesis [46,47].

In conclusion, cardiac LDH appears to be the first protein basally regulated by nitration

and not only a loss-of-function biomarker of metabolic stress. NO_2-LDH has a lower V_{max}

and the nitration is further increased under cardiac metabolic stress. Furthermore, *in vivo*

NO_2Tyr immunoreactivity not necessarily reflects Tyr nitration or solely NO scavenging.

Instead, it may indicate nitrite/MPO-dependent Trp nitration and physiological signaling.

This has wide implications on re-interpreting previous data using NO_2Tyr immunoreactivity

as a biomarker and on our understanding of nitration as a physiological protein

modification and signaling mechanism.

Methods

Detailed experimental procedures are provided in Supplementary Methods.

Word count (main text without references): 2137 words.

References

1. Sies, H. Oxidative stress: a concept in redox biology and medicine. *Redox Biol.* **4,** 180–183 (2015).

2. Beckman, J. S. & Koppenol, W. H. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am. J. Physiol. Physiol.* **271,** C1424-37 (1996).

3. Radi, R. Nitric oxide, oxidants, and protein tyrosine nitration. *Proc. Natl. Acad. Sci.* **101,** 4003–4008 (2004).

4. Armitage, M. E., Wingler, K., Schmidt, H. H. H. W. & La, M. Translating the oxidative stress hypothesis into the clinic: NOX versus NOS. *J. Mol. Med.* **87,** 1071–1076 (2009).

5. Viappiani, S. & Schulz, R. Detection of specific nitrotyrosine-modified proteins as a marker of oxidative stress in cardiovascular disease. *AJP Hear. Circ. Physiol.* **290,** H2167–H2168 (2006).

6. Shishehbor, M. H. & Hazen, S. L. Inflammatory and oxidative markers in atherosclerosis: Relationship to outcome. *Curr. Atheroscler. Rep.* **6,** 243–50 (2004).

7. Ho, E., Karimi Galougahi, K., Liu, C.-C., Bhindi, R. & Figtree, G. A. Biological markers of oxidative stress: Applications to cardiovascular research and practice. *Redox Biol.* **1,** 483–491 (2013).

8. Fernández, E., García-Moreno, J.-M., Martín de Pablos, A. & Chacón, J. May the Evaluation of Nitrosative Stress Through Selective Increase of 3-Nitrotyrosine Proteins Other Than Nitroalbumin and Dominant Tyrosine-125/136 Nitrosylation of Serum α-Synuclein Serve for Diagnosis of Sporadic Parkinson's Disease? *Antioxid. Redox Signal.* **19,** 912–918 (2013).

9. Halliwell, B. What nitrates tyrosine? Is nitrotyrosine specific as a biomarker of peroxynitrite formation in vivo? *FEBS Lett.* **411,** 157–160 (1997).

10. Ye, Y. Z., Strong, M., Huang, Z.-Q. Q. & Beckman, J. S. Antibodies that recognize

nitrotyrosine. *Methods Enzymol.* **269,** 201–209 (1996).

11.   Reiter, C. D., Teng, R.-J. & Beckman, J. S. Superoxide Reacts with Nitric Oxide to

Nitrate Tyrosine at Physiological pH via Peroxynitrite. *J. Biol. Chem.* **275,** 32460–

32466 (2000).

12.   Breusing, N. & Grune, T. Biomarkers of protein oxidation from a chemical, biological

and medical point of view. *Experimental Gerontology* (2010).

doi:10.1016/j.exger.2010.04.004

13.   Sampson, J. B., Ye, Y., Rosen, H. & Beckman, J. S. Myeloperoxidase and

Horseradish Peroxidase Catalyze Tyrosine Nitration in Proteins from Nitrite and

Hydrogen Peroxide. *Arch. Biochem. Biophys.* **356,** 207–213 (1998).

14.   Bigelow, D. J. Nitrotyrosine-modified SERCA2: a cellular sensor of reactive nitrogen

species. *Pflügers Arch. - Eur. J. Physiol.* **457,** 701–710 (2009).

15.   Knyushko, T. V., Sharov, V. S., Williams, T. D., Schöneich, C. & Bigelow, D. J. 3-

Nitrotyrosine Modification of SERCA2a in the Aging Heart: A Distinct Signature of

the Cellular Redox Environment †. *Biochemistry* **44,** 13071–13081 (2005).

16.   Li, H., Gutterman, D. D., Rusch, N. J., Bubolz, A. & Liu, Y. Nitration and Functional

Loss of Voltage-Gated K+ Channels in Rat Coronary Microvessels Exposed to High

Glucose. *Diabetes* **53,** 2436–2442 (2004).

17.   Haddad, I. Y., Zhu, S., Ischiropoulos, H. & Matalon, S. Nitration of surfactant protein

A results in decreased ability to aggregate lipids. *Am. J. Physiol. Cell. Mol. Physiol.*

270, L281-8 (1996).

18.   Ferrer-Sueta, G. *et al.* Biochemistry of Peroxynitrite and Protein Tyrosine Nitration.

Chemical Reviews (2018). doi:10.1021/acs.chemrev.7b00568

19.   Akaike, T. *et al.* Pathogenesis of influenza virus-induced pneumonia: involvement of

both nitric oxide and oxygen radicals. *Proc. Natl. Acad. Sci.* **93,** 2448–2453 (1996).

20.   Akaike, T. *et al.* 8-Nitroguanosine formation in viral pneumonia and its implication for

pathogenesis. *Proc. Natl. Acad. Sci.* **100,** 685–690 (2003).

21.   Evans, T. J. *et al.* Cytokine-treated human neutrophils contain inducible nitric oxide

synthase that produces nitration of ingested bacteria. *Proc. Natl. Acad. Sci.* **93,**

9553–9558 (1996).

22.   Hayashi, T. *et al.* Modulating role of estradiol on arginase II expression in

hyperlipidemic rabbits as an atheroprotective mechanism. *Proc. Natl. Acad. Sci.* **103,**

10485–10490 (2006).

23.   Szabo, C. *et al.* Protection against peroxynitrite-induced fibroblast injury and arthritis

development by inhibition of poly(ADP-ribose) synthase. *Proc. Natl. Acad. Sci.* **95,**

3867–3872 (1998).

24.   Graham, M. L., Janecek, J. L., Kittredge, J. A., Hering, B. J. & Schuurman, H. J. The

streptozotocin-induced diabetic nude mouse model: Differences between animals

from different sources. *Comp. Med.* (2011). doi:10.1152/ajprenal.00292.2015

25.   Berthiaume, J. M. & Wallace, K. B. Adriamycin-induced oxidative mitochondrial

cardiotoxicity. in *Cell Biology and Toxicology* (2007). doi:10.1007/s10565-006-0140-

y

26.   Peluffo, G. & Radi, R. Biochemistry of protein tyrosine nitration in cardiovascular

pathology. *Cardiovasc. Res.* **75,** 291–302 (2007).

27.   van der Vliet, A., Eiserich, J. P., Halliwell, B. & Cross, C. E. Formation of Reactive

Nitrogen Species during Peroxidase-catalyzed Oxidation of Nitrite. *J. Biol. Chem.*

272, 7617–7625 (1997).

28.   Nuriel, T., Hansler, A. & Gross, S. S. Protein nitrotryptophan: Formation, significance

and identification. *J. Proteomics* **74,** 2300–2312 (2011).

29.   Rebrin, I., Brégère, C., Kamzalov, S., Gallaher, T. K. & Sohal, R. S. Nitration of

Tryptophan 372 in Succinyl-CoA:3-Ketoacid CoA Transferase during Aging in Rat

Heart Mitochondria †. *Biochemistry* **46,** 10130–10144 (2007).

30.   Souza, J. M., Peluffo, G. & Radi, R. Protein tyrosine nitration—Functional alteration

or just a biomarker? *Free Radic. Biol. Med.* **45,** 357–366 (2008).

31.   Suzuki, T. *et al.* Nitration and nitrosation of N-acetyl-l-tryptophan and tryptophan

residues in proteins by various reactive nitrogen species. *Free Radic. Biol. Med.* **37,**

671–681 (2004).

32.   Yamakura, F. & Ikeda, K. Modification of tryptophan and tryptophan residues in

proteins by reactive nitrogen species. *Nitric Oxide* **14,** 152–161 (2006).

33.   Yamakura, F. *et al.* Nitrated and Oxidized Products of a Single Tryptophan Residue

in Human Cu,Zn-Superoxide Dismutase Treated with Either Peroxynitrite-Carbon

Dioxide or Myeloperoxidase-Hydrogen Peroxide-Nitrite. *J. Biochem.* **138,** 57–69

(2005).

34.   Yamakura, F., Matsumoto, T., Taka, H., Fujimura, T. & Murayama, K. 6-

Nitrotryptophan: a specific reaction product of tryptophan residue in human Cu, Zn-

SOD treated with peroxynitrite. *Adv. Exp. Med. Biol.* **527,** 745–9 (2003).

35.   Ikeda, K. *et al.* Detection of 6-nitrotryptophan in proteins by Western blot analysis

and its application for peroxynitrite-treated PC12 cells. *Nitric Oxide* **16,** 18–28

(2007).

36.   Alvarez, B. *et al.* Peroxynitrite-Dependent Tryptophan Nitration. *Chem. Res. Toxicol.*

9, 390–396 (1996).

37.   Herold, S. Nitrotyrosine, dityrosine, and nitrotryptophan formation from

metmyoglobin, hydrogen peroxide, and nitrite. *Free Radic. Biol. Med.* **36,** 565–579

(2004).

38.   Wang, Y. *et al.* The Nitrated Proteome in Heart Mitochondria of the db/db Mouse

Model: Characterization of Nitrated Tyrosine Residues in SCOT. *J. Proteome Res.*

9, 4254–4263 (2010).

39.   Marcondes, S., Turko, I. V & Murad, F. Nitration of succinyl-CoA:3-oxoacid CoA-

transferase in rats after endotoxin administration. *Proc. Natl. Acad. Sci.* **98,** 7146–7151 (2001).

40. Milsom, A. B., Fernandez, B. O., Garcia-Saura, M. F., Rodriguez, J. & Feelisch, M. Contributions of nitric oxide synthases, dietary nitrite/nitrate, and other sources to the formation of NO signaling products. *Antioxid. Redox Signal.* **17,** 422–32 (2012).

41. Wilks, H. M. *et al.* Designs for a broad substrate specificity keto acid dehydrogenase. *Biochemistry* **29,** 8587–8591 (1990).

42. Shoemark, D. K., Cliff, M. J., Sessions, R. B. & Clarke, A. R. Enzymatic properties of the lactate dehydrogenase enzyme from Plasmodium falciparum. *FEBS J.* **274,** 2738–2748 (2007).

43. Fan, J. *et al.* Tyrosine Phosphorylation of Lactate Dehydrogenase A Is Important for NADH/NAD+ Redox Homeostasis in Cancer Cells. *Mol. Cell. Biol.* (2011). doi:10.1128/MCB.06120-11

44. Liemburg-Apers, D. C., Willems, P. H. G. M., Koopman, W. J. H. & Grefte, S. Interactions between mitochondrial reactive oxygen species and cellular glucose metabolism. *Archives of Toxicology* (2015). doi:10.1007/s00204-015-1520-y

45. Zorov, D. B., Juhaszova, M. & Sollott, S. J. Mitochondrial Reactive Oxygen Species (ROS) and ROS-Induced ROS Release. *Physiol. Rev.* (2014). doi:10.1152/physrev.00026.2013

46. Nabeebaccus, A. A. *et al.* Nox4 reprograms cardiac substrate metabolism via protein O-GlcNAcylation to enhance stress adaptation. *JCI insight* (2017). doi:10.1172/jci.insight.96184

47. Zhang, M. *et al.* NADPH oxidase-4 mediates protection against chronic load-induced stress in mouse hearts by enhancing angiogenesis. *Proc. Natl. Acad. Sci.* (2010). doi:10.1073/pnas.1009700107

Acknowledgements

Work in A.J. laboratory was supported by BBSRC, BHF and MRC; in P.B. laboratory by

NKFIH (K123975); in H.H.H.W.S. laboratory by an ERC Advanced Investigator Grant

(294683 - RadMed), an ERC proof-of-concept grant (139-101052 - SAVEBRAIN) and the

EU Horizon 2020 programme, REPO-TRIAL. The authors would like to thank G. Jansen

for technical help, and Dr. P.M.H. Schiffers for carefully reading earlier stages of the

manuscript. We also thank the Advanced Computing Research Centre at Bristol and

BrisSynBio for access to the HPC machines BlueCrystal and BlueGem. We thank the

HECBiosim Consortium for access to the UK National HPC facility Archer.

Author information

These authors contributed equally: Cristian Nogales, Jeroen Frijhoff

Affiliations

Department of Pharmacology and Personalised Medicine, Maastricht University,

Maastricht, The Netherlands

Cristian Nogales, Ana I. Casas & Harald H.H.W. Schmidt.

Medtronic Bakken Research Center, Maastricht, The Netherlands

Jeroen Frijhoff

DiaSys Diagnostic Systems GmbH, Holzheim, Germany

Detlef Thoenges

School of Biochemistry, University of Bristol, Bristol, UK

Richard B. Sessions

Leibiniz-Institut für Analytische Wissenschaften – ISAS, Dortmund, Germany

Albert Sickmann

Rudolf-Buchheim-Institue for Pharmacology, Justus-Liebig-University, Giessen, Germany

Tatiana Nedvetskaya

School of Chemistry, Autonomous University of Querétaro. Querétaro, México

Cesar Ibarra-Alvarado

German Diabetes Research Institute, Düsseldorf, Germany

Vera Schmitz & Peter Rösen

Heinrich-Heine-University, Department Cardiovascular Physiology, Düsseldorf, Germany

Axel Goedecke

University of Nicosia Medical School, Nicosia, Cyprus

Aleksandar Jovanović

Merck Exploratory Science Center, Cambridge, Massachusetts

David H. Perlman

Cologne Cardiovascular Research Center CCRC, Cologne, Germany

Anna Klinke & Stephan Baldus

Department of Medical Chemistry, Faculty of Medicine, University of Debrecen, Debrecen,

Hungary and MTA-DE Lendület Laboratory of Cellular Metabolism, Debrecen, Hungary

Péter Bai

Department of Oncology, Microbiology and Immunology; University of Fribourg, Fribourg,

Switzerland

Csaba Szabo

University of Southampton Medical School, Southampton, UK

Martin Feelisch

Contributions

H.H.H.W.S. designed research; J.F., D.T., T.N., C.I.A., A.S., P.B., C.S., V.S., P.R., A.J.,

A.G., D.H.P., R.B.S. and M.F. performed research; A.S., A.G., A.K., S.B., D.H.P., M.F. and

H.H.H.W.S. contributed new reagents/analytic tools; J.F., D.T., D.H.P., and H.H.H.W.S.

analyzed data; and C.N., J.F., C.S., D.H.P., M.F., A.I.C. and H.H.H.W.S. wrote the paper.

Competing interests

The authors declare no conflict of interests.

Corresponding authors

Correspondence to Harald H.H.W. Schmidt or Cristian Nogales.

1 **Data availability**

2 Data available from the corresponding authors upon reasonable request.

Figures and figure legends

Figure 1. Experimental outline for an anti-NO$_2$Tyr antibody-based biomarker

detection approach. Central panel: Antibody panel screening, an anti-NO$_2$Tyr antibody

panel was tested in porcine and rat tissues under physiological conditions. Different

antibodies might recognize different nitrated proteins (green and orange antibodies) or one

antibody might detect more than one nitrated protein (blue antibody). Right panel: A 38

kDa band was especially prominent and, subsequently, purified in large mammals and

identified via MALDI-TOF as LDH. NO$_2$-LDH activity was also assessed. We identified the

nitrated residue and simulated the effect of the nitration in the protein structure and

mechanism using human LDH. Left panel: Different rodent models, as well as knock-out

mouse models were used to elucidate the effects of NO$_2$-LDH in disease, diabetes and

myocardial stress, and the enzymatic sources of reactive nitrogen species (RNS) leading

to LDH nitration, respectively.

Figure 2. Screening of different native tissue homogenates identifies a 38 kDa band

in heart as major physiologically NO$_2$Tyr-immuno positive protein. Further

purification and MALDI-MS analysis identified the 38 kDa band as LDH. (a)

Immunopositive pig and rat tissues from Fig S1 were further analyzed by SDS-PAGE

under reducing conditions (dithiothreitol) and subsequent western blot. The band pattern

was different for all three polyclonal (pAb) or monoclonal (mAb) antibodies: pAb1, mAb1

and pAb2. The most intense NO$_2$Tyr immunopositive protein band at approximately 38

kDa (p38^{nt}) was detected using pAb1 in heart. (**b**) p38^{nt} was purified from porcine heart

homogenate by affinity (Blue Sepharose) followed by ion-exchange (Q-Sepharose)

chromatography. Purification steps were tracked via protein nitrotyrosine detection using

pAb1. *(c)* Purified cardiac p38^{nt} was cut out from the gel, digested with trypsin and

analyzed by mass spectrometry and peptide mass fingerprint analysis. Most peptide ions

detected were matched to lactate dehydrogenase: 11 ions matched to LDH (■) out of a

total of 20 detected, with 5 ions matching to the next best protein assignment, malate

dehydrogenase (MDH, ☐), identifying p38^{nt} as LDH with a $p<0.05$. (**d**) LDH nitration was

assessed by quantitative western blot analysis using pAb1 (n=9). (**e**) Vmax[LDH] remained

unchanged up to an equimolar peroxynitrite:LDH ratio, and then decreased at a ratio of 10

(n=8). Data represent means ±SEM.

Figure 3. Pathological LDH nitration levels in short-term (ST) and long-term (LT)

diabetic disease models and an oxidative stress myocardial dysfunction model. (**a**)

Heart homogenates of ST and LT streptozotocin-treated diabetic rats or doxorubicin-

treated mice (DoxoR) were analyzed by SDS-PAGE and immunoblot analysis for LDH

nitration using the pAb1 antibody. In all three animal models, the relative NO_2-LDH signal

intensity was increased. In diabetes, the long-term model yielded higher values than the

short-term model, which may indicate either higher nitrosative stress, or, more likely,

accumulation of LDH nitration (**b**). The highest increase was observed in the doxorubicin

model (**c**). Data are expressed as % of untreated control animals (open bars) and

represent means ± SEM of n=6-18 experiments.

Figure 4. *In vivo* genetic validation of reactive nitrogen species (RNS) sources. (**a**)

Events leading to oxidized and nitrated LDH. RNS (represented as $NO_2^-/H_2O_2/ONOO^-$) can

nitrate LDH, resulting in decreased V_{max}. (**b**) Cardiac tissue was obtained from endothelial

nitric oxide synthase (eNOS) and myeloperoxidase (MPO) knock-out (KO), and the

respective wild-type (WT), mice. Basal nitration of LDH (p38[nt]) was analyzed by

immunoblot, and immunosignals quantified densitometrically. Data are expressed as % of

control (i.e., of the respective WT animals) and represent means ± SEM of n=3-5 animals.

NO_2-LDH depended on both eNOS and MPO.

a NO₂-LDH MALDI-TOF MS-based residue identification

b Protein structure
WT human LDH

- β-K--β-L loop
- α-H C-terminal helix
- Active-site loop
- α-2G
- Pyruvate
- NADH

c WT and T4N backbone RMSF differences per residue

Binary complex (NADH)

Ternary complex (NADH + pyruvate)

d Association of the active site with α-H (m:LH)

Binary complex

Ternary complex

e Openness of the active site loop (m:LO)

Binary complex

Ternary complex

Figure 5. Identification of LDH Trp-324 as physiologically nitrated and oxidized. The

interaction of the active site loop with the C-terminal α-helix is essential and

restricted in NO$_2$Trp-LDH. (**a**) Rat heart homogenate was subjected to 2D-PAGE and

Coomasie staining; spots were subjected to in-gel digestion and MALDI-TOF MS analysis.

Leftft panel: Mass spectrum derived from a spot corresponding to rat LDH over the range

m/z 600 to 2700. Right panel: An expansion of the same spectrum over the range m/z

1170 to 1230, showing peptide ions assigned primarily to the peptide 319-328

(319SADTLWDIQK328), and its oxidized and nitrated derivatives. Major peptide ions are

labeled with their observed m/z values and corresponding LDH mature protein amino acid

intervals, as assigned by Mascot™, and additionally by manual comparison to predicted

values calculated from the LDH sequence and oxidative and nitrative modifications. The

ions correspond to the peptide 319-328 Trp (W)-nitrated species (319-328 + NO2) and its

typical high energy decomposition product (319-328 + NO$_2$-O), as well as ions

corresponding to the characteristic family of Trp oxidation forms, hydroxytryptophan (319-

328 + Ox), didehydro-hydroxytryptophan (319-328 + Ox-2H), dihydroxytryptophan (319-

328 + 2Ox), and kynurenin (319-328 + Ox-C). Other peptide ions display single (+ Ox) and

double (+ 2Ox) oxidation, or didehydrogenation (- 2H). (**b**) Distance metrics mapped onto

the crystal structure of WT (1I0Z). Color scheme: β-K--β-K loop, dark blue; α-helix, pink;

active-site loop, green; α-2G, light blue; pyruvate, grey; NADH, cyan. Residue Trp-324 is

shown in space filling. (**c**) Plot of the differences between T4N and WT backbone RMSF

per residue (T4N – WT) in the binary complex. Structures are highlighted using the same

color scheme as previously described plus the juxtaposed loop to the C-terminal α-helix

which is highlighted in light orange. (**d** – **e**) Histograms of the distribution of distances. Blue

WT; Red T4N. Metrics m:LH (107:CB-326:CB) showing the juxtaposition of the active-site

loop and α-C-terminal helix and m:LO (103:CA-243:CB) showing the open/closed state of

the active-site loop.

Supplementary Material: Tryptophan nitration physiologically regulates cardiac

lactate dehydrogenase active site loop mobility and activity

Cristian Nogales[1,*], Jeroen Frijhoff[1], Detlef Thoenges, Richard B. Sessions, Albert

Sickmann, Tatiana Nedvetskaya, Cesar Ibarra-Alvarado, Vera Schmitz, Peter Rösen, Axel

Goedecke, Aleksandar Jovanović, David H. Perlman, Anna Klinke, Stephan Baldus, Ana I.

Casas, Péter Bai, Csaba Szabo, Martin Feelisch & Harald H.H.W. Schmidt[*].

[1]Both first authors contributed equally to this work.

[*]To whom correspondence may be addressed. Email: h.schmidt@maastrichtuniversity.nl or

c.nogales@maastrichtuniversity.nl

Supplementary Methods

Chemicals

Immunoblotting reagents and column materials were obtained from Amersham Pharmacia

Biotech (Freiburg, Germany); membranes, from Bio-Rad Laboratories (Hercules, CA, USA).

Primary antibodies against nitrotyrosine were generous gifts from L.O. Uttenthal and J.S.

Beckman or purchased from HyCult Biotechnology (Uden, Netherlands), Calbiochem-

Novabiochem GmbH/Merck Millipore (Bad Soden/Darmstadt, Germany), Alexis/Axxora

GmbH (Grünberg/Lörrach, Germany) and Upstate Inc./Millipore (Lake Placid, NY,

USA/Darmstadt, Germany). As secondary antibodies, we used horseradish peroxidase

(HRP)-conjugated goat anti-rabbit immunoglobulins for polyclonal antibodies or anti-mouse

immunoglobulins for monoclonal primary antibodies from DAKO Diagnostika GmbH/Agilent

Technologies (Hamburg, Germany). For gel loading we used Rotiload buffer from Carl Roth

GmbH + Co. KG (Karlsruhe, Germany). The Oxyblot kit was supplied by Intergen

Company/Millipore (Purchase, NY, USA/Darmstadt, Germany). Nonfat dry milk was

obtained from Nestlé Inc. Carnation (Solon, Ohio, USA); Cell culture medium M199, from

PAA Laboratories GmbH (Martinsried/Cölbe, Germany); Nitrating reagent, H_2O_2/NO_2^-, from

Alexis/Axxora GmbH; Complete Protease Inhibitor Cocktail, from Roche Diagnostics GmbH

(Mannheim, Germany). Porcine heart LDH was purchased from MyBiosource. All other

chemicals were of the highest purity grade available and purchased from either Sigma-

Aldrich Chemie GmbH (Taufkirchen, Germany) or Merck KG (Darmstadt, Germany).

Animals

Adult male Wistar Unilever rats (270-320 g) were purchased from Harlan-Winkelmann

(Borchen, Germany). eNOS and MPO KO mice were generated and bred as described

previously (19). Rats and mice were housed under standard or SPF conditions, respectively,

at a temperature of 18-20°C and a day-night rhythm of 12 h, with access to water and food

ad libitum. Porcine hearts were obtained from a local abattoir.

Nitration of proteins by peroxynitrite

If not stated otherwise, nitration was performed by continuous mixing protein and nitrating

reagent (H_2O_2/NO_2^-) in a 1:5 molar ratio in phosphate buffer pH 7.3 (15 mM KH_2PO_4, 80 mM

Na_2HPO_4) for 5 min at room temperature. Due to the strong alkaline storage conditions of

H_2O_2/NO_2^-, the pH in the samples was immediately readjusted to neutral pH with HCl. No

further purification or dialysis steps were conducted. For longer storage, samples were

aliquoted and frozen at -20°C and used within 1 month or mixed with gel loading buffer and

boiled for 10 min at 95°C for daily use. In this case, no change in signal characteristics or

intensities was observed over time.

SDS-PAGE, western and dot-blot analysis

Protein samples were separated by sodium dodecylsulfate-polyacrylamide gel

electrophoresis (SDS-PAGE) under reducing conditions using 10% polyacrylamide gels.

Alternatively, pre-cast gels (NuPAGE Novex 10% bis-Tris Midi Gel 1,0 mm x 26 well,

Invitrogen) were used according to the manufacturer's protocol with MES running buffer

XCell SureLock Midi-Cell running tanks. For mass spectrometry proteins were stained with

colloidal Coomassie blue. Proteins were immobilized on nitrocellulose membranes using

standard dot-blot or semi-dry blotting techniques (20). Efficiency of protein immobilization

was evaluated by staining the membrane in 0.5 % (w/v) Ponceau S in 3 % (w/v)

trichloroacetic acid (TCA). Membranes were treated with 5 % (w/v) nonfat dry milk in Tris-

buffered saline (TBS, 20 mM Tris/HCl pH 7.5, 150 mM NaCl) containing 0.1 % (v/v) Tween

20 (TBS-T) for 1 hour to block nonspecific protein binding. Then the membrane was

incubated overnight at 4°C with the first antibody in 5% (w/v) nonfat dry milk in TBS-T and

afterwards washed 3×10 min with TBS-T. To test the specificity for NO_2Tyr, antibodies

were pre-incubated with 3 mM NO_2Tyr. Alternatively, in order to reduce NO_2 groups of

nitrated proteins to NH_2 groups, blots were first incubated with the reducing agent dithionite

(1 mM) in 100 mM sodium borate pH 10 for 5 min at room temperature, washed 3 X 10 min

with TBS-T and then incubated with anti-NO_2Tyr antibody (41). As second antibody, we

used horseradish peroxidase (HRP)-coupled antibodies against rabbit immunoglobulins (Ig)

for polyclonal first antibodies or against mouse Ig for monoclonal first antibodies. HRP-

labeled conjugates against the first antibodies were dissolved in 5% (w/v) nonfat dry milk in

TBS-T. After washing 3 X 10 min with TBS-T the immunosignals were developed using the

ECL detection reagent by Amersham Biosciences and blots were analyzed using a Kodak

chemiluminescence Imager (Scientific Imaging System, Eastman Kodak Company, New

Haven, USA). The resolution of the imager was 752 x 582 pixels and the linear area of signal

quantification was verified by repeated measurements using increasing protein amounts.

Light signals were directly quantified employing the Kodak 1D Image analysis software

(version 3.5.2).

Screening of tissues and determination of nitrated proteins

Several tissues from rat and pig were homogenized in 10 volumes (w/v) of lysis buffer,

containing 25 mM triethanolamine HCl (TEA) pH 7, 1 mM EDTA, 5 mM dithiothreitol, 50 mM

NaCl, 10 % (v/v) glycerol and Complete^{TM} Protease Inhibitor Cocktail (Roche, Mannheim,

Germany). For long-term storage, the homogenates were shock frozen in liquid nitrogen and

stored at –20 °C. For daily use, gel loading buffer was added to the samples and heated to

95°C for 10 min. Any remaining particles were pelleted by centrifugation in a desktop

centrifuge (10000 x g, 10 min, 4°C). The samples were first screened by use of all antibodies

listed in Tab. 1 by dot blot analysis. Tissues that yielded more than 4 hits by all antibodies

were subjected to further analysis by SDS-PAGE and western blot as described above.

Characterization and isolation of a 38 kDa protein in pig heart tissue

A column packed with 2 ml of Blue Sepharose resin (Amersham Bioscience) was washed

and pre-equilibrated with 10 mL lysis buffer. Pig heart supernatant (1 mL) was applied to the

column. The flow-through was subsequently applied to 1 mL Q-Sepharose column that was

also washed and equilibrated with lysis buffer. This column was then washed again with

lysis buffer (5 mL) and the bound proteins were eluted with 2 mL of lysis buffer containing 1

M NaCl. The partially purified 38-kDa protein was mixed with gel loading buffer, boiled at

95°C for 10 min and subjected to SDS-PAGE. It was stained with colloidal Coomassie blue

(22). The stained protein band at 38 kDa was cut out, destained and enzymatically digested.

Tryptic fragments were separated by high pressure liquid chromatography (HPLC).

Fractions containing tryptic peptides were subjected to matrix-assisted laser-

desorption/ionization mass spectrometry (MALDI-MS, Bruker Reflex III™, Bruker Daltonics,

Bremen, Germany). Peptide sequences were compared with proteins in the NCBIprot

database.

2D-PAGE analysis of rat heart tissue

2D-PAGE analysis was conducted using standard conditions (42). Briefly, 7 cm

ReadyStrips™ IPG strips (BioRad) of pI 3-10 were subjected to active rehydration overnight

with heart homogenate in Destreak™ IEF buffer, supplemented with 0.02% pI 3-10

ampholites (BioRad). Isoelectric focusing was performed in a Protean™ IEF cell (Bio-Rad)

using a ramped voltage up to 4000 V, until a total of approximately 10,000 Vh had elapsed.

Strips were subjected to reduction, alkylation, and equilibration for the second dimension,

using Equilibration Buffers I & II (BioRad) for 30 min each, then SDS-PAGE, using 10%

polyacrylamide NuPage™ Novex Bis-Tris gels (Invitrogen). Gels were subsequently

Coomassie stained (Gelcode™ Blue, Pierce).

In-gel digestion of rat heart samples

In-gel digestion was conducted as previously described (43). Briefly, the well-resolved

protein spot of interest was punch-excised using wide-orifice pipette tips. Gel pieces were

destained with 100 mM ammonium bicarbonate, pH 9/50% acetonitrile, and were washed

three times with wash solution 1 (100 mM ammonium bicarbonate, pH 9), wash solution 2

(100 mM ammonium bicarbonate, pH 9/50% acetonitrile), followed by wash solution 3 (100%

acetonitrile). Gel pieces were swelled in 50 mM ammonium bicarbonate, pH 9/10%

acetonitrile, containing approximately 10 ng Trypsin Gold (Promega) and incubated

overnight at 37°C. Peptides were extracted (each time for 20 min) once with 20 mM

ammonium bicarbonate, pH 9, twice with 1% trifluoroacetic acid/50% acetonitrile, followed

by once with 100% acetonitrile. The entire extraction process was performed in duplicate,

and all extraction supernatants were pooled and spun to dryness in a SpeedVac™ (Thermo-

Savant, Waltham, MA).

MALDI-TOF MS of rat heart samples

Peptides were desalted and prepared for MS using ZipTips™ (Millipore). Mass spectra

were obtained after purified peptides were co-crystallized with the matrix 2,5-

dihydroxybenzoic acid onto AnchorChip™ targets (Bruker Daltonics, Billerica, MA) using

the dried-droplet technique (28, 44) and a Reflex IV™ MALDI-TOF MS instrument (Bruker-

Daltonics) in positive ion, reflectron mode, over the range 400-8000. The laser intensity

was adjusted to be ~3% above the threshold value for peptide ion signals. Signals from

100 to 200 laser shots were summed for each mass spectrum. External and internal

calibration was achieved to within 50 ppm using the masses of known purified peptide

standards and trypsin autolysis peptides.**MS data analysis and peptide mass**

fingerprinting

Peak lists from MALDI-TOF mass spectra (derived using MoverZ™ software, Genomic

Solutions, Ann Arbor, MI) were submitted on-line to the Mascot™ search engine (Matrix

Science, London, UK) for peptide mass fingerprinting (PMF) analysis against the SwissProt

or NCBI non-redundant protein databases, using the following restrictions: a) Rattus sp., b)

trypsin digestion with up to 3 missed cleavages, c) +/- 50 ppm error, and d) cysteine

carbamidomethylation and methionine oxidation as fixed and variable modifications,

respectively. LDH was unambiguously identified by PMF, with a Mascot™ Score of 150

(corresponding to an E value of 6×10^{-12}, the likelihood of a false assignment) and a

sequence coverage of 65% from 35 matched peptide ions, given search parameters.

Additional peptide ion assignments (as well as validation of Mascot™ assignments) were

made by manual inspection of spectra using theoretical digest and post-translational

modification mass values for LDH.

LDH activity, oxidation and nitration by H_2O_2/NO_2^-

Commercial LDH was treated with different amounts of peroxynitrite as described above.

Aliquots of H_2O_2/NO_2^- were diluted with 2 M NaOH to reach the desired concentrations

indicated in **Fig. 2**. LDH activity and kinetic parameters of exogenously nitrated LDH were

measured using a fluorescence-based assay, using excitation/emission setting of 340/470

nm, respectively. The assay used uses the fluorescent properties of NADH which can be

excited at 340 nm and emission is measured at 440 nm. The oxidized form of NADH, NAD^+,

does not have these properties. Measurements were collected every 20 seconds during 5

minutes. The final concentrations in the reaction were LDH, 2 mU/mL, NADH 0.12 mM and

Sodium pyruvate 1.13 mM. Nitration of LDH was measured as the intensity of the NO_2Tyr

immunosignal in arbitrary units.

Molecular Dynamics Methods

System setup: Two simulation systems were setup based on the crystal structure of the

ternary complex of human-heart LDH (pdb 1I0Z). One being the wild type tetramer (WT) and

the other being the tetramer with a nitro group substituted at each Trp-324 residue (T4N).

The oxamate molecules were converted to pyruvate and acpype (45) used to generate

Amber GAFF parameters for the substrate and cofactor molecules. A combination of GAFF

parameters for nitrotryptophan and analogy was used to generate a residue library entry for

a nitrotryptophan in the Gromacs AmberSB99-ildn forcefield (46). Gromacs tools were used

to add hydrogen atoms to all residues consistent with pH 7 and to protonate the catalytic

histidine (His-193). The complexes were solvated in boxes of TIP3P water containing 0.15

M sodium chloride ions. Calculations were performed at NTP under period boundary

conditions and the PME method used for long range electrostatics. The pressure was 1 Barr

and the temperature 300 K for all molecular dynamics (MD) simulations. Both systems were

relaxed by 2000 steps of steepest descent energy minimisation followed by 0.2 ns MD while

restraining the protein atoms to their initial positions.

Another pair of simulation systems were prepared in the same fashion, with the pyruvate

removed. This gave a total of four simulation systems, referred to as WT-ternary, WT-binary,

T4N-ternary and T4N-binary.

Trajectory development and analysis: Three 200 ns simulations (initialized with different

velocities) were performed for each of the four simulation systems. These simulations were

performed on the University of Bristol HPC machines BlueCrystal and BlueGem. Structures

were extracted every 5 ns from each trajectory and energy minimized. This gave 120 starting

structures for each of the four systems for further parallel trajectory development. Each new

trajectory was initialized with a fresh set of random velocities and run for 28-31 ns on the

UK National HPC machine Archer, giving around 3.5 µs combined simulation data for each

system. All simulations were run and analyzed using the Gromacs 5.1.2 suite of software

(47) and visualizations performed with VMD 1.9.2 (48) and Chimera 1.11 (49).

Molecular Dynamics Analysis

The active site loop (residues 98-110) was found to be less mobile in T4N than WT in the

binary complex (LDH+NADH), while the C-terminal helix (α-H) and the juxtaposed loop

(residues 277-287) were more mobile in T4N. The lower mobility of the active site loop is

also detectable in the ternary complex (LDH+NADH+pyruvate) albeit greatly attenuated,

while the C-terminal helix mobility is similar to the WT. The other main features in these plots

are the differences in mobility of the 205-225 loop. However, as this is one of the most

flexible parts of the protein and is not located near the active site we do not consider these

differences to impact the activity of the two proteins. By contrast, the C-terminal side of the

active site loop juxtaposes the C-terminal end of the final helix in the protein, α-H (residues

309-332). A crucial contact between these two structural elements is made by Leu-107 and

Ile-326. Hence, we define one metric reporting the association of the active site loop with α-

H via the distance between CB atoms of these residues, as m:LH. Likewise, two residues

spanning the mouth of the active site Gly-103 and Lys-243 were chosen to monitor

openness of the active site loop, or m:LO. Distance histograms show the distribution of these

distances over each of the four 3.5 µs combined trajectories. They indicate an increased

mobility in the binary complexes compared with the ternary, expressed as a broadening of

the histograms.

The distance between the active site loop and α-H (m:LH) is slightly shorter on average for

WT than T4N in the ternary complex and this difference is amplified in the binary complex

where about half of the WT loops interact more tightly with α-H than T4N. Comparison with

the second metric monitoring active site loop closure (m:LO) in the binary complex shows

that the T4N loop spends more time at a distance comparable with the ternary complex than

the WT and also is an "over-closed" state. The results of the ternary simulations are again

more similar, but a tendency towards and "over-closed" state persists.

Synthesis and Characterization of Nitrotryptophan

Nitrotryptophan (NO_2Trp) was synthesized according to Moriya, et al. (15). 20 g L-Trp and

0.25 g urea were suspended in 350 mL glacial acetic acid. Under vigorous stirring at 4°C,

10 mL 24 M nitric acid in 30 mL glacial acetic acid were added. After the suspension changed

to a yellow clean solution, it was stirred at 4°C until it changed again to a suspension. Then

an additional 10 mL fuming nitric acid in 30 mL glacial acetic acid were added drop by drop

at 15°C. After stirring overnight at room temperature, the resulting yellow precipitate was

collected by filtration and washed with glacial acetic acid. The crude product was

recrystallized in 100 mL 2 M nitric acid at temperatures below 50°C and converted to the

free amino acid in 100 mL hot water by the addition of sodium carbonate until the pH was

neutral. As analyzed by its infrared (IR) and nuclear magnetic resonance (NMR) spectra, no

4-NO_2Trp was detectable. Because 7-NO_2Trp cannot be synthesized by the action of 24 M

nitric acid on the tryptophan ring (16), the products of the synthesis with respect to UV spectrum are a mixture of 5/6-NO_2Trp. NO_2Trp crystallized as yellow needles with a melting point at 265°C (decomp.). The compound was characterized according to: UV (in 1 N NaOH) 330 nm, 380 nm; IR (KBr) ν(-NH_3^+) 3190 cm^{-1}, ν(C=C) and amide I 1600-1650 cm^{-1}, amide II and δ(-NH_3^+) 1550-1600 cm^{-1}, ν_{as}(-NO_2) 1495 cm^{-1}, ν_s(-NO_2) 1330 cm^{-1}, predominantly 5/6-NO_2 substitution at 818 cm^{-1} instead of 4-NO_2 substitution at 780 cm^{-1}; NMR (D_2O-NaOH, δ 4.7 HDO) δ 2.9-3.1 (2H, m), 3.4-3.6 (1H, m), 7.35 (1H,s), 7.5 (1H, d, J= 9 Hz), 7.7 (1H, dd, J= 2 and 9 Hz), 7.95 (1H, d, J= 2 Hz).

Animal models of disease

To examine LDH nitration under pathophysiological conditions, two *in vivo* models involving nitrative stress were assessed, including a short- and longer-term streptozotocin diabetes model in the rat, and a murine model of doxorubicin-induced cardiotoxicity.

Streptozotocin model of diabetes mellitus

At an age of 10 weeks, rats (n=12) received a single intraperitoneal injection of 60 mg/kg BW of streptozotocin (STZ) dissolved in citrate buffer 50mM pH 4,5. Age matched control rats (n=12) received vehicle injections with citrate buffer. Hyperglycemia was defined as a positive test of glucose in urine (Haemo-Glukotest, Boehringer Mannheim, Germany) and a non-fasting blood glucose level higher than 11 mmol/L (hexokinase method) one week after STZ application. Rats were sacrificed 3 (n=12) and 16 weeks (n=12) after diabetes induction by CO_2-inhalation. After thoracotomy, hearts were excised and freed of connective tissue while immersed in modified Krebs-Hepes buffer pH 7.35 containing 99 mM NaCl; 4.7 mM KCl; 1.9 mM $CaCl_2$; 1.2 mM $MgSO_4$; 25 mM $NaHCO_3$; 1 mM K_2HPO_4; 20 mM Na-Hepes; 11.1 mM D-glucose. Hearts were shock frozen in liquid nitrogen, pulverized, and homogenized in lysis buffer. For SDS-PAGE and western blot, gel loading buffer was added and protein concentrations were determined after TCA precipitation by the method of Lowry.

Doxorubicin cardiotoxicity model

Doxorubicin (DOX, Adriamycin) is a broad-spectrum anthracycline antibiotic that is used to treat a variety of cancers. However, the clinical use of DOX is limited because of its serious cardiotoxicity, possibly involving increased oxidative stress, alteration of cardiac energetic status and direct effects on DNA. Hearts from DOX-treated mice were obtained as described elsewhere (50), and homogenized and shock-frozen as described above.

Endogenous LDH nitration in eNOS-KO and MPO-KO mice

Hearts from eNOS-KO and MPO-KO mice were excised, washed in perfusion buffer, blotted dry on filter paper, weighed, and homogenized immediately in ice-cold 50 mM PBS, pH= 7.4, containing 2.5 mM EDTA, 1 mM DTT, 1% NP-40, phosphatase inhibitors (10 mM sodium fluoride, 10 mM sodium pyrophosphate, 2 mM sodium orthovanadate, and 50 mM β-glycerophosphate), and Complete™ protease inhibitor cocktail (Roche Diagnostics, Indianapolis, IN). Homogenates were incubated on ice for 30 min to facilitate solubilization, after which they were clarified by centrifugation at 12000 x g for 15 min at 4°C, aliquoted and stored at -80°C. Prior to use, samples were thawed on ice and assayed for protein concentration using the BCA method.

1 **Supplementary Tables**

Abbreviation	Specificity	Antigen	Species	Source	Reference
pAb1	anti-NO$_2$Tyr	ONOO-KLH	rabbit	L.O. Uttenthal	(33)
pAb2				J.S. Beckman	(34)
mAb1			mouse		(34)
Identical to mAb1:					
mAb2	anti-NO$_2$Tyr	ONOO-KLH	mouse	Alexis	(34)
mAb3				Upstate	(34)
Rejected:					
mAb4	anti-NO$_2$Tyr	ONOO-KLH	mouse	Calbiochem	(51)
mAb5				HyCult biotechnology	(37)
pAb3	anti-NOTrp	NOTrp-KLH	rabbit	Calbiochem	None
mAb6	anti-NOCys	NOCys	mouse	Biolabs	None

Supplementary Table 1. Commercial and non-commercial antibodies detecting

reactive nitrogen species-induced protein modifications. Poly (pAb)- and monoclonal

(mAb) antibodies from the indicated species were obtained from either commercial sources

or kindly provided by the listed researchers. All antibodies, except pAb3 and mAb6, were

raised against keyhole limpet hemocyanin (KLH) treated with 'nitration reagent', i.e.

H_2O_2/NO_2^-. For pAb3, the hapten was nitrosocysteine (NOCys); for mAb6, nitrosotryptophan

(NOTrp), and in both cases glutaraldehyde-coupled to KLH. Antibodies mAb1, mAb2 and

mAb3 behaved identically and referred to the same reference (34) , and were thus

considered identical. Antibodies mAb4, mAb5 and pAb3 were later omitted from the study

because of low titer and/or low affinity, which allowed to detect only highly abundant antigens

such as H_2O_2/NO_2^--treated bovine serum albumin (BSA), but not lower abundance

endogenously nitrated proteins in native tissues. Antibody mAb6 was also excluded from

further use because it reacted similarly with untreated and H_2O_2/NO_2^--treated BSA.

1   Consequently, we pursued the further use of the first three antibodies, pAb1, pAb2 and

2   mAb1, for the present study (see **Fig. 2**).

Antigen \ Antibody	pAb1 1:3000	pAb2 1:3000	mAb1 1:3000
NO$_2$Tyr-Block	+	+	+
NO$_2$BSA (Control)			
Pul. Artery (pig)			
Heart (pig)			
Lung (rat)			
Kidney (rat)			
Cerebellum (rat)			
Spleen (rat)			
Cortex (rat)			

Supplementary Figure 1. Anti-NO2-Tyr antibodies greatly vary in sensitivity and

specificity in native tissues. Dot blot screen of tissue homogenates of porcine pulmonary

artery and heart, and rat heart, lung, kidney, cerebellum, spleen and cortex. For each dot,

50 µg or, in the case of pulmonary artery, 100 µg of protein were blotted onto a nitrocellulose

membrane. NO$_2$Tyr-immuno-positive proteins were then detected using antibodies pAb1,

pAb2 or mAb1. NO$_2$BSA was used as a positive control; pre-incubation with 3 mM free

NO$_2$Tyr, as a negative control (+).

Supplementary Figure 2. Nitration and oxidation of LDH *in vitro* inversely correlates

with LDH activity. As LDH (0.1 nmol) was treated with increasing amounts of peroxynitrite,

total LDH activity ($\blacksquare$, left axis) was determined by the NADH-coupled photometric assay of

Warburg, as described, and all LDH samples were tested under the same conditions for the

occurrence of nitration ($\bullet$) using quantitative Western analysis with the NO_2Tyr antibody

pAb1 and oxidation ($\triangle$) using the oxyblot detection kit (both, right axes).

2 **Supplementary Figure 3.** Plots of the backbone RMSF per residue averaged over all

3 subunits of all 120 simulations. (**a**) WT-binary. (**b**) WT-ternary. (**c**) T4N-binary. (**d**) T4N-

4 ternary.