1 **Dynamism and context dependency in the diversification of the megadiverse plant genus**
2 ***Solanum* L. (Solanaceae).**
3

4 Susy Echeverría-Londoño[1,2,3*], Tiina Särkinen[4], Isabel S Fenton[1], Sandra Knapp[1], and Andy
5 Purvis[1,2]
6
7 [1]Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK
8 [2]Department of Life Sciences, Imperial College London, Ascot SL5 7QN, UK
9 [3]Department of Biology, Kenyon College, Gambier, Ohio 43022, USA
10 [4]Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh EH3 5LR, UK
11
12 *Corresponding author: Susy Echeverría-Londoño; Phone, +1(740)5046838; Email,
13 susyelo@gmail.com

14
15 Total word count main body: 5370; Introduction (721), Methods (1923), Results (959),
16 Discussion (1730), Acknowledgements (37).
17
18 Number of Figures: 3 (all in colour).
19 Number of Tables: 0
20 Supporting Information: Methods S1-S2, Fig. S1-S13, Table S1-S3.
21
22
23 **Summary**
24

• Explosive radiations have been considered one of the most intriguing diversification
patterns across the Tree of Life, but the subsequent change, movement and extinction of
the constituent species makes radiations hard to discern or understand as geological
time passes.
• We synthesised phylogenetic and distributional data for an ongoing radiation — the
mega-diverse plant genus *Solanum* L. — to show how dispersal events and past climatic
changes have interacted to shape diversification.
• We found that despite the vast diversity of *Solanum* lineages in the Neotropics, lineages
in the Old World are diversifying more rapidly. This recent explosive diversification
coincides with a long-distance dispersal event from the Neotropics, at the time when,

and to places where, major climatic changes took place. Two different groups of

Solanum have migrated and established in Australia, but only the arid-adapted lineages

experienced significant increases in their diversification, which is consistent with

adaptation to the continent's long-term climatic trend and the diversification of other arid-

adapted groups.

•   Our findings provide a clear example of how successful colonisation of new areas and

niches can – but do not always – drive explosive radiations.

Key words: Biogeography, diversification rates, plant macroevolution, Solanaceae, *Solanum*.

Introduction

The uneven distribution of taxonomic diversity among the branches of the tree of life and

geographic regions is one of the most intriguing puzzles in biology. The increasing availability of

global field study data and large phylogenies continues to reveal how some groups are highly

diverse compared with their depauperate sister groups. Even though we expect the stochastic

nature of the diversification process to produce asymmetric patterns of diversity which do not

necessarily require deterministic explanations (Raup *et al.*, 1973; Slowinski & Guyer, 1993;

Purvis, 1996), identifying significant patterns of unequal diversity (i.e., those that depart from the

null expectation) throughout the tree of life has the potential to reveal important mechanism

about the origin and maintenance of biodiversity.

One of the main hypotheses in macroevolution is that unequal diversity can arise from

significant differences in net diversification rates (i.e., speciation – extinction) among lineages

and through time and space (Stanley *et al.*, 1981; Alfaro *et al.*, 2009; Wiens, 2011, 2015). This

asymmetry in evolutionary rates is often interpreted as resulting from the presence of a

spectrum of ecological opportunities, such as the opening of niche space by the development of

new key traits, extinction of antagonists, and/or colonisation of new open areas (Simpson, 1955;

Schluter, 2000; Moore & Donoghue, 2007; Losos, 2010; Yoder *et al.*, 2010; Rabosky *et al.*,

2013). The first step to uncover the relative influence of these factors in diversification is to

decompose the macroevolutionary dynamics of a clade into its essential components ---

speciation and extinction rates. Phylogenetic Comparative Methods (PCM) provide a series of

approaches to quantifying these rates using reconstructed phylogenies along with various

stochastic models (see Stadler, 2013 and Morlon, 2014 for a detailed review of diversification

approaches). However, the results of such approaches are often limited and challenged by the

difficulty of accounting for extinct species usually not included in molecular phylogenies; these

factors are especially problematic in older clades (Liow *et al.*, 2010; Quental & Marshall, 2010;

Rabosky, 2010). Furthermore, poor species-level sampling and branch support in the underlying

phylogenies used for testing these hypotheses usually limits the power of tests of diversification

dynamics. Therefore, a focus on recently emerged clades that are species-rich, widespread and

are underpinned by well-sampled and robust phylogenies will provide us with a more accurate

picture about the macroevolutionary dynamics shaping biodiversity.

In this study, we investigate the tempo and mode of species diversification of the mega-diverse

plant genus *Solanum* L. (Solanaceae). This genus is one of the most diverse and economically

important plant groups with around 1,500 species distributed worldwide (i.e., almost the half of

the diversity of the Solanaceae family). *Solanum* contains species from all major tropical and

subtropical biomes and shows high heterogeneity of richness among its subclades. This plant

genus has been traditionally divided into two major groups, the spiny (prickly species) and non-

spiny solanums. With robust monophyletic support, the spiny solanums (known as the

Leptostemonum clade, or subgenus *Leptostemonum Bitter*) include ca. 420 species and can be

recognised by the presence of stellate trichomes, prickles, and long tapering anthers

(Vorontsova *et al.*, 2013). Spiny solanums are most diverse in the Neotropics (ca. 150 species),

but also have high diversity in Australia (ca. 130) and Africa (ca. 79). This high diversity of spiny

solanums out of the Americas contrasts with the distribution of the diversity of the non-

monophyletic group — the non-spiny solanums — which has more than 90% of its ca. 670

species distributed in the Neotropics.

As one of the top ten species-rich genera of angiosperms (Frodin, 2004)*, Solanum* is suggested

to have high diversification rates, especially in the Neotropics which contains most of its

diversity and endemism (Olmstead & Palmer, 1997; Särkinen *et al.*, 2013; Knapp & Vorontsova,

2016). However, where radiating lineages occur and how much they contribute to current

diversity and geographic differences in richness among subclades of *Solanum* has not yet been

explored. Here, we assembled the first set of complete time-calibrated and species-level

phylogenies of extant *Solanum* species (1169 species), to reconstruct diversification rates

across lineages and analyse them in a geographical context. We test the origins of the high

heterogeneity of species richness observed among subclades of *Solanum*, investigating the

relative importance of clade-specific, tree-wide and geographic variation in evolutionary rates,

as well as how these patterns are associated with historical biogeographic movements and/or

environmental changes.

Materials and Methods

Phylogenetic relationships and divergence times among the main clades of *Solanum* were

obtained from the Särkinen *et al.*, 2013 Solanaceae phylogeny. This tree was built using two

nuclear and six plastid loci from 454 *Solanum* species (ca. 34% of the extant species). This

phylogeny identified a total of 37 clades and subclades within the genus, with a vast

heterogeneity of species richness within these.

Correcting for non-random taxon sampling

The incomplete nature of the Särkinen *et al.*, 2013 phylogeny could be problematic because

non-random incomplete sampling in phylogenetic studies could significantly affect the estimate

of macroevolutionary rates (Höhna *et al.*, 2011). Despite the claims that current speciation-

extinction models account for incomplete species sampling (e.g., MEDUSA, Alfaro *et al.*, 2009;

BAMM, Rabosky, 2014; TreePar, Stadler, 2011; TESS, Höhna *et al.*, 2016b; RevBayes, Höhna

et al., 2016a), these models often assume random, even sampling of the species across the

phylogeny (i.e., each terminal in the phylogeny is assumed to have the same probability of

being sampled). However, this assumption is very often violated in empirical examples due to

geographic, temporal and taxonomic sampling biases. To account for non-random incomplete

taxon sampling in the diversification analysis, we used the stochastic polytomy resolver PASTIS

(Thomas *et al.*, 2013), to place missing species of *Solanum* in the phylogeny using taxonomic

constraints following a birth-death model. This method is a conservative approach for increasing

the sampling in phylogenies since it infers the timing of missing splits under a constant rates

birth-death model. Therefore, any inference of rate-heterogeneity in the final analysis will have

stronger support since this approach is expected to bias towards the detection of constant-rate

models of diversification (Thomas *et al.*, 2013). Every extant and accepted *Solanum* species

was assigned to one of the clades (listed in Supporting Information Table S1) using the

Särkinen *et al.*, 2013 phylogeny as a backbone. Complete trees were then generated for each

clade of *Solanum* using a combination of species with genetic data and taxonomic constraints.

For each clade, PASTIS creates an output file in a nexus format, which contains the full set of

tree constraints ready to be executed in MrBayes. Posterior distributions of phylogenies for each

clade were then inferred in MrBayes 3.2.3 (Ronquist & Huelsenbeck, 2003) using a relaxed

clock model (independent branch rates – igr prior), with the default (exponential) prior on the

distribution of branching rates. A distribution of 100 dated trees was produced containing nearly

all described species of *Solanum*, 1169 species in total (excluding those with uncertain

taxonomic position), after sampling and grafting the clades distributions. For further information

see Supporting Information Methods S1.

Diversification analysis

Several approaches have been developed to detect significant shifts in diversification across the

branches of a phylogeny. In this analysis, we mainly used the BAMM approach (Rabosky,

2014), a Bayesian framework for reconstructing evolutionary dynamics from phylogenetic trees,

which aims to provide an improvement on methods that identify heterogeneity in rates only

across specific subclades, such as MEDUSA (Alfaro *et al.*, 2009). However, other approaches

such as TESS (Höhna *et al.*, 2016b) and RevBayes (Höhna *et al.*, 2016a), were also

implemented to assess the robustness of the results obtained in BAMM.

The underlying branching process in BAMM includes the effect of time-dependence on

diversification (i.e., the age of a lineage can affect its diversification rate). It also implements a

diversity-dependent diversification model where the number of lineages in a clade may affect its

diversification rates. BAMM uses a reversible-jump Markov Chain Monte Carlo approach to

explore a larger space of parameters and candidate models of diversification. Since it follows a

Bayesian statistical framework (rather than the maximum likelihood framework used in

MEDUSA), BAMM implicitly accounts for the uncertainty in parameter estimates by providing a

distribution of marginal posterior probabilities instead of point estimates. To implement this

approach in the *Solanum* phylogeny, we used the distribution of 100 complete species trees

(generated using PASTIS, see previous section), to consider the phylogenetic uncertainty in the

estimates of diversification rates. We set the priors of speciation, extinction and the expected

number of diversification shifts (g=1) using the R package BAMMtools v 2.0.5 (Rabosky *et al.*,

2014), which identifies the priors of the diversification parameters based on the distribution of

divergence times of the phylogeny. For each of the 100 trees, an MCMC analysis was

performed with four separate runs of 20 million generations. All the analyses were run using the

C ++ BAMM command line program v 2.5.0 on the Imperial College London's High-Performance

computing cluster (http://www.imperial.ac.uk/admin-services/ict/self-service/research-

support/hpc/). We then checked for convergence of the MCMC samples making sure the effect

sample size was at least 200 for both the number of evolutionary shifts and likelihood using the

CODA R package V 0.16-1 (Plummer *et al.*, 2006). The first 25% of the samples were discarded

as burn-in.

For each of the 100 trees run in BAMM, a distribution of 1000 samples of the posterior

probabilities of diversification were created. Each sample from the posterior includes either a

single event (i.e., the diversification is described by a single time-varying process — no shifts in

diversification) or a mixture of one or more shifts and associated parameters. For each set of

posterior probabilities, we extracted the list of nodes associated with "core" rate shifts (i.e., rate

shifts with a marginal probability significantly higher than the probability expected from the prior

alone) and calculated the frequency with which these nodes are associated with significant rate

shifts across the 100 trees run in BAMM (i.e., across a distribution of 100 x 1000 trees). Finally,

we computed the mean of diversification rates for each of the species present in the Särkinen *et*

al., 2013 phylogeny across the pooled distribution of posterior samples produced in the BAMM

analysis.

Studies such as Moore *et al.*, 2016 have raised concerns about several methodological issues

in BAMM including a potentially incorrect likelihood function, the strong influence of priors on

posterior estimates, and some theoretical errors (e.g., the incorrect use of the Poisson

distribution as the error distribution of the prior number of shifts of diversification). To assess the

effects of these potential limitations on the results, we performed several sensitivity analyses

focusing on the two most important issues stated by Moore et al., 2016, (1) the prior sensitivity

of posterior distribution on the number of rate shifts, and (2) the reliability of the diversification

rate estimates. We performed these analyses using alternative approaches such as TESS

(Höhna *et al.*, 2016b) and RevBayes (Höhna *et al.*, 2016a). Further details about the

methodology and results of these analyses can be found in Supporting Information Methods S2.

Geographical patterns of diversification

Using 64,826 unique records from 1,096 taxa (see Supporting Information Methods S2 for

further details), we mapped the mean assemblage diversification rates of *Solanum* on a 1 x 1

degree map to determine which regions have accumulated or are currently accumulating a

higher or lower number of *Solanum* lineages across the globe. The mean assemblage

diversification rate was calculated as the geometric mean of all species-specific rates present in
a grid cell. We also computed a weighted version of this, dividing the mean species-specific
diversification rates by the inverse of their range size --- log of the area (m^2) occupied by each
species, to correct for the diversification rates in areas with a high occurrence of widespread
species (i.e., large ranging species have less weight in the overall diversification rates of a grid
cell).
211
212 Finally, we reconstructed lineages-through-time plots for the three-principal diversity centres for
213 *Solanum* (Neotropics, Australia and Africa) using the R package "paleotree" v 2.3 (Bapst, 2012)
214 to visualise the geographical differences in lineage accumulation dynamics among continents.
215

Ancestral range reconstruction

We investigated the historical biogeography of *Solanum* using the R package "BioGeoBEARS"
(Matzke, 2012, 2016) — BioGeography with Bayesian (and likelihood) Evolutionary Analysis in
R Scripts. This approach provides a statistical framework to compare traditional models in
biogeography such as DIVA (Dispersal-Vicariance Analysis, Ronquist, 1997), DEC (Dispersal-
Extinction-Cladogenesis, Ree *et al.*, 2005), and BayArea (Landis *et al.*, 2013). Each of these
models assume different processes to reconstruct the ancestral range of lineages. In all the
models, species ranges are allowed to change along the branches by anagenetic evolution
through two main events: dispersion (range expansion) and extinction (range contraction). The
events allowed in cladogenesis vary depending on the fitted model. For example, sympatric
speciation subset, or peripatric speciation (i.e., the ancestral range is completely inherited by
one of the daughters whereas the other inherits only a subset of the range) is only allowed in
the DEC model. However, unlike the DIVA and the BayArea models, the DEC model only allows
one daughter to inherent widespread distributions during a vicariant event, since the widespread
distribution of both daughters would assume additional events such as post-speciation dispersal
(Ree *et al.*, 2005). In addition to these classic events of biogeography, BioGeoBEARS includes
founder-event speciation (+j) which considers the influence of speciation through long-distance
dispersal, such as is common in Island-like models.
235
To avoid the effects of birth–death polytomy resolvers on the natural patterns of trait
phylogenetic structure (following Rabosky, 2015), we reconstructed the ancestral ranges of
Solanum using the time-calibrated clade credibility (MCC) from Särkinen *et al.*, 2013. Several

species were pruned from the tree as follows: (1) species which are considered widely

cultivated or with ambiguous native distribution, (2) species considered as synonyms, and

therefore duplicated in the tree, and (3) species with a low support of a direct ancestor-

descendant relationships leading to negative branches in the MCC. The final pruned phylogeny

used for subsequent analyses contained 386 species of *Solanum* with the addition of *Jaltomata*

andersonii as the outgroup.

Using the extracted distribution records, we created an occurrence matrix of *Solanum* species

split into six biogeographic areas—Africa, Australia, Indo-Pacific, Neotropics, Nearctic and

Palearctic based on the floral kingdoms defined by Cox (2001). The distribution of widespread

species was double checked and corrected for potentially recent cultivated and/or naturalised

species using the native descriptions available from http: //solanaceaesource.org/ or specific

taxonomic monographs. The maximum number of areas in the ancestral range reconstruction

was set to three to reduce the complexity and computational time in the analysis.

Three biogeographic models were fitted to the *Solanum* phylogeny and the associated

geographic distributions using BioGeoBEARS — DIVALIKE, DEC and BAYAREALIKE models.

The influence of founder-event speciation event (+j) was also included into each model,

resulting in a total of six models. An additional set of six biogeographic models were fitted using

a dispersal matrix multiplier to weight the dispersal probability of adjacent areas as 1, 0.5 and

0.001 for easy, medium, and hard dispersal. The model that best describes the empirical data

(i.e., optimal fixed model structure) was then chosen using a stepwise selection from the

candidate models ranking under the Akaike Information Criterion (AIC, Burnham & Anderson,

2002).

Once the best-fitting model was identified, we estimated the overall probabilities of the

anagenetic and cladogenetic events conditional on the model, the phylogeny, and the

geographic distributions from 100 Biogeographic Stochastic Maps (BSM) to account for the

uncertainty in the state transitions and ancestral range reconstructions (Matzke, 2016). These

stochastic maps are similar to simulations of trait change along phylogenies (Huelsenbeck *et al.*,

2003) using transition rate models (Pagel, 1999). Given the observed range data, the phylogeny

and the best fitting model of biogeographic events, the BSM simulates possible histories

constrained by the observed ancestral ranges. The ancestral state probabilities obtained under

the best fitting model are equivalent to the average of all the probabilities of the simulated

histories from the biogeographic stochastic maps (Matzke, 2016).

Results

We found significant heterogeneity in diversification rates along the branches of the *Solanum*

phylogeny. This variation in evolutionary rates has contributed to the pronounced disparities in

species richness among groups; the Old World spiny clade and the Petota subclade are

supported as the most rapid radiations (Fig. 1). The Old World clade has diversified nearly as

twice as rapidly as any other group, with approximately 0.68 lineages per million years (lineages

Myr^{-1}) as compared to the global mean of 0.25 lineages Myr^{-1} (Fig. 1). The node supporting the

crown group of the Old World clade was the node with the highest frequency of shifts found

across the 100 runs. Other groups such as the Petota subclade (from the Potato clade, Petota +

Tomato) and the node supporting the Leptostemonum group (i.e., the spiny solanums) also

show some signal of shifts in diversification but with a weaker support compared with the Old

World clade (Fig. 1, see also Supporting Information Fig. S2, S3).

The number of diversification shifts found in the BAMM analysis is largely robust to the defined

prior distribution. Supporting Information Fig. S7-S10 show that across the 100 *Solanum*

phylogenies run in BAMM, the distribution of estimated shifts (posterior samples) differs

significantly from the distribution of the expected number of shifts (prior). Although the prior

distribution (γ=1) applied in this analysis was strong and conservative (i.e., the zero-shift model

was set to be the most likely outcome), the zero-shift model was never sampled in the posterior

for any of the BAMM runs showing overwhelming evidence of the heterogeneity of shifts found

in this analysis. Moreover, the sensitivity analysis based on a subset of the data (i.e. the

Särkinen et al. 2013 phylogeny without the additional species) found that the number of shifts

was not sensitive to different priors of diversification rates (i.e., different values of γ,

0.5,1,2,10,100) as shown in Supporting Information Fig. S11. Additionally, the analysis of

diversification rates in RevBayes showed similar results to those from the BAMM analysis (see

Supporting Information Fig. S3, S4).

Although most of the diversity and endemism of *Solanum* is found in the Neotropics (ca.

70% of species, see Fig.1, 2a), the highest diversification rates are seen in lineages

mainly concentrated in Australia, Africa and the Indo-Pacific (Fig. 2b). Africa shows a

heterogeneity of rates as it contains species from groups with standard diversification rates such

as Morelloids (0.27 lineages Myr^{-1}) and African non-spiny (0.25 lineages Myr^{-1}) as well as

species from the rapidly diversifying Old World clade (0.68 lineages Myr^{-1}).

Lineages-through-time curves (Fig. 3a) confirm that most of the diversification in *Solanum*

has occurred within the Neotropics. At the global level, there is a slight acceleration in the

number of lineages of *Solanum* in the last 5 Myr, which is likely to be shaped by the

considerable increase of lineages in Australia around the same time. Australia shows an

interesting latitudinal heterogeneity of diversification rates (Fig. 3b) shaped by the distribution of

lineages of different evolutionary origins — species from the Archaesolanum subclade, which

belongs to the M1 clade in Fig. 1, are mainly distributed in the south with 0.20 lineages Myr^{-1}

whereas the Old World spiny species with 0.68 lineages Myr^{-1} have a widespread distribution.

Ancestral range reconstructions across the *Solanum* phylogeny are best explained by the

DEC M1 model (Dispersion, Extinction and cladogenesis), which allows for equal probabilities of

dispersion from the Neotropics to both Africa and Australia (Supporting Information Table S2).

Including founder-event speciation in the model did not improve the AIC values significantly (Δ

AIC = 0.5), therefore, all the historical biogeographic results, including the Biogeographic

Stochastic Mapping, in this study were based on the simpler model, DEC M1.

The genus *Solanum* appears to have risen in the Neotropics ca. 15 Myr (95% HPD 13-18), as

do the majority of its subclades (see Fig. 1). The distribution of *Solanum* appears to have

occurred mainly via within-area sympatric speciation and dispersion, with vicariance only

supported in 3% of the total events (see Supporting Information Table S3). The distribution of

Solanum in the Nearctic region occurred through at least seven dispersal events out of the

Neotropics. Although several subclades are currently distributed in the Old World, only two

subclades appear to have arisen there via historical events (as opposed to recent introductions).

These dispersal events from the Neotropics to the Old World occurred at different times and in

two different groups of *Solanum*. The first movement from the Neotropics to Africa + Australia is

likely to have occurred ca. 10 Myr (95% HPD 7-12) within the non-spiny solanums, in the crown

group of the subclades *Solanum valdiviense* + African-non-spiny + Normania + Archaesolanum

(referred to as the M1 subclade on Fig. 1). The second dispersal event occurred ca. 6 Myr (95%

HPD 5-7) within the spiny solanums in the subclade Elaeagnifolium + Old World spiny clade

(see Fig. 1). The direction of these dispersal events from the Neotropics to either Africa and

Australia is still unresolved from the stochastic mapping estimates in both the non-spiny

solanums (a 30% dispersal probability from the Neotropics to Africa, and 24% to Australia) and

the spiny solanums (a probability of 45% of dispersing from the Neotropics to Africa, and 29% to

Australia).

Overall, the Neotropics were the main source of *Solanum* movements with more than 60% of

the estimated dispersal events (Supporting Information Fig. S12, S13). The most frequent

dispersals involved movements from the Neotropics to the Nearctic with ca. 40% of the total

estimated events. Movements from the Neotropics to Australia appear to be more frequent than

those from the Neotropics to Africa (8.7 ± 1.1 vs 2.9 ± 0.9). Dispersals from Africa to Australia

are more common than those in the opposite direction (5.5 ± 1.2 vs 1.64 ± 0.9).

Discussion

In this study, we demonstrate that the great diversity of *Solanum* is likely to be the result of a

great heterogeneity in evolutionary rates among its subclades supported by at least two recent

radiations — the Old World spiny clade and the Petota subclade (Fig. 1). Previous studies such

as Whalen, 1979 and Whalen & Caruso, 1983 were unable to identify unusual rates of

speciation within *Solanum*, concluding that the great taxonomic diversity of this genus may

reflect a gradual accumulation of species in a relatively old clade (which at the time was

believed to have a Cretaceous origin, Hawkes & Smith, 1965). However, the results of these

previous studies were based on species-poor but well-known sections of *Solanum* such as

Lasiocarpa (13 species) and Androceras (12 species) which ignored the great heterogeneity of

evolutionary regimes present in other groups of the genus. Our results demonstrate the

importance of accounting for among-clade rate heterogeneity in understanding the complex

dynamics of diversification of megadiverse groups such as *Solanum* (Ricklefs, 2007; Morlon,

2014).

Despite most of the richness and endemism of *Solanum* occurring in the Neotropics (Fig. 2a),

this region has lower diversification rates than the Old World (Fig. 2b). This mismatch between

the high diversity of *Solanum* within the Neotropics and its net diversification rates could be

explained by the early and long history of diversification of *Solanum* in the Neotropics. *Solanum*

lineages within the Neotropics are older and, therefore, had more time and opportunity to

accumulate before establishing in other regions, creating the current imbalance of diversity

(McPeek & Brown, 2007; Weir & Schluter, 2007). The Neotropics is the region where *Solanum*

species experienced the greater within-area speciation events, which has also been

demonstrated as the highest within the family Solanaceae as a whole (Dupin *et al.*, 2016). The

long history of diversification and expansion of *Solanum* species within the Neotropics,

therefore, drove a greater lineage accumulation compared with other regions (Fig. 3a), making

the Neotropics the main source of diversity and origin of biogeographic movements of solanums

across the globe (Fig. 1, and Supporting Information Fig. S12, S13).

There are no obvious novel morphological or physiological traits associated with the

diversification of the Old World clade, given this group is not well defined by any single

distinctive morphological traits or combination of morphological characters (Stern *et al.*, 2011),

but instead is defined by its geographical distribution and robust monophyly (Levin *et al.*, 2006;

Weese & Bohs, 2007, 2010; Stern *et al.*, 2011; although Aubriot *et al.*, 2016 showed that six

species thought to belong to the Old World clade, based on their distribution, now belong to the

Torva clade). However, the signal of change in diversification at the base of the clade

supporting all spiny solanums in Fig. 1, indicate that the distinctive traits of spiny solanums as a

whole, such as stellate indumentum and prickles (Whalen, 1984) could have played an essential

role in the diversification of the Old World clade. In this context, the Old World clade could be an

example of an exaptive radiation where previously acquired traits – originally shaped by

different selective forces – are advantageous under a new selective regime (Simões *et al.*,

2016). Furthermore, this nested radiation of lineages of the Old World clade within spiny

solanums might be a classic example of Key confluence *sensu* Donoghue & Sanderson, 2015,

where shifts in diversification are associated with the interaction between key innovations and

extrinsic factors such as biogeographic movements and/or environmental changes, as shown in

our ancestral range reconstruction (Fig. 1).

We found that not all the *Solanum* groups distributed in the Old World (i.e., mainly in Africa,

Australia, and the Indo-Pacific) that were associated with long-dispersal events resulted in

explosive radiations such as occurred in the Old World clade. This suggests that distinct

ecological opportunities and evolutionary regimes might have shaped the diversity of spiny and

non-spiny *Solanum* in the Old World. The lineages of non-spiny solanums present in the Old

World are much less diverse than the spiny solanums and occur in different environments. In

Africa, the non-spiny solanums are less diverse despite being significantly older than the spiny

solanums; they are mostly found in forested and mesic regions (Knapp & Vorontsova, 2016). In
Australia, the diversity of non-spiny solanums is represented by the Archaesolanum group
known as the Kangaroo apples. With eight species in total (Poczai *et al.*, 2011), this group
occurs in the temperate and forested areas of the South West Pacific (e.g., Australia, Tasmania,
New Zealand and Papua New Guinea). In contrast, spiny solanums occurring in continental
Africa and Australia are much more diverse and are predominantly concentrated in arid areas
(Symon & Others, 1981; Whalen, 1984; Bean, 2004; Vorontsova *et al.*, 2013). In Africa, these
are in vegetation types such as the "Somalia-Masai regional centre of endemism" *sensu* White
(1983), whilst in Australia, the ca. 200 species of spiny solanums are predominantly distributed
in warmer and arid regions (Fig. 3b).
A comparison of temporal patterns of lineage accumulation among major regions shows a
distinctive signature of a rapid diversification within Australia at ca. 5 Myr (Fig. 3a). The timing of
diversification and dispersion of spiny solanums in the Old World, as well as their concentration
in arid and semi-arid regions of Australia (Fig. 3b), suggests a potential link between past
environmental changes and radiation of this group. A similar signature has been shown in a
wide range of groups from Australia (Harmon *et al.*, 2003; Crisp *et al.*, 2004; Crayn *et al.*, 2006;
Crisp & Cook, 2007, 2013; Rabosky *et al.*, 2007; Byrne *et al.*, 2008, 2011; Harrington *et al.*,
2012; Blom *et al.*, 2016). The dramatic climate change in Australia since its separation from
Gondwana at ca. 50 Myr is likely to have triggered the radiation of several taxa (Byrne *et al.*,
2008) and the extinction of others (Byrne *et al.*, 2011). The initial stage of aridification in the
mid-Miocene (ca. 15 Myr) and the late expansion of desert regions of Australia in the last 1-4
Myr, has led to the assembly of new biomes, such as the arid, monsoonal and alpine regions,
encouraging the establishment and expansion of arid-adapted lineages (e.g., Crisp *et al.*, 2004;
Crayn *et al.*, 2006; Rabosky *et al.*, 2007; Byrne *et al.*, 2008; Crisp & Cook, 2013). This pattern
and the congruence of the timing of diversification with other arid-adapted Australian groups
could indicate that the Old World spiny solanums established and diversified in Australia as the
arid environments expanded. The widespread distribution of Old World spiny solanums in
Australia suggests a positive effect of distribution area on diversification rates. The large
expanse of arid habitat in Australia may have created opportunities for allopatric speciation
within the biome itself (Losos & Schluter, 2000; Davies *et al.*, 2005; Rabosky *et al.*, 2007). The
successful expansion and diversification of spiny solanums in warm and arid zones in Australia,
in contrast with the slower diversification of the more mesic non-spiny clades, could indicate the
influence of features adapted to warm and dry environments on species diversification.
Moreover, the sister group of the diverse Old World clade is the species-poor Elaeagnifolium

clade, with five species confined to semi-arid regions of North and South America (Knapp &
Vorontsova, 2016), suggesting that some pre-adaptation traits to dry conditions may have
driven diversification in the Old World. This evidence suggests that the dispersal of spiny
solanums to the Old World, the influence of past environmental changes, as well as the pre-
adaptive condition of spiny solanums to colonise warm and arid regions contributed to the
explosive radiation of *Solanum* lineages in Africa and Australia.

Implications of the study

The dynamics shaping the diversity of species-rich mega-diverse genera may be more complex
than previously thought. A variety of mutually reinforcing drivers appear to have formed the
current patterns of diversity in *Solanum*, which cautions against single-cause hypotheses, such
as the influence of "key innovations" often invoked in macroevolutionary studies (Donoghue &
Sanderson, 2015). These findings also raise important questions about the reliability of global
studies that assume a single model for the evolution of character change and diversification
across large phylogenies (e.g., Zanne *et al.*, 2014), or studies that use average values of
diversity, trait distribution, or geographical distribution at generic level (e.g., Cornwell *et al.*,
2014). For instance, Beaulieu *et al.*, 2013 and Chira & Thomas, 2016 have demonstrated that
heterogeneity of diversification rates can have significant consequences for the model selection
of trait change that best fits comparative data. In general, therefore, the understanding of the
magnitude and location of shifts in diversification rates should be included as prior information in
macroevolutionary analyses, especially in analyses of trait evolution (Morlon, 2014; Chira &
Thomas, 2016), or at least models of trait evolution must not implicitly assume that such shifts
do not occur.

The results of this study also highlight the importance of context dependency in the study of
diversification dynamics (De Queiroz, 2002; Donoghue, 2005; Donoghue & Sanderson, 2015).
Our study shows that both opportunity and success in geographic movement, as well as correct
environmental setting and climatic adaptations, were necessary to spark faster diversification in
the spiny clade of *Solanum* in the Old World. Thus, focusing on effects of the combination of
events and attributes on changes in diversification could represent a more productive framework
in which to understand diversity processes (Donoghue & Sanderson, 2015). This combination
could also explain why, although many correlates of diversification rates have been reported in
the literature, few have demonstrated either generality or high explanatory power. Our results

suggest the necessity of considering more complex analyses that involve the integration of

phylogenies with other biological and Earth system data sources such as geography, climate,

historical biogeography, and physiology.

We demonstrate that *Solanum* is a remarkable study system for understanding the

diversification of plants, not only because it contains a recent and ongoing radiation, but also

because its diversity is supported by several clades with different and sometimes contrasting

evolutionary dynamics. There are still many drivers that need to be explored before we have the

whole picture of the diversification of *Solanum*, especially correlations with biogeographic

events and morphological traits, but increased taxonomic, phylogenetic, and morphological

understanding of *Solanum* now means that we are beginning to uncover many aspects of the

macroevolution of this mega-diverse plant genus, and can use it as a model for studying

diversity in widespread, species-rich groups in general.

Acknowledgements

This study was supported by the National Science Foundation (NSF) grant (to SK) 'PBI

Solanum – a world treatment' DEB-0316614, and Colciencias (Administrative department of

Science, Technology and Innovation of Colombia) who partially funded S.E.-L.'s PhD

programme.

Author Contribution

498
S.E.-L designed the study, analysed the data and wrote the first draft of the manuscript. TS

provided the phylogenetic data (reconstructed the phylogeny), distribution and taxonomic data.

ISF provided advice on the analysis and contributed to the main discussion of the

macroevolutionary analyses. SK provided the taxonomic, distribution and diversity data, and

supervised the study. AP designed and supervised the analysis. All authors reviewed the

manuscript.

References

Alfaro ME, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky DL, Carnevale G, Harmon
LJ. **2009**. Nine exceptional radiations plus high turnover explain species diversity in jawed
vertebrates. *Proceedings of the National Academy of Sciences* **106**: 13410–13414.

Aubriot X, Singh P, Knapp S. **2016**. Tropical Asian species show that the Old World clade of
`spiny solanums' (Solanum subgenus Leptostemonum pro parte: Solanaceae) is not
monophyletic. *Botanical journal of the Linnean Society. Linnean Society of London* **181**: 199–
223.

Bean AR. **2004**. The taxonomy and ecology of *Solanum* subg. Leptostemonum (Dunal) Bitter
(Solanaceae) in Queensland and far north-eastern New South Wales, Australia. *Austrobaileya*:
639–816.

Beaulieu JM, O'Meara BC, Donoghue MJ. **2013**. Identifying hidden rate changes in the
evolution of a binary morphological character: the evolution of plant habit in campanulid
angiosperms. *Systematic biology* **62**: 725–737.

Blom MPK, Horner P, Moritz C. **2016**. Convergence across a continent: adaptive
diversification in a recent radiation of Australian lizards. In: Proc. R. Soc. B. The Royal Society,
20160181.

Burnham KP, Anderson DR. **2002**. *Model selection and multi-model inference: a practical*
information-theoretic approach. Springer.

Byrne M, Steane DA, Joseph L, Yeates DK, Jordan GJ, Crayn D, Aplin K, Cantrill DJ, Cook
LG, Crisp MD, *et al.* **2011**. Decline of a biome: evolution, contraction, fragmentation, extinction
and invasion of the Australian mesic zone biota. *Journal of biogeography* **38**: 1635–1656.

Byrne M, Yeates DK, Joseph L, Kearney M, Bowler J, Williams MAJ, Cooper S, Donnellan
SC, Keogh JS, Leys R, *et al.* **2008**. Birth of a biome: insights into the assembly and
maintenance of the Australian arid zone biota. *Molecular ecology* **17**: 4398–4417.

Chira AM, Thomas GH. **2016**. The impact of rate heterogeneity on inference of phylogenetic
models of trait evolution. *Journal of evolutionary biology* **29**: 2502–2518.

Cornwell WK, Westoby M, Falster DS, FitzJohn RG, O'Meara BC, Pennell MW, McGlinn DJ, Eastman JM, Moles AT, Reich PB, *et al.* 2014. Functional distinctiveness of major plant lineages. *The Journal of ecology* **102**: 345–356.

Cox CB. **2001**. The biogeographic regions reconsidered. *Journal of biogeography* **28**: 511–523.

Crayn DM, Rossetto M, Maynard DJ. **2006**. Molecular phylogeny and dating reveals an Oligo-Miocene radiation of dry-adapted shrubs (former Tremandraceae) from rainforest tree progenitors (Elaeocarpaceae) in Australia. *American journal of botany* **93**: 1328–1342.

Crisp MD, Cook LG. **2007**. A congruent molecular signature of vicariance across multiple plant lineages. *Molecular phylogenetics and evolution* **43**: 1106–1117.

Crisp MD, Cook LG. **2013**. How was the Australian flora assembled over the last 65 million years? A molecular phylogenetic perspective. *Annual review of ecology, evolution, and systematics* **44**: 303–324.

Crisp M, Cook L, Steane D. **2004**. Radiation of the Australian flora: what can comparisons of molecular phylogenies across multiple taxa tell us about the evolution of diversity in present--day communities? *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **359**: 1551–1571.

Davies TJ, Savolainen V, Chase MW, Goldblatt P, Barraclough TG. **2005**. Environment, area, and diversification in the species-rich flowering plant family Iridaceae. *The American naturalist* **166**: 418–425.

De Queiroz A. **2002**. Contingent predictability in evolution: key traits and diversification. *Systematic biology* **51**: 917–929.

Donoghue MJ. **2005**. Key innovations, convergence, and success: macroevolutionary lessons from plant phylogeny. *Paleobiology* **31**: 77–93.

Donoghue MJ, Sanderson MJ. **2015**. Confluence, synnovation, and depauperons in plant diversification. *The New phytologist*.

Dupin J, Matzke NJ, Särkinen T, Knapp S, Olmstead RG, Bohs L, Smith SD. **2016**. Bayesian estimation of the global biogeographical history of the Solanaceae. *Journal of biogeography* **44**: 887–899.

Frodin DG. **2004**. History and concepts of big plant genera. *Taxon* **53**: 753–776.

Harmon LJ, Schulte JA, Larson A, Losos JB. **2003**. Tempo and mode of evolutionary
radiation in iguanian lizards. *Science* **301**: 961–964.

Harrington MG, Jackes BR, Barrett MD, Craven LA, Barrett RL. **2012**. Phylogenetic revision
of Backhousieae (Myrtaceae): Neogene divergence, a revised circumscription of Backhousia
and two new species. *Australian systematic botany* **25**: 404–417.

Hawkes JG, Smith P. **1965**. Continental drift: continental drift and the age of angiosperm
genera. *Nature* **207**: 48–50.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP,
Ronquist F. 2016a. RevBayes: Bayesian Phylogenetic Inference Using Graphical Models and
an Interactive Model-Specification Language. *Systematic biology*: syw021.

Höhna S, May MR, Moore BR. **2016b**. TESS: Bayesian inference of lineage diversification
rates from (incompletely sampled) molecular phylogenies in R. *Bioinformatics* **35**: 789–791.

Höhna S, Stadler T, Ronquist F, Britton T. **2011**. Inferring speciation and extinction rates
under different sampling schemes. *Molecular biology and evolution* **28**: 2577–2589.

Huelsenbeck JP, Nielsen R, Bollback JP. **2003**. Stochastic mapping of morphological
characters. *Systematic biology* **52**: 131–158.

Knapp S, Vorontsova MS. **2016**. A revision of the African Non-Spiny" Clade of Solanum
L.(Solanum sections Afrosolanum Bitter, Benderianum Bitter, Lemurisolanum Bitter,
Lyciosolanum Bitter, Macronesiotes Bitter, and Quadrangulare Bitter: Solanaceae). *PhytoKeys*
66: 1.

Landis MJ, Matzke NJ, Moore BR, Huelsenbeck JP. **2013**. Bayesian analysis of
biogeography when the number of areas is large. *Systematic biology* **62**: 789–804.

Levin RA, Myers NR, Bohs L. **2006**. Phylogenetic relationships among the 'spiny solanums'
(Solanum subgenus Leptostemonum, Solanaceae). *American journal of botany* **93**: 157–169.

Liow LH, Quental TB, Marshall CR. **2010**. When can decreasing diversification rates be
detected with molecular phylogenies and the fossil record? *Systematic biology*: syq052.

Losos JB. **2010**. Adaptive radiation, ecological opportunity, and evolutionary determinism. *The*
American naturalist **175**: 623–639.

Losos JB, Schluter D. **2000**. Analysis of an evolutionary species-area relationship. *Nature* **408**:
847.

Matzke NJ. **2012**. Founder-event speciation in BioGeoBEARS package dramatically improves
likelihoods and alters parameter inference in dispersal--extinction--cladogenesis (DEC)
analyses. *Front. Biogeogr* **4**: 210.

Matzke NJ. **2016**. Stochastic mapping under biogeographical models.

McPeek MA, Brown JM. **2007**. Clade age and not diversification rate explains species richness
among animal taxa. *The American naturalist* **169**: E97–E106.

Moore BR, Donoghue MJ. **2007**. Correlates of diversification in the plant clade Dipsacales:
geographic movement and evolutionary innovations. *The American naturalist* **170**: S28–S55.

Moore BR, Höhna S, May MR, Rannala B, Huelsenbeck JP. **2016**. Critically evaluating the
theory and performance of Bayesian analysis of macroevolutionary mixtures. *Proceedings of the*
National Academy of Sciences **113**: 9569–9574.

Morlon H. **2014**. Phylogenetic approaches for studying diversification. *Ecology letters* **17**: 508–
525.

Olmstead RG, Palmer JD. **1997**. Implications for the phylogeny, classification, and
biogeography of Solanum from cpDNA restriction site variation. *Systematic botany*: 19–29.

Pagel M. **1999**. The maximum likelihood approach to reconstructing ancestral character states
of discrete characters on phylogenies. *Systematic biology* **48**: 612–622.

Plummer M, Best N, Cowles K, Vines K. **2006**. CODA: Convergence Diagnosis and Output
Analysis for MCMC. *R News* **6**: 7–11.

Poczai P, Hyvönen J, Symon DE. **2011**. Phylogeny of kangaroo apples (Solanum subg.
Archaesolanum, Solanaceae). *Molecular biology reports* **38**: 5243–5259.

Purvis A. **1996**. Using interspecies phylogenies to test macroevolutionary hypotheses. *New*

uses for new phylogenies. Oxford University Press, Oxford: 153–168.

Quental TB, Marshall CR. **2010**. Diversity dynamics: molecular phylogenies need the fossil
record. *Trends in ecology & evolution* **25**: 434–441.

Rabosky DL. **2010**. Extinction rates should not be estimated from molecular phylogenies.
Evolution; international journal of organic evolution **64**: 1816–1824.

Rabosky DL. **2014**. Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. *PloS one* **9**: e89543.

Rabosky DL. **2015**. No substitute for real data: A cautionary note on the use of phylogenies
from birth--death polytomy resolvers for downstream comparative analyses. *Evolution;*
international journal of organic evolution **69**: 3207–3216.

Rabosky DL, Donnellan SC, Talaba AL, Lovette IJ. **2007**. Exceptional among-lineage
variation in diversification rates during the radiation of Australia's most diverse vertebrate clade.
Proceedings of the Royal Society of London B: Biological Sciences **274**: 2915–2923.

Rabosky DL, Grundler M, Anderson C, Shi JJ, Brown JW, Huang H, Larson JG, Others.
2014. BAMMtools: an R package for the analysis of evolutionary dynamics on phylogenetic
trees. *Methods in ecology and evolution / British Ecological Society*.

Rabosky DL, Santini F, Eastman J, Smith SA, Sidlauskas B, Chang J, Alfaro ME. **2013**.
Rates of speciation and morphological evolution are correlated across the largest vertebrate
radiation. *Nature communications* **4**.

Raup DM, Gould SJ, Schopf TJM, Simberloff DS. **1973**. Stochastic models of phylogeny and
the evolution of diversity. *The Journal of geology* **81**: 525–542.

Ree RH, Moore BR, Webb CO, Donoghue MJ, Crandall K. **2005**. A likelihood framework for
inferring the evolution of geographic range on phylogenetic trees. *Evolution; international journal*
of organic evolution **59**: 2299–2311.

Ricklefs RE. **2007**. Estimating diversification rates from phylogenetic information. *Trends in*
ecology & evolution **22**: 601–610.

Ronquist F. **1997**. Dispersal-vicariance analysis: a new approach to the quantification of

historical biogeography. *Systematic biology* **46**: 195–203.

Ronquist F, Huelsenbeck JP. **2003**. MrBayes 3: Bayesian phylogenetic inference under mixed
models. *Bioinformatics* **19**: 1572–1574.

Särkinen T, Bohs L, Olmstead RG, Knapp S. **2013**. A phylogenetic framework for evolutionary
study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC evolutionary biology* **13**:
214.

Schluter D. **2000**. *The ecology of adaptive radiation*. Oxford University Press.

Simões M, Breitkreuz L, Alvarado M, Baca S, Cooper JC, Heins L, Herzog K, Lieberman
BS. **2016**. The Evolving Theory of Evolutionary Radiations. *Trends in ecology & evolution* **31**:
27–34.

Simpson GG. **1955**. *Major features of evolution*. Columbia University Press: New York.

Slowinski JB, Guyer C. **1993**. Testing whether certain traits have caused amplified
diversification: an improved method based on a model of random speciation and extinction. *The*
American naturalist: 1019–1024.

Stadler T. **2011**. Mammalian phylogeny reveals recent diversification rate shifts. *Proceedings of*
the National Academy of Sciences **108**: 6187–6192.

Stadler T. **2013**. Recovering speciation and extinction dynamics based on phylogenies. *Journal*
of evolutionary biology **26**: 1203–1219.

Stanley SM, Signor PW III, Lidgard S, Karr AF. **1981**. Natural clades differ from ``random''
clades: simulations and analyses. *Paleobiology*: 115–127.

Stern S, Agra M de F, Bohs L. **2011**. Molecular delimitation of clades within New World
species of the ``spiny solanums" (Solanum subg. Leptostemonum). *Taxon* **60**: 1429–1441.

Symon DE, Others. **1981**. A revision of the genus Solanum in Australia. *Journal of the Adelaide*
Botanic Gardens **4**: 1–367.

Thomas GH, Hartmann K, Jetz W, Joy JB, Mimoto A, Mooers AO. **2013**. PASTIS: an R
package to facilitate phylogenetic assembly with soft taxonomic inferences. *Methods in ecology*

and evolution / British Ecological Society **4**: 1011–1017.

Vorontsova M, Stern S, Bohs L, Knapp S, Others. **2013**. African spiny Solanum (subgenus
Leptostemonum, Solanaceae): a thorny phylogenetic tangle. *Botanical journal of the Linnean*
Society. Linnean Society of London **173**: 176–193.

Weese TL, Bohs L. **2007**. A three-gene phylogeny of the genus Solanum (Solanaceae).
Systematic botany **32**: 445–463.

Weese TL, Bohs L. **2010**. Eggplant origins: out of Africa, into the Orient. *Taxon* **59**: 49–56.

Weir JT, Schluter D. **2007**. The latitudinal gradient in recent speciation and extinction rates of
birds and mammals. *Science* **315**: 1574–1576.

Whalen MD. **1979**. Allozyme variation and evolution in Solanum section Androceras.
Systematic botany: 203–222.

Whalen MD. **1984**. Conspectus of species groups in Solanum subgenus Leptostemonum.
Gentes Herb. (Ithaca) **12**: 179–282.

Whalen MD, Caruso EE. **1983**. Phylogeny in Solanum sect. Lasiocarpa (Solanaceae):
congruence of morphological and molecular data. *Systematic botany*: 369–380.

Wiens JJ. **2011**. The causes of species richness patterns across space, time, and clades and
the role of 'ecological limits'. *The Quarterly review of biology* **86**: 75–96.

Wiens JJ. **2015**. Faster diversification on land than sea helps explain global biodiversity
patterns among habitats and animal phyla. *Ecology letters* **18**: 1234–1241.

Yoder JB, Clancey E, Des Roches S, Eastman JM, Gentry L, Godsoe W, Hagey TJ,
Jochimsen D, Oswald BP, Robertson J, *et al.* **2010**. Ecological opportunity and the origin of
adaptive radiations. *Journal of evolutionary biology* **23**: 1581–1596.

Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG, McGlinn DJ,
O'Meara BC, Moles AT, Reich PB, *et al.* **2014**. Three keys to the radiation of angiosperms into
freezing environments. *Nature* **506**: 89–92.

Figures legends:

Fig. 1. Mean net diversification rates and historical biogeographic reconstruction across the

phylogeny of *Solanum*. For the net diversification rates, bars represent the frequency with which

nodes are associated with shifts in diversification rates across a distribution of 100 complete

phylogenies of *Solanum* (see Methods for further detail). For the historical biogeographic

reconstruction, pies at each node represent the probabilities of the ancestral range distribution;

the map indicates the regional colour code. The branches highlighted in blue represent the

dispersals events inferred in at least the 50% of Biogeographic Stochastic Maps (BSMs). The

timescale is given in Myr.

Fig. 2. Global distribution of species richness (a) and the net diversification rates of *Solanum* (b)

using 1 x 1 degree grid cells. Weighted mean lineage net diversification rates were calculated

as the mean net diversification rate for all the species present in a grid cell assemblage,

weighted by the inverse of their range size, $log_e m^2$

Fig. 3. (a) Lineages-through-time plot of *Solanum* species for the global and three regional

diversities (Neotropics, Australia, and Africa). Coloured areas around the median curves, shown

in black, represent two-tailed 95% upper and lower quantiles of the number of lineages across a

distribution of 100 complete phylogenies of *Solanum* (see Methods for further details). (b)

Distribution of the net diversification rates of *Solanum* in Australia (same units as in Fig. 2b).

0.2 Net Diversification Rates 0.7

Old World clade

Spiny

Petota

non-Spiny

M1

Species richness(log$_e$)

0   1   2   3   4   5

Weighted mean diversification rates(species Myr^{-1}/log$_e$m^2)

0.010   0.015   0.020   0.025   0.030