

## The global burden of trichiasis in 2016

Surname	Full Name	Affiliation
Flueckiger	Rebecca M Flueckiger	Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom
Courtright	Paul Courtright	Kilimanjaro Centre for Community Ophthalmology, Division of Ophthalmology, University of Cape Town, Cape Town, South Africa
Abdala	Mariamo Abdala	Ministerio da Saude, Maputo, Mozambique
Abdou	Amza Abdou	Ministère de la Santé et de la population, Niamey, Niger
Abdulnafa	Zaid Abdulnafa	Ministry of Health, Baghdad, Iraq
Al-Khatib	Tawfik K Al-Khatib	National Eye Health Programme, Ministry of Public Health and Population, Sana'a, Yemen
Amer	Khaled Amer	Ministry of Health, Egypt
Amiel	Olga Nelson Amiel	Ministerio da Saude, Maputo, Mozambique
Awoussi	Sossinou Awoussi	Ministère de la Santé, Lomé, Togo
Bakhtiari	Ana Bakhtiari	The Task Force for Global Health, Atlanta, GA, USA
Batcho	Wilfried Batcho	Ministère de la Santé, Cotonou, Benin
Bella	Assumpta Lucienne Bella	Ministère de la Santé, Yaoundé, Cameroun
Bennawi	Kamal Hashim Bennawi	Prevention of Blindness Program, Khartoum, Sudan
Brooker	Simon J Brooker	Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom; Global Health, Neglected Tropical Diseases, Bill & Melinda Gates Foundation, Seattle, USA
Chu	Brian K Chu	The Task Force for Global Health, Atlanta, GA, USA
Dejene	Michael Dejene	Michael Dejene Public Health Consultancy Services, Addis Ababa, Ethiopia
Dezoumbe	Djore Dezoumbe	Ministère de la Santé, Chad
Eishafie	Balgesa Elkheir Eishafie	Trachoma Control Program, National Ministry of Health, Khartoum, Sudan
Elvis	Aba Ange Elvis	Programme National de la Santé Oculaire et de la lutte contre l'Onchocercose, Abidjan, Côte d'Ivoire
Fabrice	Djouma Nembot Fabrice	Ministère de la Santé, Yaoundé, Cameroun
Omar	Fatma Juma Omar	Ministry of Health, Zanzibar, United Republic of Tanzania
François	Missamou François	Bureau des Maladies Oculaires, Ministère de la Santé, Kinshasa, Democratic Republic of the Congo
François	Drabo François	Programme national de lutte contre les maladies tropicales négligées (PNMTN), Ouagadougou, Burkina Faso
Garap	Jambi Garap	National Department of Health, Papua New Guinea
Gichangi	Michael Gichangi	Ministry of Health, Nairobi, Kenya
Goepogui	André Goepogui	Programme National de Lutte Contre l'Onchocercose et la Cécité et les Maladies Tropicales Négligées, Conakry, Guinea
Hammou	Jaouad Hammou	Ministry of Health, Rabat, Morocco
Kadri	Boubacar Kadri	Ministère de la Santé et de la population, Niamey, Niger
Kabona	George Kabona	Ministry of Health, Dar es Salaam, United Republic of Tanzania
Kabore	Martin Kabore	l'unité d'élimination du trachome, PNMTN, Ouagadougou, Burkina Faso
Kalua	Khumbo Kalua	Department of Ophthalmology, University of Malawi, College of Medicine, Blantyre, Malawi
Kamugisha	Mathias Kamugisha	National Institute for Medical Research, Dar es Salaam, United Republic of Tanzania
Kebede	Biruck Kebede	Federal Ministry of Health, Addis Ababa, Ethiopia
Keita	Kaba Keita	Programme National de Lutte Contre l'Onchocercose et la Cécité et les Maladies Tropicales Négligées, Conakry, Guinea
Khan	Asad Aslam Khan	Ministry of Health, Islamabad, Pakistan
Kiflu	Genet Kiflu	Federal Ministry of Health, Addis Ababa, Ethiopia
Mackline	Garae Mackline	Ministry of Health, Port Vila, Vanuatu
Macleod	Colin Macleod	Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom
Manangazira	Portia Manangazira	Department of Epidemiology and Disease Control, Ministry of Health & Child Welfare, Harare, Zimbabwe
Masika	Michael P Masika	Ministry of Health, Lilongwe, Malawi
Massangaie	Marilia Massangaie	Ministerio da Saude, Maputo, Mozambique

Mduluza	Taka Fira Mduluza	Department of Biochemistry, University of Zimbabwe, Harare, Zimbabwe
Meno	Nabicassa Meno	Ministère de la Santé, Bissau, Guinea-Bissau
Midzi	Nicholas Midzi	Department of Medical Microbiology, College of Health Sciences, University of Zimbabwe, Harare, Zimbabwe
Minnih	Abdallahi Ould Minnih	Minister de la Sante, Nouakchott, Mauritania
Mishra	Sailesh Mishra	Nepal Netra Jyoti Sangh, Kathmandu, Nepal
Mpyet	Caleb Mpyet	Sightsavers Nigeria, Kaduna, Nigeria & Department of Ophthalmology, Jos University, Jos, Nigeria
Muraguri	Nicholas Muraguri	Ministry of Health, Nairobi, Kenya
Mwingira	Upendo Mwingira	Ministry of Health, Dar es Salaam, United Republic of Tanzania
Nassirou	Beido Nassirou	Ministère de la Santé et de la population, Niamey, Niger
Ndjemba	Jean Ndjemba	Bureau des Maladies Oculaires, Ministère de la Santé, Kinshasa, Democratic Republic of the Congo
Nieba	Cece Nieba	Programme National de Lutte Contre l'Onchocercose et la Cécité et les Maladies Tropicales Négligées, Conakry, Guinea
Ngondi	Jeremiah Ngondi	RTI International, Dar es Salaam, United Republic of Tanzania
Olobio	Nicholas Olobio	Nigeria Federal Ministry of Health, Abuja, Nigeria
Pavluck	Alex Pavluck	RTI International, Washington DC, USA
Phiri	Isaac Phiri	Department of Epidemiology and Disease Control, Ministry of Health & Child Welfare, Harare, Zimbabwe
Pullan	Rachel Pullan	Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom
Qureshi	Babar Qureshi	Christian Blind Mission, Bensheim, Germany
Sarr	Boubacar Sarr	Ministère de la santé et de l'Action Sociale, Dakar, Senegal
Seiha	Do Seiha	Prevention of Blindness Programme, Ministry of Health, Phnom Penh, Cambodia
Serrano	Gloria Marina Serrano Chávez	Ministry of Health, Guatemala City, Guatemala
Sharma	Shekhar Sharma	Ministry of Health, Kathmandu, Nepal
Sisaleumsak	Siphethavong Sisaleumsak	National Ophthalmology Centre, Vientiane, Lao People's Democratic Republic
Southisombath	Khamphoua Southisombath	National Program for the Prevention of Blindness, Ministry of Health, Vientiane, Lao People's Democratic Republic
Stevens	Gretchen Stevens	Department of Information, Evidence and Research, World Health Organization, Geneva, Switzerland
Woldendrias	Andeberhan Tesfazion Woldendrias	Department of Public Health, Ministry of Health, Asmara, Eritrea
Traoré	Lamine Traoré	Ministère de la Santé, Bamako, Mali
Turyaguma	Patrick Turyaguma	Trachoma Program, Ministry of Health, Kampala, Uganda
Willis	Rebecca Willis	The Task Force for Global Health, Atlanta, GA, USA
Yaya	Georges Yaya	Ministère de la Santé Publique, Bangui, Central African Republic
Yeo	Souleymane Yeo	Programme National de la Santé Oculaire et de la lutte contre l'Onchocercose, Abidjan, Côte d'Ivoire
Zambroni	Francisco Zambroni	Ministry of Health, Guatemala City, Guatemala
Zhao	Jialiang Zhao	Department of Ophthalmology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences, Beijing, China
Solomon	Anthony W Solomon	Department of Control of Neglected Tropical Diseases, World Health Organization, Geneva, Switzerland

\*Email: [Rebecca.Flueckiger@lshtm.ac.uk](mailto:Rebecca.Flueckiger@lshtm.ac.uk)

## **Abstract**

### **Background**

Trichiasis is present when one or more eyelashes touches the eye. Uncorrected, it can cause blindness. Accurate estimates of numbers affected, and their geographical distribution, help guide resource allocation.

### **Methods**

We obtained district-level trichiasis prevalence estimates for 44 endemic and previously-endemic countries. We used (1) the most recent data for a district, if more than one estimate was available; (2) age- and sex-standardized corrections of historic estimates, where raw data were available; (3) historic estimates adjusted using a mean adjustment factor for districts where raw data were unavailable; and (4) expert assessment of available data for districts for which no prevalence estimates were available.

### **Findings**

Internally age- and sex-standardized data represented 1,355 districts and contributed 662 thousand cases (95% confidence interval [CI] 324 thousand–1.1 million) to the global total. Age- and sex-standardized district-level prevalence estimates differed from raw estimates by a mean factor of 0.45 (range 0.03–2.28). Previously non-standardized estimates for 398 districts, adjusted by  $\times 0.45$ , contributed a further 411 thousand cases (95% CI 283–557 thousand). Eight countries retained previous estimates, contributing 848 thousand cases (95% CI 225 thousand–1.7 million). New expert assessments in 14 countries contributed 862 thousand cases (95% CI 228 thousand–1.7 million). The global trichiasis burden in 2016 was 2.8 million cases (95% CI 1.1–5.2 million).

### **Interpretation**

The 2016 estimate is lower than previous estimates, probably due to more and better data; scale-up of trichiasis management services; and reductions in incidence due to lower active trachoma prevalence.

## **Author Summary**

As an individual with trichiasis blinks, the eyelashes abrade the cornea, which can lead to corneal opacity and blindness. Through high quality surgery, which involves correcting the position of the in-turned eyelid, it is possible to reduce the number of people with trichiasis. An accurate estimate of the number of persons with trichiasis and their geographical distribution are needed in order to effectively align resources for surgery and other necessary services. We obtained district-level trichiasis prevalence estimates for 44 endemic and previously-endemic countries. We used the most recently available data and expert assessments to estimate the global burden of trichiasis. We estimated that in 2016 the global burden was 2.8 million cases (95% CI 1.1–5.2 million). The 2016 estimate is lower than previous estimates, probably due to more and better data; scale-up of trichiasis management services; and reductions in incidence due to lower active trachoma prevalence.

## Background

Trachoma, a neglected tropical disease, is endemic or has recently been endemic in more than 50 countries[1]. It affects the most impoverished people of the world. Improved living standards are credited for the disappearance of trachoma from Europe and North America, but in many less developed countries, trachoma is still a public health problem, and contributes to the continued suffering and deepening of poverty of millions of people.

*Chlamydia trachomatis* is the causative organism. Repeated ocular chlamydial infection results in chronic inflammation, characterised by sub-epithelial follicles in the tarsal conjunctiva, which may be sufficiently large and numerous to meet the definition of trachomatous inflammation—follicular (TF), a key sign for assessing trachoma prevalence[2]. Over time, with repeated reinfection, scarring may develop; this scarring can eventually cause the eyelid to turn inwards in some people, resulting in eyelashes touching the globe. This is called trachomatous trichiasis (TT) and is very painful[3]. As an individual with trichiasis blinks, the eyelashes abrade the cornea, which can lead to corneal opacity and blindness[4].

The World Health Organization (WHO) endorses population based prevalence surveys (PBPSs) to guide decisions regarding implementation of interventions for trachoma elimination[5]. The typical survey design involves two-stage cluster sampling, which uses probability-proportional-to-size methodologies to select 20–30 clusters. The outputs are estimates of the prevalence of TF in children aged 1–9 years, and the prevalence of TT in adults aged 15 years and older[6].

Through high quality surgery, which involves correcting the position of the in-turned eyelid,[7] it is possible to reduce the number of people with TT. An accurate estimate of the number of persons with TT (the TT backlog) and their geographical distribution are needed in order to effectively align resources for surgery and other necessary services.

In 2009, Mariotti et al estimated the global TT backlog to be 8.2 million people[8]. This estimate was derived by summarizing a combination of published and unpublished information. First, a literature review was performed to identify published prevalence data. Where published data were not available, unpublished data were compiled from

the Eleventh (2007) Meeting of the WHO Alliance for the Global Elimination of Trachoma by 2020[9]. Where information was still missing, unpublished reports were provided by health ministries. Finally, if none of these resources were available, data were extrapolated from a proxy country believed to have common epidemiological conditions and demographic structure.

There are many uncertainties inherent in the 2009 estimate[8]. First, where PBPS data were available, the results were not systematically standardized by age and sex. This is problematic because women and the elderly are more likely than men and younger adults, respectively, to (1) be at home at the time that a house-to-house survey team calls, and (2) have TT. Second, in countries for which data were available, survey coverage was generally far from complete. Though not an invariable rule, surveys to estimate the prevalence of neglected diseases have a tendency to be done first in areas of high prevalence; for the 2009 estimate, if any data were available for a particular country, the TT prevalence figure from it was applied across the yet-to-be-mapped suspected-endemic population. Third, the use of proxy countries is extremely subjective.

In 2012, WHO collected and collated provisional 2011 country reports, and published an updated figure for the global TT backlog. The total given was 7.3 million people, but the methodologies used in each country to generate national backlog figures were not described[10]; they are likely to have been highly heterogeneous.

From December 2012 to January 2016, the Global Trachoma Mapping Project (GTMP) sought to systematically complete the global trachoma map using standardized techniques for both collecting and analysing survey data[11]. GTMP measured trachoma prevalence using gold standard PBPSs conducted at district level. People of all ages living in selected households of 1,542 districts across 29 countries were examined for trachoma, resulting in the examination of 2.6 million people[12]. GTMP analyses included standardizing trichiasis prevalence estimates against national population pyramids in an attempt to partially account for demographic differences between those examined and the national averages.

As a result of the GTMP, there are now high quality PBPS data available for most suspected-endemic areas that were previously unsurveyed. The availability of these data has catalysed the current attempt to generate a new estimate of the global trichiasis backlog.

## **Methodology**

We updated previous global estimates using the best available data, according to the following hierarchy. First, where GTMP data[11] were available, we used the age- and sex-standardized trichiasis prevalence estimates for adults in each survey. In August 2014, GTMP added examination for the presence or absence of trachomatous scarring (TS)[2] for all eyes determined to have trichiasis; prior to this, and for all non-GTMP data, trichiasis prevalence estimates used must be considered to be estimates of “all-cause trichiasis” rather than “trachomatous trichiasis”.

Second, where PBPSs had been done without the support of the GTMP, we requested raw survey data from national health ministries, or appropriate government-designated agencies. Where those data were provided by 1 March 2016, we applied the same age- and sex- standardization as was used in the GTMP[11].

Third, where PBPSs had been done but raw data were not available, prevalence estimates (whether standardized for age and sex or not) were obtained from country programs. When data had not been standardized, estimates were multiplied by the mean adjustment factor for districts in which age- and sex-standardization was possible.

Fourth, if prevalence estimates were not available, we reviewed previous estimates through desk reviews and communications with country programs. If there was adequate evidence to revise the estimates, new estimates were used; otherwise, the 2009 estimates[8] were retained.

The population data used in this analysis were derived from the UN population division (UNdata)[13] and [www.worldpop.org](http://www.worldpop.org)[14]. As trachoma is a disease primarily affecting rural populations[4], we used rural population estimates for this purpose. Rural population pyramids were obtained from UNdata. Microsoft Excel (2007) was used to organize the datasets into 5-year age bands stratified by sex for each country. The percentage of the population

within each stratum was estimated from the bands. The district level populations were derived from [www.worldpop.org](http://www.worldpop.org) raster files[14] using the zonal statistics tool in ArcGIS 10.3[15], where the sum raster value falling within each district boundary was taken as the district value. A sensitivity analysis was also performed, comparing the [www.worldpop.org](http://www.worldpop.org) population estimates and district population estimates provided by national trachoma elimination programs. The mean ratio between the national program estimates and [www.worldpop.org](http://www.worldpop.org) was 0.97. Because of this close correlation, and for the purposes of standardizing our methods, the [www.worldpop.org](http://www.worldpop.org) data were used throughout this analysis.

The statistical software package R[16] was used to perform age- and sex- standardization. First, the crude prevalence was calculated for each cluster. Second, the prevalence was standardized for each cluster by weighting the proportion of each sex-specific five-year age band observed to have trichiasis by the proportion of the adults aged 15 years and older expected to have that age and sex in that district, if available, or (if not available), nationwide. Third, the un-weighted arithmetic mean of the standardized cluster-level trichiasis proportions was taken as the district-level prevalence. Once the initial standardization was completed, bootstrapping was undertaken on the dataset for each individual district to derive 95% confidence intervals (CIs). For a district surveyed by examining individuals in  $n$  clusters, this involved bootstrap resampling (with replacement) of  $n$  clusters, over 10,000 replications. The R code is provided in the Appendix.

Where PBPSs had been done but raw data were not available, CIs were constructed by applying an approximation to the binomial distribution  $CI = p - (z \times s.e.)$  to  $p + (z \times s.e.)$ , where  $p$  is proportion of positive cases in the sample,  $s.e.$  is  $\sqrt{p(1-p)/n}$ , and  $z$  is the standard normal distribution (95% CI = 1.96). Where the sample size was not available, a sample of 1000 was assumed.

Standardized prevalence estimates and the lower and upper 95% CI bounds were multiplied by the 15-years-and-older population for the relevant district (projected to 2016) to provide an estimate of the number of persons with trichiasis.



For expert backlog assessments, confidence intervals were constructed using the relative precisions of all datasets for which bootstrapped CIs could be determined. The range of relative upper bound and lower bound precisions were trimmed to exclude 10% of the greatest and least values. The means of the remainder were applied.

Upper (and lower) CI bounds around district-level estimates, regardless of the method used for calculation, were summed to generate country-level, region-level and global estimates of the upper (and lower) CI bounds.

## **Role of the funding source and ethical review**

The funder (WHO) of this project was involved in all aspects of the work. Funders of the surveys that contributed data to the re-estimates were not involved in its design, in data analysis, data interpretation, manuscript preparation, or decisions on where, how or when to publish in the peer reviewed press. The study was approved by the Research Ethics Committee of the London School of Hygiene & Tropical Medicine (11208). The corresponding author had full access to all the data and had final responsibility for the decision to submit for publication.

## **Results**

Data from 1,355 districts in 31 countries were age- and sex- standardized and contribute an estimated 662 thousand cases (95% confidence interval (CI) 324 thousand-1.1 million) to the global total. Adjusting prevalence estimates by age and sex reduced raw district-level estimates by a mean factor of 0.45 (range 0.03–2.28). In countries where it was possible to adjust the datasets by age and sex, the country level backlog reduced by a mean factor of 0.56 (range 0.22–0.94) (Table 1). Estimates from PBPSs in 398 districts could not be standardized by age and sex because the original datasets were unavailable; the unadjusted backlogs in these districts totalled 781 thousand cases. We adjusted these estimates by multiplying them across the board by 0.45, yielding 411 thousand cases (95% CI 283-557 thousand). Based on new expert assessment of available data, it was determined that eight countries no longer have a trichiasis backlog. For six countries, local expert assessment generated revised, non-zero estimates. For one country, other published estimates[17] were used. Finally, there were seven countries for which the 2009 estimates were retained. Data included in the overall estimate had been collected from 2000–2016 (Figure 1).




Figure 1 Number of districts contributing data to the overall estimate, by year of survey

**Table 1 Estimated national-level trichiasis backlogs, 2016, with comparisons to the corresponding estimates for 2009**

WHO region	Country	2009 estimate[8] (data generally from 2007)	Source for 2016 estimate (1)	2016 estimate [year of data collection] (2) (95% confidence intervals)	Representativeness
African Region	Algeria	86,700	Retained previous estimate	86,700 (23,022-175,625)	
	Angola	no data	Expert assessment (3)	0	
	Benin	7,600	GTMP[18]	11,782 [2014-2015] (5,148-20,861)	Estimate based on 12 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	Botswana	32,900	Expert assessment (3)	0	
	Burkina Faso	32,800	PBPS – no raw data available	19,443 [2005-2015] (5,381-35,236)	Estimate based on 63 PBPS datasets covering all districts in the country
	Burundi	no data	Expert assessment (3)	0	
	Cameroon	47,200	PBPS – no raw data available	698 [2012-2014] (27-1,523)	Estimate based on 1) 35 age- and sex-adjusted PBPS datasets, plus 2) 15 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
			PBPS – raw data available	10,743 [2010-2014] (5,705-16,525)	
	Central African Republic	1,000	PBPS – no raw data available	6,539 [2011] (1,991-11,259)	Estimate based on 8 PBPS datasets; security concerns prevented the remaining 7 districts in the country where evidence indicates trachoma may be a public health problem from being surveyed
	Chad	34,300	GTMP	24,597 [2014-2015] (8,513-44,924)	Estimate based on 1) 43 age- and sex-adjusted PBPS datasets, plus 2) 14 PBPS datasets. At the time of publication 1 district where evidence indicates trachoma may be a public health problem remained to be surveyed
			PBPS – no raw data available	23,024 [2002] (15,365-30,683)	
	Congo		GTMP	0 [2015]	Estimate based on 1 age- and sex-adjusted PBPS dataset covering all districts in the country where evidence indicated that trachoma might have been a public health problem
	Côte d'Ivoire	59,900	GTMP	1,216 [2015] (176-2,582)	Estimate based on 10 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	Democratic Republic of the Congo	no data	GTMP[19]	33,333 [2014-2016] (18,369-52,870)	Estimate based on 30 age- and sex-adjusted PBPS datasets; security concerns prevented 1 district from being surveyed. At the time of publication an additional 9 districts where evidence indicates trachoma may be a public health problem remained to be surveyed.
Eritrea	42,000	GTMP	774 [2014] (274-1,308)	Estimate based on 1) 2 age- and sex-adjusted PBPS datasets, plus 2) 36 PBPS datasets; security concerns prevented the remaining 1 district in the country where evidence indicates trachoma may be a public health problem from being surveyed	
		PBPS – no raw data available	24,793 [2006-2014] (15,005-34,823)		

Ethiopia	1,272,600	Expert assessment[20-24] (4)	693,037 [2012-2016] (184,025-1,403,858)	Expert assessment based on 1) a backlog calculated from 196 age- and sex-adjusted PBPS datasets; security concerns prevented the remaining 16 districts in the country where evidence indicates trachoma may be a public health problem from being surveyed; minus 2) programmatic TT surgery output from 2012-2016
Gambia	10,500	PBPS – no raw data available	0 [2016]	PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
Ghana	3,000	PBPS – no raw data available	1,379 [2007-2008] (188-2,875)	Estimate based on 23 PBPS datasets, covering all districts in the country where evidence indicated that trachoma may have been a public health problem
Guinea	25,100	GTMP	5,523 [2014-2016] (1,141-11,834)	Estimate based on 31 age- and sex-adjusted PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
		PBPS – raw data available	24,302 [2011-2013] (13,035-38,193)	
Guinea-Bissau	16,400	PBPS – raw data available	21,255 [2005] (12,975-29,638)	Estimate based on 11 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
Kenya	306,800	PBPS – no raw data available	30,195 [2004-2012] (21,404-39,116)	Estimate based on 1) 3 age- and sex-adjusted PBPS datasets, plus 2) 12 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
		PBPS – raw data available	21,363 [2004] (13,583-31,471)	
Malawi	33,400	GTMP[25, 26]	13,446 [2013-2015] (3,302-27,937)	Estimate based on 1) 30 age- and sex-adjusted PBPS datasets, plus 2) 3 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
		PBPS – no raw data available	1,128 [2012] (227-2,124)	
		PBPS – raw data available	817 [2014] (48-1,918)	
Mali	67,600	Expert assessment (5)	13,852 (3,678-28,059)	Estimate based on 55 PBPS datasets, covering all districts in the country
Mauritania	2,500	PBPS – no raw data available	1,556 [2004-2013] (250-3,325)	Estimate based on 29 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
Mozambique	60,500	GTMP[27]	18,817 [2013-2014] (3,958-39,883)	Estimate based on 1) 99 age- and sex-adjusted PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
		PBPS – raw data available	96 [2015] (89-111)	
Namibia	6,100	Expert assessment (3)	0	
Niger	59,600	Expert assessment (8)	40,592 (9,712-83,784)	Estimate based on 49 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
Nigeria	627,300	GTMP[28-32]	193,951 [2013-2014] (111,251-297,277)	Estimate based on 1) 294 age- and sex-adjusted PBPS datasets, plus 2) 103 PBPS datasets, security concerns prevented the remaining 47 districts in the country where evidence indicates trachoma may be a public health problem from being surveyed
		PBPS – no raw data available	90,865 [2003-2014] (56,635-126,704)	
		PBPS – raw data available	3,800 [2014-2015] (1,801-6,295)	

Eastern Mediterranean Region	Senegal	129,800	GTMP	12,707 [2014] (6,823-19,364)	Estimate based on 1) 20 age- and sex-adjusted PBPS datasets, plus 2) 27 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
			PBPS – no raw data available	30,905 [2000-2012] (18,450-43,437)	
			PBPS – raw data available	3,077 [2015] (1,502-4,695)	
	South Sudan	no data	Expert assessment	81,093 (21,533-164,267)	Estimate provided by the Ministry of Health, South Sudan. To date, only a quarter of South Sudan has been mapped – this figure is likely to be an under-estimate for the country
	Togo	2,900	PBPS – no raw data available	318 [2009] (45-680)	Estimate based on 3 PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	Uganda	610,600	GTMP	680 [2014] (253-1,200)	Estimate based on 1) 39 age- and sex-adjusted PBPS datasets, plus 2) 5 PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
			PBPS – no raw data available	14,559 [2008-2012] (9,635-19,483)	
			PBPS – raw data available	65,329 [2013-2016] (39,924-149,042)	
	United Republic of Tanzania	214,800	GTMP[33, 34]	8,096 [2014] (2,491-15,225)	Estimate based on 97 age- and sex-adjusted PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
			PBPS – raw data available	63,157 [2004-2016] (28,844-87,028)	
	Zambia	8,500	GTMP	1,524 [2015] (712-2,670)	Estimate based on 1) 7 age- and sex- adjusted PBPS datasets, plus 2) 3 PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
			PBPS – no raw data available	2,168 [2010-2015] (1,289-3,047)	
	Zimbabwe	44,100	GTMP	6,765 [2014-2015] (3,133-10,988)	Estimate based on 16 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	Afghanistan	83,100	Retained previous estimate	83,100 (22,066-168,332)	
Djibouti	3,900	Expert assessment (6)	75 (20-152)		
Egypt	35,400	GTMP	35,362 [2015] (16,003-58,414)	Estimate based on 1) 4 age- and sex-adjusted PBPS datasets, plus 2) 1 PBPS dataset	
		PBPS – no raw data available	35,400 [2015] (30,596-40,204)		
Iran (Islamic Republic of)	49,300	Expert assessment (3)	0		
Iraq	43,900	Expert assessment (3)	0		
Libya	13,200	Expert assessment (7)	33,400 (8,869-67,657)		
Morocco	6,400	PBPS – no raw data available	0 [2015]	Estimate based on PBPS datasets covering all districts in the country where evidence indicated that trachoma may have been a public health problem	
Oman	600	PBPS – no raw data available	600 [2005]	Estimate based on PBPS datasets covering all districts in the	

				(41-619)	country where evidence indicated that trachoma may have been a public health problem
	Pakistan	71,700	GTMP	5,330 [2015] (1,160-10,732)	Estimate based on 1) 42 age- and sex-adjusted PBPS datasets, plus 2) 31 PBPS datasets;; security concerns prevented 15 districts from being surveyed. At the time of publication an additional 1 district where evidence indicates trachoma may be a public health problem remained to be surveyed.
			PBPS – no raw data available	23,420 [2012] (12,205-34,975)	
	Somalia	10,300	Retained previous estimate	10,300 (2,735-20,864)	
	Sudan	528,100	GTMP[35]	22,508 [2014-2015] (14,208-33,449)	Estimate based on 1) 109 age- and sex-adjusted PBPS datasets, plus 2) 22 PBPS datasets;; security concerns prevented 12 districts from being surveyed. At the time of publication an additional 1 district where evidence indicates trachoma may be a public health problem remained to be surveyed.
			PBPS – no raw data available	6,751 [2006-2013] (2,599-10,903)	
			PBPS – raw data available	36,982 [2007-2010] (7,070-80,470)	
Yemen	270,800	GTMP	5,821 [2013-2015] (896-13,165)	Estimate based on 1) 42 age- and sex-adjusted PBPS datasets;; security concerns prevented the remaining 38 districts in the country where evidence indicates trachoma may be a public health problem from being surveyed	
Region of the Americas	Brazil	58,000	Retained previous estimate	58,000 [2003-2006] (15,401-117,488)	Estimate based on PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	Colombia	no data	GTMP	48 [2015] (0-123)	Estimate based on 1) 3 age- and sex-adjusted PBPS datasets, plus 2) 6 PBPS datasets. At the time of publication an additional 3 districts where evidence indicates trachoma may be a public health problem remained to be surveyed.
			PBPS – no raw data available	23 [2003-2009] (4-44)	
	Guatemala	30	PBPS – no raw data available	543 [2011] (268-819)	Estimate based on 2 PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
Mexico	20	Expert assessment (3)	0		
South-East Asia Region	India	443,000	Retained previous estimate	443,000 (117,632-897,368)	
	Nepal	138,800	PBPS – no raw data available	25,240 [2001-2012] (16,889-33,631)	Estimate based on 1) 30 age- and sex-adjusted PBPS datasets, plus 2) 17 PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
PBPS – raw data available			3,252 [2002-2009] (462-7,323)		
Western Pacific Region	Australia	1,100	Retained previous estimate	1,100 (292,2,228)	
	Cambodia	29,200	GTMP[36]	4,999 [2014-2015] (729-10,676)	Estimate based on 14 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
	China	2,330,600	PBPS – no raw data available	71,280 (70,399-72,161)	Estimate based on PBPS datasets covering all districts in the country where evidence indicated that trachoma may have been a public health problem before 2012
	Fiji	800	PBPS using the methods of the GTMP[37] and subsequent	0 [2013]	Estimate based on 1 age- and sex-adjusted PBPS dataset covering all districts in the country where evidence indicates trachoma

		investigation[38]		may be a public health problem
Kiribati	100	GTMP	69 [2015] (30-108)	Estimate based on 1 age- and sex-adjusted PBPS dataset covering all districts in the country where evidence indicates trachoma may be a public health problem
Lao People's Democratic Republic	900	GTMP[39]	630 [2013-2015] (0-1,728)	Estimate based on 16 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
Myanmar	65,800	Retained previous estimate	65,800 (17,472-133,289)	
Nauru	0	Expert assessment (3)	0	
Papua New Guinea	5,800	GTMP[40]	156 [2015] (0-362)	Estimate based on 7 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
Solomon Islands	500	GTMP[41]	44 [2013] (4-104)	Estimate based on 1) 3 age- and sex-adjusted PBPS datasets, plus 2) 5 PBPS datasets, covering all districts in the country where evidence indicates trachoma may be a public health problem
		PBPS – no raw data available	8 [2012] (0-23)	
Vanuatu	200	GTMP[42]	48 [2014] (47-50)	Estimate based on 1 age- and sex-adjusted PBPS datasets covering all districts in the country where evidence indicates trachoma may be a public health problem
Viet Nam	210,000	Expert assessment (6)	100,000[17] (26,553-202,566)	

1) Unless otherwise specified, "Retained previous estimate" refers to the 2009 estimate by Mariotti et al[8]

2) Individuals examined in these surveys were men and women aged  $\geq 15$  years

3) Health ministry reports that there is no evidence of trichiasis being a public health problem, or that evidence indicates that trichiasis is not a public health problem

4) Estimate provided by the Federal Ministry of Health of Ethiopia; determined by calculating the backlog indicated by the most recent population-based prevalence survey in each trachoma-endemic district and subtracting from it the number of individuals with trichiasis managed by the health system since those surveys

5) Estimate provided by the Ministère de la Santé, Mali; determined by calculating the backlog indicated by the most recent population-based prevalence survey in each trachoma-endemic district and subtracting from it the number of individuals with trichiasis managed by the health system since those surveys

6) derived from information provided at the 2010 WHO Alliance for GET2020 meeting

7) Estimate informed by data from a Rapid Assessment of Avoidable Blindness (RAAB) survey

Through the inclusion of GTMP data, adjusting older datasets by age and sex, and obtaining current local expert assessment of available data, the global backlog estimate reduces from 8.2 million[8] (or 7.3 million[10]) to 2.8 million (95% CI 1.1-5.2 million) (Table 2).

In our estimates, data collected in surveys set up prior to August 2014 include all trichiasis, irrespective of the presence or absence of trachomatous conjunctival scarring. Data collected in surveys set up after August 2014 contributed data on trichiasis in eyes that also demonstrated trachomatous conjunctival scarring (or had an eyelid that could not be everted, due to presumed dense scar), thereby including only those cases of trichiasis attributable to trachoma. In both scenarios the data represent both “managed” and “unmanaged” trichiasis, irrespective of whether individuals have previously been offered corrective surgery or epilation[43].

**Table 2 Estimated region-level trichiasis backlogs, 2016, with comparisons to the corresponding estimates for 2009**

WHO Region	2009 Estimate	2016 Estimate				Ratio (2016 estimate / 2009 estimate)
		Retained previous estimate or expert assessment (95% CI)	Unstandardized (reduced by a factor of 0.45) (95% CI)	Standardized (95% CI)	Total (95% CI)	
African Region	3,846,500	915,274 (241,969-1,855,593)	247,570 (150,310-363,186)	547,151 (283,052-913,841)	1,709,995 (675,331-3,132,620)	0.45
Region of the Americas	58,050	58,000 (15,401-117,488)	566 (272-863)	48 (0-123)	58,614 (15,673-118,475)	1.01
Eastern Mediterranean Region	660,000	126,875 (33,690-257,006)	66,171 (45,441-86,702)	106,003 (39,337-196,230)	299,049 (118,467-539,937)	0.52
South-East Asia Region	485,000	443,000 (117,632-897,368)	25,240 (16,889-33,631)	3,252 (462-7,323)	471,492 (134,982-938,323)	0.97
Western Pacific Region	2,630,000	166,900 (44,318-338,083)	71,288 (70,399-72,184)	5,947 (810-13,027)	244,135 (115,527-423,294)	0.09
Total	8,248,050	1,710,049 (453,010-3,465,538)	410,835 (283,311-556,567)	662,403 (323,661-1,130,543)	2,783,285 (1,059,980-5,152,649)	0.35

## Discussion

Trichiasis remains a significant public health problem in many countries, with a global backlog estimated for 2016 at 2.8 million people, 61% of whom live in sub-Saharan Africa. While there were methodological challenges in generating the 2009 estimate, it was the best estimate that could be made with the information then available. The



considerable reduction from that estimate to the one generated here is likely to be the result of a combination of factors. First, there are now more and better data derived from rigorous surveys. Second, in many countries, there has been impressive programmatic scale up to manage TT, conducted by complex networks of governments and their partners. Third, there is likely to be an effect on the incidence of TT from the intensive efforts to reduce active trachoma prevalence in many contexts; such efforts have been ramping up in endemic countries since the World Health Assembly's 1998 commitment to global elimination of trachoma[44]. Teasing out the relative contribution of each of these factors is not possible at the present time, but regardless of cause, the reduction is welcome news for global health.

A better understanding of the backlog and distribution of trichiasis cases is necessary to effectively plan for surgical services and other components of management of individuals with trichiasis. To reduce TT prevalence in each district of each endemic country to  $<0.2\%$  in adults, which is the defined elimination threshold for TT[45], at least 2.0 million people will need to have their TT appropriately managed. From available data, it is estimated that 56% of people with trichiasis have bilateral trichiasis.

While our calculations lessen the uncertainty around the global backlog estimate, important limitations remain. Expert assessment of available data was used for 13 endemic countries, in seven of which the estimate was zero cases. Ethiopia, Mali, and Niger have undergone intensive surgical scale-up in the time since the most recent round of prevalence surveys. Because of this programmatic output, and a lack of consensus around how to counterbalance backlog reduction with new incident cases and post-surgical recurrence (both of which are inescapable, but presently difficult to accurately quantify) these countries provided results based on expert assessment. Uncertainty is greatest among the eight countries in which previous estimates were retained; these countries accounted for 848 thousand cases of trichiasis (30% of the global estimate) and further investigation is needed here. India, which accounts for 443,000 cases of trichiasis (52% of the total retained estimate) is a particular priority. A second ongoing uncertainty stems from the unavailability of raw data for 398 districts, for which, as a result, age- and sex-standardization was not possible. As noted in our calculations, there is a mean prevalence reduction against the raw prevalence estimate of  $\times 0.45$ , which was applied to previously-unstandardized district-level estimates. Third, our CIs reflect statistical uncertainty. Because statistical uncertainty is uncorrelated across countries, our method of aggregating CIs across

administrative boundaries overestimates statistical uncertainty at regional and global levels. However, there are other sources of uncertainty in addition to statistical uncertainty, so it is not possible to say whether our regional and global CIs are biased.

Ongoing lack of trichiasis (and TF) prevalence data for 235 suspected-endemic districts is due to local insecurity, and the global trachoma community stands ready to support national governments to undertake the needed mapping in those populations, when and if security conditions improve to allow safe conduct of fieldwork.

It is recognized[11] that the PBPS methodology described earlier is potentially imprecise in estimating TT against the WHO elimination target of  $\leq 0.2\%$  in adults. Thus, at district level, the uncertainty around the estimates of the TT backlog can be large. This is reflected in our CIs. Work is underway to try to develop more reliable methodologies for estimating the prevalence of trichiasis.

At present, the diagnosis of TT is based on the presence of clinical trichiasis (one or more lashes touching the globe or evidence of recent epilation of in-turned eyelashes) plus residence in a (presumed) trachoma-endemic setting. This definition is by nature somewhat circular and may need review, as it inevitably leads to classification of trichiasis as TT regardless of whether trachomatous conjunctival scarring is present or not. Trichiasis without trachomatous conjunctival scarring may be age-related or due to trauma, distichiasis, epiblepharon or other inflammatory disease[46]; the sight-threatening potential and optimal management strategies for non-trachomatous trichiasis still require further investigation[47].

Assumptions were made when adjusting the data for age and sex. First, we assumed that ages reported in surveys were accurate. However, it is reasonable to hypothesise that during a survey, individuals demonstrate a terminal digit preference when stating their age. Second, this analysis assumed that UNdata population pyramids used were representative of all districts for which survey data were available. Finally, because many studies have demonstrated a high correlation between trichiasis and increased age[48-51], we assumed zero trichiasis cases in the population aged 14 years and younger.

These calculations of national and global TT burdens are point-prevalence estimates based on data of differing vintages; they will change as additional surveys, baseline or impact, are undertaken. Other than for Ethiopia, Mali and Niger, the estimates do not take into account the number of surgeries done during the period between the most recent survey and this analysis. Nevertheless, estimates of current national TT backlog are essential for countries to appropriately allocate resources for surgical campaigns. Future district-level impact surveys in countries with trachoma elimination programmes will provide progressively improved evidence for countries to assess their residual TT burden and, ultimately, validate elimination of trichiasis as a public health problem. Baseline surveys of trachoma are still needed in a handful of countries (e.g., Egypt, Somalia and Central African Republic); these will lead to revisions in national estimates. In some settings, there will be a need to undertake trichiasis-only surveys in order to assess progress to elimination. Unfortunately, insecurity may continue to limit surveys in a few, but not all, of the settings in which previous estimates were retained. Where possible, surveys should be undertaken. Such data will help us drive towards, and demonstrate success in, the global elimination of trachoma as a public health problem.

## References

1. World Health Organization. WHO Alliance for the Global Elimination of Blinding Trachoma by the year 2020: progress report on elimination of trachoma, 2013. *Wkly Epidemiol Rec.* 2014;89(39):421-8.
2. Thylefors B, Dawson CR, Jones BR, West SK, Taylor HR. A simple system for the assessment of trachoma and its complications. *Bulletin of the World Health Organization.* 1987;65(4):477-83. PubMed PMID: 3500800; PubMed Central PMCID: PMC2491032.
3. Palmer SL, Winskell K, Patterson AE, Boubacar K, Ibrahim F, Namata I, et al. 'A living death': a qualitative assessment of quality of life among women with trichiasis in rural Niger. *International health.* 2014;6(4):291-7. doi: 10.1093/inthealth/ihu054. PubMed PMID: 25125577.
4. Mabey DC, Solomon AW, Foster A. Trachoma. *Lancet.* 2003;362(9379):223-9. doi: 10.1016/S0140-6736(03)13914-1. PubMed PMID: 12885486.
5. Solomon AW ZM, Kuper H, et al. *Trachoma control: a guide for program managers.* Geneva: World Health Organization, 2006.
6. Solomon AW, Peeling RW, Foster A, Mabey DC. Diagnosis and assessment of trachoma. *Clinical microbiology reviews.* 2004;17(4):982-1011. doi: 10.1128/CMR.17.4.982-1011.2004. PubMed PMID: 15489358; PubMed Central PMCID: PMC523557.
7. Merbs SL, Resnikoff S, Kello AB, Mariotti S, Greene G, West SK. *Trichiasis surgery for trachoma.* Geneva, Switzerland: The World Health Organization, 2015.
8. Mariotti SP, Pascolini D, Rose-Nussbaumer J. Trachoma: global magnitude of a preventable cause of blindness. *The British journal of ophthalmology.* 2009;93(5):563-8. doi: 10.1136/bjo.2008.148494. PubMed PMID: 19098034.
9. WHO Alliance for the Global Elimination of Blinding Trachoma by 2020. Report of the 11th meeting of the WHO Alliance for the Global Elimination of Blinding Trachoma. Geneva, Switzerland: 2007 2-4 April 2007. Report No.
10. World Health Organization. Global WHO Alliance for the Elimination of Blinding Trachoma by 2020. *Wkly Epidemiol Rec.* 2012;87(17):161-8.
11. Solomon AW, Pavluck AL, Courtright P, Aboe A, Adamu L, Alemayehu W, et al. The Global Trachoma Mapping Project: Methodology of a 34-Country Population-Based Study. *Ophthalmic epidemiology.* 2015;22(3):214-25. doi: 10.3109/09286586.2015.1037401. PubMed PMID: 26158580.
12. Solomon AW, Abdala M, Alemayehu W. et, al. Completing the Global Trachoma Map. Submitted.
13. UNdata: UNSD Demographic Statistics. 0.14.6 ed. UN population division.
14. Worldpop: High resolution age-structured populatin distribution maps [Internet]. GeoData Institute, University of Southampton. [cited 12 January 2016]. Available from: [www.worldpop.org](http://www.worldpop.org).
15. ArcGIS Desktop. 10.3 ed. Redlands, CA: Environmental Systems Research Institute (ESRI); 2014.
16. R Development Core Team. *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing; 2016.
17. *The end in sight: 2020 INSight.* International Coalition for Trachoma Control, 2011.
18. Bio AA, Boko PM, Dossou YA, et al. Prevalence of trachoma in northern Benin: results from 11 population-based prevalence surveys covering 26 districts. *Ophthalmic epidemiology.* 2017;in press.
19. Kilangalanga J, Ndjemba JM, Uvon PA, et al. Trachoma in the Democratic Republic of the Congo: results of 46 baseline prevalence surveys conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology.* 2017;(in press).
20. Abashawl A, Macleod C, Riang J, Mossisa F, Dejene M, Willis R, et al. Prevalence of Trachoma in Gambella Region, Ethiopia: Results of Three Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology.* 2016;23(sup1):77-83. doi: 10.1080/09286586.2016.1247875. PubMed PMID: 27918222.

21. Adamu Y, Macleod C, Adamu L, Fikru W, Kidu B, Abashawl A, et al. Prevalence of Trachoma in Benishangul Gumuz Region, Ethiopia: Results of Seven Population-Based Surveys from the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):70-6. doi: 10.1080/09286586.2016.1247877. PubMed PMID: 27918248.
22. Adera TH, Macleod C, Endriyas M, Dejene M, Willis R, Chu BK, et al. Prevalence of and Risk Factors for Trachoma in Southern Nations, Nationalities, and Peoples' Region, Ethiopia: Results of 40 Population-Based Prevalence Surveys Carried Out with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):84-93. doi: 10.1080/09286586.2016.1247876. PubMed PMID: 27918229.
23. Bero B, Macleod C, Alemayehu W, Gadisa S, Abajobir A, Adamu Y, et al. Prevalence of and Risk Factors for Trachoma in Oromia Regional State of Ethiopia: Results of 79 Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(6):392-405. doi: 10.1080/09286586.2016.1243717. PubMed PMID: 27820657.
24. Sherief ST, Macleod C, Gigar G, Godefay H, Abraha A, Dejene M, et al. The Prevalence of Trachoma in Tigray Region, Northern Ethiopia: Results of 11 Population-Based Prevalence Surveys Completed as Part of the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):94-9. doi: 10.1080/09286586.2016.1250917. PubMed PMID: 27918232.
25. Kalua K, Chisambi A, Chinyanya D, Kamwendo Z, Masika M, Willis R, et al. Completion of Baseline Trachoma Mapping in Malawi: Results of Eight Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):32-8. doi: 10.1080/09286586.2016.1230224. PubMed PMID: 27726469.
26. Kalua K, Phiri M, Kumwenda I, Masika M, Pavluck AL, Willis R, et al. Baseline Trachoma Mapping in Malawi with the Global Trachoma Mapping Project (GTMP). *Ophthalmic epidemiology*. 2015;22(3):176-83. doi: 10.3109/09286586.2015.1035793. PubMed PMID: 26158575; PubMed Central PMCID: PMC4673584.
27. Abdala M, Singano CC, Willis R, et al. The epidemiology of trachoma in Mozambique: results of 96 population-based prevalence surveys. *Ophthalmic epidemiology*. in press.
28. Muhammad N, Mpyet C, Adamu MD, William A, Umar MM, Goyol M, et al. Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys. *Ophthalmic epidemiology*. 2016;23(sup1):46-54. doi: 10.1080/09286586.2016.1250918. PubMed PMID: 27918227.
29. Adamu MD, Mpyet C, Muhammad N, Umar MM, Muazu H, Olamiju F, et al. Prevalence of Trachoma in Niger State, North Central Nigeria: Results of 25 Population-Based Prevalence Surveys Carried Out with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):63-9. doi: 10.1080/09286586.2016.1242757. PubMed PMID: 27918223.
30. Mpyet C, Muhammad N, Adamu MD, Muazu H, Umar MM, Abdull M, et al. Prevalence of Trachoma in Bauchi State, Nigeria: Results of 20 Local Government Area-Level Surveys. *Ophthalmic epidemiology*. 2016;23(sup1):39-45. doi: 10.1080/09286586.2016.1238945. PubMed PMID: 27846369.
31. Mpyet C, Muhammad N, Adamu MD, Muazu H, Umar MM, Goyol M, et al. Prevalence of Trachoma in Katsina State, Nigeria: Results of 34 District-Level Surveys. *Ophthalmic epidemiology*. 2016;23(sup1):55-62. doi: 10.1080/09286586.2016.1236975. PubMed PMID: 27775463.
32. Mpyet C, Muhammad N, Adamu MD, Muazu H, Umar MM, Alada J, et al. Trachoma Mapping in Gombe State, Nigeria: Results of 11 Local Government Area Surveys. *Ophthalmic epidemiology*. 2016;23(6):406-11. doi: 10.1080/09286586.2016.1230633. PubMed PMID: 27726459.
33. Mwingira UJ, Kabona G, Kamugisha M, Kirumbi E, Kilembe B, Simon A, et al. Progress of Trachoma Mapping in Mainland Tanzania: Results of Baseline Surveys from 2012 to 2014. *Ophthalmic epidemiology*. 2016;23(6):373-80. doi: 10.1080/09286586.2016.1236974. PubMed PMID: 27775455; PubMed Central PMCID: PMC4673584.

34. Omar FJ, Kabona G, Abdalla KM, Mohamed SJ, Ali SM, Ame SM, et al. Baseline Trachoma Surveys in Kaskazini A and Micheweni Districts of Zanzibar: Results of Two Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(6):412-7. doi: 10.1080/09286586.2016.1235206. PubMed PMID: 27775454; PubMed Central PMCID: PMC5116987.
35. Elshafie BE, Osman KH, Macleod C, Hassan A, Bush S, Dejene M, et al. The Epidemiology of Trachoma in Darfur States and Khartoum State, Sudan: Results of 32 Population-Based Prevalence Surveys. *Ophthalmic Epidemiol*. 2016;23(6):381-91. doi: 10.1080/09286586.2016.1243718. PubMed PMID: 27841721.
36. Meng N, Seiha D, Thorn P, Willis R, Flueckiger RM, Dejene M, et al. Assessment of Trachoma in Cambodia: Trachoma Is Not a Public Health Problem. *Ophthalmic epidemiology*. 2016;23(sup1):3-7. doi: 10.1080/09286586.2016.1230223. PubMed PMID: 27726472.
37. Macleod CK, Butcher R, Mudaliar U, Natutusau K, Pavluck AL, Willis R, et al. Low Prevalence of Ocular Chlamydia trachomatis Infection and Active Trachoma in the Western Division of Fiji. *PLoS neglected tropical diseases*. 2016;10(7):e0004798. doi: 10.1371/journal.pntd.0004798. PubMed PMID: 27404379; PubMed Central PMCID: PMC4942140.
38. Macleod C, Yalen C, Butcher R, Mudaliar U, Natutusau K, Rainima-Qaniuci M, et al. Eyelash Epilation in the Absence of Trichiasis: Results of a Population-Based Prevalence Survey in the Western Division of Fiji. *PLoS neglected tropical diseases*. 2017;11(1):e0005277. doi: 10.1371/journal.pntd.0005277. PubMed PMID: 28114364; PubMed Central PMCID: PMC5256864.
39. Southisombath K, Sisalermsak S, Chansan P, Akkhavong K, Phommala S, Lewallen S, et al. National Trachoma Assessment in the Lao People's Democratic Republic in 2013-2014. *Ophthalmic epidemiology*. 2016;23(sup1):8-14. doi: 10.1080/09286586.2016.1236973. PubMed PMID: 27846362.
40. Ko R, Macleod C, Pahau D, Sokana O, Keys D, Burnett A, et al. Population-Based Trachoma Mapping in Six Evaluation Units of Papua New Guinea. *Ophthalmic epidemiology*. 2016;23(sup1):22-31. doi: 10.1080/09286586.2016.1235715. PubMed PMID: 27893297.
41. Sokana O, Macleod C, Jack K, Butcher R, Marks M, Willis R, et al. Mapping Trachoma in the Solomon Islands: Results of Three Baseline Population-Based Prevalence Surveys Conducted with the Global Trachoma Mapping Project. *Ophthalmic epidemiology*. 2016;23(sup1):15-21. doi: 10.1080/09286586.2016.1238946. PubMed PMID: 27937043.
42. Taleo F, Macleod CK, Marks M, Sokana O, Last A, Willis R, et al. Integrated Mapping of Yaws and Trachoma in the Five Northern-Most Provinces of Vanuatu. *PLoS neglected tropical diseases*. 2017;11(1):e0005267. doi: 10.1371/journal.pntd.0005267. PubMed PMID: 28118354.
43. World Health Organization. Report of the 3rd global scientific meeting on trachoma. Johns Hopkins University, Baltimore, MA, 19-20 July 2010: Geneva: World Health Organization; 2010.
44. Global elimination of blinding trachoma. 51st World Health Assembly; May 16, 1998; Geneva: World Health Assembly; 1998.
45. World Health Organization. Validation of elimination of trachoma as a public health problem. WHO/HTM/NTD/2016.8. Geneva: World Health Organization, 2016.
46. Rajak SN, Habtamu E, Weiss HA, Bedri A, Gebre T, Bailey RL, et al. The clinical phenotype of trachomatous trichiasis in Ethiopia: not all trichiasis is due to entropion. *Investigative ophthalmology & visual science*. 2011;52(11):7974-80. doi: 10.1167/iops.11-7880. PubMed PMID: 21896855; PubMed Central PMCID: PMC3219424.
47. World Health Organization Alliance for the Global Elimination of Trachoma by 2020. Second Global Scientific Meeting on Trachomatous Trichiasis; Cape Town, 4-6 November 2015 (WHO/HTM/NTD/2016.5): Geneva: World Health Organization; 2016.
48. Ngondi J, Gebre T, Shargie EB, Graves PM, Ejigsemahu Y, Teferi T, et al. Risk factors for active trachoma in children and trichiasis in adults: a household survey in Amhara Regional State, Ethiopia.

Transactions of the Royal Society of Tropical Medicine and Hygiene. 2008;102(5):432-8. doi: 10.1016/j.trstmh.2008.02.014. PubMed PMID: 18394663.

49. Burton MJ, Bowman RJ, Faal H, Aryee EA, Ikumapayi UN, Alexander ND, et al. The long-term natural history of trichomatous trichiasis in the Gambia. *Investigative ophthalmology & visual science*. 2006;47(3):847-52. doi: 10.1167/iovs.05-0714. PubMed PMID: 16505016.

50. Jones BR. The prevention of blindness from trachoma. *Transactions of the ophthalmological societies of the United Kingdom*. 1975;95(1):16-33. PubMed PMID: 775692.

51. Schemann JF, Laffly D, Sacko D, Zephak G, Malvy D. Trichiasis and geoclimatic factors in Mali. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2007;101(10):996-1003. doi: 10.1016/j.trstmh.2007.05.015. PubMed PMID: 17658570.