Multiobjective Strain Design: A Framework for Modular Cell Engineering

Sergio Garcia[1,2] and Cong T. Trinh[1,2*]

[1]Department of Chemical and Biomolecular Engineering, University of Tennessee, Knoxville,

TN

[2]Center for Bioenergy Innovation, Oak Ridge National Laboratory, Oak Ridge, TN

[*]Corresponding author: Tel: 865-974-2181. Email: ctrinh@utk.edu.

ABSTRACT

Diversity of cellular metabolism can be harnessed to produce a large space of molecules. However, development of optimal strains with high product titers, rates, and yields required for industrial production is laborious and expensive. To accelerate the strain engineering process, we have recently introduced a modular cell design concept that enables rapid generation of optimal production strains by systematically assembling a modular cell with an exchangeable production module(s) to produce target molecules efficiently. In this study, we formulated the modular cell design concept as a general multiobjective optimization problem with flexible design objectives derived from mass action. We developed algorithms and an associated software package, named ModCell2 to implement the design. We demonstrated that ModCell2 can systematically identify genetic modifications to design modular cells that can couple with a variety of production modules and exhibit a minimal tradeoff among modularity, performance, and robustness. Analysis of the modular cell designs revealed both intuitive and complex metabolic architectures enabling modular production of these molecules. We envision ModCell2 provides a powerful tool to guide modular cell engineering and sheds light on modular design principles of biological systems.

Keywords

INTRODUCTION

Engineering microbial cells to produce bulk and specialty chemicals from renewable and sustainable feedstocks is becoming a feasible alternative to traditional chemical methods that rely on petroleum feedstocks (1). However, only a handful of chemicals, out of the many possible molecules offered by nature, are industrially produced by microbial conversion, mainly because the current strain engineering process is laborious and expensive for profitable biochemical production (2). Thus, innovative technologies to enable rapid and economical strain engineering are needed to harness a large space of industrially-relevant molecules (3).

The modular organization of biological systems has been a source of inspiration for synthetic biology and metabolic engineering (4, 5). Modular pathway engineering breaks down target pathways into tractable pathway modules that can be finely tuned for optimal production of desirable chemicals (6, 7). Harnessing combinatorial pathways (e.g., fatty acid biosynthesis, reverse beta oxidation, polyketide or isoprenoid biosynthesis) is one excellent example of modular metabolic engineering. These pathways contain metabolic similarity (or combinatorial characteristics) such as a group of common specific enzymes capable of catalyzing linear reaction steps (8) and/or elongation cycles (9-11) and hence are capable of producing a large library of unique molecules (12). Since these molecules are derived from a common precursor metabolite(s), the optimal production strains often share common genotypes and phenotypes, and hence, the costly strain optimization process is only performed once for these molecules. Remarkably, this advantageous strain optimization strategy can be applied even for production of molecules derived from different precursors, using the concept of modular cell (ModCell) design (2, 13, 14).

With the arrival of steady-state, constraint-based stoichiometric models of cellular

metabolism, various computational algorithms have been developed to guide strain engineering

(15-17). These methods have featured the design of strains capable of growth-coupled product

synthesis (*GCP*), enabling adaptive laboratory evolution of these designed strains to enhance

product titers, rates, and yields (14, 18-20). Two approaches on growth-coupled production have

been formulated ─ one based on the coexistence of maximum growth and product synthesis rates

during the exponential growth phase (21) and the other based on the obligate requirement of

optimal product synthesis in any growth phase (22). The distinction between these two types of

growth coupling are also referred to weak coupling (*wGCP*) and strong coupling (*sGCP*) (16,

23).

Development of most strain design algorithms has been focused on overproduction of

only one target molecule. The first algorithm proposed for modular cell design compatible for

overproduction of multiple target molecules is MODCELL (13), which guided several

experimental studies (14, 24-27). It works by generating *sGCP* strain designs for each target

product based on elementary mode analysis (28), and then comparing the design strategies of

different products to identify common genetic modifications among them. A similar approach

was adapted in a subsequent work (29). For MODCELL to find optimal solutions for multiple

target products, it requires: 1) enumerating all possible designs above a predefined minimum

product yield and with minimal reaction deletion sets for each production network, which might

lead to a large number of solutions for each network and hence make the problem

computationally intractable, and 2) the resulting designs for all products must be compared to

identify common interventions, which is a computationally-hard, set-covering problem. Thus, the

current enumerative approach of MODCELL might become intractable very quickly, especially

for large-scale metabolic networks and potentially generate non-optimal designs, i.e., requiring

more knock-outs than necessary or including fewer products than possible.

In this study, we generalized the concept of modular cell design and addressed the

computational limitation of implementing it. We developed a novel computational platform

(ModCell2), based on multiobjective optimization and analysis of mass action of cellular

metabolism, to guide the design of modular cells for large-scale metabolic networks. We

demonstrated that ModCell2 can systematically identify genetic modifications to design modular

cells that can couple with a variety of production modules and exhibit a minimal tradeoff among

modularity, performance, and robustness. By analyzing these designs, we further revealed both

intuitive and complex metabolic architectures enabling modularity in modular cell and

production modules required for efficient biosynthesis of target molecules.

METHODS

Design principles of modular cell engineering

In the conventional strain engineering approach, a parent strain is genetically modified to yield

an optimal production strain to make only a target product. To produce each new molecule, the

design-build-test cycles of strain engineering must be repeated, which is laborious and expensive

(Figure 1). To minimize the cycles, modular cell engineering is formulated by genetically

transforming a parent strain into a modular (chassis) cell that must be assembled with

exchangeable modules to create optimal production strains (13). A modular cell is designed to

contain core metabolic pathways shared across designed optimal production strains.

Exchangeable modules are production pathways designed to synthesize desirable chemicals. A

combination of a modular cell and a production module(s) is required to balance redox, energy,

and precursor metabolites for sustaining cellular metabolism during growth and/or stationary

phases and exhibiting only desirable phenotypes. Practically, modular cell engineering can be

applied to monocultures and polycultures, where a production module(s) can be embedded in a

modular cell and activated by intracellular and/or extracellular cues such as light and/or signaling

molecules.

Multiobjective strain design framework for modular cell engineering

For modular cell engineering, we seek to design a chassis cell compatible with as many

production modules as possible to achieve only desirable production phenotypes while requiring

minimal genetic modifications. Since all production modules must leverage cellular resources of

the modular cell (e.g. precursor metabolites, cofactors, and energy), they form competing

objectives. Therefore, the framework of modular cell engineering can be formulated as a

multiobjective optimization problem, named ModCell2, as described below.

$\text{maximize}_{over\ y_j,\ z_{jk}}\ \ \boldsymbol{F} = \left(f_1, f_2, \dots, f_{|K|}\right)^T$ [1]

subject to

$f_k \in \arg max\{\sum_{j \in J_k} c_{jk} v_{jk}$

subject to

$\sum_{j \in J_k} S_{ijk} v_{jk} = 0, for\ all\ i \in I_k$ [2]

$l_{jk} \leq v_{jk} \leq u_{jk}, for\ all\ j \in J_k$ [3]

$l_{jk} d_{jk} \leq v_{jk} \leq u_{jk} d_{jk}, for\ all\ j \in C\ and\ C\ \square\ J_k$ [4]

$where\ d_{jk} = y_j\ V\ z_{jk}\}, for\ all\ k\ \square\ K$

$z_{jk} \leq (1 - y_j), for\ all\ j \in C, k \in K$ [5]

$\sum_{j \in C}(1 - y_j) \leq \alpha$ [6]

$\sum_{j \in C} z_{jk} \leq \beta_k, \text{for all } k \square K$ [7]

where i, j, and k are indices of metabolite i, reaction j, and production network k, respectively; f_k

is a design objective for network k; c_{jk} represents the cellular objective for reaction j in network

k associated with a design objective defined in eqns. 8-10; v_{jk} (mmol/g DCW/h) is metabolic flux

of reaction j bounded by l_{jk} and u_{jk} in network k, respectively; y_j and z_{jk} are binary design

variables for deletion reaction j and module reaction j in network k, respectively; α and β_k are

design parameters for deletion and module reactions, respectively; S_{ijk} is a stoichiometric

coefficient of metabolite in reaction j of network k; and C (eqn. 4) is the candidate reaction set

(Supplementary File S1). The goal of the optimization problem is to simultaneously maximize

all design objectives f_k.

Steady-state mass balance constraint of cellular metabolism. Quasi steady-state flux

balance of cellular metabolism (eqn. 2) is used as metabolic constraints for eqn. 1 (30, 31). A

model corresponding to each modular production strain (i.e. production network k) will be

derived from a parent strain (i.e. parent network) by adding necessary reactions (e.g., a

production module) to produce a target molecule. A feasible flux distribution for each production

network is described by mass balance (eqn. 2) and reaction flux bounds (eqns. 3 and 4). For a

given production network, the phenotypic space can be illustrated by the gray area that is

projected onto the two-dimensional space spanned by product synthesis and growth rates (Figure

2).

Design variables. In our formulation for modular cell engineering, we introduced two

design variables – binary reaction deletions (y_j) inherent to the modular cell and module-specific

reaction insertions (z_{jk}) (eqn. 4). These variables can be experimentally manipulated to constrain

the desirable phenotypes of production strains as shown in Figure 2. Specifically, $y_j = 0$ if

reaction j is deleted from the modular cell; otherwise, $y_j = 1$. Deleting metabolic reactions

removes undesired functional states of the network and leaves those with high design objectives.

Likewise, $z_{jk} = 1$ if reaction j is present in the production network k; otherwise, $z_{jk} = 0$. These

module reactions are endogenous reactions removed from the parent network (eqn. 5), but are

added back to a specific production module to enhance the compatibility of a modular cell. The

maximum number of reaction deletions (α) and module-specific reaction insertions (β_k) are user-

defined parameters.

Design objectives. To generalize ModCell2 design, we allow three different types of

design objectives (f_k, eqn. 1) that determine production phenotypes for each production network.

Depending on the application, a phenotype can be designed to be weak coupling (*wGCP*), strong

coupling (*sGCP*), and/or non-growth production (*NGP*) (Figure 2). The constrained phenotypic

spaces based on these design objectives are shown in color; any point within these spaces is a

feasible physiological state of the cell that can be represented by a metabolic flux distribution.

The *wGCP* design seeks to achieve a high product rate at maximum growth rate (Figure

2A). The *wGCP* design objective, f_k^{wGCP} ($\in [0, 1]$), is calculated as follows:

161
$$f_k^{wGCP} = \frac{v_{Pk}^{\mu}}{v_{Pmaxk}^{\mu}}$$
[8]

where v_{Pk}^{μ} is the minimum synthesis rate of the target product P at the maximum growth rate for

production network k and v_{Pmaxk}^{μ} is the maximum synthesis rate of P (Supplementary File S1).

This *wGCP* design formulation is equivalent to RobustKnock (32) or OptKnock with a tilted

objective function (21, 33, 34). In eqn. 8, f_k^{wGCP} is scaled from 0 to 1 for proper comparison

among products. The *wGCP* design is appropriate for applications where growth rate is not

limited by the nutrients, and the product is formed during the growth phase.

The *sGCP* design seeks to achieve a high product rate not only at optimal growth rate but

also during non-growth phase (Figure 2B). The *sGCP* design objective, f_k^{sGCP} ($\in$ [0, 1]), is

calculated as follows:

171
$$f_k^{sGCP} = \frac{v_{Pk}^{\mu}}{v_{P_{max}k}^{\mu}} \cdot \frac{v_{Pk}^{\bar{\mu}}}{v_{P_{max}k}^{\bar{\mu}}}$$
[9]

where $v_{Pk}^{\bar{\mu}}$ and $v_{P_{max}k}^{\bar{\mu}}$ are the minimum and maximum product formation rates for production

network k in the stationary phase, respectively (Supplementary File S1). The *sGCP* design

objective is comparable to the one implemented in MODCELL (13). Different from *wGCP*,

sGCP requires high product synthesis rate for any growth phase. However, the additional

constraint of optimal product synthesis during the stationary phase requires more genetic

manipulations or specific experimental conditions (e.g., anaerobic growth condition, supply of

intermediate metabolites). Both *wGCP* and *sGCP* designs enable fast growth selection to attain

the optimum product rates by adaptive laboratory evolution (35, 36).

The *NGP* design aims to maximize the minimum product rate during the non-growth

phase by eliminating carbon fluxes directed to biomass synthesis (Figure 2C). The *NGP* design

objective, f_k^{NGP} ($\in$ [0, 1]), is calculated as follows:

183
$$f_k^{NGP} = \frac{v_{Pk}^{\bar{\mu}}}{v_{P_{max}k}^{\bar{\mu}}}$$
[10]

While the *NGP* design is not suitable for growth selection, it can be derived from a *wGCP* (or

sGCP) design by imposing additional genetic modifications. Practically, *NGP* design strains can

be activated during cell culturing using a regulatory genetic circuit to toggle switch between

production phases.

Design solutions. Optimal solutions for eqn. 1 are a Pareto set (***PS***) that correspond to

design variables, including reaction deletions (y_j) and module reaction insertions (z_{jk}). Each

solution constitutes a design of a modular cell:

$$PS := \{\mathbf{x} \in \Omega: \nexists \mathbf{t} \in \Omega, \mathbf{F(t)} \prec \mathbf{F(x)}\} \tag{11}$$

Here, $\mathbf{F(t)} \prec \mathbf{F(x)}$ means $\mathbf{F(t)}$ *dominates* $\mathbf{F(x)}$ if and only if $f_i(\mathbf{t}) \geq f_i(\mathbf{x})$ for all i, and $\mathbf{F(t)}$

differs from $\mathbf{F(x)}$ in at least one entry. The feasible space of design variables, Ω, is defined by

the problem constraints (eqn. 2-7, also see Supplementary File S1). Phenotypes of modular cells

will be the image of the Pareto set in the objective space, known as the Pareto front (***PF***):

$$PF := \{\mathbf{F(x)}: \mathbf{x} \in PS\} \tag{12}$$

For the multiobjective strain design framework, the input parameters include α (eqn. 6), β_k (eqn.

7), and the production networks as input metabolic models. Each model contains a production

module to produce one target chemical. The output is a Pareto set (genetic modifications) and its

respective Pareto front (desirable production phenotypes). For a special case with no trade-off

among the design objectives, an optimal solution, named a utopia point, exists where each

objective achieves its maximum value. The multiobjective strain design formulation presented is

general and can be applied to design modular cells for any organism.

Algorithm and implementation

ModCell2 algorithm. To solve the multiobjective optimization problem for modular cell

engineering, we used multiobjective evolutionary algorithms (MOEAs) (37). MOEAs were

selected because they can efficiently handle linear and non-linear problems and do not require

preferential specification of design objectives (38). MOEAs start by randomly generating a

population of individuals (a vector of design variables), each of which is mapped to a design

objective vector (i.e., a fitness vector). In ModCell2 (Supplementary File S1), the objective

values of an individual are calculated by solving the linear programming problems for each

production network. Next, individuals are shuffled to generate an offspring, from which the most

fit individuals are kept. This process was repeated until the termination criteria was reached, for

instance, either the solutions cannot be further improved or the simulation time limit is reached.

216   *ModCell2 implementation.* To streamline the modular cell design, we developed the

217 ModCell2 software package based on three core classes (Figure S1 in Supplementary File S2).

218 The Prodnet class parses and pre-processes production network models, and computes

219 production phenotypes. The MCdesign class serves as an interface between the MOEA

220 optimization method and metabolic models. Finally, the ResAnalysis class loads the Pareto set

221 computed by MCdesign and identifies the most promising modular cell designs.

222   The code was written in MATLAB 2017b (The Mathworks Inc.) using the function

gamultiobj() from the MATLAB Optimization Toolbox that implements the NSGA-II algorithm

(39) to solve the multiobjective optimization problem. The solution and analysis methods were

parallelized using the MATLAB Parallel Computing Toolbox. The linear programs to calculate

metabolic fluxes were solved using the GNU Linear Programming Kit (GLPK). The COBRA

toolbox (40, 41) and F2C2 0.95b (42) were also used for COBRA model preprocessing and

manipulation.

229   *Metabolic models*. In our study, we used three parent models including i) a small

metabolic network to illustrate the modular cell design concept (13), ii) a core metabolic network

of *Escherichia coli* to compare the performance of ModCell2 with respect to the conventional

single-product strain design strategy and the first-generation modular cell design platform

MODCELL (13), and iii) a genome-scale metabolic network of *E. coli* (i.e., iML1515 (33)) for

biosynthesis of a library of endogenous and heterologous metabolites, including 4 organic acids,

6 alcohols, and 10 esters (Figures S2 in Supplementary File S2) (8, 22, 25, 43-49).

Simulation protocols. Anaerobic conditions were imposed by setting oxygen exchange

fluxes to be 0, and the glucose uptake rate was constrained to be at most 10 mmol/gCDW/h, as

experimentally observed for *E. coli*. When using the genome-scale model iML1515 to simulate

wGCP designs, the commonly observed fermentative products (acetate, CO_2, ethanol, formate,

lactate, succinate) were allowed for secretion as described elsewhere (50). For simulation of

sGCP and *NGP* designs, the glucose uptake rate was fixed (i.e., -10 mmol/gCDW/h); otherwise,

the flux is not active during the no-growth phases, resulting in the product synthesis rate of 0

regardless of genetic manipulations. To compare ModCell2 with Optknock, we applied the

OptKnock algorithm with a titlted objective function (51) to generate *wGCP* designs for each

production network, using the open-source algebraic modeling language Pyomo (52). The MILP

problems were solved using CPLEX 12.8.0 with a time limit of 10,000 seconds set for each

product. ModCell2 is provided as an open-source software package and is freely available for

academic research. The software package and documentation can be download via either

web.utk.edu/~ctrinh or Github https://github.com/TrinhLab.

Analysis methods for design solutions

Compatibility. The compatibility, C ($\Box$ Z^+), of a design is defined as the number of

products that are coupled with a modular cell and has objective values above a specified cutoff

value θ. As a default, we set $\theta = 0.6$ for the *wGCP* and *NGP* design objectives and $\theta = 0.36$

(0.6^2) for the *sGCP* design objective. For example, a *wGCP* design for 3 products that has the

design objective values of 0.4, 0.9, and 0.6 has a compatibility of 2, given a cutoff value of $\theta \geq$

0.6.

Compatibility difference and loss. Robustness is the ability of a system to maintain its

function against perturbations, and hence is very important of designing biological and

engineered systems (53). To evaluate the robustness of modular cell designs, we defined two

metrics, the compatibility difference (*CD*) and compatibility loss (CL $\Box$ [0,1]) as follows:

$$CD = C_{initial} - C_{final} \hspace{3cm} [13]$$

$$CL = \frac{C_{initial} - C_{final}}{C_{initial}} \hspace{3cm} [13]$$

where $C_{initial}$ and C_{final} are the compatibilities of a modular cell design before and after a single

reaction deletion, respectively. The value *CD* > 0 (or *CL* > 0) means the modular gains fitness

while *CD* < 0 (or *CL* < 0) means that it loses its fitness. In the analysis, we did not consider

essential and blocked reactions for our single-deletion analysis; for instance, there are only 1139

potential reaction deletions in the iML1515 model.

Metabolic switch design. A metabolic switch design is a modular cell that can possess

multiple production phenotypes (i.e., *wGCP*, *sGCP*, and *NPG*), activated by an environmental

stimulus (e.g. metabolites, lights). The metabolic switch design is enforced to have a set of

reaction (gene) deletions in one production phenotype to be a subset of the other, for instance,

$\{\mathbf{y}_{wGCP}\} \subseteq \{\mathbf{y}_{NPG}\}$, $\{\mathbf{y}_{wWCP}\} \subseteq \{\mathbf{y}_{sGCP}\}$. The metabolic switch design is beneficial for multiphase

fermentation configurations that enable flexible genetic modification and implementation.

Specifically, the metabolic switch design can exhibit the *wGCP* phenotype during the growth

phase and the *NPG* (or *sGCP*) phenotype during the stationary phase. The metabolic switches

can be implemented using the genetic switchboard (54).

RESULTS AND DISCUSSION

Illustrating ModCell2 for modular cell design of a simplified network

An example parent network, adapted from (13), was used to illustrate ModCell2 (Figure 3A).

Inputs for the multiobjective optimization problem include i) three production networks (Figure

3B), comprising of one endogenous production module (module 1) and two heterologous

production modules (modules 2 and 3) and ii) design parameters (Figure 3C), containing design

objective type, maximum number of deletion reactions (α), and maximum number of module

reactions (β_k). The output of ModCell2 generated the Pareto set and the corresponding Pareto

front for modular cell designs (Figure 3D). The 2-D plots of product yields versus growth rates

presented the feasible phenotypic spaces of the wildtype (gray area) and the designed strain (blue

area).

Using various α and β_k values, ModCell2 simulation generated three $wGCP\text{-}\alpha\text{-}\beta_k\text{-}d$, four

$sGCP\text{-}\alpha\text{-}\beta_k\text{-}d$ designs, and three $NGP\text{-}\alpha\text{-}\beta_k\text{-}d$ designs, where d is the design solution index

(Figure 3D). For instance, by setting $\alpha = 3$ and $\beta_k = 0$, we found three $sGCP$ designs including

sGCP-3-0-1, *sGCP-3-0-2*, and *sGCP-3-0-3*. The first design *sGCP-3-0-1* has a compatibility of 2

(given $\theta \ \square \ 0$) with the design objective values of 0.42 and 0.97 for the products P2 and P3,

respectively. In contrast, the *sGCP-3-0-2* and *sGCP-3-0-3* designs have compatibilities of only 1

with the design objectives of 0.63 for P2 and 0.45 for P1, respectively.

Based on all designs, we can clearly see the trade-offs for optimization of different

products for $\beta_k = 0$. However, setting $\beta_k \geq 1$ helps increase the compatibility of a modular cell

with different production modules. In addition, we found that the Pareto front collapses into a

utopia point as seen in the *wGCP-1-1-1*, *sGCP-3-1-1*, and *NGP-3-1-1* designs. For instance, the

modular cell, *sGCP-3-1-1*, is compatible with all three products. The three corresponding

optimal production strains can couple growth and product formation during the growth phase.

During the stationary phase, these strains produce the products at maximum theoretical yields. In

theory, a universal modular cell always exists, provided that enough reaction deletions and

module reactions are used. It might be more tractable to construct such a modular cell from a

synthetic minimal cell using the bottom-up approach. However, construction of a universal

modular cell from a host organism (e.g., *E. coli*, *S. cerevisiae*) using the top-down approach will

require a significantly large number of genetic modifications, that might be challenging.

Comparing ModCell2 designs with first-generation MODCELL and single product designs

ModCell2 can generate more and better designs than the first-generation modular cell

design platform. To evaluate the algorithms and performance of ModCell2, we directly

compared it with MODCELL (13) in two case studies, using the same core model of *E. coli* for

production of five alcohols (ethanol, propanol, isopropanol, butanol, and isobutanol) and 5

derived butyrate esters (ethyl butyrate, propyl butyrate, isopropyl butyrate, butyl butyrate, and

isobutyl butyrate) from glucose (Figure 4A).

In the first case study, we fixed the reaction module, i.e. $\beta_k = 2$ for ethanol

dehydrogenase (FEM5) and ethanol export reaction (TRA1), in ModCell2 to emulate the same

input as MODCELL (Supplementary File S3). The results showed that ModCell2 generated all

the designs with the same *sGCP* objective values like MODCELL (Figure 4B, 4C, 4D) together

with other alternative solutions (Supplementary File S3). Interestingly, ModCell2 only required 5

and 6 reaction deletions as opposed to 7 and 7 for the *sGCP-5-0-5* and *sGCP-5-0-6* designs,

respectively. By setting the maximum reaction deletions to $\alpha \geq 6$, ModCell2 could find better

design solutions with fewer deletion reaction requirement and higher objective values

(Supplementary File S3).

In the second case study, we used the same model configuration but did not enforce the

module reactions. By setting $\alpha = 5$ and $\beta_k = 1$, we found the *sGCP-5-1-8* design that is

compatible with all products and achieves the same objective values for products found in *sGCP-*

5-0-5, *sGCP-5-0-6*, and *sGCP-5-0-2* (Figure 4E). The desirable phenotypic spaces can be further

constrained for many products if α is increased from 5 to 6 (Figure 4F). Remarkably, by setting

$\alpha = 8$ and $\beta_k = 2$, we found a utopia point design, *sGCP-8-2-9*, without any trade-off among

design objectives (Figure 4G). This utopia point design could not be achieved with $\alpha < 8$

regardless of any β_k value.

Overall, the results demonstrate that ModCell2 can efficiently compute the Pareto front

of modular cell designs. It can find better designs with fewer reaction deletion and module

reaction requirements, improve design objective values, and enhance compatibility.

ModCell2 can identify designs with more compatibility than the conventional single-

product designs. To evaluate if the conventional, single-product design strategy is suitable for

modular cell engineering, we first used OptKnock to generate *wGCP* designs for the same 10

target molecules independently with various allowable reaction deletions ($\alpha = 2, 3, ..., 7$).

Likewise, we employed ModCell2 to produce *wGCP* designs using the same α and various β. To

directly compare OptKnock and ModCell2 solutions, we calculated the *wGCP* design objective

values for all products based on each OptKnock solution (Supplementary File S3). As expected,

our result showed that ModCell2 and OptKnock designs have the same highest objective values

for each product (Figure 5A). However, several OptKnock solutions were always dominated by

ModCell2 solutions in all parameter configurations (Figure 5B). With $\alpha \geq 4$, ModCell2 could

identify *wGCP-α-1* designs with the maximum compatibility of 10, while the best OptKnock

designs only achieved the highest compatibility of 5 (Figure 5C, 5D).

349   Overall, ModCell2 can generate modular cells compatible with the maximum number of

350   modules and achieve high objective values. Single-product designs might not be compatible with

351   a large number of products, and the solutions might be far from Pareto optimality.

353   **Exploring emergent features of modular cell design using an *E. coli* genome-scale network**

354   *Modcell2 can design modular cells using a large-scale metabolic network.* To

demonstrate that ModCell2 can be applied for a genome scale metabolic network, we tested it to

generate *wGCP* designs for 20 target molecules with $\alpha = 4$ and various β_k (Supplementary File

S4). The α value was chosen because with 4 deletions, OptKnock could identify single product

designs with objectives above 60% of the theoretical maximum (Supplementary File S5). With

$\beta_k = 0$, ModCell2 could identify modular cell designs with compatibility of 17, for example, the

wGCP-4-0-50 design featuring deletion of ACALD (acetaldehyde dehydrogenase, *adhE*),

ACKr/PTAr (acetate kinase, *ack;* phosphotransacetylase, *pta*), GLYAT (glycine C-

acetyltransferase, *kbl*), and LDH_D (lactate dehydrogenase, *ldhA*) (Figure 6A, 6D,

Supplementary File S4). By analyzing all *wGCP-4-β_k-d* designs (257 total for $\beta_k = 0$, 1, 2, and

3), we found that the ethanol and D-lactate production modules are most compatible with all

modular cell designs (Figure 6A, 6C, Supplementary File S4). Among reaction deletions, *LdhA*

(86% of designs), *Pta* (38%), and *AdhE* (25%) are the most frequent deletion reactions (Figure

6B). This finding is consistent with a comprehensive survey of metabolic engineering

publications (55) showing that these deleted reactions appeared in most of *E. coli* engineered

369 strains for production of fuels and chemicals. The result supports the potential use of modular

370 cell engineering to systematically build modular platform strains.

371   ***ModCell2 designs can capture combinatorial characteristics of production modules.*** To

372 evaluate whether ModCell2 could capture the combinatorial properties among production

373 modules, we analyzed the Pareto front of *wGCP-4-0-d* that have a total of 58 designs.

374 Hierarchical clustering of this Pareto front revealed certain products with similar objective values

375 across solutions, such as ethyl esters and butyrate esters (Figure 6A). These products together

376 were compatible with different modular cells and exhibited metabolic similarity in their

377 production modules. Thus, ModCell2 could generate designs that capture the combinatorial

378 properties useful for modular cell engineering.

379   ***ModCell2 can identify highly compatible modular cells.*** Analysis of compatibility shows

380 that certain modular cells can couple with production modules that may not exhibit the

381 combinatorial properties (Figure 6D). However, there exists a tradeoff between the number of

382 feasible designs and degree of compatibility. Some modular cell designs are compatible with up

383 to 17 out of 20 products, for instance, the most compatible design, *wGCP-4-0-48*, featuring

384 deletions of ACALD (*adhE*), ACKr/PTAr (*ack*, *pta*), GND (phosphogluconate dehydrogenase,

385 *gnd*) and LDH_D (*ldhA*) (Supplementary File S4). An alternative design *wGCP-4-0-48-*

386 *alternative* also exists where deletion of G6PDH2r (gluose-6-phosphate dehydrogenase, *zwf*) is

387 replaced by that of GND, the first step in the oxidative pentose phosphate pathway. The gene

388 deletions in the design *wGCP-4-0-48-alternative* are a subset of the modular *E. coli* strain

389 TCS095, whose modular properties have recently been validated experimentally (14).

390   To determine if modular cell design is a viable alternative to single-product design, we

391 also analyzed a potential tradeoff between design performance and modularity by comparing the

maximum value of each objective across all solutions in the Pareto front and the single-product

design optima. If production modules exhibit competing phenotypes, a modular cell will not

achieve the same performance in all modules as a single-product design strain. Analysis of the

most compatible design *wGCP-4-0-48-alternative* showed that it could achieve objectives within

4% of the single-product optima in 14 products and within 10% in 3 products (Figure 6E). This

result indicates that it is feasible to identify highly compatible modular cell designs without a

significant tradeoff between performance and modularity.

Analysis of potential tradeoff between robustness and modularity can identify

conserved metabolic features. To evaluate the robustness of modular cells, we analyzed the

compatibility change (*CD*) of *wGCP-4-0* designs with compatibilities of 4 or greater (Figure S3

in Supplementary File 2). Remarkably, the result shows that only 7.5% of potential reaction

deletions were detrimental to the robustness of modular cells while the large remaining portion

did not affect *CD* values. Out of the 85 reactions whose deletion affected compatibility, only a

few appeared consistently across the designs. For instance, deletion of TPI (triose-phosphate

isomerase, *tpi*) led to an average compatibility loss of 95%, inactivating most modular cell

designs. Based on flux variability analysis, TPI must operate in the forward direction by

converting glycerone phosphate (dhap) to glyceraldehyde-3-phosphate (g3p) to drive sufficient

flux through glycolysis and hence preventing synthesis of undesired byproducts (D-lactate or

1,2-propanediol) from dhap. Likewise, deletion of carbon dioxide and water transport and

exchange reactions caused compatibility loss across all designs. Pyruvate carboxylase (PPC) is

an important reaction to channel carbon flux through the Krebs cycle (56), and hence, deletion of

PPC reduces compatibility in most modular cell designs with an average *CL* of 43%.

While some reaction deletions are critical for modular cell robustness, others are

associated with specific products. For example, deletion of PDH (pyruvate dehydrogenase

complex, *lpd/aceEF*) removes compatibility in all butanol-derived designs, indicating PFL

(pyruvate formate lyase, *pfl*) is not an appropriate route. To make heterologous butanol-derived

molecules under anaerobic conditions, FDH (NADH-dependent formate dehydrogenase, *fdh*) is

required in butanol-derived modules where enzymatic reaction pairs of PFL and FDH could

substitute PDH known to be anaerobically inhibited.

Overall, analysis of tradeoff between modularity and robustness can identify not only the

conserved metabolic features of modular cells but also potential bottlenecks in specific

production modules.

Enabling metabolic switch among different design objectives using ModCell2

The ability to dynamically control growth and production phases can potentially enhance

product titers, rates, and yields. For instance, two-phase fermentation can be employed where

growth phase is optimized for biomass synthesis and stationary phase for chemical production

(57). Using ModCell2, we investigated the feasibility to design optimal strains to toggle switch

desirable production phenotypes.

To design a *wGCP→NPG* metabolic switch, we first used our reference *wGCP* design as

a parent strain (Figure 7A) and then employed ModCell2 to identify the most compatible

wGCP→NPG designs. With 5 additional deletions, we could find *wGCP→*NPG designs that

encompass both *wGCP* and *NPG* phenotypes, for instance, the *sup-NGP-5-0-23* design featuring

deletion of PGI (glucose-6-phosphate isomerase, *pgi*), MDH (malate dehydrogenase, *mdh*),

ASPT (L-aspartase, *aspT*), Tkt2 (transketolase, *tktB*), and ATPS4rpp (ATP synthase, *atp*) (Figure

7B). The deletion reactions in the *wGCP→NPG* designs appear in both catabolic (PGI,

ATPS4tpp) and anabolic (ASP, TKT2) processes, responsible for growth disruption and direction

of carbon flow to the biosynthesis of target products.

Likewise, we used ModCell2 to design a *wGCP→sGCP* metabolic switch. We identified

the most compatible *wGCP→sGCP* designs with 6 additional deletions, for instance, the *sup-*

sGCP-6-0-39 design featuring the deletion of MGSA (Methylglyoxal synthase, *mgsA*), ALCD2x

(alcohol dehydrogenase, *adhE*), PFL, MDH, FADRx (FAD reductase, *fadI*), and GLUDy

(NADP$^+$dependent glutamate dehydrogenase, *gdhA*) (Figure 7C). Different from the

wGCP→NGP metabolic switch, all deletions in the *wGCP→sGCP* designs are involved the

elimination of biosynthesis pathways of undesirable byproducts.

While it is feasible to metabolically switch among different production phenotypes, it not

only requires more reaction deletions but also reduces the product compatibility. For instance,

the *wGCP→sGCP* and *wGCP→NGP* designs are only compatible with 5 products while the

wGCP parent design have a compatibility of 17 out of 20 products with 4 deletions. The main

reason is that both *wGCP→sGCP* and *wGCP→NGP* designs must eliminate all possible

redundant pathways that result in biosynthesis of undesirable byproducts.

CONCLUSION

In this study, we developed a multiobjective strain design platform for modular cell

engineering. With a new developed algorithm and computational platform, ModCell2 enables

flexible design of modular cells that can couple with production modules to exhibit desirable

production phenotypes. In comparison to the first-generation strain design platform, ModCell2

can handle large-scale metabolic networks and identify better solutions that require fewer genetic

modifications and exhibit more product compatibility. Different from the conventional single-

product strain design, ModCell2 can find solutions that are Pareto optimal with negligible

tradeoffs among modularity, performance, and robustness. We envision ModCell2 is a useful

tool to implement modular cell engineering and fundamentally study modular designs in natural

and synthetic biological systems.

ACKNOWLEDGEMENTS

This research was financially supported in part by the NSF CAREER Award

(NSF#1553250) and the DOE subcontract Grant (DE-AC05-000R22725) by the Center of

Bioenergy Innovation (CBI), the U.S. Department of Energy Bioenergy Research Center funded

by the Office of Biological and Environmental Research in the DOE Office of Science. The

funders had no role in the study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

REFERENCES

1. Nielsen J & Keasling Jay D (2016) Engineering Cellular Metabolism. *Cell* 164:1185-1197.

2. Trinh CT & Mendoza B (2016) Modular cell design for rapid, efficient strain engineering toward industrialization of biology. *Current Opinion in Chemical Engineering* 14:18-25.

3. Connelly T, *et al.* (2015) Industrialization of Biology: A Roadmap to Accelerate the Advanced Manufacturing of Chemicals.

4. Sauro HM (2008) Modularity defined. *Molecular systems biology* 4.

5. Purnick PEM & Weiss R (2009) The second wave of synthetic biology: from modules to systems. *Nature reviews Molecular cell biology* 10:410-422.

6. Yadav VG, De Mey M, Giaw Lim C, Kumaran Ajikumar P, & Stephanopoulos G (2012) The future of metabolic engineering and synthetic biology: Towards a systematic practice. *Metabolic Engineering* 14:233-241.

7. Biggs BW, De Paepe B, Santos CNS, De Mey M, & Ajikumar PK (2014) Multivariate modular metabolic engineering for pathway and strain optimization. *Current opinion in biotechnology* 29:156-162.

8. Rodriguez GM, Tashiro Y, & Atsumi S (2014) Expanding ester biosynthesis in Escherichia coli. *Nat Chem Biol* 10:259-265.

9. Cheong S, Clomburg JM, & Gonzalez R (2016) Energy-and carbon-efficient synthesis of functionalized small molecules in bacteria using non-decarboxylative Claisen condensation reactions. *Nature biotechnology* 34:556-561.

496   10.   Tseng H-C & Prather KL (2012) Controlled biosynthesis of odd-chain fuels and
497   chemicals via engineered modular metabolic pathways. *Proceedings of the National*
498   *Academy of Sciences* 109(44):17925-17930.

499   11.   Xu P*, et al.* (2013) Modular optimization of multi-gene pathways for fatty acids
500   production in *E. coli*. *Nat Commun* 4:1409.

501   12.   Ng CY, Chowdhury A, & Maranas CD (2016) A microbial factory for diverse chemicals.
502   *Nat Biotech* 34:513-515.

503   13.   Trinh CT, Liu Y, & Conner DJ (2015) Rational design of efficient modular cells.
504   *Metabolic engineering* 32:220-231.

505   14.   Wilbanks B, Layton D, Garcia S, & Trinh C (2017) A Prototype for Modular Cell
506   Engineering. *ACS Synthetic Biology*:acssynbio.7b00269.

507   15.   Long MR, Ong WK, & Reed JL (2015) Computational methods in metabolic engineering
508   for strain design. *Current opinion in biotechnology* 34:135-141.

509   16.   Machado D & Herrgård MJ (2015) Co-evolution of strain design methods based on flux
510   balance and elementary mode analysis. *Metabolic Engineering Communications* 2:85-92.

511   17.   Chowdhury A, Zomorrodi AR, & Maranas CD (2015) Bilevel optimization techniques in
512   computational strain design. *Computers & Chemical Engineering* 72:363-372.

513   18.   Fong SS*, et al.* (2005) In silico design and adaptive evolution of Escherichia coli for
514   production of lactic acid. *Biotechnology and Bioengineering* 91:643-648.

515   19.   Trinh CT & Srienc F (2009) Metabolic engineering of Escherichia coli for efficient
516   conversion of glycerol to ethanol. *Applied and Environmental Microbiology* 75(21):6696-
517   6705.

20.   Otero JM, *et al.* (2013) Industrial systems biology of Saccharomyces cerevisiae enables

novel succinic acid cell factory. *PLoS One* 8:e54144.

21.   Burgard AP, Pharkya P, & Maranas CD (2003) Optknock: a bilevel programming

framework for identifying gene knockout strategies for microbial strain optimization.

Biotechnology and Bioengineering 84:647-657.

22.   Trinh CT, Unrean P, & Srienc F (2008) Minimal Escherichia coli Cell for the Most

Efficient Production of Ethanol from Hexoses and Pentoses. *Applied and Environmental*

Microbiology 74(12):3634-3643.

23.   Klamt S & Mahadevan R (2015) On the feasibility of growth-coupled product synthesis

in microbial strains. *Metabolic engineering* 30:166-178.

24.   Wierzbicki M, Niraula N, Yarrabothula A, Layton DS, & Trinh CT (2016) Engineering

an Escherichia coli platform to synthesize designer biodiesels. *Journal of biotechnology*

224:27-34.

25.   Layton DS & Trinh CT (2014) Engineering modular ester fermentative pathways in

Escherichia coli. *Metabolic Engineering* 26:77-88.

26.   Layton DS & Trinh CT (2016) Expanding the modular ester fermentative pathways for

combinatorial biosynthesis of esters from volatile organic acids. *Biotechnology and*

bioengineering.

27.   Layton DS & Trinh CT (2016) Microbial synthesis of a branched-chain ester platform

from organic waste carboxylates. *Metabolic Engineering Communications* 3:245-251.

28.   Trinh CT, Wlaschin A, & Srienc F (2009) Elementary mode analysis: a useful metabolic

pathway analysis tool for characterizing cellular metabolism. *Applied Microbiology and*

Biotechnology 81:813-826.

29.   Jouhten P*, et al.* (2016) Yeast metabolic chassis designs for diverse biotechnological

542   products. *Scientific reports* 6.

543   30.   Price ND, Papin JA, Schilling CH, & Palsson BO (2003) Genome-scale microbial in

544   silico models: the constraints-based approach. *Trends in Biotechnology* 21:162-169.

545   31.   Palsson BØ (2015) Systems biology: constraint-based reconstruction and analysis.

546   (Cambridge University Press).

547   32.   Tepper N & Shlomi T (2010) Predicting metabolic engineering knockout strategies for

548   chemical production: accounting for competing pathways. *Bioinformatics* 26:536-543.

549   33.   Feist AM*, et al.* (2010) Model-driven evaluation of the production potential for growth-

550   coupled products of Escherichia coli. *Metabolic engineering* 12:173-186.

551   34.   Yang L, Cluett WR, & Mahadevan R (2011) EMILiO: a fast algorithm for genome-scale

552   strain design. *Metabolic engineering* 13:272-281.

553   35.   Fong SS*, et al.* (2005) In silico design and adaptive evolution of Escherichia coli for

554   production of lactic acid. *Biotechnology and bioengineering* 91(5):643-648.

555   36.   Trinh C & Srienc F (2009) Metabolic engineering of *Escherichia coli* for efficient

556   conversion of glycerol to ethanol. *Appl Environ Microbiol* 75(21):6696 - 6705.

557   37.   Coello Coello CA (2002) Theoretical and numerical constraint-handling techniques used

558   with evolutionary algorithms: a survey of the state of the art. *Computer Methods in*

559   *Applied Mechanics and Engineering* 191(11):1245-1287.

560   38.   Marler RT & Arora JS (2004) Survey of multi-objective optimization methods for

561   engineering. *Structural and multidisciplinary optimization* 26:369-395.

562   39.   Deb K, Pratap A, Agarwal S, & Meyarivan T (2002) A fast and elitist multiobjective

563   genetic algorithm: NSGA-II. *Evolutionary Computation, IEEE Transactions on* 6:182-

564   197.

565   40.   Schellenberger J, *et al.* (2011) Quantitative prediction of cellular metabolism with

566   constraint-based models: the COBRA Toolbox v2. 0. *Nature protocols* 6:1290-1307.

567   41.   Heirendt L, *et al.* (2017) Creation and analysis of biochemical constraint-based models:

568   the COBRA Toolbox v3.0.

569   42.   Larhlimi A, David L, Selbig J, & Bockmayr A (2012) F2C2: a fast tool for the

570   computation of flux coupling in genome-scale metabolic networks. *BMC Bioinformatics*

571   13:57.

572   43.   Shen CR, *et al.* (2011) Driving forces enable high-titer anaerobic 1-butanol synthesis in

573   Escherichia coli. *Applied and environmental microbiology* 77:2905-2915.

574   44.   Atsumi S, Hanai T, & Liao JC (2008) Non-fermentative pathways for synthesis of

575   branched-chain higher alcohols as biofuels. *Nature* 451:86-89.

576   45.   Tseng H-C & Prather KLJ (2012) Controlled biosynthesis of odd-chain fuels and

577   chemicals via engineered modular metabolic pathways. *Proceedings of the National*

578   *Academy of Sciences* 109:17925-17930.

579   46.   Yim H, *et al.* (2011) Metabolic engineering of Escherichia coli for direct production of 1,

580   4-butanediol. *Nature chemical biology* 7:445-452.

581   47.   Akita H, Nakashima N, & Hoshino T (2016) Pyruvate production using engineered

582   Escherichia coli. *AMB Express* 6(1):94.

583   48.   Niu D, *et al.* (2014) Highly efficient L-lactate production using engineered Escherichia

584   coli with dissimilar temperature optima for L-lactate formation and cell growth.

585   *Microbial cell factories* 13(1):78.

586   49.   Yu JL, Xia XX, Zhong JJ, & Qian ZG (2014) Direct biosynthesis of adipic acid from a

587   synthetic pathway in recombinant *Escherichia coli. Biotechnology and bioengineering*

588   111(12):2580-2586.

589   50.   von Kamp A & Klamt S (2017) Growth-coupled overproduction is feasible for almost all

590   metabolites in five major production organisms. *Nature communications* 8:15956.

591   51.   Maranas CD & Zomorrodi AR (2016) Optimization Methods in Metabolic Networks.

592   52.   Hart WE, *et al.* (2012) *Pyomo-optimization modeling in python* (Springer).

593   53.   Kitano H (2004) Biological robustness. *Nature Reviews Genetics* 5(11):826.

594   54.   Callura JM, Cantor CR, & Collins JJ (2012) Genetic switchboard for synthetic biology

595   applications. *Proceedings of the National Academy of Sciences* 109(15):5850-5855.

596   55.   Winkler JD, Halweg-Edwards AL, & Gill RT (2015) The LASER database: Formalizing

597   design rules for metabolic engineering. *Metabolic Engineering Communications* 2:30-38.

598   56.   Coomes MW, Mitchell BK, Beezley A, & Smith TE (1985) Properties of an Escherichia

599   coli mutant deficient in phosphoenolpyruvate carboxylase catalytic activity. *Journal of*

600   *bacteriology* 164(2):646-652.

601   57.   Klamt S, Mahadevan R, & Hädicke O (2018) When Do Two☐Stage Processes

602   Outperform One☐Stage Processes? *Biotechnology journal* 13(2):1700539.

 Figure legends

Figure 1: Comparison between the conventional single-product strain design and modular cell

engineering. In the conventional approach, each target product requires to go through the

iterative optimization cycle. The modular cell engineering approach exploits common

phenotypes associated with high product titers, rates, and yields; and hence, the strain

optimization cycle only needs to be performed once for multiple products, which helps reduce

the cost and time of strain development.

Figure 2: Graphical representation of phenotypic spaces for different strain design objectives

including **(A)** weak growth coupling (*wGCP*), **(B)** strong growth coupling (*sGCP*), and **(C)** no-

growth production (*NPG*). v_{Pk}^{μ} is the minimum product formation rate at the maximum growth

rate for production network k, and $v_{P_{max}k}^{\mu}$ is the maximum product secretion rate attainable. $v_{Pk}^{\bar{\mu}}$

and $v_{P_{max}k}^{\bar{\mu}}$ are the minimum and maximum product formation rates for production network k

during the stationary phase, respectively.

Figure 3: Illustration of ModCell2 workflow and analysis including **(A)** parent model, **(B)**

production modules, **(C)** design parameters, and **(D)** simulation output for Pareto set and Pareto

front based on design input.

Figure 4: The 2-D metabolic phenotypic spaces of different *sGCP* designs using the core

metabolic model. **(A)** Metabolic map, **(B)** *sGCP-5-0-5* design, **(C)** *sGCP-5-0-6* design, **(D)**

sGCP-5-0-2 design, **(E)** *sGCP-5-1-8* design, **(F)** *sGCP-6-1-10* design, and **(G)** *sGCP-8-2-9*

design. For each panel, the gray and blue areas correspond to the phenotypic spaces of the

wildtype and the optimal production strain, respectively.

Figure 5: Comparison of strain design by OptKnock and Modcell2. (**A**) A correlation between

the maximum objective values for each product generated by OptKnock and the equivalent

values attained by ModCell2. Each point is colored based on the number of reaction deletions,

with warmer colors corresponding to more reaction deletions. (**B**) A comparison between the

Optknock objective vectors with at most 7 reaction deletions and the representative ModCell2

objective vector, *wGCP-7-1* which dominates them. Each color circle represents a pair of

dominating *wGCP* design and dominated OptKnock solution (Supplementary File S3). (**C**)

Maximum compatibility of OptKnock designs (blue), *wGCP* designs ($\beta_k = 0$, orange), wGCP

designs ($\beta_k = 1$, yellow). (**D**) Compatibility distribution of Optknock ($\alpha = 7$, orange) and *wGCP-*

7-1 (blue).

Figure 6: (A) Pareto front of *wGCP-4-0-d*. The columns correspond to different designs labeled

by their design index, *d*, where the rows correspond to different products. (**B**) Frequency of the

top deletion reactions. (**C**) Product compatibility distribution across designs. (**D**) Design

compatibility. (**E**) Tradeoff between modularity and performance. The bars correspond to the

maximum objective values attainable for each product whereas the blue line represent the

objective values of the *wGCP-4-0-48alternative* design.

Figure 7: Production phenotypes of (**A**) the wild type (gray) and the representative, highly-

compatible design *wGCP-4-0-48-alternative* (blue), (**B**) the *wGCP→NPG* design, *sup-NGP-5-0-*

23, and (**C**) the *wGCP→sGCP* design, *sup-sGCP-6-0-39*.

Figure 1

Features	Conventional strain engineering	Modular cell engineering
Parent strain		
Modular cell	Absent	Optimized common production phenotypes
Exchangeable modules	1	Multiple
Optimal production strains		Module1 ... Module 2 ... Module 3
Design-build-test cycle	Repeated for every new product	One time for many products

656   **Figure 2**

658
659
660

Objective	α	β_k	d	Reaction deletions	Module reactions			f_k			Prod. yield vs growth		
					P1	P2	P3	P1	P2	P3	P1	P2	P3
wGCP	1	0	1	R10				0.45	0.00	0.00			
wGCP	1	0	2	R9				0.00	0.63	0.97			
wGCP	1	1	1	R9	R9			0.45	0.63	0.97			
sGCP	3	0	1	R8, R9, R11				0.00	0.42	0.97			
sGCP	3	0	2	R6, R9, R10				0.00	0.63	0.00			
sGCP	3	0	3	R6, R8, R11				0.45	0.00	0.00			
sGCP	3	1	1	R6, R9, R11	R9		R6	0.45	0.63	0.97			
NGP	3	0	1	R6, R11				1.00	0.00	0.00			
NGP	3	0	2	R4, R9, R11				0.00	1.00	1.00			
NGP	3	1	1	R4, R9, R11	R9			1.00	1.00	1.00			

667   **Figure 4**

Figure 5

Figure 6

Figure 7