Parameter inference with analytical propagators for stochastic models of autoregulated gene expression

Frits Veerman[1], **Nikola Popović**[1,*]**, and Carsten Marr**[2]

[1]University of Edinburgh, School of Mathematics, Edinburgh, EH9 3FD, United Kingdom
[2]Helmholtz Zentrum München, Institute of Computational Biology, 85764 Neuherberg, Germany
[*]Nikola.Popovic@ed.ac.uk

ABSTRACT

Stochastic gene expression in regulatory networks is conventionally modelled via the Chemical Master Equation (CME). As explicit solutions to the CME, in the form of so-called propagators, are oftentimes not readily available, various approximations have been proposed. A recently developed analytical method is based on a separation of scales that assumes significant differences in the lifetimes of mRNA and protein in the network, allowing for the efficient approximation of propagators from asymptotic expansions for the corresponding generating functions. Here, we showcase the applicability of that method to simulated data from a 'telegraph' model for gene expression that is extended with an autoregulatory mechanism. We demonstrate that the resulting approximate propagators can be applied successfully for Bayesian parameter inference in the non-regulated model; moreover, we show that, in the extended autoregulated model, autoactivation or autorepression may be refuted under certain assumptions on the model parameters. Our results indicate that the method showcased here may allow for successful parameter inference and model identification from longitudinal single cell data.

1 Introduction and background

Gene expression in regulatory networks is an inherently stochastic process[1]. Mathematical models typically take the form of a Chemical Master Equation (CME), which describes the temporal evolution of the probabilities of observing specific states in the network[2]. Recent advances in single-cell fluorescence microscopy[3–7] have allowed for the generation of experimental longitudinal data, whereby the fluorescence intensity of mRNA or protein abundances in single cells is measured. While most common models assume the availability of protein abundance data, abundances of mRNA may equally be of interest, depending on the model[8]. Here, we focus on abundances of protein, which we assume to be measured at regular sampling intervals Δt. A typical data set, denoted by Q, thus consists of protein abundances n_i at $N+1$ different points in time; see Figure 1A. We can group these abundances into transitions $n_i \to n_{i+1}$; cf. Figure 1B. A model-derived propagator $P_{n_{i+1}|n_i}(\Delta t, \Theta)$ allows for the calculation of the probabilities of such transitions for some set of model parameters Θ. Summing over all these probabilities for all N observed transitions, we can calculate the log-likelihood $L(\Theta)$ of that particular parameter set as

$$L(\Theta) = \sum_{i=0}^{N-1} \log P_{n_{i+1}|n_i}(\Delta t, \Theta), \tag{1.1}$$

which can be evaluated over a range of values for the model parameters to yield a 'log-likelihood landscape', the maximum of which corresponds to the most likely parameter set Θ subject to the measured data set Q. Assessing the likelihood of a model by evaluating the associated propagator for the observed transitions in a time-lapse experiment is a feasible, established approach that has been successfully applied previously[6,9,10]. Due to the complex nature of the underlying regulatory networks, explicit expressions for $P_{n_{i+1}|n_i}$ are difficult to obtain in general. Hence, a variety of approximations have been proposed, which can be either numerical[10,11] or analytical[12,13], to cite but a few examples. Here, we apply the analytical method recently developed by the current authors[14], which was based on ideas presented by Popović, Marr & Swain[13], to obtain fully time-dependent approximate propagators; an outline of the method is given in Section 2.

Our aim in the present article is to demonstrate the applicability of these propagators, as well as to evaluate their performance in the context of Bayesian parameter inference for synthetic data. Specifically, we showcase the resulting inference procedure for a family of stochastic gene expression models. First, in Section 3, we consider a model that incorporates DNA 'on'/'off' states ('telegraph model'); see also the work of Raj *et al.*[15] and Shahrezaei & Swain[9]. Subsequently, in Section 4, that model is extended with an autoregulatory mechanism, whereby protein influences its own production through an autocatalytic reaction. In Section 5, we summarise our results and present an outlook to future research; finally, in Appendix A, we collate the analytical formulae that underly our inference procedure for the family of models showcased here.

Figure 1. (A) Simulated time series of protein abundance n, with measurements at times t_i and sampling interval Δt. (B) Histogram of the frequency k_j of transitions $(n_0 \rightarrow n)_j$, inferred from a longer time series with 100 transitions.

2 Method

2.1 Calculation of propagators

Our method[14] is based on an analytical approximation of the probability generating function that is introduced for analysing the CME corresponding to the given gene expression model. Propagators can be calculated from the generating function via the Cauchy integral formula, which implies

$$P_{n_{i+1}|n_i}(\Delta t, \Theta) = \frac{1}{2\pi i} \oint_\gamma \frac{F(z; \Delta t, n_i, \Theta)}{z^{n_{i+1}+1}} \, dz; \tag{2.1}$$

here, $F(z; \Delta t, n_i, \Theta)$ is the generating function of the (complex) variable z, which additionally depends on the sampling time Δt, the protein abundance n_i, and the model parameter set Θ. The integration contour γ is a closed contour in the complex plane around $z = 0$. The choice of contour is arbitrary; however, it can have a significant effect on computation times and on the accuracy of the resulting integrals[16]. Here, we choose γ to be a regular 150-sided polygon approximating a circle of radius 0.8 that is centred at the origin in the complex plane, which results in a 'hybrid analytical-numerical' procedure for the evaluation of $P_{n_{i+1}|n_i}$.

2.2 Parameter inference

The parameter inference procedure proposed here can be divided into the following steps:

1) Data binning. The simulated data Q is presented as a time series $\{n_i\}$, $0 \leq i \leq N$, which yields N transitions $n_i \rightarrow n_{i+1}$. Generically, some of these transitions occur more than once. For computational efficiency, we bin the data accordingly to create a binned data set $\widehat{Q} = \left\{ \left(k_j, (n_0 \rightarrow n)_j \right) \right\}$, with $1 \leq j \leq \widehat{N}$ for $\widehat{N} \leq N$, where k_j is the frequency of the transition $(n_0 \rightarrow n)_j$; see also Figure 1. Here, $\widehat{N}$ denotes the number of different transitions observed in the data, that is, the size of the binned data set $\widehat{Q}$. We emphasise that binning can substantially accelerate parameter inference, in particular for near-stationary processes.

2) Marginalisation. Frequently, some of the involved species in a model are not observed, and hence have to be marginalised over. In the models discussed in Sections 3 and 4, we assume that protein is measured, while mRNA remains unobserved. Marginalisation over unobserved species is usually carried out on the transition probabilities in (2.1). However, since the marginalisation procedure is linear, it commutes with the Cauchy integral. Introducing the linear 'marginalisation operator' $\mathbb{M}$, we may write

$$\mathbb{M} P_{n_{i+1}|n_i}(\Delta t, \Theta) = \frac{1}{2\pi i} \oint_\gamma \frac{\mathbb{M} F(z; \Delta t, n_i, \Theta)}{z^{n_{i+1}+1}} \, dz, \tag{2.2}$$

where $\mathbb{M}$ now acts on the generating function F. Therefore, given the analytical approximation for F resulting from our method[14], we define

$$\widehat{F}(z; \Delta t, n_i, \widehat{\Theta}) = \mathbb{M} F(z; \Delta t, n_i, \Theta), \tag{2.3}$$

where $\widehat{\Theta} \subset \Theta$ is the subset of parameters that remain after the marginalisation procedure has been applied. Note that $\widehat{F}$ is still a fully analytical, general expression which depends on the as yet unspecified values of its arguments.

3) Evaluation. We choose a set $\widehat{\Theta}_0$ of numerical values for the parameters in $\widehat{\Theta}$. Moreover, we specify the integration contour γ, which we discretise as described in 2) to approximate the Cauchy integral in (2.1) by a finite sum. Suppose that the contour γ is discretised as $\{\zeta(l)\}$, with $0 \le l \le L$ and $\zeta(0) = \zeta(L)$; then, the integral of a function G along γ is approximated as

$$\oint_\gamma G(z)\,\mathrm{d}z \approx \sum_{l=0}^{L-1} G(\zeta(l))\,\Delta\zeta(l), \quad \text{with } \Delta\zeta(l) = \zeta(l+1) - \zeta(l). \tag{2.4}$$

Now, for every transition $(n_0 \to n)_j$ in the binned data set $\widehat{Q}$, we evaluate $\widehat{F}$, as given in (2.3), for the chosen parameter values $\widehat{\Theta}_0$ along the discretised contour. We hence obtain the array

$$\left\{ \frac{1}{2\pi i} \frac{\widehat{F}(\zeta(l); \Delta t, (n_0)_j, \widehat{\Theta}_0)}{\zeta(l)^{(n)_j + 1}} \Delta\zeta(l) \right\} \quad \text{for } 0 \le l \le L-1 \text{ and } 1 \le j \le \widehat{N}, \tag{2.5}$$

which we sum over l to find

$$p_j(\widehat{\Theta}_0, \Delta t) = \sum_{l=0}^{L-1} \frac{1}{2\pi i} \frac{\widehat{F}(\zeta(l); \Delta t, (n_0)_j, \widehat{\Theta}_0)}{\zeta(l)^{(n)_j + 1}} \Delta\zeta(l) \tag{2.6}$$

as the approximate value of the propagator for the transition $(n_0 \to n)_j$.

4) Calculation of the log-likelihood. To calculate the log-likelihood of the parameter subset $\widehat{\Theta}_0$, we substitute the approximate propagators p_j, as defined in (2.6), into (1.1) to obtain

$$L(\widehat{\Theta}_0) = \sum_{j=1}^{\widehat{N}} k_j \log p_j(\widehat{\Theta}, \Delta t). \tag{2.7}$$

3 Showcase 1: The telegraph model

To demonstrate our parameter inference procedure, we consider a stochastic gene expression model that incorporates DNA 'on'/'off' states ('telegraph model')[9,15]:

$$
\begin{array}{lll}
D & \underset{k_1}{\overset{k_0}{\rightleftharpoons}} D^* & \text{(DNA activation/deactivation),} \\
D^* & \overset{v_0}{\to} D^* + M & \text{(transcription of DNA to mRNA),} \\
M & \overset{v_1}{\to} M + P & \text{(translation of mRNA to protein),} \\
M & \overset{d_0}{\to} \emptyset & \text{(decay of mRNA),} \\
P & \overset{d_1}{\to} \emptyset & \text{(decay of protein).}
\end{array}
\tag{3.1}
$$

In recent work[14], we presented an analytical method for obtaining explicit, general, time-dependent expressions for the generating function associated to the CME that arises from the model in (3.1). A pivotal element of the application of that method to (3.1) is the assumption that the protein decay rate d_1 is notably smaller than the decay rate d_0 of mRNA, which implies that the parameter $\varepsilon := \frac{d_1}{d_0}$ is small; hence, the associated generating function is *approximated* to a certain order $O = k$, corresponding with a theoretical accuracy that is proportional to ε^k. For more details on the resulting approximation, we refer to Appendix A.

To obtain synthetic data, we simulate the model in (3.1) using Gillespie's stochastic simulation algorithm[17], for fixed values of the (rescaled) parameters

$$\kappa_0 := \frac{k_0}{d_1} = 1.3, \ \kappa_1 := \frac{k_1}{d_1} = 1.2, \ \lambda := \frac{v_0}{d_1} = 3.3, \ \mu := \frac{v_1}{d_0} = 2.85, \ \varepsilon := \frac{d_1}{d_0} = 0.1, \text{ and } d_1 = 1 \tag{3.2}$$

on the time interval $0 \le t \le 10$, and we measure the protein abundance n with a fixed sampling interval Δt. As our method assumes that Δt is of order ε, cf. again Appendix A, we set $\Delta t = \varepsilon = 0.1$, which yields $N = 100$ transitions. Finally, we consider random initial states, with mRNA and protein numbers chosen uniformly between 0 and 10, and we assume DNA to be in the 'on' state with probability $\frac{\kappa_0}{\kappa_0 + \kappa_1}$. Based on the simulated measurement data, we perform the parameter inference procedure described in Section 2. As the data consists of protein abundances only, and as propagators for the model in (3.1) depend on abundances of both mRNA and protein, we marginalise over mRNA abundance assuming a steady-state distribution

Figure 2. Log-likelihood landscapes inferred from a simulation of the telegraph model in (3.1) with $N = 100$ transitions and parameter values as in (3.2). (A) Leading (zeroth) order approximation. (B) First order approximation; true value (cross) versus maximum likelihood estimate (MLE; dot).

reported by Raj *et al.*[15]; that distribution coincides with the steady-state limit of the associated fully time-dependent distribution considered by Veerman, Marr & Popović[14]. We assume that the values of κ_0, κ_1, ε, and d_1 are known, and calculate the log-likelihood in (2.7) for varying λ and μ. We scan these two parameters in the range $\{10^{-3} \leq \lambda \leq 10^3,\ 10^{-3} \leq \mu \leq 10^2\}$, using a logarithmically spaced grid of 50×40 grid points. Figure 2 shows the resulting log-likelihood landscapes and, in particular, a comparison of the performance of the leading (zeroth) order approximation for the generating function, see Figure 2A, with that of the first order approximation in Figure 2B.

To quantify the performance of the method developed by Veerman, Marr & Popović[14] for parameter inference, we compare four different scenarios:

(a) Parameter values as in (3.2), with sampling interval $\Delta t = \varepsilon = 0.1$ on the time interval $0 \leq t \leq 10$, corresponding to $N = 100$ transitions, which is the original setup that yields the results shown in Figure 2.

(b) As in (a), with the time interval increased to $0 \leq t \leq 100$, which yields $N = 1000$ transitions.

(c) As in (a), with $\varepsilon = 0.01$; the sampling interval is decreased accordingly to $\Delta t = \varepsilon = 0.01$; measurements are taken on the time interval $0 \leq t \leq 1$, which yields $N = 100$ transitions.

(d) As in (a), with $\mu = 28.5$.

For each scenario, we infer the most likely values of the parameters λ and μ, for increasing approximation order O. The inferred values of λ and μ are compared to the 'true' values λ_{true} and μ_{true}, where we consider relative errors to quantify the performance of our inference procedure. The results of that comparison are shown in Figure 3. The accuracy of inference for λ clearly increases when the approximation order O is increased from 0 to 1; the increase in accuracy from $O = 1$ to $O = 2$ is obfuscated by grid size effects. A ten-fold increase in the number of transitions (b) increases the accuracy of the leading order approximation, while a ten-fold increase in the value of μ_{true} (d) decreases the accuracy of the leading order approximation. For μ, there is no noticeable increase in accuracy with the approximation order O, within the parameter grid used. However, the accuracy of inference for μ increases overall when the number of transitions is increased (b), the small parameter ε is decreased (c), or the value of μ_{true} is increased (d).

4 Showcase 2: An autoregulated telegraph model

We extend the telegraph model in (3.1) with an autoregulatory mechanism, where the DNA activation rates are influenced by the presence of protein. Autoregulation is modelled in a catalytic manner, via one of the two following reactions:

$$D + P \xrightarrow{a_P} D^* + P \qquad \text{(autoactivation through protein)}, \qquad (4.1a)$$

$$D^* + P \xrightarrow{r_P} D + P \qquad \text{(autorepression through protein)}. \qquad (4.1b)$$

Figure 3. Relative error $\Delta_r(x) := \frac{x-x_{\text{true}}}{x_{\text{true}}}$ of the inferred parameters λ (A) and μ (B), for increasing approximation order O; for $O = k$, the propagators $P_{n_{i+1}|n_i}$ are approximated up to and including terms of order ε^k. (a) $N = 100$ transitions, $\varepsilon = 0.1$, $\lambda_{\text{true}} = 3.3$, and $\mu_{\text{true}} = 2.85$. (b) $N = 1000$ transitions, other parameters as in (a). (c) $\varepsilon = 0.01$, number of transitions N and other parameters as in (a). (d) $\mu_{\text{true}} = 28.5$, number of transitions N and other parameters as in (a).

The above pair of autoregulation mechanisms was introduced by Hornos *et al.*[18], and implemented e.g. by Iyer-Biswas & Jayaprakash[19]; see Section 5 for a discussion of the physical validity of these mechanisms.

To assess the performance of our parameter inference procedure, we fix the parameter values as in (3.2). Again, we marginalise over mRNA abundance assuming a steady-state distribution; see Section 3. We generate six data sets, as follows:

(A) Simulate the model in (3.1) without autoregulation ('null model'; $a_P = r_P = 0$) on the time interval $0 \le t \le 10$, which yields $N = 100$ transitions.

(B) As in (A), with the time interval increased to $0 \le t \le 100$, which yields $N = 1000$ transitions.

(C) Simulate the extended model $\{(3.1),(4.1a)\}$ with autoactivation for $a_P \delta = 0.3$ on the time interval $0 \le t \le 10$, which yields $N = 100$ transitions.

(D) As in (C), with the time interval increased to $0 \le t \le 100$, which yields $N = 1000$ transitions.

(E) Simulate the extended model $\{(3.1),(4.1b)\}$ with autorepression for $r_P \delta = 0.3$ on the time interval $0 \le t \le 10$, which yields $N = 100$ transitions.

(F) As in (E), with the time interval increased to $0 \le t \le 100$, which yields $N = 1000$ transitions.

Every data set consists of 10 runs of equal length.

Generating functions for the autoregulated extension, by (4.1), of the telegraph model in (3.1) have been derived in the theoretical companion article[14] to the current work, under the assumption that the autoregulation rate a_P or r_P is small compared to the protein decay rate d_1. That assumption implies that the ratios $\frac{a_P}{d_1} := \alpha_P \delta$ and $\frac{r_P}{d_1} := \rho_P \delta$ are small.

Parameter inference now proceeds as follows. We fix a data set, and take a single run from that set. For that run, we determine the likelihood of the autoactivated model in $\{(3.1),(4.1a)\}$, varying $0 \le \alpha_P \delta \le 1$; likewise, we determine the likelihood of the autorepressed model in $\{(3.1),(4.1b)\}$, varying $0 \le \rho_P \delta \le 1$. The likelihood L of the autoregulated extension is then compared with the likelihood L_0 of the non-regulated model in (3.1); as before, N denotes the number of data points, where L is defined as in (2.7), that is, we simply evaluate the probability of the observed transitions given the model under consideration, after marginalisation over mRNA. The likelihood difference $L - L_0$ quantifies the evidence for that model. We repeat the above procedure for all 10 runs in the data set, and determine the mean and standard deviation; the outcome is illustrated in Figure 4. We observe that 1000 transitions suffice to correctly refute autorepression in (B,D), and to correctly refute autoactivation in (F). In the case of 100 transitions, no conclusion can be drawn from (A) and (C), while (E) correctly refutes autoactivation; however, the small difference between L and L_0, with $|L - L_0| < 1$, indicates low significance.

5 Discussion

In the present article, we showcase a parameter inference procedure that is based on a recently developed analytical method[14] which allows for the efficient numerical approximation of propagators via the Cauchy integral formula on the basis of asymptotic series for the underlying generating functions. The resulting hybrid analytical-numerical approach reduces the need

Figure 4. Parameter inference for the extended autoregulated model {(3.1),(4.1)} on the basis of various types of synthetic data, where the performance of the model is quantified via the likelihood difference $L - L_0$. On the vertical axis, $L - L_0$ indicates the difference between the likelihood of the autoactivated or autorepressed model and that of the non-regulated model in (3.1); the higher the value of $L - L_0$, the more likely the associated model is. In each panel, the solid curve indicates the mean values based on 10 model runs; dashed curves indicate the uncertainty (one standard deviation). On the horizontal axis, the strength of autoregulation is measured by $\alpha_P \delta$ (increasing to the right) or $\rho_P \delta$ (increasing to the left). (A) Data generated from the null (non-regulated) model in (3.1), with $N = 100$ transitions. (C) Data generated from the model in (3.1), with autoactivation as in (4.1a), for $\alpha_P \delta = 0.3$ and $N = 100$ transitions. (E) Data generated from the model in (3.1), with autorepression as in (4.1b), for $\rho_P \delta = 0.3$ and $N = 100$ transitions. (B,D,F) as (A,C,E), but with $N = 1000$ transitions; note that the vertical axis has a different scaling. All other model parameters were assumed to be known.

for computationally expensive simulations; moreover, due to its perturbative nature, it is highly applicable over relatively short time scales, such as occur naturally in the calculation of the log-likelihood in (1.1).

We model protein expression with a simple stochastic simulation algorithm[17]; the resulting protein abundances are in the lower range of experimentally measured values[20, 21], which does not, however, limit our proof of principle. In addition, in the presented application of our approach, we assume that some of the model parameters are known, that protein abundances can be measured at regular sampling intervals, and that there is no noise. We then present results for synthetic data in a family of models for stochastic gene expression from the literature under the central assumption that lifetimes of protein are significantly longer than those of mRNA, which introduces a small parameter ε and, hence, a separation of scales. An extensive discussion of the validity of our assumption that ε is small can be found elsewhere[9, 14]; see also Section 3. The assumed smallness of ε is crucial to the underlying analytical method, as introduced by Veerman, Marr & Popović[14]. In addition, our approach is specifically attuned to the time variability of the expression process, in the sense that we assume the sampling interval Δt to be small, as well; cf. again Section 3. It is thus ideally suited to describing transient dynamics far from steady state, as is evident in the bursting behaviour seen in Figure 1A.

In Section 3, we discuss a simple ('telegraph') gene expression model without autoregulation, showing that our approach can successfully infer relevant model parameters. Unlike in previous work[22], the underlying implementation avoids potential bias due to zero propagator values and large initial protein numbers through the use of 'implicit' series expansions in ε; see Appendix A for an in-depth argument.

In Section 4, we perform a model comparison in an autoregulated extension of the telegraph model. We consider three types of gene regulation: autoactivation, autorepression, and no regulation of DNA activity (null model). For each type, we simulate protein abundance data with 100 and 1000 transitions, respectively. Throughout, we find that 100 data points are insufficient to reject model hypotheses with our approach. With 1000 data points, however, we can successfully reject the non-regulated and the autorepressed model for simulated data from an autoactivated model, in which case we can even infer the correct order of the autoactivation parameter. For simulated autorepression, we can reject the model with autoactivation, but not the non-regulated model. Our approach fails to identify the correct model for data from a non-regulated model for 1000 transitions, where the autoactivated model is clearly, but wrongly, favoured. We believe that more research is needed into the sources of these discrepancies in dependence on both model parameters and the order of our approximation.

In both showcases, we observe a trade-off between the accuracy of inference versus the required computation time. Computation times seem to increase exponentially with the approximation order, at least for the setup realised in this article. For practical purposes, we hence propose an algorithm whereby the fastest, leading order approximation is used to obtain an initial estimate for the underlying model parameters; that estimate can then be improved by including higher order corrections, resulting in a much more computationally efficient procedure.

It is insightful to compare our results with other recent work on parameter inference in regulated gene expression models. In work by Feigelman et al.[10], three models for regulated gene expression with a slightly different structure compared to the models studied in the present article were simulated and inferred via a particle filtering-based inference procedure that employs genealogical information of dividing cells. Interestingly, positive and negative autoregulation could be successfully rejected there for data that was simulated from a no-feedback model. However, the no-feedback model could not be rejected for data originating from the corresponding models with positive or negative feedback. From that comparison with Feigelman et al.[10], we conclude that the structure of the data, the intensity of regulatory feedback, and the chosen inference procedure together will influence the extent of insight which can be obtained from an inference approach that is based on stochastic models of gene expression.

We emphasise that our analytical method is not restricted to the specific models showcased here. Our aim in the present article is to demonstrate the applicability of that method, and to investigate its performance, rather than to assess the biological validity of a given model. Minimally, our approach can be extended to recent, physiologically more relevant modifications of the telegraph model with autoregulation[18] by Grima, Schmidt & Newman[23] and Congxin et al.[24]; another feasible alternative model can be obtained by introduction of a refractory state[25]. Ideally, we would like to test a variety of stochastic gene expression models against a given set of measurement data. Current computational approaches struggle to provide propagators for models with more than one regulated species, which can often only be approximated even in that simple scenario. The principal advantage of our hybrid approach is that propagators can be evaluated in a computationally efficient manner, via a combination of mathematical analysis and numerical integration[14]; other approaches rely either on the calculation of propagators based on direct numerical simulation of the underlying model – which is computationally demanding – or on the assumption that symbolic derivatives of the generating function are explicitly known, which only holds for specific models of relatively low complexity[13].

The input for our propagator-based approach are the abundances of the involved species, viz. of protein. Thus, we assume that absolute protein numbers are measured, which is in practice hampered by an unknown scaling factor between the observed fluorescence and the corresponding abundances, and by noise. While various suggestions for inferring that factor have been

made[5, 26, 27], and while a linear scaling is customarily assumed[6, 11], an accurate experimental determination remains extremely challenging.

Finally, the showcases presented in this article are based on synthetic data that was generated *in silico*; in the future, we plan to consider experimental data, such as can be found in work by Suter *et al.*[6].

Acknowledgements

The authors thank Ramon Grima and Peter Swain (both University of Edinburgh) for valuable comments and suggestions. This work has been supported by the Leverhulme Trust, through Research Project Grant RPG-2015-017 ('The nature of gene expression: model selection and parameter inference').

Author Contributions Statement

All authors conceived the experiments; F.V. conducted the experiments; F.V. and C.M. analysed the results; F.V. and N.P. wrote the manuscript. All authors reviewed the manuscript.

Additional Information

Competing interests
The authors declare no competing interests.

References

1. Elowitz, M., Levine, A., Siggia, E. & Swain, P. Stochastic gene expression in a single cell. *Science* **297**, 1183–1186, DOI: 10.1126/science.1070919 (2002).

2. van Kampen, N. G. *Stochastic processes in physics and chemistry*, vol. 888 of *Lecture Notes in Mathematics* (North-Holland Publishing Co., Amsterdam-New York, 1981).

3. Crane, M., Clark, I., Bakker, E., Smith, S. & Swain, P. A microfluidic system for studying ageing and dynamic single-cell responses in budding yeast. *PLoS One* **9**, e100042, DOI: 10.1371/journal.pone.0100042 (2014).

4. Filipczyk, A. *et al.* Network plasticity of pluripotency transcription factors in embryonic stem cells. *Nat. Cell Biol.* **17**, 1235–1246, DOI: 10.1038/ncb3237 (2015).

5. Hoppe, P. *et al.* Early myeloid lineage choice is not initiated by random PU.1 to GATA1 protein ratios. *Nature* **535**, 299–302, DOI: 10.1038/nature18320 (2016).

6. Suter, D. *et al.* Mammalian genes are transcribed with widely different bursting kinetics. *Science* **332**, 472–474, DOI: 10.1126/science.1198817 (2011).

7. Zenklusen, D., Larson, D. & Singer, R. Single-RNA counting reveals alternative modes of gene expression in yeast. *Nat. Struct. & Mol. Biol.* **15**, 1263–1271, DOI: 10.1038/nsmb.1514 (2008).

8. Janicki, S. *et al.* From silencing to gene expression: real-time analysis in single cells. *Cell* **116**, 683–698, DOI: 10.1016/S0092-8674(04)00171-0 (2004).

9. Shahrezaei, V. & Swain, P. Analytical distributions for stochastic gene expression. *Proc. Natl. Acad. Sci. United States Am.* **105**, 17256–17261, DOI: 10.1073/pnas.0803850105 (2008).

10. Feigelman, J. *et al.* Analysis of cell lineage trees by exact bayesian inference identifies negative autoregulation of nanog in mouse embryonic stem cells. *Cell Syst.* **3**, 480–490, DOI: 10.1016/j.cels.2016.11.001 (2016).

11. Zechner, C., Unger, M., Pelete, S., Peter, M. & Koeppl, H. Scalable inference of heterogeneous reaction kinetics from pooled single-cell recordings. *Nat. Methods* **11**, 197–202, DOI: doi:10.1038/nmeth.2794 (2013).

12. Schnoerr, D., Sanguinetti, G. & Grima, R. Approximation and inference methods for stochastic biochemical kinetics—a tutorial review. *J. Phys. A* **50**, 093001, 60, DOI: 10.1088/1751-8121/aa54d9 (2017).

13. Popović, N., Marr, C. & Swain, P. A geometric analysis of fast-slow models for stochastic gene expression. *J. Math. Biol.* **72**, 87–122, DOI: 10.1007/s00285-015-0876-1 (2016).

14. Veerman, F., Marr, C. & Popović, N. Time-dependent propagators for stochastic models of gene expression: an analytical method. *J. Math. Biol.* DOI: 10.1007/s00285-017-1196-4 (2017).

15. Raj, A., Peskin, C., Tranchina, D., Vargas, D. & Tyagi, S. Stochastic mRNA synthesis in mammalian cells. *PLoS Biol.* **4**, 1707–1719, DOI: 10.1371/journal.pbio.0040309 (2006).

16. Bornemann, F. Accuracy and stability of computing high-order derivatives of analytic functions by Cauchy integrals. *Foundations Comput. Math.* **11**, 1–63, DOI: 10.1007/s10208-010-9075-z (2011).

17. Gillespie, D. Exact stochastic simulation of coupled chemical reactions. *The J. Phys. Chem.* **81**, 2340–2361, DOI: 10.1021/j100540a008 (1977).

18. Hornos, J. *et al.* Self-regulating gene: An exact solution. *Phys. Rev. E: statistical, nonlinear, biological, soft matter physics* **72**, 051907, DOI: 10.1103/PhysRevE.90.051907 (2005).

19. Iyer-Biswas, S. & Jayaprakash, C. Mixed Poisson distributions in exact solutions of stochastic autoregulation models. *Phys. Rev. E: statistical, nonlinear, biological, soft matter physics* **90**, 052712, DOI: 10.1103/PhysRevE.90.052712 (2014).

20. Schwanhausser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473**, 337–342, DOI: 10.1038/nature10098 (2011).

21. Milo, R. What is the total number of protein molecules per cell volume? a call to rethink some published values. *Bioessays* **35**, 1050–1055, DOI: 10.1002/bies.201300066 (2013).

22. Feigelman, J., Popović, N. & Marr, C. A case study on the use of scale separation-based analytical propagators for parameter inference in models of stochastic gene regulation. *J. Coupled Syst. Multiscale Dyn.* **3**, 164–173, DOI: 10.1166/jcsmd.2015.1074 (2015).

23. Grima, R., Schmidt, D. & Newman, T. Steady-state fluctuations of a genetic feedback loop: An exact solution. *J. Chem. Phys.* **190**, 035104, DOI: 10.1063/1.4736721 (2012).

24. Congxin, L., Cesbron, F., Oehler, M., Brunner, M. & Höfer, T. Frequency modulation of transcriptional bursting enables sensitive and rapid gene regulation. *Cell Syst.* **6**, 409–423, DOI: 10.1016/j.cels.2018.01.012 (2018).

25. Zoller, B., Nicolas, D., Molina, N. & F., N. Structure of silent transcription intervals and noise characteristics of mammalian genes. *Mol. Syst. Biol.* **11**, 823, DOI: 10.15252/msb.20156257 (2015).

26. Rosenfeld, N., Perkins, T., Alon, U., Elowitz, M. & Swain, P. A fluctuation method to quantify in vivo fluorescence data. *Biophys. J.* **91**, 759–766, DOI: 10.1529/biophysj.105.073098 (2006).

27. Bakker, E. & Swain, P. Estimating numbers of fluorescent molecules in single cells by analysing fluctuations in photobleaching. *bioRxiv* **272310**, DOI: 10.1101/272310 (2018).

<div style="text-align:center">

APPENDIX TO

PARAMETER INFERENCE WITH ANALYTICAL PROPAGATORS FOR STOCHASTIC MODELS OF AUTOREGULATED GENE EXPRESSION

FRITS VEERMAN, NIKOLA POPOVIĆ, AND CARSTEN MARR

</div>

A. ANALYTICAL DETAILS

The hybrid analytical-numerical approach developed by Veerman, Marr & Popović [2] introduces a probability generating function which transforms the CME corresponding to a given stochastic gene expression model into a system of partial differential equations. Explicit expressions for the solutions to these equations are obtained using dynamical systems techniques, in combination with perturbation theory. The values of the associated propagators are recovered through a numerically efficient implementation of the Cauchy integral in (2.1); see also Section 2.

A.1. **Approximate generating functions.** The generating functions used to approximate propagators in the present article, cf. Section 2, are derived via the analytical method presented by Veerman, Marr & Popović [2]. For the telegraph model in Section 3, the leading order generating function $\widehat{F}_0$ is given by

$$(\text{A.1}) \qquad \widehat{F}_0(z; \Delta t, n_0, \varepsilon, \lambda, \mu, \chi) = \left[1 - (1-z)\mathrm{e}^{-\Delta t}\right]^{n_0} {}_1\mathrm{F}_1\left[\kappa_0, \kappa_0 + \kappa_1, \varepsilon\lambda\frac{1 - f(z)}{f(z)}\left(1 - \mathrm{e}^{-f(z)\frac{\Delta t}{\varepsilon}}\right)\right];$$

here, $\chi = \frac{\kappa_1}{\kappa_0 + \kappa_1}$ and $f(z) = 1 + \mu(1-z)\mathrm{e}^{-\Delta t}$. All parameters have been rescaled according to (3.2). The generating function has been marginalised over mRNA, assuming a steady-state distribution reported by Raj et al. [1]. Analogously, the first order approximation $\widehat{F}_1$ of the generating function is given by

(A.2)

$$\widehat{F}_1(z; \Delta t, n_0, \varepsilon, \lambda, \mu, \chi) = \left[1 - (1-z)\mathrm{e}^{-\Delta t}\right]^{n_0}\left[1 + \varepsilon(1-\chi)\lambda\frac{1 - f(z)}{f(z)}\mathrm{e}^{-f(z)\frac{\Delta t}{\varepsilon}}\left(\frac{1 - \mathrm{e}^{f(z)\frac{\Delta t}{\varepsilon}}}{f(z)} + \frac{\Delta t}{\varepsilon}\mathrm{e}^{f(z)\frac{\Delta t}{\varepsilon}}\right)\right]$$

$$\times {}_1\mathrm{F}_1\left[\kappa_0, \kappa_0 + \kappa_1, \varepsilon\lambda\frac{1 - f(z)}{f(z)}\left(1 - \mathrm{e}^{-f(z)\frac{\Delta t}{\varepsilon}}\right)\right]\left\{1 + \varepsilon\left[\frac{1}{f(z)^2} + \frac{1}{f(z)}\frac{\Delta t}{\varepsilon} + (1 - f(z))\frac{\Delta t^2}{2\varepsilon^2}\right]\right\}\right].$$

For the autoregulated model discussed in Section 4, the same expressions for the generating functions are used; however, χ now depends on the autoregulatory mechanism according to

$$(\text{A.3}) \qquad \chi = \begin{cases} \dfrac{\kappa_1}{\kappa_0 + \kappa_1} & \text{(no autoregulation)}, \\[1em] \dfrac{\kappa_1}{\kappa_0 + \kappa_1 + \alpha_P \delta\, n_0} & \text{(autoactivation)}, \\[1em] \dfrac{\kappa_1 + \rho_P \delta\, n_0}{\kappa_0 + \kappa_1 + \rho_P \delta\, n_0} & \text{(autorepression)}. \end{cases}$$

A.2. **'Implicit' expansions.** It is important to note that neither the leading order generating function in (A.1) nor the first order approximation given by (A.2) are expressed as asymptotic series in powers of ε, as would be expected on the basis of the perturbative approach taken by Veerman, Marr & Popović [2]. The underlying reasoning can be summarised as follows.

First, in the derivation of the generating functions, it was assumed that the sampling time Δt was small, i.e. of order ε; note that this assumption is satisfied in all numerical simulations shown in the current article, where $\Delta t = \varepsilon$ throughout. Thus, we can write

$$(\text{A.4}) \qquad \Delta t = \varepsilon \Delta s.$$

With the above scaling for Δt, an expansion of $\widehat{F}_0$ and $\widehat{F}_1$, as defined in (A.1) and (A.2), respectively, into asymptotic series in ε to the appropriate order yields

(A.5) $\qquad \widehat{F}_0 = z^{n_0},$

(A.6) $\qquad \widehat{F}_1 = z^{n_0}\left\{1 + \varepsilon(1-z)\left[\dfrac{n_0\Delta s}{z} - \dfrac{(1-\chi)\lambda\mu}{1+\mu(1-z)}\left(\dfrac{\mu(1-z)\left\{1-\mathrm{e}^{[-(1+\mu(1-z)]\Delta s}\right\}}{1+\mu(1-z)} + \Delta s\right)\right]\right\}.$

From (A.5), one can immediately conclude that

(A.7) $$\oint_\gamma \widehat{F}_0 = \delta_{n,n_0},$$

where $\delta_{n,n_0} = 1$ if $n = n_0$ and $\delta_{n,n_0} = 0$ otherwise. From the series for $\widehat{F}_1$ in (A.6), we see that we can write

(A.8) $$\widehat{F}_1(z) = z^{n_0}\left\{1 + \varepsilon\sum_{k=-1}^{\infty} f_{1,k}z^k\right\},$$

with appropriately chosen coefficients $f_{1,k}$; hence, it follows that

(A.9) $$\oint_\gamma \widehat{F}_1 = 0 \quad \text{if } n_0 > n+1.$$

More generally, an expansion of the generating function to order k in ε will yield

(A.10) $$\oint_\gamma \widehat{F}_k = 0 \quad \text{if } n_0 > n+k.$$

From these observations, we conclude that decreasing transitions $(n_i \to n_{i+1})$, where $n_i > n_{i+1} + k$, will be assigned a probability that is identically zero. Hence, if such transitions *are* present in the data, the model is ruled out immediately, as our perturbative approach excludes the possibility that decreasing transitions can occur. One can understand this phenomenon by considering the definition of the small parameter ε as the ratio of the protein decay rate d_1 over the mRNA decay rate d_0. A leading order approximation of $\varepsilon = 0$ is thus equivalent to taking $d_1 \to 0$ which, in turn, implies that protein does not decay at all, since (natural) protein decay is the only reaction in (3.1) that can decrease protein abundance. By the same reasoning, a straightforward expansion of the generating function to order ε^k will restrict the model to transitions $(n_i \to n_{i+1})$, where $n_{i+1} - n_i \geq -k$. It would follow that either the order O of the method would be limited from below by the data, leading to high-order expansions in ε and, hence, to increased computation times, or that the method could only be applied to a subset of the given data, which would introduce a bias.

Lastly, an asymptotic expansion such as (A.6) implicitly assumes that all parameters and variables in the model are of order 1 in ε. For the series expansion of $\widehat{F}_1$ in (A.6), that assumption would significantly restrict the range of λ; in comparison, in Figure 2, likelihood values for λ up to order ε^{-3} are calculated. More importantly, the above assumption would restrict the range of n_0, implying that only a subset of the data – with sufficiently low protein numbers – could be used as input for parameter inference.

We emphasise that none of these difficulties occur with the expressions in (A.1) and (A.2), where the expansion order in ε is expressed 'implicitly' in the respective functional forms of $\widehat{F}_0$ and $\widehat{F}_1$.

References

[1] A. Raj, C.S. Peskin, D. Tranchina, D.Y. Vargas, and S. Tyagi. Stochastic mRNA synthesis in mammalian cells. *PLoS Biology*, 4(10):1707–1719, 2006.

[2] F. Veerman, C. Marr, and N. Popović. Time-dependent propagators for stochastic models of gene expression: an analytical method. *Journal of Mathematical Biology*, DOI: 10.1007/s00285-017-1196-4, 2017.