Title: Empowering conservation practice with efficient and economical genotyping from poor

quality samples using mPCRseq

Authors: MeghanaNatesh[1,2 #], Ryan W. Taylor[3,4,#], Nathan Truelove[5],Elizabeth A. Hadly[3],
Stephen Palumbi[3,5], Uma Ramakrishnan[1#*] and Dmitri Petrov[3#*]

Author Affiliations:

[1]National Centre for Biological Sciences, TIFR, Bellary Road, Bangalore - 560065, India
[2]Sastra University, Tirumalaisamudram, Thanjavur – 613401, India

[3]Department of Biology, Stanford University, Stanford, CA - 94305, USA

[4]End2End Genomics LLC, Davis, CA

[5]Hopkins Marine Station, Stanford University, Pacific Grove, CA – 93950, USA

#: Equal contribution
12
13   email addresses: meghanan@ncbs.res.in, ryan@ryantaylor.net, truelovn@stanford.edu,
14   hadly@stanford.edu, spalumbi@stanford.edu, uramakri@ncbs.res.in, dpetrov@stanford.edu

15   Running title: Genetic data to empower conservation
16
17   Keywords: conch, conservation genetics, endangered species, monitoring, new methods, next
18   generation sequencing, non-invasive samples, SNPs, tigers
19
20   Type of article: Policy Perspectives
21
22   Number of words in the abstract:*196*
23   Number of words in the manuscript:*2548*
24   Number of references:*12*
25   Number of figures and tables: *Figure 3, Tables 0*
26

27   *Correspondence to: Uma Ramakrishnan, Associate Professor, Wellcome/DBT India Alliance
28   Senior Fellow, National Centre for Biological Sciences, TIFR, Bangalore, 560065, India, +91-80-
29   23666030, uramakri@ncbs.res.in
30   and Dmitri Petrov, Professor, Department of Biology, 371 Serra St., Stanford University
31   Stanford, CA 94305-5020, USA, +1-(650) 736 1169, dpetrov@stanford.edu

32
33

Abstract: *Genetic tools are widely used in conservation but expansion to genomic scale data*

that are rapid and cost effective has been slow. Most genomic tools are developed for high

quality DNA sources from lab or medical settings. So far, genetic data from market or field

settings assess easily amplified mitochondrial DNA or a few microsatellites. Here we use

multiplex PCR with low quality DNA from feces, hair, and cooked samples to quickly provide

multi-locus data from many SNPs. We demonstrate the wide range of potential applications

through tools to monitor individual wild tigers and track commercial trade in Caribbean queen

conch. 100 SNPs from degraded tiger samples identified individuals, discerned close relatives,

and detected population differentiation. 62 SNPs from conch fritters and field collected samples

identified individuals, tested for close kin and detected population structure. Our study provides

proof of concept for a rapid, simple, cost-effective, and scalable method, a framework that can

be applied to other conservation scenarios previously limited by low quality DNA samples. These

approaches provide a critical advance for wildlife monitoring and forensics, open the door to

field-ready testing, and will strengthen the use of science in policy decisions and wildlife trade.

Introduction

Stemming the tide of global species decline requires continuous monitoring and nimble,

adaptive management to promote species recovery. Effective monitoring requires, minimally,

the ability to identify and track presence of a species and ultimately to track specific individuals

and to determine their familial relationships. Integrating local data across species range

provides the ability to monitor global threats from population range reduction to illegal wildlife

trade.

In principle, all of these goals can be achieved via genotyping at a modest number of loci

such as microsatellites or single nucleotide polymorphisms (SNPs). Given that endangered

species tend to be rare and elusive, the necessary approach must be able to accommodate non-

invasive sources of DNA such as feces, shell, feathers, hair, and saliva that yield impure, mixed,

and/or extremely small amounts of degraded DNA. Moreover, market samples generated by

wildlife trade may be processed, cooked, dried, or mixed with other species, again providing

low quality and often mixed DNA. In contrast, the current approaches tend to require large

amounts of high quality, pure DNA from the target species most readily obtained by

immobilizing or sacrificing an animal (Fitak, Naidu, Thompson, & Culver, 2016; Kraus et al.,

2015) or demand expensive and generally inefficient enrichment strategies (Chiou & Bergey,

2018; Snyder-Mackler et al., 2016) to enhance the quantity of the target DNA. As a result, these

approaches are unsuitable, unaffordable, or too low throughput in most conservation settings.

The tiger (*Panthera tigris*) is one of the most endangered carnivores and despite drastic

habitat reduction, the remaining ~5,000 wild tigers inhabit over 14 countries. Their wide

distribution makes it critical that locally collected data is comparable across their range. While

photographic mark-recapture is commonly used to monitor tigers, capture rates are poor in

low-density habitats. Neither can photographs always be matched with confiscated tiger parts.

Instead, genotyping of scat or hair, along with rapid forensic testing of confiscated skins or

other traded parts can verify species, individual identity, and source populations.

Although formerly abundant, the queen conch (*Strombus gigas*) was listed in CITES

Appendix II in 1990. Trade was further restricted in 2004 due to declining stocks. Florida is

currently the largest market for queen conch seafood products, consuming 82% of international

trade. Matching the populations of origin of queen conch in Florida markets with international

regulations is crucial to recover this formerly lucrative fishery. The most direct access to

imported conch is from the hundreds of restaurants in Florida, necessitating techniques that

would allow genotyping from the most abundant menu item, fried conch fritters.

Here we demonstrate that a multiplex PCR approach (Campbell, Harmon, & Narum,

2015) followed by next-generation sequencing satisfies all the requirements necessary for

inexpensive, fast, and easy genotyping of low-quality samples. We illustrate the power of this

method for these two endangered species in very divergent conservation contexts and real-life

settings: genotypes from fecal, shed hair, and saliva samples found on the killed prey from wild

Indian tigers and from CITES-regulated Caribbean queen conch imported to the US and sold in

fried fritters.

Methods

We designed primers for 192 SNP loci from a pool of 50,000 SNPs discovered through

whole genome sequencing of 75 tigers of captive and wild origin from *P. t. tigris, P. t. jacksoni,*

P. t. altaica, and P. t. sumatrae subspecies (see supplementary materials). The pool consisted of

SNPs that were polymorphic in all subspecies (> 10% MAF), or differentiated subspecies from

each other (see supplementary materials). We used the publicly available program Primer3 to

design primer pairs (amplicon size 50-90 bp, primer size 17-25 bp, and Tm of 60-61ºC).The

resulting 192 primers were pooled for use in a single amplification of all target SNPs for each

tiger sample. The initial primer set was tested on fecal and tissue samples from captive tigers

(Table S1). The test set included fecal and tissue samples from the same individuals, multiple

subspecies, samples left to weather for up to two days, and a serially diluted tissue DNA

sample. Tiger DNA was quantified (see supplementary materials) and all samples were

genotyped in triplicate. SNPs were then tested on a range of non-invasive samples from across

India (representative photographs in Fig. 1b, sample list in Table S2). Multiple locations and

differing tiger landscapes, two sympatric carnivore species, the leopard (*Panthera pardus*) and

Indian wild dog or dhole *(Cuon alpinus*) were genotyped. Fecal samples, saliva and shed hair

were included.

The tiger example builds on a large number of SNPs ascertained from multiple fully

sequenced genomes. Often such resources are not available for endangered species, although

this is changing rapidly. For queen conch, we sought to identify SNPs from the transcriptomes

of 83 tissue samples from six populations in the Caribbean including Florida, collected from

fishing operations or non-lethally by research permit (see Table S3). Following methods

paralleling those used for tigers, we then genotyped at 192 SNPs an additional 237 conch (from

14 populations) and as well as 42 putatively conch samples isolated from 15 conch fritters

purchased at several Miami restaurants (see supplementary materials).

Results

SNP genotyping for wild tigers

Of the 192 SNPs, 126 produced the most consistent results (see supplementary materials for details, Fig. S1, Table S4), and this panel of 126 SNP panel had a high overall genotyping success rate (Fig. 2). An average of 95 SNPs (range: 4 - 114) were successfully typed across all samples. When tiger DNA was > 0.01ng/ul (irrespective of sample type) an average of 105 SNPs (range: 48 - 114) were typed.12 SNPs did not amplify for over 50% of samples and were excluded from subsequent analysis. The dhole sample had very poor amplification success (mean across replicates <10 SNPs) as expected. Leopards had very high amplification success (mean: 110 SNPs), but the SNPs were monomorphic and identical across two individuals (mean: 0.03% mismatches). All tiger samples in contrast had distinct genotypes and could be readily distinguished from the leopard samples.

Replications of 'within-sample' and 'within individual-between tissue types' provided important insights on the precision and accuracy of SNP identification and its efficacy for non-invasive sampling. The proportion of matches between tissue types for the same individual (n=5) was very high (precision, mean 0.97; range 0.91 - 0.99), showing that our approach resulted in nearly identical genotypes for different sample types (feces, saliva, hair). Sample replicates (within-sample comparisons) were also highly concordant and had a high proportion of matches (precision, mean: 0.957, range: 0.757 - 1.0). Among these, 77 samples genotyped at least 60% of SNPs had even higher concordance across replicates (mean proportion of matches: 0.969, range: 0.86-1.0), with only 5 samples showing values less than 0.95. These results are in sharp contrast to the high error rates observed for microsatellite genotypes from other non-

invasive sampling techniques (Coates et al., 2011). Our probability of misidentifying two

different individuals as the same is vanishingly low (pID = 1.6E-22 assuming a population of

siblings).

Observed relatedness values (Fig. 3) and clustering of samples (Fig. S2) further highlight

our ability to detect biological patterns. Our data had two pairs of samples with known

relationships (zoo samples) while the remaining were sampled from different populations in the

wild. The known parent offspring and sibling pairs had relatedness values close to the expected

value of 0.5 (Figure 3). Observed pair-wise relatedness was higher within than between known

genetic clusters or populations (see Natesh et al., 2017). As expected, relatedness between

individuals from a small, isolated population (NW, Figure 3) was high. Even with the few

individuals in our pilot dataset presented here, wild individuals fell into three genetic

clusters/populations (Figure S2), similar to previous data (Natesh et al., 2017). For small

isolated populations (such as NW) future genotyping could help identify whether individuals are

also inbred, and if such inbreeding has fitness consequences through estimates of individual

reproductive success. Should genetic supplementation of this population be required, SNP data

from different tiger landscapes could help identify possible sources.

SNPs genotyping for queen conch

Our second example illustrates that this multiplex PCR approach can be applied even if

background genetic information is not as complete (as with tiger), and again highlights use in

difficult samples: in this case fried conch fritters. We identified 480,962 SNPs from the

sequenced transciptomes. Possibly because we based our primer design on a transcriptome

rather than a genome, our SNP success was lower for conch than tigers. Approximately half the

192 conch primer pairs failed to provide data. For the best 62 SNPs success was about the same

as for tigers (average 86%, Fig. S3). Replicate samples shared 99%-100% of their alleles, but we

also found several genetically identical samples among small processed conch pieces collected

from fishing boats. As with genetic re-sampling of tigers in the wild, discovery of repeated

individuals in a sample can inform overall fishery numbers. There were no obvious close

relatives among the samples, a result expected for low sample sizes and very large populations

(Fig. S3). The data also inform basic information important in design of marine protected areas.

Outlying islands (Aruba, St. Eustatius) were genetically differentiated from the central

Caribbean and Florida (F_{ST} = 0.037, 0.048 respectively, Table S5). Samples from the same island

group/on the same coast (e.g. Florida, Bahamas) were not differentiated. These patterns

parallel a recent survey of queen conch with microsatellites (Truelove et al., 2017).

While the conch isolated from deep fried fritters showed lower success rates than conch

sampled by fishermen or from biopsies, the success rates were high enough to allow individual

identification and initial population comparison. Twenty-three of the fritter conch did not

amplify with any primer, or produced data for less than 18 SNPs, and we suspect they were not

Queen conch. A further 8 fritter conch showed poor SNP amplification (average 41%) but could

be confirmed as Queen conch samples. The most successful 17 conch from fritters showed

success rates comparable to those of fresh samples (77% vs 86%) (Fig. S3). Even with this

preliminary data set, we can determine that the fritter samples showed lower (average) genetic

identity to Florida populations (average 78.6 – 80.5 alleles shared, Fig. S6)compared to the

Puerto Rico or St. Eustatius (81.1-82.3, Fig. S6). Positive identification of populations for

imported samples will require greater geographic sampling, a larger SNP panel, and a better

genotype map of potential source locations. For example, assignment testing successfully

places 65% of Bahamas conch samples back into that population, but this is not yet high enough

for precise population discovery. Nevertheless, we can already suggest that the Florida Keys

and Nassau are less likely to be the populations of origin for the fritters we genotyped, and

show that high resolution nuclear data can be readily obtained even from deeply processed

commercial samples and used to address key conservation needs.

Discussion

Our results on these small, pilot datasets provide proof of concept for multiplex SNP

genotyping of non-invasive and processed market samples from two species with vastly

different physiologies, biologies, ecologies, and conservation challenges. This approach satisfies

the key requirements for a practical conservation application. It is inexpensive to scale up, we

estimate that implementation costs (for upto 1000 SNPs) can be as low as $5 per sample. The

initial costs for primer design might include full-genome sequencing, which is becoming more

and more affordable. The cost of a genome of reasonable quality, even for fairly large

mammalian genomes, in the range of a few thousand dollars (Armstrong et al., 2017). In

addition, primers based on RADSeq data or RNAseq data provide cheaper solutions for very

large genomes. The largest development cost is of primer synthesis, which is expected to go

down as high-throughput oligo synthesis is becoming a large-scale, commercial commodity

service. Lab equipment required of the approach is minimal (centrifuge, thermocycler,

pipettes), and field-based sequencers are allowing the use of this method anywhere in the field.

Increasing the number of SNPs beyond a few hundred can provide additional

information. For instance, even with a few thousand SNPs it should be possible to start fully

imputing large regions of the genome in cases where full genomes of individuals in the pedigree

can be obtained. Moreover, strategic design of primers that would amplify clusters of closely

located SNPs should allow detection of very recent inbreeding by detecting long contiguous

runs of homozygosity (Kirin et al., 2010). Some of the SNPs could be located in such close

proximity that they generate minihaplotypes that have been shown to be particularly useful in

pedigree reconstruction (Baetscher, Clemento, Ng, Anderson, & Garza, 2018). The SNP panels

can be further designed to allow simultaneous species, individual, and diet identification for

clusters of species, such as, for instance, all large carnivores and all common species in their

diet in India or sub-Saharan Africa.

Most importantly, our data show that multiplex PCR is successful for degraded, cooked,

mixed, or small and low-concentration samples, making it an ideal tool for monitoring

individuals in populations under rigors of field conditions or commercial markets. In the future,

when populations maybe very small and isolated, we may need the ability to impute whole

genomes, construct full pedigrees, and conduct genome-wide association studies for

detrimental or advantageous traits. In principle, this method will allow such advances.

Effective monitoring of individuals, populations and species is critical to designing rapid

conservation action and management. In the future, conservation will depend critically on such

tools. Small, isolated populations will require inbreeding management, genotype tracking for

population assessment, and data to assess when individuals should be introduced. Populations

across fragmented landscapes may require assisted migration, especially in the context of

climate change and the range shifts it demands. Minimizing illegal wildlife trade will require

enforcement based on genetically identified sources of trafficked animals. Rapid genetic

monitoring of endangered species from commonly occurring non-invasive samples will provide

a data-pathway to species recovery. We believe that multiplex PCR presents an example of such

rapid, accessible, cheap and efficient technology that will make this possible.

References

Armstrong, E. E., Taylor, R. W., Prost, S., Blinston, P., Der, E. Van, Madzikanda, H., … Petrov, D.

(2017). Entering the era of conservation genomics: Cost-effective assembly of the African

wild dog genome using linked long reads. *BioRxiv*.

http://doi.org/http://dx.doi.org/10.1101/195180.

Baetscher, D. S., Clemento, A. J., Ng, T. C., Anderson, E. C., & Garza, J. C. (2018).

Microhaplotypes provide increased power from short-read DNA sequences for relationship

inference. *Molecular Ecology Resources*, *18*(2), 296–305. http://doi.org/10.1111/1755-

0998.12737

Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by sequencing

(GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing.

Molecular Ecology Resources, *15*(4), 855–867. http://doi.org/10.1111/1755-0998.12357

Chiou, K. L., & Bergey, C. M. (2018). Methylation-based enrichment facilitates low-cost,

noninvasive genomic scale sequencing of populations from feces. *Scientific Reports*, *8*(1),

1–10. http://doi.org/10.1038/s41598-018-20427-9

Coates, B. S., Bayles, D. O., Wanner, K. W., Robertson, H. M., Hellmich, R. L., & Sappington, T.

W. (2011). The application and performance of single nucleotide polymorphism markers

for population genetic analyses of lepidoptera. *Frontiers in Genetics*, *2*(JUNE), 1–10.

http://doi.org/10.3389/fgene.2011.00038

Fitak, R. R., Naidu, A., Thompson, R. W., & Culver, M. (2016). A New Panel of SNP Markers for

the Individual Identification of North American Pumas. *Journal of Fish and Wildlife*

Management, *7*(1), 13–27. http://doi.org/10.3996/112014-JFWM-080

Kirin, M., McQuillan, R., Franklin, C. S., Campbell, H., Mckeigue, P. M., & Wilson, J. F. (2010).

Genomic runs of homozygosity record population history and consanguinity. *PLoS ONE*,

5(11), 1–7. http://doi.org/10.1371/journal.pone.0013996

Kraus, R. H. S., vonHoldt, B., Cocchiararo, B., Harms, V., Bayerl, H., Kühn, R., … Nowak, C. (2015).

A single-nucleotide polymorphism-based approach for rapid and cost-effective genetic

wolf monitoring in Europe based on noninvasively collected samples. *Molecular Ecology*

Resources, *15*(2), 295–305. http://doi.org/10.1111/1755-0998.12307

Natesh, M., Alta, G., Nigam, P., Jhala, Y. V, Zachariah, A., Borthakur, U., & Ramakrishnan, U.

(2017). Conservation priorities for endangered Indian tigers through a genomic lens.

Scientific Reports, *7*(1), 1–11. http://doi.org/10.1038/s41598-017-09748-3

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. a R., Bender, D., … Sham, P. C.

(2007). PLINK: a tool set for whole-genome association and population-based linkage

analyses. *American Journal of Human Genetics*, *81*(3), 559–75.

http://doi.org/10.1086/519795

Snyder-Mackler, N., Majoros, W. H., Yuan, M. L., Shaver, A. O., Gordon, J. B., Kopp, G. H., …

Tung, J. (2016). Efficient Genome-Wide Sequencing and Low Coverage Pedigree Analysis

from Non-invasively Collected Samples. *Genetics*, *203*(June), 699–714.

http://doi.org/10.1534/genetics.116.187492

Truelove, N. K., Box, S. J., Aiken, K. A., Blythe-Mallett, A., Boman, E. M., Booker, C. J., … Stoner,

A. W. (2017). Isolation by oceanic distance and spatial genetic structure in an

overharvested international fishery. *Diversity and Distributions*, *23*(11), 1292–1300.

http://doi.org/10.1111/ddi.12626

Acknowledgements: We thank San Diego Zoo Global, Conservation Society of California at

Oakland Zoo, Oakland, CA, San Francisco Zoo, the Bronx Zoo, WCS Russia program, El Paso Zoo,

Performing Animal Welfare Society, Wildlife Conservation Society India Program, Centre for

Wildlife Studies, India, Wildlife Conservation Trust, Himanshu Chhattani, Anubhab Khan, Prachi

Thatte, Aditya Joshi, Arun Zachariah, Erica Calcagno and Jackie Gai for tiger samples. We thank

the Forest Departments of Rajasthan, Karnataka and Madhya Pradesh and the American Zoo

Association, Tiger SSP for permissions. Awadhesh Pandit and NCBS next generation genomics

facility helped with Illumina sequencing. We thank Steve Box, and Steve Canty for logistical help

and support in framing conservation questions for conch. **Funding:** R43HG009482, Wildlife

Conservation Trust grant to UR, Fulbright Nehru Academic exchange Program (UR) and

Wellcome Trust/DBT India Alliance senior fellowship (UR), Summit Foundation and the Global

Genome Initiative at the Smithsonian Institution (to NT and SRP and Steve Box) supported

conch work. Author Contributions: MN, RWT, DP, UR, EAH, SP and NT designed the study. MN,

RWT and NT conducted lab work and analyses. MN, RWT, DP, UR, EAH, SP wrote the paper.

Competing interests: Ryan W. Taylor and End2End Genomics LLC have received NIH small

business funding (R43HG009482) to develop molecular tools for the study of non-model

species. The other authors declare that we have no competing interests. **Data Availability:** On

publication, the data will be made available on the Dryad repository.

Figures

Figure 1a: A schematic showing the protocol and approximate time taken for each step.

Figure 1b: A panel of photographs showing the conservation setting for tiger and conch genomics. On the left, the top panel shows a tiger defecating (courtesy Himanshu Chattani), the middle panel a tiger scrape mark on a tree associated with hair samples (courtesy Kaushal Patel). On the right, the top panel shows a conch emerging from its shell and the middle a fisherman holding his catch.

Figure 2: SNP typing success (in triplicate) for various tiger samples and controls. The concentration of each sample is shown on the right. Empty cells in a particular row correspond to SNPs that could not be genotyped. Filled cells indicate successfully typed SNPs, and the color indicates sample type.

Figure 3: Pairwise relatedness (PI_HAT; (Purcell et al., 2007)) captures known relationships, and provides insights into wild tiger relatedness. 'Self' are pairwise comparisons among the three replicate analyses of each sample, as well as comparisons between different sample types for the same individual. Our dataset includes one pair of siblings (Sibs) and one parent-offspring pair (P-O). Sibs and P-O relationships were provided by zoo records. W-P are within population relatedness between samples from previously identified populations (Natesh et al., 2017). B-P are between population relatedness. Values in W-P and B-P are from field samples, or samples from wild caught captive tigers, where the relatedness is unknown. The inset shows sampling locations of the wild samples.