A data-driven interactome of synergistic genes improves network based cancer outcome prediction

Amin Allahyar:

Department of Genetics, Center for Molecular Medicine, University Medical Center Utrecht, Utrecht

University, 3584 CG Utrecht, The Netherlands.

Delft Bioinformatics Lab, Faculty of Electrical Engineering, Mathematics and Computer Science, Delft

University of Technology, 2628 CD Delft, The Netherlands.

a.allahyar@tudelft.nl

Joske Ubels:

Department of Genetics, Center for Molecular Medicine, University Medical Center Utrecht, Utrecht

University, 3584 CG Utrecht, The Netherlands.

Skyline DX, Marconistraat 16, 3029 AK Rotterdam.

Department of Hematology, Erasmus MC Cancer Institute, Rotterdam.

J.Ubels@skylinedx.com

Jeroen de Ridder (Corresponding author):

Department of Genetics, Center for Molecular Medicine, University Medical Center Utrecht, Utrecht

University, 3584 CG Utrecht, The Netherlands.

J.deRidder-4@umcutrecht.nl

Abstract

Robustly predicting outcome for cancer patients from gene expression is an important challenge

on the road to better personalized treatment. Network-based outcome predictors (NOPs), which

considers the cellular wiring diagram in the classification, hold much promise to improve

performance, stability and interpretability of identified marker genes. Problematically, reports on

the efficacy of NOPs are conflicting and for instance suggest that utilizing random networks

performs on par to networks that describe biologically relevant interactions. In this paper we turn

the prediction problem around: instead of using a given biological network in the NOP, we aim to

identify the network of genes that truly improves outcome prediction. To this end, we propose

SyNet, a gene network constructed ab initio from synergistic gene pairs derived from survival-

labelled gene expression data. To obtain SyNet, we evaluate synergy for all 69 million pairwise

combinations of genes resulting in a network that is specific to the dataset and phenotype under

study and can be used to in a NOP model. We evaluated SyNet and 11 other networks on a

compendium dataset of >4000 survival-labelled breast cancer samples. For this purpose, we used

cross-study validation which more closely emulates real world application of these outcome

predictors. We find that SyNet is the only network that truly improves performance, stability and

interpretability in several existing NOPs. We show that SyNet overlaps significantly with existing

gene networks, and can be confidently predicted (~85% AUC) from graph-topological descriptions

of these networks, in particular the breast tissue-specific network. Due to its data-driven nature,

SyNet is not biased to well-studied genes and thus facilitates post-hoc interpretation. We find that

SyNet is highly enriched for known breast cancer genes and genes related to e.g. histological

grade and tamoxifen resistance, suggestive of a role in determining breast cancer outcome.

Keywords: Cancer outcome prediction; Network based models; Cross-study performance

Author Summary

Cancer is caused by disrupted activity of several pathways. Therefore, outcome predictors

analyze patient's expression profiles from perspective of gene groups collected from interactomes

(e.g. protein interaction networks). These Network based Outcome Predictors (NOPs) hold

potential to facilitate identification of dysregulated pathways and delivering improved prognosis.

Nonetheless, recent studies revealed that compared to classical models, neither performance nor

consistency can be improved using NOPs.

We argue that NOPs can only perform well under guidance of suitable networks. The commonly

used networks may miss associations specially for under-studied genes. Additionally, these

networks are often generic with low resemblance to perturbations that arise in cancer.

To address this issue, we exploit ~4100 samples and infer a disease specific network called SyNet

linking synergistic gene pairs that collectively show predictivity beyond individual performance of

genes.

Using identical datasets, we show that a NOP yields superior performance merely by considering

groups of genes in SyNet. Further, NOP performance severely reduces if SyNet nodes are

shuffled, confirming relevance of SyNet links.

Due to simplicity of our approach, this framework can be used for any phenotype of interest. Our

findings represent the value of network-based models and crucial role of interactome in their

performance.

68 1 Introduction

Metastases at distant sites (e.g. in bone, lung, liver and brain) is the major cause of death in

breast cancer patients [1]. However, it is currently difficult to assess tumor progression in these

patients using common clinical variables (e.g. tumor size, lymph-node status, etc.) [2]. Therefore,

for 80% of these patients chemotherapy is prescribed [3]. Meanwhile, randomized clinical trials

showed that at least 40% of these patients survive without chemotherapy and thus unnecessarily

have to suffer from the toxic side effect of this treatment [3,4]. For this reason, substantial efforts

have been made to derive molecular classifiers that can predict clinical outcome based on gene

expression profiles obtained from the primary tumor at the time of diagnosis [5,6].

An important shortcoming in molecular classification is that 'cross-study' generalization is often

poor [7]. This means that prediction performance decreases dramatically when a classifier trained

on one patient cohort is applied to another one [8]. Moreover, the gene signatures found by these

classifiers vary greatly, often sharing only few or no genes at all [9–11]. This lack of consistency

casts doubt on whether the signatures capture true 'driver' mechanisms of the disease or rather

subsidiary 'passenger' effects [12].

Several reasons for this lack of consistency have been proposed, including small sample size

[11,13,14], inherent measurement noise [15] and batch effects [16,17]. Apart from these technical

explanations, it is recognized that traditional models ignore the fact that genes are organized in

pathways [18]. One important cancer hallmark is that perturbation of these pathways may be

caused by deregulation of disparate sets of genes which in turn complicates marker gene

discovery [19,20].

To alleviate these limitations, the classical models are superseded by Network-based Outcome

Prediction (NOP) methods which incorporate gene interactions in the prediction model [21]. NOPs

have two fundamental components: aggregation and prediction. In the aggregation step, genes

that interact, belong to the same pathway or otherwise share functional relation are aggregated

(typically by averaging expressions) into so called "meta-genes" [22]. This step is guided by a

secondary data source describing gene-gene interactions such as cellular pathway maps or

protein-protein interaction networks. In the consequent prediction step, meta-genes are selected

and combined into a trained classifier, similar to a traditional classification approach. Several

NOPs have been reported to exhibit improved discriminative power, enhanced stability of the

classification performance and signature and better representation of underlying driving

mechanisms of the disease [18,23–25].

In recent years, a range of improvements to the original NOP formulation has been proposed. In

the prediction step, various linear and nonlinear classifiers have been evaluated [26,27].

Problematically, the reported accuracies are often an overestimation as many studies neglected

to use cross-study evaluation scheme which more closely resembles the real world application of

these models [7]. Also for the aggregation step which is responsible for forming meta-genes from

gene sets several distinct approaches are proposed, such as clustering [23] and greedy

expansion of seed genes into subnetworks [18]. Moreover, in addition to simple averaging,

alternative means by which genes can be aggregated, such as linear or nonlinear embeddings,

have been proposed [17,28]. Most recent work combines these steps into a unified model [8,29].

Despite these efforts and initial positive findings, there is still much debate over the utility of NOPs

compared to classical methods, with several studies showing no performance improvement

[21,30,31]. Perhaps even more striking is the finding that utilizing a permuted network [31] or

aggregating random genes [10] performs on par with networks describing true biological

relationships. Several meta-analyses attempting to establish the utility of NOPs have appeared

with contradicting conclusions. Notably, Staiger et al. compared performance of nearest mean

classifier [32] in this setting and concluded that network derived meta-genes are not more

predictive than individual genes [21,31]. This is in contradiction to Roy et al. who achieved

improvements in outcome prediction when genes were ranked according to their t-test statistics

compared to their pagerank property [33] in PPI network [28,34]. It is thus still an open question

whether NOPs truly improve outcome prediction in terms of predictive performance, cross-study

robustness or interpretability of the gene signatures.

A critical - yet often neglected - aspect in the successful application of NOPs is the contribution

of the biological network. In this regard, it should be recognized that many network links are

unreliable [35,36], missing [37] or redundant [38] and considerable efforts are being made to

refine these networks [37,39–41]. In addition, many links in these networks are experimentally

obtained from model organisms and therefore may not be functional in human cells [42–44].

Finally, most biological networks capture only a part of a cell's multifaceted system [45]. This

incomplete perspective may not be sufficient to link the wide range of aberrations that may occur

in a complex and heterogeneous disease such as breast cancer [46,47]. Taken together, these

issues raise concerns regarding the extent to which the outcome predictors may benefit from

inclusion of common biological networks in their models.

In this work, we propose to construct a network *ab initio* that is specifically designed to improve

outcome prediction in terms of cross-study generalization and performance stability. To achieve

this, we will effectively turn the problem around: instead of using a given biological network, we

aim to use the labelled gene expression datasets to identify the network of genes that truly

improves outcome prediction (see **Figure 1** for a schematic overview).

Figure 1. Schematic overview of SyNet inference and NOP training. For every 69 million

combinations of gene pairs **(a)** we compute three criteria including synergy (S_{ij}, purple),

average AUC (M_{ij}, pink), and correlation (C_{ij}, blue) **(b)**. These three criteria form a three-

dimensional space **(c)** from which Fitness (F_{ij}) can be calculated for each pair. Top pairs (green

dots) in this space collectively form SyNet **(d)**. SyNet is subsequently used in a NOP **(e)**, in

which the links in SyNet guide the construction of "meta-genes". Within a NOP groups of genes

are formed **(f)** and then integrated into meta-genes (typically using averaging) **(g)**. The

constructed meta-genes are then used as regular features to train standard classifiers **(h)**. The

phenotype of interest is patient outcome (i.e. 5-year survival).

Our approach relies on the identification of *synergistic gene pairs*, i.e. genes whose joint

prediction power is beyond what is attainable by both genes individually [48]. To identify these

pairs, we employed grid computing to evaluate all 69 million pairwise combinations of genes. The

resulting network, called SyNet, is specific to the dataset and phenotype under study and can be

used to successfully infer a NOP model.

To obtain SyNet, and allow for rigorous cross-study validation, a dataset of substantial size is

required. For this reason, we combined 14 publicly available datasets to form a compendium

encompassing 4129 samples. To the best of our knowledge, the data combined in this study

represents the largest breast cancer gene expression compendium to date. Further, to ensure

unbiased evaluation, sample assignments in the inner as well as the outer cross-validations folds

are kept equal across all assessments throughout the paper.

In the remainder of this paper, we will demonstrate that integrating genes based on SyNet

provides superior performance and stability of predictions when these models are tested on

independent cohorts. In contrast to previous reports, where shuffled versions of networks also

performed well, we show that the performance drops substantially when SyNet links are shuffled,

suggesting its connections are truly informative. We further evaluate the content and structure of

SyNet by overlaying it with known genesets and existing networks, revealing marked enrichment

for known breast cancer prognostic markers. While overlap with existing networks is highly

significant, the majority of direct links in SyNet is absent from these networks. Interestingly, they

can be reliably predicted from existing networks when more complex topological descriptions are

taken into account. Taken together, our findings suggest that compared to generic gene networks,

phenotype specific networks, which are derived directly from labeled data, can provide superior

performance while at the same time revealing valuable insight into etiology of breast cancer.

2 Results

2.1 SyNet improves NOP performance

We first evaluated NOP performance for three existing methods (Park, Chuang and Taylor) and the GL when supplied with a range of networks, including generic networks, tissue specific networks and SyNet. The AUC of the four NOPs, presented in **Figure 2**, clearly demonstrates that SyNet improves the performance of all NOPs, except for the Park method in which performs on par using the Corr network. None of the existing NOPs truly outperform the classifier Lasso (i.e. baseline; rightmost bar in **Figure 2)** which is the top performing regular classifier in our evaluations; see **S3** for details).

Figure 2. Performance comparison of NOPs for 4 methods and 12 networks including SyNet. Each bar represents average performance in terms of the AUC. The rightmost bar represents the performance of standard Lasso which considers all individual genes as features (i.e. no network). Error bars represent the standard deviation across 10 repeats.

The GL clearly outperforms all other methods, in particular when it exploits the information contained in SyNet. This corroborates our previous finding [8] that existing methods which construct meta-genes by averaging are suboptimal (see **S1** for a more extensive analysis). The GL using the Corr network also outperforms the baseline model, albeit non-significantly (p~0.6), which is in line with previous reports [23]. It should be noted that across all these experiments an identical set of samples is used to train the models so that any performance deviation must be due to differences in (i) the set of utilized genes or (ii) the integration of the genes into meta-genes. In the next two sections, we will investigate these factors in more details.

2.2 SyNet provides feature selection capabilities

Networks only include genes that are linked to at least one other gene. As a result, networks can provide a way of ranking genes based on the number and weight of their connections. One explanation for why NOPs can outperform regular classifiers is that networks provide an a priori gene (feature) selection [31]. To test this hypothesis and determine the feature selection capabilities of SyNet, we compare classification performances obtained using the baseline classifier (i.e. Lasso) that is trained using enclosed genes in each network. While this classifier performs well (see **S3** for details), it cannot exploit information contained in the links of given network. So any performance difference must be due to the genes in the network. The number of genes in the network was varied by thresholding the weighted edges in the network and removing unconnected genes. The edge weight threshold and the Lasso regularization parameter were determined using a cross-validated grid search (see **S5** for details). **Figure 3** provides the optimal performances for 12 distinct networks along with number of genes used in the final model (i.e. genes with non-zero Lasso coefficients). We also included the baseline model where all genes are utilized to train Lasso classifier (rightmost bar).

Figure 3. Performance comparison between networks when interconnections are ignored and genes contained in each network are utilized to train Lasso. **a.** Performance (AUC) of Lasso classification using individual genes in 12 networks. Numbers below each bar represent the median number of non-zero coefficients after training Lasso across 10 repeats and 14 folds. **b.** Stability of identified signatures measured by overlap between identified gene sets using Jaccard index. X and y-axis represent average and standard deviation of the Jaccard index measured across 10 repeats and 14 folds.

The results presented in **Figure 3.a** demonstrate that SyNet is the only network that performs significantly better than the baseline model which is trained on all genes. Interestingly, we observe

that SyNet is the top performing network while utilizing a comparable number of genes to other

networks. The second-best network is the Corr network. Compared to the Corr network, SyNet

additionally had access to label information, making it also specific to the class label under study.

It should be noted that the data on which SyNet and the Corr networks are constructed are

completely independent from the validation data on which the performance is based due to our

multi-way cross-validation scheme (see Methods and **S5**). We conclude that dataset specific

networks, in particular SyNet which also exploits label information, provides a meaningful feature

selection that is beneficial for classification performance.

Our result show that none of the tissue specific networks outperform the baseline. Despite the

modest performance, it is interesting to observe that performance for these networks increases

as more relevant tissues (e.g. breast and lymph node networks) are utilized in the classification.

Additionally, we observe that tissue specific networks do not outperform the generic networks.

This may be the result of the fact that generic networks predominantly contain broadly expressed

genes with fundamental roles in cell function which may still be relevant to survival prediction. A

similar observation was made for GWAS where SNPs in these widely-expressed genes can

explain a substantial amount of missed heritability [59].

In addition to classifier performance, an important goal of employing NOPs is to identify stable

gene signatures, that is, the same genes are selected irrespective of the study used to train the

models. Gene signature stability is necessary to confirm that the identified genes are independent

of dataset specific variations and are true biological drivers of the disease. To measure the

signature consistency, we assessed the overlap of selected genes across all repeats and folds

using the Jaccard Index. **Figure 3.b** shows that, using genes included in SyNet, Lasso identifies

more similar genes across folds and studies compared to other networks. Surprisingly, despite

the fact that the expression data from which SyNet is inferred changes in each classification fold,

the signature stability for SyNet is markedly better than for generic or tissue specific networks, in

which the genes are fixed. Therefore, our results demonstrate that synergistic genes in SyNet

truly aid the classifier to robustly select signatures across independent studies.

2.3 SyNet connections are beneficial for NOP

The ultimate goal of employing NOPs compared to classical models that do not use network

information is to improve prognosis prediction by harnessing the information contained in the links

of the given network. Therefore, we next aimed to assess to what extend also connections

between the genes, as captured in SyNet and other networks, can help NOPs to improve their

performance beyond what is achievable using individual genes. As before, we utilized identical

datasets (in terms of genes and samples) in inner and outer cross-validation loops to train all four

NOPs as well as the baseline model which uses Lasso. As a result, in our comparison, the only

difference between each NOP and the regular Lasso is the grouping of genes based on their

connectivity in the networks.

Our results, presented in **Figure 4.a**, clearly demonstrate that SyNet is the only network to achieve

superior prognostic prediction for unseen patients from an independent cohort. It outperforms all

other tested networks, including relevant tissue specific networks. Moreover, we find that using

SyNet also provides one of the largest gains in performance over its Lasso counterpart (**Figure**

4.b, x-axis), indicating that the links between genes in SyNet truly aid classification performance

beyond what is obtained as a result of the feature selection capabilities.

Figure 4. Performance of NOP models trained using SyNet compared to a shuffled version of this

network. **a.** Bars indicate average performance of models across repeats and error bars denote

the corresponding standard deviation. Solid bars represent average performance of models

trained using SyNet. Dashed bars denote performance of the same model using randomized

SyNet, i.e. the same genes are present but randomly connected. **b.** Improvement of performance

(x-axis) and stability (in terms of the standard deviation of the AUC; y-axis) compared to the

baseline model. Square and circle markers represent performance obtained using genes (i.e.

Lasso) and the network (i.e. GL), respectively.

To confirm that NOP performance using SyNet is the result of the network structure, we also

applied the GL to a shuffled version of SyNet (**Figure 4.a**). We observe a substantial deterioration

of the AUC supporting the conclusion that not only the genes, but also links contained in SyNet

are relevant to achieve good prediction. Moreover, this observation rules out that the GL by itself

is able to provide enhanced performance compared to standard Lasso. Result of similar

assessment for Corr network is given in (Figure **S12**).

Another important property of an outcome predictor is to yield comparable performance

irrespective of dataset used for training the model (i.e. performance stability). This is a highly

desirable quality, as concerns have been raised regarding the highly variable performances of

breast-cancer classifiers applied to different cohorts [7,60]. **Figure 4.b** clearly demonstrates that

a NOP model guided by SyNet not only provide superior performance, it also offers improved

stability of the classification performance.

We further compared performances when each network contains equal number of links and

observe a similar trend in performance (see **S7** for details). We also considered the more complex

Sparse Group Lasso (SGL), which offers an additional level of regularization. No substantial

difference between GL and SGL performance was found (see **S8** for details). Likewise, we did

not observe substantial performance differences when the number of genes, group size and

regularization parameters were simultaneously optimized in a grid search (see **S9** for details).

Together, these findings can be considered as the first unbiased evidence of true classification

performance improvement in terms of average AUC and classification stability by a NOP.

2.4 Gene enrichment analysis for SyNet

Many curated biological networks suffer from bias since genes with well-known roles are the

subject of more experiments and thus get more extensively and accurately annotated [61]. Post-

hoc interpretation of the features used by NOPs, often by means of an enrichment analysis, will

therefore be affected by the same bias. SyNet does not suffer from such bias, as its inference is

purely data driven. Moreover, since SyNet is built based on gene pairs that contribute to the

prediction of clinical outcome, we expect that the genes included in the network not only relate to

breast cancer; they should play a role in determining how aggressively the tumor behaves, how

advanced the disease is or how well it responds to treatment.

To investigated the relevance of genes contained in SyNet in the development of breast cancer

and, more importantly, clinical outcome, we ranked all pairs according to their median Fitness

(F_{ij}) across 14 studies and selected the top 300 genes (encompassing 3544 links). This cutoff

was frequently chosen by the Lasso as the optimal number of genes in SyNet (see section 3.1).

Figure 5 depicts a visualization of this network revealing three main subnetworks and a few

isolated gene pairs. We performed functional enrichment of all genes using Ingenuity Pathway

Analysis (IPA) [62], and functional enrichment of the subcomponents of the three large

subnetworks using Gene Set Enrichment Analysis (GSEA) [63].

Figure 5. Visualization of SyNet. SyNet consists of three main subnetworks (**a**, **b** and **c**) and five

separated gene pairs (**d**). Node size represents degree of node and link thickness indicates

fitness of the corresponding pair. **a.** The largest subnetwork encompassing 223 genes is enriched

for histologic grade of invasive breast cancer tumors. **b.** The second subnetwork is directly

connected to the first cluster and contains risk factors for developing breast cancer. **c.** The third

cluster is enriched for genes upregulated in normal-like subtype of breast cancer. **d.** Out of five

pairs, only TFF3 and TFF1 pair is enriched for genes up-regulated in early primary breast tumors.

IPA reveals that out of 300 genes in SyNet, 287 genes have a known relation to cancer (2e-

06<p<1e-34) of which 222 are related to reproductive system disease (2e-06<p<1e-34).

Furthermore, the top five upstream regulators of genes in SyNet (orange box, **Figure 5**) are

CDKN1A, E2F4, RABL6, TP53 and ERBB2, all of which are well known players in the

development of breast cancer [64–68]. The mean degree of the 300 genes in the network build

by SyNet is 24, but there are 12 genes which have a degree of 100 or above: ASPM [69], BUB1

[70], CCNB2 [71], CDKN3 [72], CENPA [73], DLGAP5 [74], KIF23 [75], MCM10 [76], MELK [77],

RACGAP1 [78], TTK [79] and UBE2C [80]. All these genes play a vital role in progression through

the cell cycle and mitosis, by ensuring proper DNA replication, correct formation of the mitotic

spindle and proper attachment to the centromere.

In addition to a clear involvement of genes linked to breast cancer generically, we also find clear

indications that the genes in SyNet are relevant to clinical outcome and prognosis of the disease.

For instance, the most highly enriched cluster (green cluster **Figure 5**) is found to be associated

to histological grade of the tumor. The histological grade, which is based on the morphological

characteristics of the tumor, has been shown to be informative for the clinical behavior of the

tumor and is one of the best-established prognostic markers [81]. Notably, the blue cluster is

enriched for genes involved in tamoxifen resistance, one of the important treatments of ER-

positive breast cancer.

Two other sub-clusters (yellow and purple in **Figure 5**), contain genes from distinctly different

biological processes than the main cluster. In these clusters we also observe clear hub genes:

SLC7A7 and CD74 in the yellow and ACKR1 and MFAP4 in the purple cluster. ACKR1 is a

chemokine receptor involved in the regulation of the bio-availability of chemokine levels and

MFAP4 is involved in regulating cell-cell adhesion. The recruitment of cells, as regulated by

chemokines, and reducing cell-cell adhesion both play an important role in the process of

metastasis. CD74 has also been linked to metastasis in triple negative breast cancer [82].

Metastasis, and not the primary tumor, is the main cause of death in breast cancer [3].

IPA highly significantly identifies the SyNet genes as upstream regulators of canonical pathways

implicated in breast cancer (**Figure 5**), such as Cell Cycle Control of Chromosomal Replication

(8e-18), Mitotic Roles of Polo-Like Kinase (4e-15), Role of CHK Proteins in Cell Cycle Checkpoint

Control (6e-12), Estrogen-mediated S-phase Entry (2e-11), and Cell Cycle: G2/M DNA Damage

Checkpoint Regulation (5e-10). Although all cancer cells deregulate cell cycle control, the degree

of dysregulation may contribute to a more aggressive phenotype. For instance, it is recognized

that the downregulation of certain checkpoint regulators are related to a worse prognosis in breast

cancer [83,84].

In summary, SyNet predominantly appears to contain genes relevant to two main processes in

the progression of breast cancer: increased cell proliferation and the process of metastasis.

Although many genes have not previously been specifically linked to breast cancer prognosis,

their role in regulating different stages of replication and mitosis points to a genuine biological role

in the progression and prognosis of breast cancer.

2.5 Similarity of SyNet to existing biological networks

We next sought to investigate the similarity between SyNet and existing biological networks that directly or indirectly capture biological interactions. To enable a comparison with networks of different sizes, we compare the observed overlap (both in terms of genes as well as links) to the distribution of expected overlap obtained by shuffling each network 1000 times. Overlap is determined for varying network sizes by thresholding the link weights such that a certain percentage of genes or links remains. Results are reported in terms of a z-score in **Figure 6**.

Figure 6. Similarity of existing biological networks to SyNet in terms of genes (**a.**) and links (**b.**). The x-axis represents the percentage of top gene/links used, the y-axis the z-score of observed vs. expected number of gene/links. The z-score is calculated by relating the observed number of SyNet gene/links that are present in existing biological networks to the expected distribution. To calculate the expected distribution, genes in biological networks are shuffled.

Figure 6.a shows that for the majority of networks a significantly higher than expected number of SyNet genes is contained in the top of each network. The overlap is especially pronounced for the tissue specific networks, in particular the Breast and Lymph node specific network, supporting our observation that SyNet contains links that are relevant for breast cancer. The enrichment becomes even more significant when considering the overlap between the links (**Figure 6.b**). In this respect, SyNet also is clearly most similar to the Breast and Lymph node specific networks. We confirmed that these enrichments are not only driven by the correlation component of SyNet by repeating this analysis with the correlation component removed (see **S10**). It should moreover be noted that, although a highly significant overlap is observed, the vast majority of SyNet genes and links are not present in the existing networks, explaining the improved performance obtained with NOPs using SyNet. Specifically, out of the 300 genes in SyNet, only 142 are contained within

the top 25% of genes (n=1005) in the Breast specific network, and 151 in the top 25% of genes

(n=1290) in the Lymph node specific network. Similarly, out of the 3544 links in SyNet, only 1182

are contained within the top 25% of links (n=12500) in the Breast specific network, and 617 in the

top 25% (n=12500) of the Lymph node specific network (see **S11** for details).

2.6 Higher order structural similarity of SyNet and existing biological networks

In addition to direct overlap, we also aimed to investigate if genes directly connected in SyNet

may be indirectly connected in existing networks. To assess this for each pair of genes in SyNet,

we computed several topological measures characterizing their (indirect) connection in the

biological networks. We included degree (**Figure 7.a**), shortest path (**Figure 7.b**) and jaccard

(**Figure 7.c**) (see supplementary text for details). To produce an edge measure from degree and

pagerank (which are node based), we computed the average degree and pagerank of genes in a

pair respectively. Furthermore, we produced an expected distribution for each pair by computing

the same topological measures for one of the genes and another randomly selected gene. The

results from this analysis supports our previous observation that the information contained in the

links of SyNet is substantially - yet partially - overlapping with the information in the existing

networks. Notably, the similarity increases for networks of increased relevance to the tissue in

which the gene expression data is measured (i.e. breast tissue).

Figure 7. Comparison of three topological measures calculated over biological networks. Each

color represents a network. Gray boxes represent the same topological measures calculated on

the shuffled network.

2.7 Predicting SyNet links from biological networks

Encouraged by the overlap with existing biological networks, we next asked whether links in

SyNet can be predicted from the complete collection of existing networks. To this end we

characterized each gene-pair by a set of 12 graph-topological measures that describe local and

global network structure around each gene-pair. In addition to the degree, shortest path and

Jaccard, we included several additional graph-topological measures including direct link, page

rank (with four betas), closeness centrality, clustering coefficient and eigenvector centrality (see

supplementary text for details). While converting node-based measures to edge based measures,

in addition to using the average, we also used the difference between the score for each gene in

the pair, similar to our previous work [85]. We applied these measures to all 10 networks in our

collection yielding a total of 210 features. The gene-pairs are labeled according to their presence

or absence in SyNet. Inspection of this dataset using the t-SNE [57] reveals that the links in SyNet

occupy a distinct part of the 2D embedding obtained (**Figure 8.a**).

Figure 8. Characterizing SyNet links by a range of graph topological measures. **a.** t-SNE

(unsupervised) visualization of the combined 180 topological measures. Each dot represents one

gene pair. Green dots indicate SyNet links while gray markers represent an equal number of

random pairs. **c.** Performance of Lasso model trained over all topological measure for different

networks and all networks combined (rightmost bar).

We trained a Lasso and assessed classification performance in a 50-fold cross validation scheme

where in each fold 1/50 of pairs in SyNet is kept hidden and the rest of pairs is utilized to train the

classifier. To avoid information leakage in this assessment, we removed gene pairs from the

training set in case one of the genes is present in the test set. Based on this analysis we find that

a simple linear classifier can reach ~85% accuracy in predicting the synergistic gene relationships

from SyNet (**Figure 8.b**, rightmost bar). The contribution from generic networks is markedly

smaller than for the tissue specific networks. In particular the networks relevant to breast cancer

are highly informative, to the extent that combining multiple networks no longer improves

prediction performance. Further investigation of feature importance revealed that the page rank

topological measure was commonly used as a predictive marker. Apparently, while direct overlap

between SyNet and existing networks is modest, the topology of the breast specific networks is

highly informative for the links contained in SyNet. This corroborates findings from Winter et al. in

which the page rank topological measure was proposed to identify relevant genes in outcome

prediction [33,34,86].

464 **3 Discussion and Future work**

Although the principle of using existing knowledge of the cellular wiring diagram to improve

performance, robustness and interpretability of gene expression classifiers appears attractive,

contrasting reports on the efficacy of such approach have appeared in literature [21,28,31,34].

Consensus in this field has particularly been frustrated by an evaluation of a limited set of sub-

optimal classifiers [21,23,28,34], small sample size [18,24,26], or the use of standard K-fold cross-

validation instead of cross-study evaluation schemes [24,26]. For this reason, it remained unclear

if network-based classification, and in particular network-based outcome prediction, is beneficial.

Here, we present a rigorously cross-validated procedure to train and validate Group Lasso-based

NOPs using a variety of networks, in particular also tissue specific networks, which have not been

evaluated in the context of NOPs before.

Based on our analyses we conclude that none of the existing networks achieve improved

performance compared to using properly regularized classifiers trained on all genes. In this work

we therefore present a novel gene network, called SyNet, which is computationally derived

directly from the labeled dataset. The links in SyNet connect synergistic gene pairs. We followed

a cross-validation procedure in which the inference of SyNet and validation of its utility in a NOP

is strictly separated. We find that SyNet-based NOPs yields superior performance with higher

stability across the folds compared to both the baseline model trained on all genes as well as

models that use other existing gene networks. We therefore conclude that networks can improve

outcome prediction, but only if this network is dataset specific and inferred from labeled training

data. The Corr network, which is also dataset specific but not inferred from labeled data, also

improved performance but much less than using SyNet.

A major benefit of SyNet over manually curated gene networks is that its inference is purely data

driven, and therefore not biased to well-studied genes. Post-hoc interpretation of the genes

selected by a NOP that utilized SyNet is therefore expected to provide a more unbiased

interpretation of the important molecular players underlying breast cancer and patient survival.

Analysis of the genes contained in SyNet shows strong enrichment for genes with known

relevance to breast cancer. More importantly, the largest subcomponent of SyNet is strongly

linked to patient prognosis because it includes many genes with a known relation to the

histological grade of the tumor.

To investigate if SyNet captures known biological gene interactions, we extensively compared

SyNet with existing networks. We find highly significant overlaps between links, indicating that

SyNet connects genes that also have a known biological interaction. Despite this significant

overlap, the majority of the SyNet links are not recapitulated by direct links in the existing

networks. However, we find that accurate predictions of links in SyNet are possible if more

complex graph topological descriptions of the indirect connections in the existing networks are

used. Interestingly, accurate predictions are only obtained when using the breast specific

networks. Apparently, although the information contained in SyNet is very similar to other gene

interaction networks, they can only be exploited by the classifier if they are connected in a certain

way. This might explain why using existing biological networks in NOPs directly is unsuccessful

and why using graph topological measures has been successful in identifying relevant genes in

outcome prediction [33,34,86].

4 Materials and Methods

Inferring a synergistic network (SyNet)

We hypothesized that, in order to improve outcome prediction by network-based classification,

interconnections in the network should correspond to gene pairs for which integration yields a

performance beyond what is attainable by either of the individual genes (i.e. synergy).

Accordingly, we formulated the synergy S_{ij} between gene i and gene j as

$$S_{ij} = \frac{A_{ij}}{Max(A_i, A_j)}$$

where A_i, A_j and A_{ij} respectively represent the Area Under Curve (AUC) of gene i, the AUC of

gene j and the AUC of meta-gene ij formed by aggregation of gene i and gene j. Meta-gene

formation is carried out by a linear regression model which demonstrated superior performance

in our experiments (see **S1** for details). Cross-validation performance of the linear regression (see

section on Cross validation design for details) is obtained and the median of 65 AUCs (13 folds

and 5 repeats) is used as the final score A_{ij} for each pair. The AUC of the individual genes (i.e.

A_i and A_j) is obtained in a similar fashion.

Defining the synergy as a function of AUC yields a phenotype (label) specific measure which

effectively ignores extraneous relationships between gene pairs that are not relevant in outcome

prediction. The synergy measure S_{ij} depends on the performance of individual genes where

poorly performing genes tend to achieve higher degree of synergy compared to two predictive

genes (see **S2** for corresponding analysis). In order to account for this effect, the average AUC

of individual genes is included as a second criterion. Furthermore, our preliminary tests confirmed

previous findings [8,23,49], that integrating highly correlated genes (which reduces meta-gene

noise) may improve survival prediction. For this reason, we added correlation of pairs as a third

criterion. To combine these three measures, each measure is normalized and combined into an

overall fitness score F_{ij} for gene pair ij:

$$F_{ij} = -\sqrt{(1 - \overline{S_{ij}})^2 + (1 - \overline{M_{ij}})^2 + (1 - \overline{C_{ij}})^2}$$

Here, M_{ij} and C_{ij} represent mean AUC and absolute spearman correlation of gene i and j

respectively. Bars above letters indicate that the corresponding values are normalized to the [0 ,

1] interval. Employing the Dutch grid infrastructure, we quantified the fitness for all 69 million

possible pairs of genes (n=11748). **Figure 1.c** visualizes the fitness of all pairs in a three-

dimensional space. Finally, the top 50,000 pairs with highest fitness are considered as SyNet.

Expression Data

Accurately estimating survival risk and identifying markers relevant for progression of a complex

disease such as breast cancer requires a large number of samples [11]. To this end, samples

from METABRIC [50] (n=1981) are combined with 12 studies collected in ACES [21] (n=1606) as

well as samples from the TCGA breast invasive carcinoma dataset [51] (n=532) (see

supplementary text for details). Collectively, these datasets, spanning 14 distinct studies, form a

compendium encompassing 4129 samples. To the best of our knowledge, the data combined in

this paper represents the largest breast cancer gene expression compendium to date. As a result,

our compendium should capture a large portion of the biological heterogeneity among breast

cancer patients, as well as technical biases originating from the variability in platforms and study-

specific sample preparations [52]. This variability will assist the trained models to achieve better

generalization which is crucial in real world application of the final classification model [9,13,53].

To correct for technical variations that may arise during the library preparation, initially the

expression data within each study is quantile normalized and then batch-effect corrected using

Combat [54] where the outcome of patients was modeled as an additional covariate to maintain

the variance associated with the prognostics. This procedure was shown to perform well among

many batch effect removal methods [55,56]. Successful removal of batch effects were confirmed

using t-SNE visualization [57] (See **S4** for details). The label for each patient corresponds to

overall survival time (or recurrence free survival if available) with respect to a 5-year threshold

(good vs. poor outcome).

Regular classifiers and Network based prediction models

Ascertaining the relevance of networks in outcome prediction should be performed using a robust

predictor capable of providing adequate performance in prognostic prediction. Previous

assessments in this regard have been limited to only few classifiers [21,23,28,34]. To identify the

optimal predictor, we have compared performance of wide range of linear and nonlinear classifiers

(see **S3** for details). Supporting our previous findings [8], this evaluation demonstrates that simple

linear classifiers outperform the more complex ones, with the regularized linear classifier (Lasso)

reaching the highest AUC. This classifier supports both classical and well as network based

prediction by its derivative called Group Lasso (GL) [58]. The GL is structurally analogous to

standard Lasso with the exception of the way in which the regularization is performed; Lasso

applies regularization to genes while GL enforces selection of groups of genes (See

supplementary text for details). In order to incorporate network information in the GL, similar to

our previous work [8], each gene in the corresponding network is considered as seed gene and

together with its K neighbors the group structure provided to the GL. Priority of neighbor selection

is determined by edge weights between each neighbor and corresponding seed gene. The

hyperparameters for each classifier (e.g. K in the GL) are determined by means of a grid search

in the inner cross validation loop (see **S5** for schematic overview).

For comparison, we include three well-known NOPs in our analysis. Park et al. utilized hierarchical

clustering to group highly correlated genes [23]. Each group is summarized into a meta-gene by

averaging the expression profile of the genes in that group. These meta-genes are then employed

as regular features to train a Lasso classifier. The optimal cluster size for hierarchical clustering

is identified by iterative application of Lasso in an inner cross-validation. Chuang et al. employs a

greedy search to define subnetworks [18]. This is done by iteratively expanding a sub-network

initiated from a seed gene guided by a supervised performance criterion which halts when

performance no longer increases (in the training set). After groups are formed, the meta-genes

are constructed by averaging expression of each gene within each group similar to Park et al.

Finally, Taylor et al. focus on hubs (i.e. highly connected genes, degree>5) in a network [24]. To

identify dysregulated subnetworks, the change in correlation between each hub and its direct

neighbors across two classes of outcome (poor vs. good) is assessed. Meta-genes are formed

from candidate subnetworks similar to the procedure employed by Park et al.

Networks

In addition to SyNet, we considered a total of 12 other publicly available networks, including

generic networks (HumanInt, BioPlex, BioGRID, IntAct and STRING), tissue specific networks

[43] (brain, kidney, ovary, breast, lymph node) and a correlation network (Corr) which was

previously shown to be an effective network in outcome prediction [8,23]. To the best of our

knowledge, our study is the first to evaluate tissue specific networks in the context of NOPs. To

maintain a reasonable network size, we utilized only the top 50,000 links (based on the link weight)

in each network (similar to number of links in SyNet). For the only unweighted network, HumanInt

[37], all interactions (n=~14k) were included and links were weighted according to the average

degree of the two interacting genes. Moreover, a randomized version of each network is

constructed by shuffling nodes in the network which destroys the biological information of the links

while preserving the overall network structure (see supplementary text for full details on

preparation of networks).

Cross validation design

In order to ascertain if network information truly aids outcome prediction, the evaluation should

be based on a rigorous cross-validation that closely resembles the real-world application of these

models. To this end, we perform cross-study validation in order to mimic a realistic situation in

which a classifier is applied to data from a different hospital than it was trained on [7]. Briefly, one

study is taken out for validation of the final performance (outer loop test set). SyNet inference and

NOP training are carried out on the 13 remaining studies (outer loop training set). Within each

fold of the outer loop training set, again one study is left out to obtain the inner loop test set and

the rest of studies for inner loop training set. The inner loop training set is sub sampled (with

replacement) to 70% and regression is performed for every genes as well as gene pairs (identical

set of samples are used across all genes and pairs). The AUC scores (A_i, A_j and A_{ij}) are

calculated on the inner loop test set. This is repeated 5 times. To train a NOP for this fold, a new

inner loop training set is formed by redrawing 70% of the samples from the outer loop training set.

This set is also used to infer correlation network. To assess the final performance of the NOP the

outer loop test set is used (see **S5** for a detailed schematic). Our initial experiments showed a

large variation of performance across studies (see **S6** for details). To prevent this variation from

influencing our comparisons, assignment of samples to folds in both inner and outer cross-

validation loops are kept identical across all comparisons throughout the paper. We used Area

Under the ROC Curve (AUC) as the main measure of performance in this paper.

Declarations

Availability of data and material: The normalized and batch effect removed cohort (excluding

METABRIC which requires access approval through synapse.org) can be downloaded from [87].

Furthermore, from [87], the full matrix of SyNet in binary format as well as top gene pairs (including

their tri-score used to calculate their fitness) is available for download in tab-delimited format.

Finally, all scripts used for preparation of data and figures in this manuscript are available for

download from [88]. To ensure the full reproducibility of our results, the indices utilized for training

and testing of all models (including inner and outer cross-validation) is also provided.

Competing interests: The authors declare there are no competing interests.

Funding: Part of computations required for this work was carried out on the Dutch national e-

infrastructure (e-infra160001) with the support of the SURF Foundation.

Authors' contributions: AA collected, analyzed the data and prepared the manuscript. JU

interpreted the results and prepared the manuscript, JdR designed the experiments, prepared

and reviewed the manuscript. All authors read and approved the final manuscript.

References

1. Fantozzi A, Christofori G. Mouse models of breast cancer metastasis. Breast Cancer Res

[Internet]. 2006;8. Available from: http://dx.doi.org/10.1186/bcr1530

2. Shapiro CL, Recht A. Side Effects of Adjuvant Treatment of Breast Cancer. N Engl J Med.

2001;344:1997–2008.

3. Weigelt B, Peterse JL, van't Veer LJ. Breast cancer metastasis: markers and models. Nat

Rev Cancer. 2005;5:591–602.

4. Cardoso F, van't Veer LJ, Bogaerts J, Slaets L, Viale G, Delaloge S, et al. 70-Gene Signature

as an Aid to Treatment Decisions in Early-Stage Breast Cancer. N Engl J Med. 2016;375:717–

29.

5. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AAM, Mao M, et al. Gene expression

profiling predicts clinical outcome of breast cancer. Nature. 2002;415:530–6.

6. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AAM, Voskuil DW, et al. A gene-

expression signature as a predictor of survival in breast cancer. N Engl J Med. 2002;347:1999–

2009.

7. Bernau C, Riester M, Boulesteix A-L, Parmigiani G, Huttenhower C, Waldron L, et al. Cross-

study validation for the assessment of prediction algorithms. Bioinformatics. 2014;30:i105–12.

8. Allahyar A, de Ridder J. FERAL: network-based classifier with application to breast cancer

outcome prediction. Bioinformatics. 2015;31:i311–9.

9. Ransohoff DF. Opinion: Bias as a threat to the validity of cancer molecular-marker research.

Nat Rev Cancer. 2005;5:142–9.

10. Venet D, Dumont JE, Detours V. Most random gene expression signatures are significantly

associated with breast cancer outcome. PLoS Comput Biol. 2011;7:e1002240.

11. Ein-Dor L, Zuk O, Domany E. Thousands of samples are needed to generate a robust gene

list for predicting outcome in cancer. Proc Natl Acad Sci U S A. 2006;103:5923–8.

12. Akavia UD, Litvin O, Kim J, Sanchez-Garcia F, Kotliar D, Causton HC, et al. An integrated

approach to uncover drivers of cancer. Cell. 2010;143:1005–17.

13. Kim S-Y. Effects of sample size on robustness and prediction accuracy of a prognostic gene

signature. BMC Bioinformatics. 2009;10:147.

14. Hua J, Tembe WD, Dougherty ER. Performance of feature-selection methods in the

classification of high-dimension data. Pattern Recognit. 2009;42:409–24.

15. Bryant PA, Smyth GK, Robins-Browne R, Curtis N. Technical Variability Is Greater than

Biological Variability in a Microarray Experiment but Both Are Outweighed by Changes Induced

by Stimulation. PLoS One. 2011;6:e19556.

16. Parker HS, Leek JT. The practical effect of batch on genomic prediction. Stat Appl Genet

Mol Biol. 2012;11:Article 10.

17. Alcaraz N, List M, Batra R, Vandin F, Ditzel HJ, Baumbach J. De novo pathway-based

biomarker identification. Nucleic Acids Res. 2017;45:e151.

18. Chuang H-Y, Lee E, Liu Y-T, Lee D, Ideker T. Network-based classification of breast cancer

metastasis. Mol Syst Biol. 2007;3:140.

19. Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. Cell. 2011;144:646–

74.

20. Hanahan D, Weinberg RA. The hallmarks of cancer. Cell. 2000;100:57–70.

21. Staiger C, Cadot S, Györffy B, Wessels LFA, Klau GW. Current composite-feature

classification methods do not outperform simple single-genes classifiers in breast cancer

prognosis. Front Genet [Internet]. 2013;4. Available from:

http://dx.doi.org/10.3389/fgene.2013.00289

22. Cheng W-Y, Ou Yang T-H, Anastassiou D. Biomolecular events in cancer revealed by

attractor metagenes. PLoS Comput Biol. 2013;9:e1002920.

23. Park MY, Hastie T, Tibshirani R. Averaged gene expressions for regression. Biostatistics.

Oxford University Press; 2007;8:212–27.

24. Taylor IW, Linding R, Warde-Farley D, Liu Y, Pesquita C, Faria D, et al. Dynamic modularity

in protein interaction networks predicts breast cancer outcome. Nat Biotechnol. 2009;27:199–

204.

25. Zhang W, Chien J, Yong J, Kuang R. Network-based machine learning and graph theory

algorithms for precision oncology. npj Precision Oncology [Internet]. 2017;1. Available from:

http://dx.doi.org/10.1038/s41698-017-0029-7

26. Popovici V, Chen W, Gallas BG, Hatzis C, Shi W, Samuelson FW, et al. Effect of training-

sample size and classification difficulty on the accuracy of genomic predictors. Breast Cancer

Res. 2010;12:R5.

27. Wessels LFA, Reinders MJT, Hart AAM, Veenman CJ, Dai H, He YD, et al. A protocol for

building and evaluating predictors of disease state based on microarray data. Bioinformatics.

2005;21:3755–62.

28. Roy J, Winter C, Schroeder M. Meta-analysis of Cancer Gene Profiling Data. Methods in

Molecular Biology. 2016. p. 211–22.

29. Dutkowski J, Ideker T. Protein networks as logic functions in development and cancer. PLoS

Comput Biol. 2011;7:e1002180.

30. Cun Y, Fröhlich H. Network and data integration for biomarker signature discovery via

network smoothed T-statistics. PLoS One. 2013;8:e73074.

31. Staiger C, Cadot S, Kooter R, Dittrich M, Müller T, Klau GW, et al. A Critical Evaluation of

Network and Pathway-Based Classifiers for Outcome Prediction in Breast Cancer. PLOS ONE

[Internet]. 2012;7. Available from:

http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0034796

32. Alpaydin E. Introduction to Machine Learning. MIT Press; 2014.

33. Winter C, Kristiansen G, Kersting S, Roy J, Aust D, Knösel T, et al. Google goes cancer:

improving outcome prediction for cancer patients by network-based ranking of marker genes.

PLoS Comput Biol. 2012;8:e1002511.

34. Roy J, Winter C, Isik Z, Schroeder M. Network information improves cancer outcome

prediction. Brief Bioinform. Oxford University Press; 2014;15:612–25.

35. Cusick ME, Yu H, Smolyar A, Venkatesan K, Carvunis A-R, Simonis N, et al. Literature-

curated protein interaction datasets. Nat Methods. 2009;6:39–46.

36. von Mering C, Krause R, Snel B, Cornell M, Oliver SG, Fields S, et al. Comparative

assessment of large-scale data sets of protein-protein interactions. Nature. 2002;417:399–403.

37. Rual J-F, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, Li N, et al. Towards a

proteome-scale map of the human protein-protein interaction network. Nature. 2005;437:1173–

8.

38. Mahdavi MA, Lin Y-H. False positive reduction in protein-protein interaction predictions

using gene ontology annotations. BMC Bioinformatics. 2007;8:262.

39. Rolland T, Taşan M, Charloteaux B, Pevzner SJ, Zhong Q, Sahni N, et al. A proteome-scale

map of the human interactome network. Cell. 2014;159:1212–26.

40. Huttlin EL, Ting L, Bruckner RJ, Gebreab F, Gygi MP, Szpyt J, et al. The BioPlex Network: A

Systematic Exploration of the Human Interactome. Cell. 2015;162:425–40.

41. Huttlin EL, Bruckner RJ, Paulo JA, Cannon JR, Ting L, Baltier K, et al. Architecture of the

human interactome defines protein communities and disease networks. Nature. 2017;545:505–

9.

42. Greene CS, Troyanskaya OG. Chapter 2: Data-driven view of disease biology. PLoS

Comput Biol. 2012;8:e1002816.

43. Greene CS, Krishnan A, Wong AK, Ricciotti E, Zelaya RA, Himmelstein DS, et al.

Understanding multicellular function and disease with human tissue-specific networks. Nat

Genet. 2015;47:569–76.

44. Yeger-Lotem E, Sharan R. Human protein interaction networks across tissues and diseases.

Front Genet. 2015;6:257.

45. Zhang S, Jin G, Zhang X-S, Chen L. Discovering functions and revealing mechanisms at

molecular level from biological networks. Proteomics. 2007;7:2856–69.

46. Kotlyar M, Pastrello C, Sheahan N, Jurisica I. Integrated interactions database: tissue-

specific view of the human and model organism interactomes. Nucleic Acids Res.

2016;44:D536–41.

47. de Anda-Jáuregui G, Velázquez-Caldelas TE, Espinal-Enríquez J, Hernández-Lemus E.

Transcriptional Network Architecture of Breast Cancer Molecular Subtypes. Front Physiol.

2016;7:568.

48. Watkinson J, Wang X, Zheng T, Anastassiou D. Identification of gene interactions

associated with disease from gene expression data using synergy networks. BMC Syst Biol.

2008;2:10.

49. Wu G, Stein L. A network module-based method for identifying cancer prognostic

signatures. Genome Biol. 2012;13:R112.

50. Curtis C, Shah SP, Chin S-F, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and

transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature.

2012;486:346–52.

51. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast

tumours. Nature. 2012;490:61–70.

52. Allott EH, Geradts J, Sun X, Cohen SM, Zirpoli GR, Khoury T, et al. Intratumoral

heterogeneity as a source of discordance in breast cancer biomarker classification. Breast

Cancer Res. 2016;18:68.

53. Kourou K, Exarchos TP, Exarchos KP, Karamouzis MV, Fotiadis DI. Machine learning

applications in cancer prognosis and prediction. Comput Struct Biotechnol J. 2015;13:8–17.

54. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using

empirical Bayes methods. Biostatistics. 2007;8:118–27.

55. Müller C, Schillert A, Röthemeier C, Trégouët D-A, Proust C, Binder H, et al. Removing

Batch Effects from Longitudinal Gene Expression - Quantile Normalization Plus ComBat as Best

Approach for Microarray Transcriptome Data. PLoS One. 2016;11:e0156594.

56. Chen C, Grennan K, Badner J, Zhang D, Gershon E, Jin L, et al. Removing batch effects in

analysis of expression microarray data: an evaluation of six batch adjustment methods. PLoS

One. 2011;6:e17238.

57. van der Maaten L, van der Maaten L, Hinton G. Visualizing non-metric similarities in multiple

maps. Mach Learn. 2011;87:33–55.

58. Yuan M, Lin Y. Model selection and estimation in regression with grouped variables. J R

Stat Soc Series B Stat Methodol. 2006;68:49–67.

59. Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From Polygenic to

Omnigenic. Cell. 2017;169:1177–86.

60. Castaldi PJ, Dahabreh IJ, Ioannidis JPA. An empirical assessment of validation practices for

molecular classifiers. Brief Bioinform. 2011;12:189–202.

61. Khatri P, Sirota M, Butte AJ. Ten years of pathway analysis: current approaches and

outstanding challenges. PLoS Comput Biol. 2012;8:e1002375.

62. Krämer A, Green J, Pollard J, Tugendreich S. Causal analysis approaches in Ingenuity

Pathway Analysis. Bioinformatics. 2013;30:523–30.

63. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-wide expression

profiles. Proc Natl Acad Sci U S A. 2005;102:15545–50.

64. Khaleel SS, Andrews EH, Ung M, DiRenzo J, Cheng C. E2F4 regulatory program predicts

patient survival prognosis in breast cancer. Breast Cancer Res. 2014;16:486.

65. Gasco M, Shami S, Crook T. The p53 pathway in breast cancer. Breast Cancer Res

[Internet]. 2002;4. Available from: http://dx.doi.org/10.1186/bcr426

66. Wei C-Y, Tan Q-X, Zhu X, Qin Q-H, Zhu F-B, Mo Q-G, et al. Expression of CDKN1A/p21

and TGFBR2 in breast cancer and their prognostic significance. Int J Clin Exp Pathol. e-Century

Publishing Corporation; 2015;8:14619.

67. Tan M, Yu D. Molecular Mechanisms of ErbB2-Mediated Breast Cancer Chemoresistance.

Advances in Experimental Medicine and Biology. 2007. p. 119–29.

68. Montalbano J, Jin W, Sheikh MS, Huang Y. RBEL1 is a novel gene that encodes a

nucleocytoplasmic Ras superfamily GTP-binding protein and is overexpressed in breast cancer.

J Biol Chem. 2007;282:37640–9.

69. Fish JL, Kosodo Y, Enard W, Pääbo S, Huttner WB. Aspm specifically maintains symmetric

proliferative divisions of neuroepithelial cells. Proc Natl Acad Sci U S A. 2006;103:10438–43.

70. Skoufias DA, Andreassen PR, Lacroix FB, Wilson L, Margolis RL. Mammalian mad2 and

bub1/bubR1 recognize distinct spindle-attachment and kinetochore-tension checkpoints. Proc

Natl Acad Sci U S A. National Academy of Sciences; 2001;98:4492–7.

71. Draetta G, Luca F, Westendorf J, Brizuela L, Ruderman J, Beach D. cdc2 protein kinase is

complexed with both cyclin A and B: Evidence for proteolytic inactivation of MPF. Cell.

1989;56:829–38.

72. Nalepa G, Barnholtz-Sloan J, Enzor R, Dey D, He Y, Gehlhausen JR, et al. The tumor

suppressor CDKN3 controls mitosis. J Cell Biol. Rockefeller University Press;

2013;jcb.201205125.

73. Foltz DR, Jansen LET, Black BE, Bailey AO, Yates JR 3rd, Cleveland DW. The human

CENP-A centromeric nucleosome-associated complex. Nat Cell Biol. 2006;8:458–69.

74. Tsou A-P, Yang C-W, Huang C-YF, Yu RC-T, Lee Y-CG, Chang C-W, et al. Identification of

a novel cell cycle regulated gene, HURP, overexpressed in human hepatocellular carcinoma.

Oncogene. 2003;22:298–307.

75. Pavicic-Kaltenbrunner V, Mishima M, Glotzer M. Cooperative assembly of CYK-

4/MgcRacGAP and ZEN-4/MKLP1 to form the centralspindlin complex. Mol Biol Cell.

2007;18:4992–5003.

76. Ricke RM, Bielinsky A-K. Mcm10 Regulates the Stability and Chromatin Association of DNA

Polymerase-α. Mol Cell. 2004;16:173–85.

77. Nakano I, Paucar AA, Bajpai R, Dougherty JD, Zewail A, Kelly TK, et al. Maternal embryonic

leucine zipper kinase (MELK) regulates multipotent neural progenitor proliferation. J Cell Biol.

2005;170:413–27.

78. Lee K-Y, Esmaeili B, Zealley B, Mishima M. Direct interaction between centralspindlin and

PRC1 reinforces mechanical resilience of the central spindle. Nat Commun. 2015;6:7290.

79. Fisk HA, Mattison CP, Winey M. Human Mps1 protein kinase is required for centrosome

duplication and normal mitotic progression. Proc Natl Acad Sci U S A. 2003;100:14875–80.

80. Hao Z, Zhang H, Cowell J. Ubiquitin-conjugating enzyme UBE2C: molecular biology, role in

tumorigenesis, and potential as a biomarker. Tumour Biol. 2012;33:723–30.

81. Rakha EA, Reis-Filho JS, Baehner F, Dabbs DJ, Decker T, Eusebi V, et al. Breast cancer

prognostic classification in the molecular era: the role of histological grade. Breast Cancer Res.

2010;12:207.

82. Greenwood C, Metodieva G, Al-Janabi K, Lausen B, Alldridge L, Leng L, et al. Stat1 and

CD74 overexpression is co-dependent and linked to increased invasion and lymph node

metastasis in triple-negative breast cancer. J Proteomics. 2012;75:3031–40.

83. Catzavelos C, Bhattacharya N, Ung YC, Wilson JA, Roncari L, Sandhu C, et al. Decreased

levels of the cell-cycle inhibitor p27Kip1 protein: Prognostic implications in primary breast

cancer. Nat Med. 1997;3:227–30.

84. Craig C, Wersto R, Kim M, Ohri E, Li Z, Katayose D, et al. A recombinant adenovirus

expressing p27Kip1 induces cell cycle arrest and loss of cyclin-Cdk activity in human breast

cancer cells. Oncogene. 1997;14:2283–9.

85. Hulsman M, Dimitrakopoulos C, de Ridder J. Scale-space measures for graph topology link

protein network architecture to function. Bioinformatics. 2014;30:i237–45.

86. Newman MEJ. Analysis of weighted networks. Phys Rev E. American Physical Society;

2004;70:056131.

87. A phenotype specific network of synergistic genes. SyNet.deRidderLab.nl. Accessed 12

April 2018.

88. Pipeline for detecting synergistic pairs from gene expression data:

github.com/UMCUGenetics/SyNet. Accessed 12 April 2018.