Causal inference for spatial constancy across whole-body motion

Florian Perdreau[1]*, James Cooke[1], Mathieu Koppen[1], and W. Pieter Medendorp[1]

[1] Radboud University, Donders Institute for Brain, Cognition & Behaviour, Nijmegen, Netherlands

*corresponding authors: Florian Perdreau (f.perdreau@donders.ru.nl)

Running title: Causal inference for spatial constancy across whole-body motion

Keywords: causal inference, spatial updating, self-motion, visuo-vestibular integration

Abstract

The brain can estimate the amplitude and direction of self-motion by integrating multiple sources of sensory information, and use this estimate to update object positions in order to provide us with a stable representation of the world. A strategy to improve the precision of the object position estimate would be to integrate this internal estimate and the sensory feedback about the object position based on their reliabilities. Integrating these cues, however, would only be optimal under the assumption that the object has not moved in the world during the intervening body displacement. Therefore, the brain would have to infer whether the internal estimate and the feedback relate to a same external position (stable object), and integrate and/or segregate these cues based on this inference – a process that can be modeled as Bayesian Causal inference. To test this hypothesis, we designed a spatial updating task across passive whole-body translation in complete darkness, in which participants (n=11), seated on a vestibular sled, had to remember the world-fixed position of a visual target. Immediately after the translation, a second target (feedback) was briefly flashed around the estimated "updated" target location, and participants had to report the initial target location. We found that the participants' responses were systematically biased toward the position of the second target position for relatively small but not for large differences between the "updated" and the second target location. This pattern was better captured by a Bayesian causal inference model than by alternative models that would always either integrate or segregate the internally-updated target position and the visual feedback. Our results suggest that the brain implicitly represents the posterior probability that the internally updated estimate and the sensory feedback come from a common cause, and use this probability to weigh the two sources of information in mediating spatial constancy across whole-body motion.

Author Summary

A change of an object's position on our retina can be caused by a change of the object's location in the world or by a movement of the eye and body. Here, we examine how the brain solves this problem for spatial updating by assessing the probability that the internally-updated location during body motion and observed retinal feedback after the motion stems from the same object location in the world. Guided by Bayesian causal inference model, we demonstrate that participants' errrors in spatial updating depend nonlinearly on the spatial discrepancy between internally-updated and reafferent visual feedback about the object's location in the world. We propose that the brain implicitly represents the probability that the internally updated estimate and the sensory feedback come from a common cause, and use this probability to weigh the two sources of information in mediating spatial constancy across whole-body motion.

Introduction

Motor acts have immediate consequences for the sensory input. For example, a saccadic eye movement across the visual scene temporarily suppresses visual processing [1] and alters the retinal image [2]. Nevertheless, the brain retains correspondence between the presaccadic and postsaccadic scenes – called visual stability – by dissociating these changes in retinal input from those due to changes of the visual scene itself [3].

To do so, it has been suggested that the brain uses an internal forward model that, based on a copy of the saccadic motor command, predicts the postsaccadic scene, which can then be compared with the actual feedback of the postsaccadic scene [4,5]. However, this evaluation process is not flawless because both signals, i.e., the predicted and the actual feedback, are noisy [6]. The optimal strategy for the brain to cope with such uncertainty is through statistically weighting the evidence that the predicted and the actual feedback reflect the same scene or not. This strategy is known as Bayesian causal inference [7].

Recently, we provided evidence for this strategy using the saccadic suppression of displacement task [8], testing how participants judge the presaccadic location of a visual object that shifted during a saccade [9]. Following the rules of Bayesian causal inference, integration was strong when predicted and actual feedback represented spatially close target locations (as if they had a common cause), but weakened with larger spatial differences, depending on the precision of these signals [9].

While the saccadic system has provided evidence for Bayesian causal inference, it is not trivial that this mechanism is also applied to retain visual stability in other motion conditions. Saccades are rapid, self-generated movements that result in an abrupt alteration of the visual scene [10], and more critically in a selective suppression of visual information [11]. Therefore, a mechanism that predicts the reafferent visual information based on motor commands (via

forward models) may be a prerequisite for visual updating across saccades [4,5]. In contrast to

saccades, passively induced motions, such as riding a car, induce slow and progressive changes

of the visual input, and do not have corresponding motor commands that could be used to

predict the visual consequences of self-motion. Given these differences, it is not clear how the

brain deals with passive self-motion when the environment remains visible.

During passive self-motion, the brain must rely on vestibular and other sensory signals to infer

the motion [12–16]. Various studies suggested that there is a clear compensation for passive

self-motion in the updating of visual space, although compensation is not always perfect [17–

20]. Other studies have shown that this compensation is severely compromised when the

vestibular system is lesioned, indicating that vestibular signals weight significantly into visual

space updating [21].

Despite these insights, it is important to point out that most of these studies operationalized

visual updating by measuring how the brain, in darkness, keeps track of remembered target

locations during the motion, in which reliance on self-motion feedback in updating is necessary.

In heuristic terms, these self-motion updates may be superfluous in natural settings, where the

visual world remains continuously available, uninterrupted by the motion [22].

Here, we ask whether the brain applies Bayesian causal inference in the processing of self-

motion-based visual updates and actual visual feedback signals, or whether it simply derives

heuristic, suboptimal solutions to achieve visual stability during passive self-motion, e.g., by

relying on visual feedback alone.

To address this question, we designed a spatial updating task across passive whole-body

translation, in which participants, seated on a vestibular sled, had to remember the world-fixed

position of a visual target and report its location after the intervening body displacement.

Critically, in contrast to previous studies, the target was briefly presented again at the end of

the displacement (as actual visual feedback), but shifted relative to the updated target location.

In line with the predictions of Bayesian causal inference, we found that our participants'

responses were systematically biased to the actual visual feedback, depending on its spatial

discrepancy with the updated location. Our data could not be accounted for by a standard

optimal integration model that integrates the internal update and actual feedback irrespectively

of their spatial discrepancy, or by reliance on either one of these signals. Our findings suggest

that the brain explicitly represents the causal structure in multiple signal integration for visual

stability across whole-body motion.

Methods

Participants

11 participants took part in the present study (mean age = 27.3 yrs (SE = 2.4), 7 males). All

subjects had normal or corrected-to-normal vision and had no known vestibular or neurological

disorders. The present study was approved by the Ethics Committee of the Faculty of Social

Sciences of the Radboud University, Nijmegen. Every participant gave written informed

consent prior to participating in the experiment.

Apparatus

Participant's displacement was operated by a custom-made sled, consisting of a chair mounted

on an 800-mm track (see [16] for more details). The sled was powered by a linear motor

(TB15N, Technotion, Almelo, The Netherlands) and controlled by a Kollmorgen S700 drive

(Danaher, Washington, DC). The movements of the sled were controlled with a precision better

than 0.034 mm, 2 mm/s and 150 mm/s2. Participants were seated with their interaural axis

aligned with the direction of the sled motion. Head movements were restricted by an ear-fixed

mold and a chin rest so that participants' eyes were kept at a distance of 1.47 m orthogonal to

an OLED screen of size 1234x676 mm (55EA8809-ZC, LG, Seoul, South Korea). The screen

had a refresh rate of 60 Hz. It was placed in front of the sled, aligned with its center (see Fig

1A). A black cardboard frame was mounted on the screen to prevent any residual illumination

that could make the screen edges visible.

A linear guide was mounted on the sled, in front of the participants and at the level of their

thoracic diaphragm. By moving a slider on this guide, subjects controled the horizontal motion

of a cross-hair cursor displayed on the screen. Position of the slider was continuously tracked

at 200 Hz using two Optotrak Certus systems (NDI, Northern Digital Instruments, Waterloo,

Canada). The experiment and setup were controlled using software written in Python 2.7.

A

B

Figure 1**. Setup and procedure**. (A) Experimental setup. Participants were seated on a vestibular sled. They viewed stimuli on a screen and could control a cursor using a linear guide mounted to the seld. (B) Procedure. A test trial started after the participant had moved a cursor (green cross) to the homing position (yellow square). Next, a target (red disc) was flashed for 300 ms, whose world-centered location had to be remembered. Then, the participant was displaced by 40 cm within 1 s, after which a probe target appeared for 50 ms, shifted relative to the updated target position. Finally, the green cursor reappeared and the participant had to position it at the remembered world-centered location of the target.

Procedure

We designed a task that addresses spatial updating across whole-body passive translation. The task comprised two kinds of trials, *update-only trials* and *test trials*. In the update-only trials, participants had to remember the world-fixed position of a target, briefly flashed prior to the motion, and report its location after the motion. Previous work has shown imperfect spatial updating for passive translation in complete darkness [17,23] and the update-only trials were used to determine the updating gain per participant (see below). The test trials were identical to the update-only trials, with the exception that at the end of the motion, but prior to the participant's response, the target was briefly displayed again. The location of this probe target varied, but was centered on the internally-updated target location, as estimated on the basis of the update-only trials.

The details of the task are shown in Fig 1B. At the beginning of each trial, participants were passively moved to the homing position of the sled, either at -200 mm or at +200 mm relative to the screen's center, depending on whether the trial was to test updating across rightward or leftward translation. Then, participants were presented with a 20x20 mm green cursor on the screen, along with a 20x20mm yellow square (cursor homing position) and a gray body-fixed fixation dot (radius: 3.5 mm) that participants had to fixate throughout the trial (Fig 1B). Using the linear guide, participants had to bring the cursor onto the homing position such that both disappeared, which triggered the onset of the target (red disc, radius: 12.5 mm), presented for 300 ms, at one of five possible locations (-100, -50, 0, 50, 100 mm relative to screen center). At target offset, participants were passively moved sideways by 40 cm to the left or to the right for a duration of 1 s with a minimum-jerk velocity profile (peak velocity: 0.7 m/s, peak acceleration: 2.2 m/s²). In the test trials, at the end of the motion a probe target was briefly flashed for 50 ms with one of eight possible shifts (-228, -80, -28, -8, 8, 28, 80, 228 mm) relative to the internally updated target location, which was estimated by a preceding block of update-

only trials. Finally, in both kinds of trials the cursor reappeared and a brief sound cued the participant to position the cursor at the initially remembered, world-centered location of the target. Participants had 2.5 s to provide their response. If no response was detected within the time limit, the trial was repeated later during the experiment. After the participant had given his/her response, a new trial started, testing updating across motion in the opposite direction. To keep participants motivated and focused on good performance, a message was displayed after every 20 trials showing the average error of the last 20 trials. If the average error was smaller than at the previous message, it was displayed in green (red otherwise). Every participant was instructed to aim for a green feedback signal as often as possible.

The experiment started with a block of 80 update-only trials (eight replications of the five target positions for two movement directions). For each such trial we computed a *motion updating gain* of $1 - \frac{r-s}{D}$, with s the actual target position, r the response position and D the signed motion amplitude ($D < 0$ for leftward, $D > 0$ for rightward motion) of the trial (all in screen coordinates). This gain equals unity for a perfect motion update and zero if any motion update is lacking. A participant's updating gain α was determined as his/her average updating gain across all trials of the update-only block and this α was then used to compute his/her internally updated target position in the test trials. Actually, to allow for asymmetric updating, separate updating gains were computed for leftward vs. rightward motion.

Next, the experiment's *test block* started, consisting of a mix of test trials and update-only trials. Each participant performed 640 test trials (8 replications of 2 motion directions x 5 target locations x 8 probe target shifts). In addition, the test block contained 160 update-only trials, 66% of them randomly interleaved in the first half of the test block and the remaining in the second half. These trials were included to check that the internal update estimate of the preceding update-only block was still valid and to help maintain it [24]. The outcomes of these

185   trials were also used to progressively update the participant's gain parameter α as the overall

186   average updating gain across all his/her update-only trials.

187   Target locations and probe target shifts used in the experiment were selected based on pilots

188   and simulations to ensure model recovery.

Data analysis

Behavior

For every trial, a response was validated if the cursor reached a velocity of 5 cm/s or less for

300 ms (in screen coordinates). Response error was then computed as the difference between

the validated response position and the target position and expressed in mm. Subsequent offline

data and statistical analyses were performed using MATLAB (2015b) and R (3.3.2; [25]).

Individual updating gains α were derived as described above and computed separately for

leftward and rightward self-motion directions. To determine whether the individual updating

gain could be modeled by a single parameter irrespective of self-motion direction we performed

a paired samples t test on leftward vs. rightward updating gains.

For the test trials, we compared individual response error and variability across conditions by

testing linear mixed models using the lme4 R package [26] with motion direction (left, right)

and probe target shift (8 levels ranging from -228 to +228 mm) as predictors. In order to

compare models, linear and quadratic trends of the probe target effect were investigated.

Overall, threshold for statistical significance was defined as 5%.

For the purpose of plotting only, participants' data were remapped to a rightward body-

displacement.

Model

In this study's main task a trial started with flashing a visual target. Next, the participant was moved sideways and at the end of the motion a probe was flashed either at or with an horizontal deviation from the internally-updated target position (determined using the update-only trials; see Procedure). Throughout the trial a body-fixed fixation cross was present. The participant was then required to indicate the world-fixed position of the target presented before the motion. The purpose was to investigate whether in this situation of passive self-motion and uninterrupted visual input the brain solves the position updating task by combining the available memory and sensory information, on the one hand the internally-updated position of the pre-motion target, denoted m, and on the other hand the post-motion probe target position, denoted v, in a statistically optimal fashion, i.e., according to a causal Bayesian inference mechanism [7,9]. The ideas of this approach are now summarized informally, but more details can be found in the supplemental material.

The causal Bayesian inference model is principally probabilistic: both update m and visual probe percept v are considered to be contaminated by noise and represented as probability distributions, taken to be Gaussian. In addition, the model involves a *prior* distribution, also Gaussian, representing the participant's a priori beliefs about target position, independent of trial information. According to the model, on each trial two hypotheses are considered: one being that m and v have a common cause (here: the probe was displayed at the correct internally-updated target position), the other that they have distinct causes (the probe was displaced relative to the internally-updated location).

Under the first hypothesis (v gives 'true' information) the optimal way to combine the m, v, and prior distributions is by Bayesian integration, resulting in a Gaussian distribution with intermediate mean and higher precision, with precision defined as inverse variance. To be precise, the mean of the integration is the average of the m and v and prior means, each weighted

by its own precision, while the integration precision is the sum of the m, v, and prior precisions.

Under the second hypothesis (v is from displaced probe) the optimal way to proceed is simply

ignore v and just integrate m and the prior. This is called segregation.

To optimally apply the distributions for the two hypotheses, the integration distribution for a

correctly positioned probe, and the segregation distribution for a displaced probe, the

probability of the probe being displaced or not is still needed. The model assumes the participant

has a prior probability for the probe being correctly positioned, which on each trial is combined

with the m and v information of that trial to result in the corresponding posteriori probability.

(Qualitatively: the less overlap between m and v, the more evidence for displaced v; again, for

the equations see supplemental material). The final model distribution is the mixture of the

integration and segregation distributions, each weighted by the posteriori probability of the

corresponding hypothesis.

In fitting this model, we have to decide about specifications of the various distributions

involved. Priors are regarded as free parameters: the prior probability p_C of correct probe

position and the mean π and variance σ_π^2 of the prior for pre-movement target position. The

distribution v for the visually presented probe is supposed to be accurate with variance σ_v^2

treated as a free parameter.

The distribution m for the internally-updated position of the pre-motion target cannot be

assumed to be accurate: it has been established that, under the conditions of our experiment,

passive self-motion amplitude is underestimated [16]. We allow for such underestimation by

introducing a gain factor α. For a pre-motion target position s in body coordinates, the correct

update after a movement with amplitude $\mathbf{D}$ would be $s - \mathbf{D}$. Assuming underestimation of

distance $\mathbf{D}$ by a factor α, however, the distribution m is not centered on $s - \mathbf{D}$, but on $s +$

$(1 - \alpha) * D$. For each participant this gain α was estimated in the first block of update-only

trials, which estimate was then used and repeatedly updated in the experiment. The precision
of m, on the other hand, contributes another free parameter σ_m^2.

Figure 2**. Model's variables and predictions**. (A) Variables: given a target was initially
presented at the location s, the task of the observer was to estimate its position after having been
passively moved over a distance D. To do so, the observer can use two sources of information:
the internally updated target location m and the probe target position v. However, because the
self-motion is underestimated, the internally updated target location m would not actually be
centered on s but rather on $s + (1 - \alpha) * D$. (B) Models' predictions for response error (left)
and variability (right): The causal inference model (green) predicts that localization response is
biased toward the probe target location (black horizontal lines) for relatively small spatial
discrepancies between the probe target (horizontal gray bars) and the internal update m (red
dashed horizontal line) and becomes aligned with m for larger discrepancies. Accordingly,
response variability should be greatest for intermediate discrepancies while decreasing as the
probe target and the internal update get optimally integrated (small discrepancies) or segregated
(large discrepancies). Alternative model are depicted in orange (optimal integration), red (visual

271 feedback only) and light blue (memory-only), see text for further explanation. Because, in these
272 models, m and v are always either integrated or segregated according to these alternative
273 models, the corresponding predicted response variability should not depend on the probe shift.
274 Parameters used to generate these predictions are: $\sigma_v = 25 \ mm$, $\sigma_\pi = 400$ mm, $\sigma_m = 60 \ mm$,
275 $\alpha = 0.5$, $p(C) = 0.25$, $\pi = 0$.

Alternative models

The Bayesian causal inference model is a true 'ideal observer' model. All available information

is used in a statistically optimal way, including computation of the posterior probe displacement

probability for determining the weights in the mixture model. Arguably, this theoretical

benchmark is hard to attain in real life and the brain might instead resort to some approximating

heuristic. One possibility is that it does not compute probabilities for the two competing

hypotheses, correct or displaced probe position, but deterministically chooses one or the other,

i.e., always integrates or always segregates, respectively. The first alternative, forced fusion,

has been previously suggested for saccadic eye-movements [6,27]. It is an extreme case of the

causal inference mixture model with 0-1 weights. Similarly, choosing to always segregate

corresponds to the opposite 0-1 weighting in the mixture model.

A third possible heuristic is segregation with roles reversed: just process the directly given

visual cue and do not consider memory updates. The plausibility of this approach derives from

the fact that during passive whole-body movements the brain lacks the possibility to predict the

consequence of self-motion due to the absence of motor commands. The usefulness of keeping

track of memory and applying updates is not clear for naturalistic situations, where continuous

visual feedback is available. For the update-only trials without visual feedback the memory

model is retained, based on the assumption that, in the absence of other, more precise

information, the brain is able to spatially update an object based on an internal estimate of self-

motion.

Next to the causal inference model also these three simpler models, one integration and two

segregation models, were fitted and the model fits were compared.

Models predictions

Fig 2B shows the predictions of the respective models of the response error (left panel) and response variability (right panel). Generally, the memory-only model (light blue line) corresponds to a flat line (with as intercept the weighted average of one minus the updating gain times motion amplitude and the participant's prior position, weighted by memory-update and prior precision, respectively). The visual-only model (dark red line) predicts a straight line with slope close to one (visual precision divided by sum of visual and prior precisons). The optimal integration model (orange line) is represented by a straight line with an intermediate slope, determined by the relative precisions of the visual probe target perception, internal memory update, and prior position. The causal inference (green line) prediction is very similar to the integration model close to zero probe target shift (reflecting a high weight for common cause, thus integration), but its slope diminishes in absolute value and may even reverse sign, curving back to the horizontal memory-only axis, reflecting the growing weight for the segregate-memory branch of the mixture distribution with increasing disceprancy of the visual feedback signal.

Furthermore, the causal inference model also makes testable predictions about response variability (Fig 2B, right panel). The integration branch of this mixture model, which minimizes variability, has a high weight (common cause probability) close to zero probe target shift and this weight decreases with growing spatial discrepancy, i.e., with growing shift amplitude. Therefore, our participants' response variability should decrease as the spatial discrepancy between the expected and actual feedback decreases. In contrast, the weights predicted by the optimal integration model as well as by the segregation models (memory-only and visual-only) do not depend on the actual spatial discrepancy between the internally-updated target position and the probe position. Therefore, as predicted by these models, response variability as a function of probe displacement follows a flat line whom intercept corresponds depends on the

precision of the internal update, of the perception of the probe location and of the prior position

(optimal integration), or on the precision of the internal update and of the prior position

(memory-only), or on the precision of the perception of the probe location and of the prior

position (visual-only).

Model fitting and evaluation

The causal inference model has six free parameters: the variances σ_m^2 and σ_v^2 of the $\boldsymbol{m}$ and $\boldsymbol{v}$

distributions, the mean π and variance σ_π^2 of the target prior, the updating gain α, and the prior

probability p_C of correct probe position. The last parameter has no role in the integration model

and the visual-only segregation model, for which five parameters are left. The memory-only

model has again one parameter less, the variance of the ignored distribution, σ_v^2, for a total of

four parameters. All these models were fit to 1D localization data from both update-only and

test trials, which were composed of 8 displacement sizes, 5 targets locations and 2 self-motion

directions. Given there is no visual feedback during updating-only trials, we assumed subjects

used the memory-model in these conditions to compute their estimate. Thus, the best-fitting

parameters set were optimized for both updating-only and test trials.

Parameters were fit by maximum likelihood estimation. Lacking an analytical solution for the

mixture model likelihood, this likelihood was sampled by taking 10,000 draws from each of the

$\boldsymbol{m}$ and $\boldsymbol{v}$ distributions. Although the segregation and integration processes possess closed-form

likelihoods, we used the same simulation approach throughout for consistency. From the

discrete draws, a likelihood function was obtained by kernel density estimation (KDE, [28]).

We chose a Gaussian kernel with bandwidth h determined by Silverman's rule of thumb ($h = (4/3n)^{1/5}\hat{\sigma} \cong 0.168\,\hat{\sigma}$, with $\hat{\sigma}$ the sample SD; Silverman, 1986). This KDE approach has been

used before [29], but alternatively the model draws can be binned into a likelihood histogram

[7,9] or the likelihood can be approximated without sampling through numerical integration

[30] or linearization [31]. To our knowledge, no study has explicitly compared these

approximations.

Likelihood optimization was performed numerically using Bayesian Adaptive Direct Search

(BADS, [32]). BADS requires specification of upper and lower bounds as well as plausible

upper and lower bounds; the bounds we used can be found in the supplemental material (Table

S1). The upper bound for the visual noise parameter (σ_v) was set to 100 mm based on the largest

visual noise level estimated in a control condition (see Table S4 in supplemental material). For

every model and participant we computed 100 fits using random parameter initializations and

selected the best from these. Models' goodness of fit was assessed using the root mean square

error (RMSE). For model comparison the Bayesian Information Criterion (BIC, [33]) was used.

To validate our fitting procedure, we performed both a parameter and model recovery analysis

to ensure parameters and models can be inferred well using our experimental design and

analysis pipeline (see Tables S2 and S3 in supplemental material).

Results

The present study aimed at determining the inference mechanism used by the brain to estimate

the position of an object after a passive, whole-body translation. More specifically, we were

interested in examining whether the brain would only rely on the visual feedback, given it is

continuously available during self-motion, or also consider the expected sensory feedback, and

then either just use the latter, or always integrate both sources of feedback, or weighting these

two possibilities according to the causal inference model. To do so, we designed a spatial

updating task across whole-body passive translation where our participants' task was to

remember the world-fixed position of a target displayed prior to the translation and then shown

again as a probe at the end of the displacement, at a location shifted relative to the internally

updated target position. We were particularly interested in examining the effect of the probe

shift on the response bias and variability.

Estimation of updating gain

For each target position, the internally updated position around which to present the probe target

was determined based on the individual updating gain parameter α, during the experiment

estimated and updated as the participant's average across all preceding update-only trials. On

average, participants had an updating gain of 0.47 (SE= 0.05). The updating gains between

leftward and rightward motion were not significantly different ($\alpha_{left} = 0.47, \alpha_{right} =$

$0.46, t(10)=0.451$, CI(95%)=[-0.018, 0.027], p=0.662), so individual updating gains were

captured by a single parameter in our models.

Response error and variability

Fig 3A shows the response error of a typical participant plotted as a function of probe shift from

the internally-updated target position (response error of all participants is shown in Fig S2 in

supplemental material). While data are replicated from panel to panel, each panel illustrates a

specific model prediction. Fig 3B shows the same plots for the group data. While three models

predict the relationship to be linear with either a null (memory-only) or positive (visual-only

and integration) probe shift slope, causal inference posits a curvilinear relationship with slopes

decreasing from zero shift outwards, possibly reversing direction. In a linear mixed model

analysis of the response error data, with motion direction and probe target shift as predictors,

this opposed curvature (concave down for positive, concave up for negative probe shifts) was

modeled by a 'signed' quadratic component: the probe shift values squared with a sign reversal

added for negative shifts. Motion direction had no main or interaction effect (all $p > 0.38$).

Beside a positive linear effect ($\chi^2(1) = 10.23, p = 0.001$), probe shift also revealed a significant

($\chi^2(1) = 5.37, p = 0.020$) quadratic effect (with the sign consistent with the CI model). The

mixed model required random effects for intercept and both linear and quadratic shift

components.

Figure 3. Response error for an individual participant (A) and the group average (B). (A)
Response error plotted as a function of the spatial discrepancy between the internally-updated
target location and the probe target location together with models' predictions. Small blue dots
represent individual responses, whereas white-filled dots and error bars present the average
response and its standard deviation for a specific condition. Solid, orange lines show models'
predictions and colored areas display the standard deviation of the predicted response
distribution. Colored dashed line represents the average internally-updated target position,
whereas the gray dashed line stands for the target position and the black horizontal bars indicate
the probe target position. (B) average participants response error plotted as a function of probe
target displacement. Error bars represent the standard error of the mean (S.E.M). Each dot is
the average response of one participant. Solid orange line and shaded area represent the average
model prediction and the standard error of the mean respectively. Horizontal dashed gray line
depicts the target position, whereas the red dashed line and the solid, black lines indicate the
average internally-updated target and the probe target positions respectively.

Fig 4A shows the response variability (standard deviation of response error) of an example

participant along with our models predictions (one model per panel. Response variability of all

participants is shown in Fig S3 in supplemental material). Similarly, Fig 4B presents the same

plots at the group level. These variability data were also subjected to a mixed model analysis

with motion direction and a linear plus (ordinary) quadratic probe target component as

predictors. Again, no main or interaction effect for motion direction was found (all $p > 0.86$).

Target shift showed no linear ($\chi^2(1) = 0.24$, $p = 0.621$), but a higly significant ($\chi^2(1) = 62.47$, p

$< 10^{-14}$) concave-up quadratic effect, as predicted by the CI model. Here, the random part

consisted of a random intercept only.

**Figure 4. Average response variability for an individual participant (A) and the group
average (B).** (A) Individual participant's response variability (standard deviation of response
error) and models' predictions (solid, colored lines) plotted as a function of the spatial
discrepancy between the internally-updated target location and the probe target location, as well
as for "updating-only" trials (data plotted in gray area). (B) Average participants response
variability plotted as a function of target displacement relative to the internally-updated
location. The Causal inference model is the only of the tested models that captures the effect of
spatial discrepancy on our participants response variability.

Model fits and evalution

Bayesian causal inference predicts that the memory-based updated target location and the visual feedback are integrated in proportion to the posterior probability that they refer to a same target location (common cause). We tested this hypothesis against the predictions made by three alternative models. One of these is an optimal integration model that combines the memory update and visual feedback based on their respective precisions regardless of their spatial discrepancy. The other two models rely on the heuristic of using just one of both sources: a memory-only model disregarding the probe target and a visual-only model disregarding the internal update. Predictions of the four models are outlined in Fig2B (see Methods).

To quantitatively compare the predictions of our four models at the individual level, we computed model fits per participant. Root-mean-square error (RMSE) computed on each participant's average response and the model predictions in every condition suggests that our participants' data were best described by the Causal inference model [RMSE = 12.74 mm, SE = 2.55 mm], followed by the optimal integration model [RMSE= 19.73 (2.54) mm], the memory-only model [RMSE= 37.53 (13.86) mm] and the visual-only model [RMSE= 74.68 (10.00) mm]. A similar analysis applied to our participants' response variability revealed that the variability pattern (Figs 3B and 4B) clearly could only be captured by the Causal Inference model's prediction [RMSE= 12.26 (2.72) mm], followed by the optimal integration model [RMSE=12.76 (2.00) mm], the "memory only" model [RMSE=26.94 (10.12) mm], and the visual only model [RMSE=30.91 (SE = 6.59)].

Figure 5. BIC differences between the Bayesian causal inference model and the three alternative models (optimal integration, memory only and visual only). Negative differences would suggest stronger evidence for the alternative model than for the causal inference model and an absolute difference smaller than 10 can be judged uninformative.

While this fit measure does not take the number of free parameters used by the models into account, we also computed the Bayesian Information criterion (BIC) for each individual model fit. Averaged across participants the causal inference model clearly outperformed the integration model ($\Delta BIC = 294.78$, SE = 93.25), the memory-only model ($\Delta BIC = 594.87$, SE = 272.06), and the visual-only model ($\Delta BIC = 1220.94$, SE = 201.87). Given that differences in BIC larger than 20 are considered strong evidence for one model against the other [34,35], this suggests overwhelming evidence for the causal inference model compared to the three alternative models (see Fig 5). Table 1 lists the best fit parameters of the causal inference model and parameter fits of all the models are presented in the supplemental material (Table S1).

Participant	σ_v (mm)	σ_π (mm)	α	σ_m (mm)	$p(C)$	π (mm)
s1	25.74	364.78	0.41	32.50	0.17	93.75
s2	81.26	106.94	-0.02	199.98	0.06	-3.41
s3	10.02	144.71	0.44	78.51	1.00	17.66
s4	28.72	128.58	1.04	61.06	0.24	-22.26
s5	99.90	80.31	0.05	55.66	0.68	3.08
s6	8.47	193.40	0.41	34.57	0.68	27.50
s7	3.45	170.35	0.70	72.13	0.37	-25.28
s8	100.00	76.61	-0.08	52.40	0.38	24.97
s9	40.18	291.16	0.97	183.61	1.00	-11.44
s10	100.00	76.93	-0.10	77.43	1.00	-3.78
s11	9.92	115.81	0.62	46.86	0.89	-8.67
Median	28.72	128.58	0.41	61.06	0.68	-3.41
Mean	46.15	159.05	0.40	81.34	0.59	8.37
S.E.M	12.26	28.05	0.12	17.14	0.11	10.04

Table 1: Best parameters recovered by the Causal inference model for every participant.

Discussion

Because images from world-centered objects shift on our retina when the eyes move in space,

either by eye rotation or body motion, the brain must compensate for our motion in order to

keep a spatially constant representation of the world. This spatial updating mechanism involves

that the brain internally updates remembered object locations based on the estimated self-

motion. To explain spatial constancy across saccadic eye movements, we recently suggested

that the brain relies on a causal inference mechanism [9], showing that the updated target

location after saccades and the new visual feedback are integrated and/or segregated based on

the posterior probability that they refer to the same position in the world. However, passive

self-motion differs from saccadic eye-movements in that visual feedback is typically

continously available and spatial updating cannot rely on the efference copy of the motor

commands. Therefore, the present study aimed at determining whether the brain would call on

Bayesian causal inference or whether it would use a simpler heuristic, for instance solely relying

on the available visual feedback for spatial updating after passive whole-body motion. In line

with the prediction made by Bayesian causal inference, our results show that responses were

biased toward the probe location, with proportionally stronger bias for small probe

displacements and a relatively smaller bias for the larger displacements. We conclude that, in

order to maintain spatial constancy across passive body motion, the brain would weigh the

internally-updated target position and the visual feedback based on the posterior probability

that they refer to a common position in the world.

As quantitative support for support this conclusion, we fitted a Bayesian Causal inference

model and compared its predictions to those of three alternative models: an optimal integration

model and a visual-only as well as a memory-only segregation model. The response patterns of

our participants could be better captured by the Causal inference model compared to these

alternative models, both in terms of quality of fit and of model evidence as measured by the

RMSE and the Bayesian Information Criterion. The fitted prior on the probability of having a

common cause was on average about 0.54, but could vary across participants, with some

participants being closer to the prediction of an optimal integration model ($p(C) \approx 1$, n = 3)

and others being closer to a segregation model ($p(C) \approx 0$, n = 1). Such variability has also been

reported in a previous study [9], perhaps related to effects of instruction and experimental

context.

Another important prediction of causal inference is that response variability should decrease

non-linearly as target and probe are optimally integrated, that is, as its spatial difference of the

probe target with the internally-updated target position is smaller. Again, this response

variability pattern was observed in most of our participants.

It is important to point out that our data cannot be used to determine whether our participants

were using a true Bayes-optimal causal inference strategy or a suboptimal approximation of it

[36]. More specifically, because the posterior probability of having a common cause partially

depends on how these two cues can be spatiotemporally discriminated, this would depend on

how precise their representation is. Our previous study investigating the role of causal inference

in spatial constancy across saccades manipulated the sensory reliability of the post-saccadic

probe target by varying its presentation duration [9]. It was found that for longer presentation

durations, which results in a more precise probe target representation, participants' responses

were more biased toward the probe location for small displacement (stronger integration) but

also more quickly disregarded as the size of the displacement increased. This result suggest that

observers could obey a true Bayes-optimal decision strategy in spatial constancy tasks.

While our results suggest updating responses rely on a Bayesian causal inference, they cannot

disentangle the different possible implementations of the decision strategy used in causal

inference. Here, we only considered causal inference as a model averaging approach: both

integration and segregation estimates are combined and weighted based on the posterior

probability of having a common cause in a particular trial. Alternative decision strategies have

been suggested as well and have been compared in previous studies, including model selection

[37], probability matching [9,38] and coupling prior [39,40]. Nonetheless, model averaging

most of the time outperformed these alternative decision strategies in predicting participants

response in cue-combination and unity judgment tasks [7,9,30,37].

Previous research has investigated spatial updating across body translation in complete

darkness using psychophysical procedure where participants have to compare the position of a

probe target to a reference [17,19]. Despite the similarity with our test trials where a probe

target was shown at the end of the translation, the aim of these studies was in fact closer to the

goal of our updating-only trials: measuring how an observer updates an object position based

on vestibular cues only. A critical difference with our test trials lies in the role played by the

probe target and the way it can be used. Comparing the probe position to the target position

necessarily implies to segregate these positions and to regard them as being generated by

different causes (compare one object vs. another). In contrast, we used an estimation task where

participants had to report the location of the target, possibly combining the probe location as an

additional source of information. In contrast to using a 2-AFC task as previously described, our

estimation procedure allows us to determine the way visual feedback regarding the target can

be used in order to more precisely update the target location.

Our spatial updating task involved passive linear body-translation in complete darkness with

restricted eye-movements due to the body-fixed fixation target. This allowed us to better control

the types of sources of information that the brain could use in order to estimate the translation

amplitude and consequently update the target location. Optic flow was not available at all.

Therefore, the amplitude estimate could mostly be derived from the integration of canals and

otoliths signals about the angular and linear acceleration of the head in space respectively

[15,41]. Increasing the number of sensory and motor sources about self-motion, reflecting more

ecological situations, however, would not rule out the use of causal inference. Integration of

these multiple sources of information would actually result in a more precise estimate of the

updated target location, which would then be better discriminated from the sensory feedback of

the target. The sharpening of the sensory cue estimates should finally impact on the posterior

probability of having a common cause and, consequently, on the weighting of integration and/or

segregation of these estimates. Interestingly, this causal inference could take place at multiple

stages of processing: e.g. related to the multisensory cue-combination to estimate self-motion,

and related to the integration of the sensory feedback in order to improve the estimate of the

target location or to detect an external change in the object.

We used a computational approach to describe and predict human updating behavior. Previous

theoretical work suggested that Bayesian optimal integration could be supported by the linear

combination of neural population activity that can be approximated by Poisson-like

distributions [42,43]. In line with this suggestion, it has been shown in macaques that MSTd

neurons compute the weighted sum of their inputs, the weights of which were varying according

to motion coherence, which was a manipulation of cue reliability [44]. Similar evidence for

neuronal correlates of optimal integration has been found in studies involving visuo-vestibular stimuli [45]. More recently, an audio-visual cue-combination study combined with fMRI found evidence for a possible cortical hierarchy implementing causal inference [46]. At the bottom of the hierarchy, primary sensory areas encode their preferred stimulus (segregation estimates), whereas optimal integration of the sensory cues and the encoding of the uncertainty about their causal structure would occur higher in the hierarchy (posterior intraparietal sulcus and anterior intraparietal sulcus respectively). This study, however, considered Bayesian causal inference in a task that involves the combination of unisensory cues. In contrast, spatial updating involves the combination of a sensory feedback and an internal, amodal estimate of the expected sensory feedback itself derived from multisensory sources of information. Therefore, it remains to be determined whether this cortical hierarchy would also support Bayesian causal inference for spatial constancy. Interfering techniques, such as TMS, could be used in order to test whether it affects how participants integrate sensory cues given their spatial disparity in spatial updating tasks.

Acknowledgments

This research was supported by a VICI grant to W.P.M.

Author contributions

Conceived and designed the experiments: FP JC WPM. Performed the experiments: FP. Analyzed the data: FP JC MK WPM. Wrote the paper: FP JC MK WPM.

References

1. Ross J, Burr D, Morrone MC. Suppression of the magnocellular pathway during saccades. Behavioural Brain Research. Elsevier; 1996. pp. 1–8. doi:10.1016/0166-4328(96)00012-5

2. Campbell FW, Wurtz RH. Saccadic omission: Why we do not see a grey-out during a

saccadic eye movement. Vision Res. Pergamon; 1978;18: 1297–1303.

doi:10.1016/0042-6989(78)90219-5

3. Helmholtz H von. Handbuch der physiologischen Optik [Handbook of physiological

optics]. Leipzig : Leopold Voss, 1867; 1 v.~; in 8.; DCCC.4.353. Hamburg, Germany:

Voss; 1867.

4. Cavanaugh J, Berman RA, Joiner WM, Wurtz RH. Saccadic Corollary Discharge

Underlies Stable Visual Perception. J Neurosci. Society for Neuroscience; 2016;36: 31–

42. doi:10.1523/JNEUROSCI.2054-15.2016

5. Wurtz RH. Neuronal mechanisms of visual stability. Vision Res. Pergamon; 2008;48:

2070–2089. doi:10.1016/j.visres.2008.03.021

6. Niemeier M, Crawford JD, Tweed DB. Optimal transsaccadic integration explains

distorted spatial perception. Nature. Nature Publishing Group; 2003;422: 76–80.

doi:10.1038/nature01435

7. Körding KP, Beierholm U, Ma WJ, Quartz S, Tenenbaum JB, Shams L. Causal inference

in multisensory perception. PLoS One. 2007;2. doi:10.1371/journal.pone.0000943

8. Bridgeman B, Hendry D, Stark L. Failure to detect displacement of the visual world

during saccadic eye movements. Vision Res. 1975;15: 719–722. doi:10.1016/0042-

6989(75)90290-4

9. Atsma J, Maij F, Koppen M, Irwin DE, Medendorp WP. Causal Inference for Spatial

Constancy across Saccades. PLoS Comput Biol. 2016;12: 1–20.

doi:10.1371/journal.pcbi.1004766

10.   Ross J, Morrone MC, Goldberg ME, Burr DC. Changes in visual perception at the time

600   of saccades. Trends Neurosci. 2001;24: 113–121. doi:10.1016/S0166-2236(00)01685-4

601   11.   Burr DC, Morrone MC, Ross J. Selective suppression of the magnocellular visual

602   pathway during saccadic eye movements. Nature. 1994;371: 511–513.

603   doi:10.1038/371511a0

604   12.   Lappe M, Bremmer F, Van Den Berg AV. Perception of self-motion from visual flow.

605   Trends Cogn Sci. 1999;3: 329–336. doi:10.1016/S1364-6613(99)01364-9

606   13.   Fetsch CR, Turner AH, DeAngelis GC, Angelaki DE. Dynamic re-weighting of visual

607   and vestibular cues during self-motion perception. J Neurosci. Society for Neuroscience;

608   2009;29: 15601–15612. doi:10.1523/JNEUROSCI.2574-09.2009

609   14.   Ter Horst AC, Koppen M, Selen LPJ, Medendorp WP. Reliability-based weighting of

610   visual and vestibular cues in displacement estimation. PLoS One. 2015;10.

611   doi:10.1371/journal.pone.0145015

612   15.   Berthoz A, Israel I, Georges-Francois P, Grasso R, Tsuzuku T. Spatial memory of body

613   linear displacement: what is being stored? Science. 1995;269: 95–98.

614   doi:10.1126/science.7604286

615   16.   Clemens IAH, Selen LPJ, Pomante A, MacNeilage PR, Medendorp WP. Eye Movements

616   in Darkness Modulate Self- Motion Perception. eNeuro. 2017;4: 1–12.

617   doi:10.1523/ENEURO.0211-16.2016

618   17.   Clemens IAH, Selen LPJ, Koppen M, Medendorp WP. Visual stability across combined

619   eye and body motion. J Vis. 2012;12: 8–8. doi:10.1167/12.12.8

18.   Bresciani J-PP, Gauthier GM, Vercher J-LL, Blouin J. On the nature of the vestibular

control of arm-reaching movements during whole-body rotations. Exp Brain Res.

Springer-Verlag; 2005;164: 431–441. doi:10.1007/s00221-005-2263-4

19.   Tramper JJ, Medendorp WP. Parallel updating and weighting of multiple spatial maps

for visual stability during whole body motion. J Neurophysiol. American Physiological

Society; 2015;114: 3211–3219. doi:10.1152/jn.00576.2015

20.   Klier EM, Hess BJM, Angelaki DE. Human Visuospatial Updating After Passive

Translations In Three-Dimensional Space. J Neurophysiol. 2008;99: 631–637.

doi:10.1152/jn.01091.2007.Human

21.   Wei M, Li N, Newlands SD, Dickman JD, Angelaki DE. Deficits and recovery in

visuospatial memory during head motion after bilateral labyrinthine lesion. J

Neurophysiol.   American   Physiological   Society;   2006;96:   1676–1682.

doi:10.1152/jn.00012.2006

22.   Gibson JJ. The Perception of the Visual World [Internet]. Carmichael L, editor. Boston:

Houghton Mifflin; 1950.

23.   Israël I, Chapuis N, Glasauer S, Charade O, Berthoz A. Estimation of passive horizontal

linear whole-body displacement in humans. J Neurophysiol. American Physiological

Society; 1993;70: 1270–1273.

24.   Mackrous I, Simoneau M. Improving spatial updating accuracy in absence of external

feedback.   Neuroscience.   IBRO;   2015;300:   155–162.

doi:10.1016/j.neuroscience.2015.05.024

25.   R Development Core Team. R: a language and environment for statistical computing.

Citeseer; 2016.

26.   Bates DM, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using

644   lme4. J Stat Softw. 2015;67: 1–48. doi:10.1177/009286150103500418

645   27.   Vaziri S, Diedrichsen J, Shadmehr R. Why Does the Brain Predict Sensory

646   Consequences of Oculomotor Commands? Optimal Integration of the Predicted and the

647   Actual   Sensory   Feedback.   J   Neurosci.   2006;26:   4188–4197.

648   doi:10.1523/JNEUROSCI.4747-05.2006

649   28.   Botev ZI, Grotowski JF, Kroese DP. Kernel density estimation via diffusion. Ann Stat.

650   2010;38: 2916–2957. doi:10.1214/10-AOS799

651   29.   De Winkel KN, Katliar M, Bülthoff HH. Causal inference in multisensory heading

652   estimation.   Lappe   M,   editor.   PLoS   One.   Springer;   2017;12:   1–20.

653   doi:10.1371/journal.pone.0169676

654   30.   Acerbi L, Dokka K, Angelaki DE, Ma WJ. Bayesian Comparison of Explicit and Implicit

655   Causal Inference Strategies in Multisensory Heading Perception. bioRxiv. 2017; 150052.

656   doi:10.1101/150052

657   31.   De Winkel KN, Katliar M, Diers D, Bülthoff HH. Causal Inference in the Perception of

658   Verticality. Sci Rep. 2018;8: 12. doi:10.1038/s41598-018-23838-w

659   32.   Acerbi L, Ma WJ. Practical Bayesian Optimization for Model Fitting with Bayesian

660   Adaptive Direct Search. In: Guyon I, Luxburg U V, Bengio S, Wallach H, Fergus R,

661   Vishwanathan S, et al., editors. Advances in Neural Information Processing Systems 30.

662   Curran Associates, Inc.; 2017. pp. 1836–1846. doi:https://doi.org/10.1101/150052

663   33.   Raftery AE. Bayesian Model Selection in Social Research. Sociol Methodol. American

664   Sociological Association; 1995;25: 111–163. doi:10.2307/271063

665   34.   Kass RE, Raftery AE. Bayes factors. J Am Stat Assoc. 1995;90: 773–795.
666   doi:10.1080/01621459.1995.10476572

667   35.   Efron B, Gous A. Scales of Evidence for Model Selection: Fisher versus Jeffreys. IMS
668   Lect Notes - Monogr Ser. 2001;38: 210–246.

669   36.   Ma WJ. Organizing probabilistic models of perception [Internet]. Trends in Cognitive
670   Sciences. 2012. pp. 511–518. doi:10.1016/j.tics.2012.08.010

671   37.   Rohe T, Noppeney U. Sensory reliability shapes perceptual inference via two
672   mechanisms. J Vis. The Association for Research in Vision and Ophthalmology;
673   2015;15: 22. doi:10.1167/15.5.22

674   38.   Wozny DR, Beierholm UR, Shams L. Probability matching as a computational strategy
675   used in perception. PLoS Comput Biol. Public Library of Science; 2010;6: e1000871.
676   doi:10.1371/journal.pcbi.1000871

677   39.   Ernst MO. A Bayesian view on multimodal cue integration. Hum body Percept from
678   Insid out. 2006;131: 105–131. doi:10.1016/j.cub.2011.11.039

679   40.   Garzorz IT, MacNeilage PR. Visual-Vestibular Conflict Detection Depends on Fixation.
680   Curr Biol. Elsevier; 2017;27: 2856–2861.e4. doi:10.1016/j.cub.2017.08.011

681   41.   Angelaki DE, Cullen KE. Vestibular System: The Many Facets of a Multimodal Sense.
682   Annu Rev Neurosci. 2008;31: 125–150. doi:10.1146/annurev.neuro.31.060407.125555

683   42.   Ma WJ, Beck JM, Latham PE, Pouget A. Bayesian inference with probabilistic
684   population codes. Nat Neurosci. Nature Publishing Group; 2006;9: 1432–1438.
685   doi:10.1038/nn1790

43.   Ma WJ, Beck JM, Pouget A. Spiking networks for Bayesian inference and choice. Curr

Opin Neurobiol. 2008;18: 217–222. doi:10.1016/j.conb.2008.07.004

44.   Morgan ML, DeAngelis GC, Angelaki DE. Multisensory Integration in Macaque Visual
Cortex   Depends   on   Cue   Reliability.   Neuron.   2008;59:   662–673.
doi:10.1016/j.neuron.2008.06.024

45.   Fetsch CR, Pouget A, DeAngelis GC, Angelaki DE. Neural correlates of reliability-based
cue weighting during multisensory integration. Nat Neurosci. Nature Publishing Group;
2011;15: 146–154. doi:10.1038/nn.2983

46.   Rohe T, Noppeney U. Cortical Hierarchies Perform Bayesian Causal Inference in
Multisensory Perception. PLoS Biol. 2015;13: 1–18. doi:10.1371/journal.pbio.1002073

47.   Silverman B. Density estimation for statistics and data analysis. Chapman and Hall.
Chapman and Hall; 1986. doi:10.2307/2347507

48.   Kontsevich LL, Tyler CW. Bayesian adaptive estimation of psychometric slope and
threshold. Vision Res. 1999;39: 2729–2737. doi:10.1016/S0042-6989(98)00285-5

49.   Prins N. The psi-marginal adaptive method: How to give nuisance parameters the
attention they deserve (no more, no less). J Vis. The Association for Research in Vision
and Ophthalmology; 2013;13: 1–17. doi:10.1167/13.7.3

50.   Prins N, Kingdom FAA. Palamedes: Matlab routines for analyzing psychophysical data.
2009.

51.   Ernst MO, Banks MS. Humans integrate visual and haptic information in a statistically
optimal   fashion.   Nature.   Nature   Publishing   Group;   2002;415:   429–433.
doi:10.1038/415429a