Original Article

Growth trajectories better reflect the influence of soil variation than functional traits in

a widespread pioneer Neotropical tree

Running title: **Soil-related phenotypic variability for a widespread tree species**

Sébastien Levionnois[1*], Niklas Tysklind[2], Eric Nicolini[3], Bruno Ferry[4], Valérie Troispoux[2],

Gilles Le Moguedec[5], Hélène Morel[6], Clément Stahl[2], Sabrina Coste[7], Henri Caron[2, 8], Patrick

Heuret[2, 5*]

1. CNRS, UMR EcoFoG, AgroParisTech, CIRAD, CNRS, INRA, Université des Antilles,

Université de Guyane, 97310 Kourou, France.

2. INRA, UMR EcoFoG, AgroParisTech, CIRAD, CNRS, INRA, Université des Antilles,

Université de Guyane, 97310 Kourou, France.

3. CIRAD, UMR AMAP, CIRAD, CNRS, INRA, IRD, Université de Montpellier, 34398

Montpellier, France.

4. AgroParisTech, UMR SILVA, AgroParisTech, INRA, 54280 Champenoux, France.

5. INRA, UMR AMAP, CIRAD, CNRS, INRA, IRD, Université de Montpellier, 34398

Montpellier, France.

6. CIRAD, UMR EcoFoG, AgroParisTech, CIRAD, CNRS, INRA, Université des Antilles,

Université de Guyane, 97310 Kourou, France.

7. Université de Guyane, UMR EcoFoG, AgroParisTech, CIRAD, CNRS, INRA, Université

des Antilles, Université de Guyane, 97310 Kourou, France.

8. INRA, UMR BIOGECO, INRA, Université de Bordeaux, Cestas, France

*Authors for correspondence:

Sébastien Levionnois, PhD student

Postal adress: UMR EcoFog, Campus Agronomique, Avenue de France

BP709 – 97387 Kourou Cedex

French Guiana

Tel: +33 760 26 73 83

Email: sebastien_levionnois@live.fr

Patrick Heuret, PhD

Postal adress: UMR AMAP, TA- A-51/PS2 Boulevard de la Lironde

34398 Montpellier Cedex 5 (France)

French Guiana

Tel: +33 467 61 55 14

Email: patrick.heuret@inra.fr

Author contributions

PH designed and led the project. PH, SL, EN, VT, HM, NT, CS, SC, BF and HC measured

tree architecture and functional traits. BF and VT described soils characteristics. SL, PH and

GL performed data analysis. SL wrote the manuscript with contributions from PH and NT.

All authors contributed critically to the drafts and gave final approval for publication.

ABSTRACT

1. Trait-environment relationships have been described at the community level across tree

species. However, whether interspecific trait-environment relationships are maintained at the

intraspecific level is yet unknown.

2. We examined phenotypic variability for 16 functional leaf (dimensions, nutrient,

chlorophyll) and wood traits (density) across two soil types, Ferralitic Soil (FS) vs. White

Sands (WS), on two sites for 70 adult trees of *Cecropia obtusa* Trécul (Urticaceae) in French

Guiana. *Cecropia* is a widespread pioneer Neotropical genus that generally dominates forest

early successional stages. To understand how soil types impact resource-use through the

processes of growth and branching, we examined the architectural development with a

retrospective analysis of growth trajectories.

3. Functional trait responses to soil types were weak, as only two traits -leaf residual water

content and K content- showed significant differences across soil types. Soil effects were

stronger on growth trajectories and tree architectural development, with WS trees having the

slower growth trajectories -mediated through smaller internode length-, smaller trunk heights

and diameters for a given age, and less numerous branches across their lifespan.

4. The analysis of growth trajectories based on architectural analysis improved our ability to

detect the effects of soil types compared to measured functional traits. Intraspecific variability

is higher for growth trajectories than functional traits for *C. obtusa*, revealing the higher

sensitivity of the architectural approach in comparison to the functional approach in the case

of *C. obtusa*. Soil-related response of *Cecropia* functional traits is not homologous to the

community-level environmental filtering, suggesting that the effects of the acting ecological

processes are different between the two levels.

Key words: architecture, *Cecropia obtusa*, ferralitic/white-sand soils, functional traits,

growth trajectory, intraspecific variability

INTRODUCTION

Trait-based community ecology seeks to predict the processes of assemblage and maintenance of plant communities over time and space (McGill *et al.* 2006). The key questions in this field are (i) the identification of ecological processes determining community composition (McGill *et al.* 2006; Shipley *et al.* 2016), and (ii) the role of intraspecific variability (ITV) in community assemblages, and to what extent ITV can be ignored by using species-level functional trait means (Violle *et al.* 2012; Shipley *et al.* 2016). Trait-based approaches have improved our understanding of the role of ecological processes in community assemblage. Environmental filtering drives community assemblage through the interaction of individuals with the abiotic environment (Kraft *et al.* 2015): Physiologically challenged individuals are eliminated, so that the breadth of functional trait values is predicted to be small (i.e. functional trait under-dispersion) within local communities. Another process, niche differentiation is based on the interaction of neighbouring individuals, and incorporates the effects of both resource competition and shared predators (Uriarte *et al.* 2004). For species co-existence, they cannot share exactly the same niche, such that evenness of functional trait value distribution is predicted to be high, leading to functional trait over-dispersion within local communities. Both ecological processes, environmental filtering and niche differentiation, have been demonstrated for various habitats and landscapes, with environmental filtering tending to be more pervasive (Kraft *et al.* 2008; Swenson and Enquist 2009; Paine *et al.* 2011; HilleRisLambers *et al.* 2012).

ITV has long been ignored, or at least underestimated, in trait-based community ecology (Violle *et al.* 2012; Shipley *et al.* 2016). This has been the case for studies investigating ecological processes of community assemblages (Schamp *et al.* 2008; Kraft *et al.* 2008; Swenson and Enquist 2009); but see Paine *et al.* (2011). ITV may allow a species to thrive in several communities. First, displaying a large ITV would allow a species to fit a large abiotic spectrum, since there is a higher probability that the required functional trait values compatible with the habitat fall into the possible range of functional trait values of the species. In such a case, the success of individuals in terms of environmental filtering is promoted. Second, displaying a large ITV allows different ways to avoid functional trait similarity with neighbours, and contributes to the niche differentiation among habitats. Recent efforts for incorporating ITV into trait-based community ecology have validated these hypotheses (Fridley and Grime 2010; Jung *et al.* 2010, 2014; Lepš *et al.* 2011; Kraft *et al.* 2014). Moreover, studies have demonstrated that incorporating ITV increases the predictive

power of models for species interactions, trait-environment relationships, and ecosystem

productivity (Jung *et al.* 2010; Paine *et al.* 2011).

Generalist species are particularly useful to understand the role of ITV and

environmental filtering in the structuring of community composition. Generalist species are

defined here as species able to thrive within a larger range of abiotic conditions than most of

species, and generally regarding one kind of condition, i.e. topographic, edaphic, light... First,

generalist species tend to display large ITV, and by definition they inhabit large ecological

spectrums (Sides *et al.* 2014). Generalist species offer the opportunity to test hypotheses

regarding how ecological processes act at the intraspecific level, how functional traits are

mediated regarding the ecological processes, and if they do so in the same way than at the

interspecific level. Second, better knowledge on how ITV of functional traits is structured

should help to better decide if and how ITV must be accounted for in trait-based community

ecology, especially for generalist species, which are often regionally widespread and abundant

(Holt *et al.* 2002; Borregaard and Rahbek 2010; Boulangeat *et al.* 2012).

The Amazon rainforest has been a rich study field for investigating key questions on

trait-based ecology, such as relationships of functional traits with environmental gradients

(Kraft *et al.* 2008). Edaphic gradients have been particularly studied to disentangle drivers of

spatial distribution of species and functional traits over the Amazon basin (Sabatier *et al.*

1997; Clark *et al.* 1999; Stropp *et al.* 2011; Allié *et al.* 2015).The contrast between white-sand

(WS) *versus* ferralitic soils (FS) has been repeatedly used for explaining Amazonian spatial

species diversity turnover. WS are mainly quartz soils, representing 3% to 5% of soils in the

Amazon basin and exist as island-like spots in a matrix of other soils such as FS (Adeney *et*

al. 2016; Fine and Baraloto 2016).WS are poor in mineral nutrients and acidic, with low-

usable water reserves and poor nitrogen mineralization, especially in comparison with

common FS. The environmental filtering ensued by the FS-WS heterogeneity has strong

impacts on species distribution patterns at the community level (Stropp *et al.* 2011; ter Steege

et al. 2013; Daly *et al.* 2016; Fine and Baraloto 2016), as well as affecting functional traits.

WS flora is characterised by a convergence towards a conservative functional strategy of

nutrient acquisition because of the scarce nutrient availability and severe water stress (Grubb

and Coomes 1997; Patiño *et al.* 2009; Fyllas *et al.* 2009; Fine *et al.* 2010; Fortunel *et al.* 2012;

Fortunel, Paine, *et al.* 2014; Fortunel, Ruelle, *et al.* 2014; Fine and Baraloto 2016). These

imply higher leaf mass area (LMA), higher wood density, smaller seeds, and lower leaf

nutrient contents associated with higher nutrient use efficiency (Fine and Baraloto 2016), in

comparison with other soil types such as FS.

Many generalist species are able to grow on either end of the FS-WS gradient (Fine

and Baraloto 2016). A pervasive soil response has been demonstrated for the tree species

Protium subserratum Engl. (Burseraceae), where Fine *et al.*, found significant differences on

the chemical traits associated to herbivory resistance between individual growing on FS and

WS, paralleling the environmental filtering acting at the interspecific level (Fine *et al.* 2013).

However, we do not know how more commonly used, morphological, and easy-to-measure

functional traits (e.g. LMA, leaf area, leaf thickness, wood density…), largely used in trait-

based plant community ecology, vary at the intraspecific level between FS and WS, and if the

environmental response of these functional traits mirrors the environmental filtering acting at

the community level (Fortunel, Paine, *et al.* 2014).

Phenotypic adjustment to abiotic factors could occur in different manners according to

the plant compartment (i.e. root, trunk, or leaf), the function (assimilation, mechanical

stability, conduction…) or the life-history strategy (growth vs survival). For instance,

functional traits associated to resource acquisition (e.g. leaf and root traits) could vary

independently of functional traits related to resource use (e.g. growth, defence). Moreover,

Paine *et al.* (2011) have shown that leaf-level traits are more responsive to environmental

filtering than stem traits among communities. Easy-to-measure functional traits commonly

used in trait-based ecology are appropriate to capture a snap-shot image of the resource-

acquisition strategy (Baraloto *et al.* 2010), but fail to take into account the growth strategy,

which integrates the long term response of the individual to its environment. Phenotypic

adjustments could also occur at the whole plant structure level, namely, its architecture.

Numerous studies have shown that trees associated to stressful and limiting habitats (e.g. low

light exposure, low water availability) exhibit prostrated and frail plant-level structure, as also

low branching potential (Charles-Dominique *et al.* 2009, 2012; Stecconi *et al.* 2010). Limiting

habitats, such as WS, are physical boundaries in terms of available energy, water and nutrients

per unit of time for a given plant. Even if functional traits associated to resource acquisition

do not vary, the resource scarcity could have an effect on whole plant structure all along the

life of an individual. Here, we combine functional trait approaches with a whole-tree

developmental approach to understand tree phenotypic responses and the interaction between

ITV and environment. With a whole-tree developmental approach, we can consider the

development of the trunk for instance, described as a sequence of repetitive elementary units

(e.g. internode, growth unit, annual shoot), universal for vascular plants, and the accumulation

and fluctuation of growth, branching, and flowering processes through a tree's lifespan

(Heuret *et al.* 2006; Guédon *et al.* 2007; Taugourdeau *et al.* 2012). Here, we aim to elucidate

the role of ITV in functional traits and growth patterns in allowing species to thrive in

different environments by studying the Amazon rainforest genus *Cecropia,* composed of

hyperdominant pioneer tree species, critical in the recovery of Amazon forests.

We focus on *Cecropia obtusa* Trécul (Urticaceae), a widespread Guiana shield

generalist species, capable of growing on both FS and WS, and displaying perennial growth

marks, which allow for an analysis of life history based on architecture analysis (growth,

branching, flowering) through time, making *C. obtusa* a model species for tree architecture

and growth (Heuret *et al.* 2002; Zalamea *et al.* 2008; Mathieu *et al.* 2012; Letort *et al.* 2012).

We measured commonly used functional leaf and wood traits, coupled with architectural

development and growth trajectory analyses for *C. obtusa* individuals from two sites with

both soil types in French Guiana. We aim to answer the following questions:

(i)   Is the soil-response of *C. obtusa* mediated by changes in functional traits (i.e.

resource acquisition) or architectural development (i.e. resource use through the

process of growth)?

(ii)   Is the effect of environmental filtering on functional traits the same at the

intraspecific and interspecific levels?

MATERIALS AND METHODS

Terms and definitions

In this study, we use the term of "functional trait" according to the definition of Violle *et al.*

(2007), as any morphological, physiological, or phenological trait which impact fitness

indirectly via their effects on growth, survival, or reproduction. But in this study, functional

traits specifically refers to easy-to-measure traits, generally and massively measured in trait-

based ecology, and sometimes referred as soft traits (Violle *et al.* 2007): e.g. leaf area, leaf

mass area, wood density… These traits are generally measured at a specific given moment of

the plant's life, disconnected from the developmental trajectory, and ignoring potential

ontogenetical effects on the trait value. That is why we oppose functional traits to

architectural traits in our study. Architectural traits are defined here as morphological traits

directly related to growth and branching processes, and that can expressed as longitudinal

data: internode length, annual shoot length… according to plant height, or plant age, or node

ranking. We also used whole-tree-level traits, defined as traits capturing whole-tree features

of architecture such as tree height, the number of branches, the number of branching orders…

Such traits are generally harder to measure than soft traits we refer as functional traits in our

study, and are not expressible as longitudinal data as our architectural traits. Nonetheless, the

reader can keep in mind that our architectural and whole-tree-level traits fundamentally

remain functional traits according to the original definition of Violle *et al.* (2007). The goal of

the use of this specific terminology in the context of our study is to contrast the architectural

approach and related measurements, which are not so common in trait-based ecology.

Study species: Why C. obtusa *is an appropriate tree model species?*

C. obtusa has several characteristics that allow the retrospective construction of a tree's past

growth. The growth of *C. obtusa* is continuous (no cessation of elongation) and monopodial

(no death of meristem), the tree is made of a set of axes, where each one is composed of an

ordered, linear, and repetitive succession of phytomers (i.e. the set of a node, an internode, a

leaf, and its axillary buds; Fig. S1). Leaves are stipulated, with an enveloping stipule named

calyptra which has a protective function (Fig. S1). At the leaf establishment, the calyptra

sheds leaving a characteristic ring scar delimiting the associated internode, and usable as a

permanent growth marker. The 10-day stable phyllochron (i.e. rhythm of leaf production)

associated with such permanent growth marker allows for the retrospective analysis of tree

growth and development, covering the tree's lifespan (Heuret *et al.* 2002; Zalamea *et al.*

2012).

There are three lateral buds in the axil of each leaf (Fig. S1). The central bud is

vegetative and can develop into a new axis. The two others are proximal lateral buds of the

vegetative central one and can develop into inflorescences The inflorescences leave

permanent scars after shedding, allowing the retrospective analysis of tree's lifespan

flowering events. The same retrospective analysis is possible with branching events since the

presence of past branches remains visible.

Study site

Two sampling sites were selected in French Guiana: (1) Counami, along the Counami forestry

road (N5.41430°, W53.17547°, geodesic system WGS84); and (2) Sparouine, along the

national road 5 (N5.27566°, W54.20048°). The warm and wet tropical climate of French

Guiana is seasonal due to the north-south movement of the Inter-Tropical Convergence Zone.

Annual rainfall is 3,041 mm year^{-1} and annual mean air temperature is 25.7 °C at Paracou

experimental station (Gourlet-Fleury *et al.* 2004) situated nearly at 30 km and 150 km to the

east of Counami and Sparouine sites respectively. There is one long dry season lasting from

mid-August to mid-November, during which rainfall is < 100 mm month^{-1}. The two studied

sites (Counami and Sparouine) are characterised by rainfall differences (Fig. S2). Counami

shows higher levels of rainfall and higher contrasts between the long rainy and the long dry

seasons. For each of the two sites, two micro-localities are identified corresponding to two

soil types: ferralitic soils (FS) and white-sand soils (WS). Local sites were chosen to be well

drained and on upper slopes.

Plant material, study conception, and sampling

Individuals had grown in clearings and formed a secondary forest where they are the

dominant species together with *C. sciadophylla*. A total of 70 trees were selected in

September and December 2014 respectively for Counami and Sparouine sites: 32 in Counami

and 38 in Sparouine. Soil samples were taken at the same time for pedological analysis. On

the Counami site, where individuals are widely spaced, a soil sample was taken at the basis of

each individual tree. On the Sparouine site, where individuals where clustered, 9 soil samples

were taken, as each soil sample was representative of 4-6 individuals located no further than

30m from the soil sample spot.

As *C. obtusa* is dioecious, only pistillate (i.e. female) trees were felled to avoid

potential sex-related variability in the measured functional traits. Trees were not felled

according to the same scheme in the two sites. Trees were preselected to have as close as

possible comparable diameters at breast height (DBH), and age was estimated with binoculars

according to the method described by (Zalamea *et al.* 2012). By counting the number of

internodes we were able to estimate the age of trees as each internode is produced in 10 days

(Heuret *et al.* 2002; Zalamea *et al.* 2012). In Sparouine, all individuals correspond to a single

colonisation pulse on both soil types: all individuals have similar age (7-10 years), with DBH

of 11.94 to 25.70 cm, and heights of 13.85 to 23.20 m (Fig. S3). Both soil types were

represented by 19 individuals and all individuals were felled and measured between the 14[th]

and the 19[th] of September 2015. Thus, season-, size-, and age-related effects on functional

traits are controlled for soil and individual comparisons.

The experimental design at Counami was different. The forestry road was opened

gradually, and therefore the age of the trees differed according to the road section (Zalamea *et*

al. 2012). All individuals assigned to WS at Counami were selected at a single small WS

patch located 6 km after the entrance of the road. WS trees represented a single colonisation

pulse and were of similar age (14-16 years, except one significantly older with 22.8 years

old), with DBH from 6.21 to 15.18 cm, and heights from 10.27 to 16.18 m, (Fig. S3). It was

not possible to choose trees on FS on a single restricted area because of the perturbation of

soil structure by the logging machines and because we excluded trees on down slopes.

Consequently, FS trees were sampled between km 6 to 11 of the forestry road and included

different cohorts with different ages (7-23 years), DBH of 9.55 to 22.44 cm, and heights of

12.16 to 22.63 m (Fig. S3). Thirteen and nineteen individuals were sampled on FS and WS

respectively. Counami trees were felled at different dates, from September 2014 to April

2016. The contrasted protocol was chosen to study seasonal and ontogenetic effect on leaf

traits, but the results of such analysis will not be addressed here. No seasonal effects on leaf

traits were detected, and ontogenetic effects on functional trait were standardised, as

presented in the *Statistical analyses* part.

Soil properties

Pedological analyses included granulometry, moisture content, pH, organic matter content,

and contents of exchangeable cations (Appendix S1, with detailed abbreviations). The

complete sampling procedure is described in the Appendix S1. Exchangeable cations were

analysed divided by cation-exchange capacity (CEC) to avoid correlations between the former

and the latter. We also calculated a soil index of fertility as: $Soil_{index} = \frac{K+Ca+Mg+Na}{CEC}$.

The *a priori* classification of soil types (FS-*versus*-WS) was confirmed by pedological

analyses of the soil properties within each site. The described pattern of soil properties is

congruent with that reported in the literature (Adeney *et al.* 2016; Fine and Baraloto 2016a).

WS consist of a large proportion of coarse sand with high Ca:CEC (calcium on CEC) and C:N

(carbon on nitrogen) ratios. FS consist of a large proportion of clay and silt with high

moisture, N, C, MO P_{tot} (total potassium) contents and a high Al:CEC (aluminium on CEC)

ratio. Based on water availability, N content, and $soil_{index}$, the site fertility can be ordered as

COU-FS > SPA-FS > COU-WS = SPA-WS. Sparouine WS are characterised by higher

H:CEC and Fe:CEC ratio than Counami WS. The related results are presented in Appendix

S1. Within sites, the difference between soil types is more contrasted in Counami than in

Sparouine.

Architectural and functional traits

For all individuals, we measured a suite of architectural traits at phytomer and whole-tree

levels to characterise growth, branching and flowering dynamics, and the resulting tree

architecture. Retrospective analysis of development allows us to consider tree developmental

trajectories as growth performance traits (i.e. the height-age relationship). Such approach

considers the development of the trunk only (i.e. it does not include the complexity of
branching events) described as a sequence of phytomers. Three variables were measured for
each phytomer: (1) internode length (2) vegetative bud state coded as: 0 for not developed or
aborted; 1 for developed, present or pruned, (3) inflorescence bud state coded as: 0 for no
inflorescence; 1 for pruned or present inflorescences. Features for bud states are treated as
binary values: presence or absence. As suggested by (Davis 1970), (Heuret *et al.* 2002), and
(Zalamea *et al.* 2008), we analysed periodical fluctuations in internode length, which are
driven by seasonal variations of rainfall, (Zalamea et al., 2013), as well as the rhythmic
disposition of inflorescences and branches to infer the past development of the tree, and
model its growth dynamic (section statistical analysis and Appendix S2).

As a first step, the fluctuation of internode length allowed us to estimate (i) the
growth representing a single year as the shortest internodes are associated with the peak of the
dry season, (ii) the age in days after germination of any internode along the trunk, and (iii) the
yearly average time taken by the tree to produce an internode (i.e. the phyllochron).

As a second step, to understand how the trees are modifying their growth strategy in
the two types of soils, we looked at (i) variations of phyllochron, internode length, and annual
shoot length over time, and (ii) contribution of the number of internode vs internode length in
the annual shoot length variation (See Appendix S2 for the followed methodology).

As a third step, we looked at how these different potential growth strategies (i.e.
number vs. length of internodes) drive the cumulative tree height over time, namely the
growth performance. Finally, to study space-foraging performance and reproductive
performance we analysed the cumulative branching and flowering over time. The measured
and estimated traits presented as longitudinal sequences, are shown in Table 1.

Whole-tree-level traits was also measured (Table 1). Functional traits were measured at the
leaf level (Table 2) as proxies of leaf resource capture, while trunk wood specific gravity was
measured as indicator of stem transport and storage capacity (Baraloto *et al.* 2010). We
measured leaf-level traits for only one leaf per individual: either the third or the fourth leaf
positioned under the apex of the A1 axis. In this way, potential effects of plant spatial
structure and senescence on variation of leaf-level traits are controlled. Leaf lifespan along the
A1 axis was estimated for each tree by counting the number of leaves on a given axis and
multiplying it by the known mean phyllochron (10 days, Heuret et al. 2002). The complete
sampling procedure for functional traits is described in Appendix S3.

Statistical analyses

Topology of trees and the different pedological, whole-tree-level, and functional features

associated with each repetitive unit are coded in sequences in Multi-scale Tree Graph format

(MTG; Godin & Caraglio, 1998; Godin, Costes, & Caraglio, 1997).

Statistical analyses relative to developmental trajectories were conducted with

AMAPmod (*op. cit*), now integrated in the OpenAlea platform, re-engineered and named

'VPlants' (Pradal *et al.* 2013), and the R programming language (R Core Team 2018). We relied

on autocorrelation coefficients on internodes (length, branch presence, inflorescence

presence) to confirm an annual periodicity on the stand level (i.e. soil x site) for growth,

branching, and flowering processes. Methods and results regarding the analysis of

autocorrelation coefficients are presented in Appendix S4. To analyse fluctuations of

internode length, we used a method of time series analysis relying on a decomposition

principle of signals, described as follows. The different sources of variation, such as long-

term changes at low-level frequency vs short-term changes at high-level frequency

fluctuations, are identified and filtered (Guédon *et al.* 2007). Firstly, we calculated a moving

average to extract the trend of internode length sequences in a similar way as (Zalamea *et al.*

2008). Having extracted the trend, we looked at local fluctuations by examining the residuals.

Residuals were generated by dividing for each internode, its length by its moving average

(Appendix S2 for details The analysis of residuals allowed the identification of the limits of

the long dry season in September/October for successive years, since shorter internodes are

elongated during this period as shown for *C. obtusifolia* Bertol. (Davis, 1970), *C. peltata* L.,

and *C. sciadophylla* Mart (Zalamea *et al.* 2013). Delimitation of annual growth for each

individual allowed the estimation of a mean phyllochron for each year according to the node

rank (Appendix S2). Knowing the phyllochron allowed the conversion of the rank node to a

temporal scale, namely the age. Finally, by considering the length or the number of nodes

elongated between two successive dry seasons, we estimated the annual shoot length (Table

1). Growth strategies are studied as (i) variations of phyllochron, internode length, and annual

shoot length over time, and (ii) contribution of the number of internode vs internode length in

the annual shoot length variation. Significant differences in architectural traits (i.e. internode

length, phyllochron, AS length, number of internodes per annual shoot) between FS and WS

were identified based on a confidence interval at 95% around the mean trajectory of the

considered architectural trait. A mean trajectory was calculated and plotted for each soil type

within each site.

To test the effect of soil type on the variability of growth trajectories –which are

longitudinal data by nature-, we tested the correspondence of distribution of (i) soil types,

with that of (ii) clusters defined by statistical signatures of growth trajectories. The clusters

were characterised with a clustering method on the generated longitudinal data (Table 1), with

the *kml* R package (Genolini and Falissard 2009). It is a classification method based on an

implementation of "k-means", itself based on a minimization function of distances among

trajectories. For each trait, 100 simulations were used, and decisions are based on the

Calinski-Harabasz criterion. The optimal number of clusters corresponds to a maximisation of

the Calinski-Harabasz criterion. The dependency of defined clusters on soil types is evaluated

with a Pearson's chi-squared test. Analyses relative to soil, whole-tree-level, and functional

trait data are realised in R language. Potential effects of seasonality and ontogeny on leaf and

whole-tree-level trait variation for Counami trees were tested with a multiple regression

analysis. No season-related effects were detected based on a total of 70 leaves. The effects of

ontogeny were filtered for each functional trait only when a significant signal was detected

based on a linear regression analysis. Principal Component Analysis (PCA) on soil properties

and functional traits were conducted with the *ade4* (Chessel *et al.* 2004) and *Factoextra*

(Kassambara and Mundt 2016) R packages. The effect of soil on functional and whole-tree-

level traits was tested with linear mixed-effect models (LMER), with the soil gradient

modelled by tree coordinates along the first axis (45,4%) of the soil PCA (Appendix S1). Soil

and tree age were set as fixed effects, and site as a random effect. A comparison of factorial

coordinates of individuals was conducted for each axis based on a nested-ANOVA and a *post-*

hoc Tukey's HSD test.

RESULTS

Developmental approach: architecture and growth trajectory

Significant differences in trajectories of architectural traits between FS and WS were

visualised through plotted confidence intervals around the mean trajectory (Fig. 1). Internode

length was significantly shorter for WS in comparison to FS in Counami (Fig. 1c) for the first

5 years only. These first 5 years corresponded to the ontogenetic stage with the longest

internodes. No difference in internode length was found in Sparouine between FS and WS

(Fig. 1e). Clusters of internode length trajectories significantly matched soil type distributions

in Counami ($P < 0.01$), but not in Sparouine ($P > 0.05$; Fig. 1d,f). Phyllochron –and the

dependant variable, the number of nodes per annual shoot-, was not significantly different

between FS and WS for either site (Fig. 1a; Fig. S4a,c). Clusters of the trajectories of the

number of internodes per annual shoot trajectories significantly matched soil type

distributions in Sparouine ($P < 0.01$), but not in Counami ($P > 0.05$; Fig. S4b,d). Annual shoot

length was significantly shorter for WS in comparison to FS in Counami (Fig. S4e) for the

first 5 years only. No difference in annual shoot length was found in Sparouine between FS

and WS (Fig. S4g). Clusters of annual shoot length trajectories significantly matched soil type

distributions in Counami (P < 0.01), but not in Sparouine (P > 0.05; Fig. S4f,h).

For both sites, there was a pattern for FS trees to be higher than WS trees for a given

age (Fig. 1g,i). Within the Counami site, FS trees covered the largest variation of growth

trajectories and reached the highest height (Fig. 1g). Within the Sparouine site, WS trees

covered a larger variation of trajectories and the WS highest trees were as tall as the highest

FS ones (Fig. 1i). For both sites, it was possible to identify two main growing phases. The

phases were differentiated by variations in growth rates over the tree's lifespan. The first

phase covered the first 5-7 years, except for FS Counami trees where it was the first 9-10

years. The second growing phase was defined by a slower growth rate, which remained

constant for all individuals. For both sites, cluster of tree height trajectories significantly

matched soil type distribution (P < 0.05; Fig. 1h,j).

The analysis of the cumulated number of pairs of inflorescences on the trunk indicated

that there was no significant difference between FS and WS for both sites based on

confidence intervals (Fig. 2a, b). In Counami trees, there was a significant difference in the

cumulated number of branches of the trunk between FS and WS after 5-6 years old (Fig. 2c).

In Sparouine trees there was no significant difference in the cumulated number of branches on

the trunk between FS and WS (Fig. 2d).

A significant effect of soil was identified for tree height, DBH, the branching order,

and the height of the first flowering and first branching (P < 0.05; Table 3;LMER), with all

whole-tree-level traits increasing in FS.

Characterisation of functional traits

The first and second axes of the PCA for functional traits explained 47.0 % of the inertia (Fig.

3a). The first axis (28.5 %) is driven by $C:N_{leaf}$, L_{pet}, A_{pet}, A_{leaf} and N_{leaf}. The second axis

(18.5 %) is driven by $H_2O_{res,leaf}$, and K_{leaf}. Conditions (i.e. soil types x sites) were

differentiated along the first axis (Fig. 3b; P < 0.05; ANOVA) with significant differences (i)

between FS and WS within Counami and (ii) between Counami and Sparouine when only

considering FS. Conditions were more strongly differentiated along the first axis (P < 0.001;

ANOVA) with Counami trees in WS differing from Sparouine trees in both FS and WS.

Significant effect of soil was detected for leaf residual water content and leaf K content (P <

0.05; Table 4), with lower residual water content but higher K content for FS trees.

DISCUSSION

To our knowledge, our study is the first incorporating both, tree architectural development and functional traits, in relation with the environment. It is also the first to quantify functional traits for an Amazonian generalist species regarding soil types: FS and WS. Our pedological analysis confirmed strong contrasts in soil characteristics between FS and WS, opening the possibility of soil-related phenotypic response. The soil-response of functional traits was rather weak, whereas the soil-response of architectural development was rather strong and dominated the phenotypic response of *C. obtusa* to the WS-FS contrast, leading to soil-response strategy modulating resource use (mediated by growth) rather than resource acquisition (mediated by functional traits).

Soil-response is mediated by architectural development, not functional traits

Our results suggest that phenotypic response to soil change is mediated by the architectural development rather than functional traits. A striking result is that the soil was not a driver of the variation of measured leaf and wood traits. Only two leaf traits were responsive to soil types: leaf residual water content and leaf K content. The residual water content, which is not a commonly used functional trait, is indicative of the capacity of leaf tissues to retain water through osmotic adjustments (Bartlett *et al.* 2012). The residual moisture content was positively correlated to K content (*results not shown*; P < 0.001; R² = 0.210), which plays a central role in the maintenance of osmotic integrity of cells and tissues (Marschner 1995). Such correlation between residual water content and soil type suggests that edaphic water stress is one of the primary factors underlying the FS-WS gradient, further shaping the phenotypic response, especially for functional traits related to hydraulics and drought tolerance. This is coherent with our pedological analysis that indicates water availability as highly determinant in the first axis of the soil PCA, underlying the FS-WS gradient (Appendix S1).

Such weak functional trait response was unexpected. Two non-mutually exclusive reasons can be explored to explain why only two functional traits responded to changes in soil type.

(1) There are relevant functional traits we did not consider in our study. It has been shown that water availability is the leading climate driver of Amazonian rainforest tree growth (Wagner *et al.* 2012). Water relation and drought-resistance traits, such as drought-induced vulnerability to embolism and stomatal sensitivity, leaf turgor

loss point, root depth, crown area to sapwood area ratio, may have played a central role in ensuring growth and survival on the different soil types (Urli *et al.* 2013; Anderegg *et al.* 2016; O'Brien *et al.* 2017; Adams *et al.* 2017; Eller *et al.* 2018). Differences in Amazonian soil characteristics can also impact the root system properties (Freschet *et al.* 2017), including mycorrhizal fungi associations. For instance, it has been shown that ectomycorrhizal mutualisms are much more common on WS (Roy *et al.* 2016), and several studies suggest that ectomycorrhizal species may be better able to acquire nutrients (Reich 2014).

(2) The different plant strategies, or life-histories, can be defined along two important strategic axes of plant functioning: the resource acquisition (e.g. photosynthesis, soil nutrients absorption) axis and the resource use (e.g. growth, defence and secondary metabolites) axis (Reich 2014). The functional traits (i.e. leaf and wood traits) measured here are related to resource acquisition, and poorly captured how resources are used. Trees may not necessarily respond to WS resource scarcity by modifying functional traits related to the acquisition axis. But instead, the reduced resources assimilated in a given time may be translated into reduced resource use possibilities. Since plants are organisms with undetermined development, growth remains one of the largest carbon and nutrients sink across lifespan. Thus, growth may be a component of an adaptive response to resource scarcity. Deciphering growth processes and strategies, and quantifying their variations, could represent an opportunity for studying changes along the resource use axis, in relation to the environment.

An analysis of growth trajectory based on architectural development analysis is a useful tool for the quantification of the resource use strategy. The autocorrelation function at the stand level confirmed a high degree of periodicity across all individuals for growth, flowering, and branching processes (Appendix S4). With the analysis of internode fluctuations, this periodicity has been shown to be annual, and further allowed to shift on a temporal scale and to conduct our retrospective analysis of architecture. We clearly showed that soil types impacted the overall growth trajectory (i.e. cumulated tree height according to age) for both sites, with WS trees having the lowest trajectories. For any given age, WS trees were always smaller, due to resource scarcity. However, such pattern is less noticeable on Sparouine trees. The site difference could be explained by (i) the less pronounced contrasts between FS and WS in Sparouine as shown by our pedological analysis (Appendix S1), and (ii) the rainier dry season in Sparouine (Fig. S2). Under the assumption that the interaction

between WS and water scarcity during the dry season is deleterious for tree growth, this may also explain the generally strongest growth trajectories in Sparouine in comparison to Counami.

Regarding the growth strategy, soil type showed a significant effect on both internode length and annual shoot length in Counami, but not in Sparouine. When the soil effect was strong enough, the differences in annual shoot length between soil types corresponded mainly to variations in internode length rather than variations in number of nodes per annual shoot. Reducing the number of nodes per annual shoot would imply the increase of the phyllochron, thus reducing the number of leaves produced per year. Such mechanism would critically affect tree carbon balance, as the contribution of a given leaf to the carbon balance is disproportionate in comparison to most of species: *C. obtusa*'s leaves are large (1,000-5,000 mm², Levionnois *et al.*, *data not published*) but few (100-600 leaves, Table 3). Similarly, (Zalamea *et al.* 2013) found no difference in phyllochron between *C. sciadophylla* from two distanced locations with contrasting rainfall. The architectural analysis also shows that WS trees in Counami had significantly fewer cumulated branches, and lower branching order, than those in FS.WS trees in Counami have, therefore, reduced space and light foraging capacities, decreasing their competitiveness. Because flowering is synchronous on all crown axes (Heuret et al., 2003), the energetic production cost of inflorescences and seeds is exponentially related to the number of main branches. Therefore, WS trees in Counami must also have comparatively reduced reproductive and dispersive abilities, leading to a reduced overall fitness compared to their FS conspecifics, under the assumption that FS and WS trees form a unique population.

Our results were not in agreement with Borges *et al.* (2019), who applied a similar approach by comparing functional traits for an Asteraceae generalist tree species growing in savanna and cloud forests in a single site in south-eastern Brazil (the study was conducted on a same site, with no distance or climatic effects on functional traits). They found contrasting functional trait responses between the two habitats for a set of functional traits related to resource acquisition and storage (i.e. leaf area and thickness, LMA, wood density), such that savanna individuals were more resource conservative (i.e. high wood density and LMA, thick and small leaves) than those from cloud forest. The discrepancy between the two studies indicates that the type of phenotypic response (i.e. resource acquisition vs resource use strategies) for generalist species is not uniform across species, and may vary depending on its functional type (e.g. evergreen vs deciduous, pioneer vs late-successional, light-demanding vs shade-tolerant), the nature of the resource heterogeneity between habitats (e.g. light, water,

soil nutrients), or the degree of habitat divergence.Such non-agreement between our results

and those from Borges *et al.* (2019), stresses the importance of considering biological

diversity and its interaction with environmental heterogeneity. One environmental constraint

may strongly regulate a particular strategic dimension of plant functioning but not the other

(e.g. resource acquisition vs. use axes), and some functional traits (e.g. leaf traits) may be

particularly sensitive to certain environmental constraints, as revealed by Paine *et al.* (2011).

Our study exemplifies the complexity of incorporating ITV in studying ecological processes,

and how ITV of different functional traits are not evenly responsive to abiotic factors.

However, we demonstrated the potential gains of incorporating architectural analysis in plant

community ecology, particularly at the intraspecific level.

Why Cecropia *functional response is not homologous to the environmental filtering at the*

community level?

At the community level in the Amazon, edaphic contrasts lead to strong environmental

filtering mediated by functional traits. WS flora is characterised by a convergence in

functional traits, particularly towards a conservative strategy (Grubb and Coomes 1997;

Patiño *et al.* 2009; Fyllas *et al.* 2009; Fine *et al.* 2010; Fortunel *et al.* 2012; Fortunel, Paine, *et*

al. 2014; Fortunel, Ruelle, *et al.* 2014; Fine and Baraloto 2016). It implies high LMA, high

WSG, and low leaf nutrient contents associated with high water use efficiency (i.e. high

photosynthetic assimilation to stomatal conductance ratio) for WS tree species (Fine and

Baraloto 2016), instead of FS tree species. Fine & Baraloto (2016) highlighted how WS are

limiting for plant development, due to resource scarcity, and how resource scarcity in WS is

an abiotic factor selecting for functional trait optima diverging from functional trait optima

found on FS. However, the intraspecific *Cecropia* functional trait response to environmental

heterogeneity did not parallel the environmental filtering operating on functional traits at the

community level (i.e. high LMA and WSG on WS, low LMA and WSG on FS…). This

suggests potential contrasting effects of ecological processes, such as environmental filtering

and biotic interactions, operating at the interspecific level and the *C. obtusa* intraspecific

level.

Regarding environmental filtering, as discussed above, other sets of functional traits

(e.g. hydraulic and drought-resistance traits) rather than those we measured could be key

determinants for the establishment on WS. Moreover, as discussed above, variation in growth

strategy could be the leading parameter bypassing constraints related to WS. Regarding biotic

interactions, the studies of (Fine *et al.* 2004, 2006) suggest that herbivory pressure could be a

primary driver of ecological speciation and diversification within genus on WS, because of

higher costs of tissue lost associated with resource-poor habitats. The resource scarcity selects

for structures with long lifespan, and resistant to herbivory pressure. Conversely, *Cecropia*

trees are characterised by short lifespans and high growth rates, in relation to their pioneering

and competitive strategy, which is in contradiction with a conservative strategy privileging

long lifespan. Under such hypothesis, selection for light competitiveness would be prevalent

on selection for a conservative strategy. This would explain why functional traits such as

LMA and WSG are not impacted by soil types as demonstrated by our study. In order to

achieve herbivory resistance, three types of defence can be produced: structural, chemical,

and mutualistic. Here again, the non-dependence of functional traits such as LMA and WSG

on soil type suggest that structural defences are not required to respond and to allow *Cecropia*

trees to grow on WS. Chemical traits related to herbivory resistance have been shown to vary

between FS and WS for the generalist tree species *Protium subserratum* (Burseraceae; (Fine

et al. 2013)). Chemical traits related to herbivory pressure, and the herbivory pressure in

itself, are unknown for *Cecropia* trees; but functional traits related to herbivory avoidance

could play an important role in the strategy required to allow *C. obtusa*'s generalist behaviour,

and need further investigations. Finally, *Cecropia* species are also famous for their mutualism

with the *Azteca* ant species, where ants offer a protection against visitors by biting (Schupp

1986; Dejean *et al.* 2009). During field work, we observed ant occupancy on both sites and on

both soil types, suggesting an undisturbed interaction between ant and host plants.

CONCLUSION

Our study demonstrated that commonly measured traits, related to resource acquisition

strategies, are not systematically responsive to contrasting habitats. Other aspects of plant

functioning such as resource use strategies (through architectural development) can rather

mediate such responses. Our study raises concerns about negative results when investigating

environmental filtering based on commonly measured functional traits like LMA, leaf

thickness, or WSG. Environmental filtering can occur on other dimensions of plant

functioning. As architectural analysis has brought insights on environmental filtering at the

intraspecific level, such approach could also be applied to the process of niche differentiation,

especially regarding intra- and interspecific competition.

ACKNOWLEDGEMENTS

We especially thank Jean-Yves Goret, Saint-Omer Cazal and Audin Patient for their major

contributions in field work and collecting data. We thank Julie Bossu, Coffi Belmys Cakpo,

Jocelyn Cazal, Aurélie Cuvelier, Bruno Clair, Aurélie Dourdain, Alexandre Haslé de Barral,

Solène Happert, Marie Hartwig, Clément Jouaux, Yohann Legraverant, Anabelle Mercrette,

Ariane Mirabel, Pascal Petronelli, Laurent Risser, Dylan Taxile, Camille Ziegler and Lore

Verryckt for their assistance with field work and measurement of leaf traits. We gratefully

thank Isabelle Maréchaux for critical and valuable comments on the manuscript. We thank

Anna Deasey for English proof reading. We thank the ONF for access to forestry roads and

the sampling. We also thank the USRAVE (INRA-COFRAC) of Bordeaux and the EEF lab

(INRA) of Nancy for measurements of leaf nutrients. S.L. was supported by a doctoral

fellowship from CEBA. This study benefited from an *Investissement d'Avenir* grant managed

by the *Agence Nationale de la Recherche* (CEBA, ref. ANR-10-LABX-0025).

LITERATURE CITED

Adams HD, Zeppel MJB, Anderegg WRL, *et al.* 2017. A multi-species synthesis of
physiological mechanisms in drought-induced tree mortality. *Nature Ecology and Evolution*.

Adeney JM, Christensen NL, Vicentini A, Cohn-Haft M. 2016. White-sand Ecosystems in
Amazonia. *Biotropica* 48: 7–23.

Allié E, Pélissier R, Engel J, *et al.* 2015. Pervasive Local-Scale Tree-Soil Habitat Association
in a Tropical Forest Community. *PLOS ONE* 10: e0141488.

Anderegg WRL, Klein T, Bartlett M, *et al.* 2016. Meta-analysis reveals that hydraulic traits
explain cross-species patterns of drought-induced tree mortality across the globe. *Proceedings*
of the National Academy of Sciences of the United States of America 113: 5024–5029.

Baraloto C, Timothy Paine CE, Poorter L, *et al.* 2010. Decoupled leaf and stem economics in
rain forest trees. *Ecology Letters* 13: 1338–1347.

Bartlett MK, Scoffoni C, Sack L. 2012. The determinants of leaf turgor loss point and
prediction of drought tolerance of species and biomes: a global meta□analysis. *Ecology*
Letters 15: 393–405.

Borges ER, Prado-Junior J, Santana LD, *et al.* 2019. Trait variation of a generalist tree species
(Eremanthus erythropappus, Asteraceae) in two adjacent mountain habitats: savanna and
cloud forest. *Australian Journal of Botany* 66: 640–646.

Borregaard MK, Rahbek C. 2010. Causality of the relationship between geographic
distribution and species abundance. *The Quarterly Review of Biology* 85: 3–25.

Boulangeat I, Lavergne S, Es JV, Garraud L, Thuiller W. 2012. Niche breadth, rarity and
ecological characteristics within a regional flora spanning large environmental gradients.
Journal of Biogeography 39: 204–214.

Charles-Dominique T, Edelin C, Bouchard A. 2009. Architectural strategies of Cornus sericea, a native but invasive shrub of Southern Quebec, Canada, under an open or a closed canopy. *Annals of Botany*: mcp273.

Charles-Dominique T, Edelin C, Brisson J, Bouchard A. 2012. Architectural strategies of Rhamnus cathartica (Rhamnaceae) in relation to canopy openness. *Botany* 90: 976–989.

Chessel D, Dufour A-B, Thioulouse J. 2004. The ade4 package - I□: One-table methods. *R News* 4: 5–10.

Clark DB, Palmer MW, Clark DA. 1999. Edaphic Factors and the Landscape-Scale Distributions of Tropical Rain Forest Trees. *Ecology* 80: 2662–2675.

Daly DC, Silveira M, Medeiros H, Castro W, Obermüller FA. 2016. The White-sand Vegetation of Acre, Brazil. *Biotropica* 48: 81–89.

Davis RB. 1970. Seasonal differences in intermodal lengths in Cecropia trees; a suggested method for measurement of past growth in height. *Turrialba*.

Dejean A, Grangier J, Leroy C, Orivel J. 2009. Predation and aggressiveness in host plant protection: a generalization using ants from the genus Azteca. *Naturwissenschaften* 96: 57–63.

Eller C, de V. Barros F, R.L. Bittencourt P, Rowland L, Mencuccini M, S. Oliveira R. 2018. Xylem hydraulic safety and construction costs determine tropical tree growth. *Plant, Cell & Environment*: n/a-n/a.

Fine PVA, Baraloto C. 2016a. Habitat Endemism in White-sand Forests: Insights into the Mechanisms of Lineage Diversification and Community Assembly of the Neotropical Flora. *Biotropica* 48: 24–33.

Fine PVA, García-Villacorta R, Pitman NCA, Mesones I, Kembel SW. 2010. A floristic study of the white-sand forests of Peru. *Annals of the Missouri Botanical Garden* 97: 283–305.

Fine PVA, Mesones I, Coley PD. 2004. Herbivores Promote Habitat Specialization by Trees in Amazonian Forests. *Science* 305: 663–665.

Fine PVA, Metz MR, Lokvam J, *et al.* 2013. Insect herbivores, chemical innovation, and the evolution of habitat specialization in Amazonian trees. *Ecology* 94: 1764–1775.

Fine PVA, Miller ZJ, Mesones I, *et al.* 2006. The growth–defense trade-off and habitat specialization by plants in amazonian forests. *Ecology* 87: S150–S162.

Fortunel C, Fine PVA, Baraloto C. 2012. Leaf, stem and root tissue strategies across 758 Neotropical tree species. *Functional Ecology* 26: 1153–1161.

Fortunel C, Paine CET, Fine PVA, Kraft NJB, Baraloto C. 2014. Environmental factors predict community functional composition in Amazonian forests. *Journal of Ecology* 102: 145–155.

Fortunel C, Ruelle J, Beauchêne J, Fine PVA, Baraloto C. 2014. Wood specific gravity and anatomy of branches and roots in 113 Amazonian rainforest tree species across environmental gradients. *New Phytologist* 202: 79–94.

Freschet GT, Valverde-Barrantes OJ, Tucker CM, *et al.* 2017. Climate, soil and plant functional types as drivers of global fine-root trait variation. *Journal of Ecology* 105: 1182–1196.

Fridley JD, Grime JP. 2010. Community and ecosystem effects of intraspecific genetic diversity in grassland microcosms of varying species diversity. *Ecology* 91: 2272–2283.

Fyllas NM, Patiño S, Baker TR, *et al.* 2009. Basin-wide variations in foliar properties of Amazonian forest: phylogeny, soils and climate. *Biogeosciences* 6: 2677–2708.

Genolini C, Falissard B. 2009. KmL: k-means for longitudinal data. *Computational Statistics* 25: 317–328.

Godin C, Caraglio Y. 1998. A Multiscale Model of Plant Topological Structures. *Journal of Theoretical Biology* 191: 1–46.

Godin C, Costes E, Caraglio Y. 1997. Exploring plant topological structure with the AMAPmod software: an outline.

Gourlet-Fleury S, Guehl JM, Laroussine O. 2004. *Ecology and management of a neotropical rainforest□: lessons drawn from Paracou, a long-term experimental research site in French Guiana*. Paris: Elsevier.

Grubb PJ, Coomes DA. 1997. Seed mass and nutrient content in nutrient-starved tropical rainforest in Venezuela. *Seed Science Research* 7: 269–280.

Guédon Y, Caraglio Y, Heuret P, Lebarbier E, Meredieu C. 2007. Analyzing growth components in trees. *Journal of Theoretical Biology* 248: 418–447.

Heuret P, Barthélémy D, Guédon Y, Coulmier X, Tancre J. 2002. Synchronization of growth, branching and flowering processes in the South American tropical tree Cecropia obtusa (Cecropiaceae). *American Journal of Botany* 89: 1180–1187.

Heuret P, Meredieu C, Coudurier T, Courdier F, Barthélémy D. 2006. Ontogenetic trends in the morphological features of main stem annual shoots of Pinus pinaster (Pinaceae). *American Journal of Botany* 93: 1577–1587.

HilleRisLambers J, Adler PB, Harpole WS, Levine JM, Mayfield MM. 2012. Rethinking Community Assembly through the Lens of Coexistence Theory. *Annual Review of Ecology, Evolution, and Systematics* 43: 227–248.

Holt AR, Gaston KJ, He F. 2002. Occupancy-abundance relationships and spatial distribution: A review. *Basic and Applied Ecology* 3: 1–13.

Jung V, Albert CH, Violle C, Kunstler G, Loucougaray G, Spiegelberger T. 2014. Intraspecific trait variability mediates the response of subalpine grassland communities to extreme drought events. *Journal of Ecology* 102: 45–53.

Jung V, Violle C, Mondy C, Hoffmann L, Muller S. 2010. Intraspecific variability and trait-
based community assembly. *Journal of Ecology* 98: 1134–1140.

Kassambara A, Mundt F. 2016. Factoextra□: Extract and Visualize the Results of
Multivariate Data Analyses.

Kraft NJB, Adler PB, Godoy O, James EC, Fuller S, Levine JM. 2015. Community assembly,
coexistence and the environmental filtering metaphor. *Functional Ecology* 29: 592–599.

Kraft NJB, Crutsinger GM, Forrestel EJ, Emery NC. 2014. Functional trait differences and
the outcome of community assembly: an experimental test with vernal pool annual plants.
Oikos 123: 1391–1399.

Kraft NJB, Valencia R, Ackerly DD. 2008. Functional Traits and Niche-Based Tree
Community Assembly in an Amazonian Forest. *Science* 322: 580–582.

Lepš J, Bello F de, Šmilauer P, Doležal J. 2011. Community trait response to environment:
disentangling species turnover vs intraspecific trait variability effects. *Ecography* 34: 856–
863.

Letort V, Heuret P, Zalamea P-C, Reffye PD, Nicolini E. 2012. Analysing the effects of local
environment on the source-sink balance of Cecropia sciadophylla: a methodological approach
based on model inversion. *Annals of Forest Science* 69: 167–180.

Marschner H. 1995. 8 - Functions of Mineral Nutrients: Macronutrients In: *Mineral Nutrition*
of Higher Plants (Second Edition). London: Academic Press, 229–312.

Mathieu A, Letort V, Cournède P h., Zhang B g., Heuret P, de Reffye P. 2012. Oscillations in
Functional Structural Plant Growth Models. *Mathematical Modelling of Natural Phenomena*
7: 47–66.

McGill BJ, Enquist BJ, Weiher E, Westoby M. 2006. Rebuilding community ecology from
functional traits. *Trends in Ecology & Evolution* 21: 178–185.

O'Brien MJ, Engelbrecht BMJ, Joswig J, *et al.* 2017. A synthesis of tree functional traits
related to drought□induced mortality in forests across climatic zones. *Journal of Applied*
Ecology.

Paine CET, Baraloto C, Chave J, Hérault B. 2011. Functional traits of individual trees reveal
ecological constraints on community assembly in tropical rain forests. *Oikos* 120: 720–727.

Patiño S, Lloyd J, Paiva R, *et al.* 2009. Branch xylem density variations across the Amazon
Basin. *Biogeosciences* 6: 545–568.

Pradal C, Coste J, Boudon F, Fournier C, Godin C. 2013. *OpenAlea 2.0: Architecture of an*
integrated modeling environment on the web. Finnish Society of Forest Science.

R Core Team. 2018. *R: A language and environment for statistical   computing*. Vienna,
Austria: R Foundation for Statistical Computing.

Reich PB. 2014. The world-wide 'fast–slow' plant economics spectrum: a traits manifesto.
Journal of Ecology 102: 275–301.

Roy M, Schimann H, Braga-Neto R, *et al.* 2016. Diversity and Distribution of Ectomycorrhizal Fungi from Amazonian Lowland White-sand Forests in Brazil and French Guiana. *Biotropica* 48: 90–100.

Sabatier D, Grimaldi M, Prévost M-F, *et al.* 1997. The influence of soil cover organization on the floristic and structural heterogeneity of a Guianan rain forest. *Plant Ecology* 131: 81–108.

Schamp BS, Chau J, Aarssen LW. 2008. Dispersion of traits related to competitive ability in an old-field plant community. *Journal of Ecology* 96: 204–212.

Schupp EW. 1986. Azteca protection of Cecropia: ant occupation benefits juvenile trees. *Oecologia* 70: 379–385.

Shipley B, Bello FD, Cornelissen JHC, Laliberté E, Laughlin DC, Reich PB. 2016. Reinforcing loose foundation stones in trait-based plant ecology. *Oecologia* 180: 923–931.

Sides CB, Enquist BJ, Ebersole JJ, Smith MN, Henderson AN, Sloat LL. 2014. Revisiting Darwin's hypothesis: Does greater intraspecific variability increase species' ecological breadth? *American Journal of Botany* 101: 56–62.

Stecconi M, Puntieri JG, Barthélémy D. 2010. An architectural approach to the growth forms of Nothofagus pumilio (Nothofagaceae) along an altitudinal gradient. *Botany* 88: 699–709.

ter Steege H, Pitman NCA, Sabatier D, *et al.* 2013. Hyperdominance in the Amazonian Tree Flora. *Science* 342: 1243092.

Stropp J, Sleen PV der, Assunção PA, Silva AL da, Steege HT. 2011. Tree communities of white-sand and terra-firme forests of the upper Rio Negro. *Acta Amazonica* 41: 521–544.

Swenson NG, Enquist BJ. 2009. Opposing assembly mechanisms in a Neotropical dry forest: implications for phylogenetic and functional community ecology. *Ecology* 90: 2161–2170.

Taugourdeau O, Dauzat J, Griffon S, Sabatier S, Caraglio Y, Barthélémy D. 2012. Retrospective analysis of tree architecture in silver fir (Abies alba Mill.): ontogenetic trends and responses to environmental variability. *Annals of Forest Science* 69: 713–721.

Uriarte M, Condit R, Canham CD, Hubbell SP. 2004. A spatially explicit model of sapling growth in a tropical forest: does the identity of neighbours matter? *Journal of Ecology* 92: 348–360.

Urli M, Porté AJ, Cochard H, Guengant Y, Burlett R, Delzon S. 2013. Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree Physiology* 33: 672–683.

Violle C, Enquist BJ, McGill BJ, *et al.* 2012. The return of the variance: intraspecific variability in community ecology. *Trends in Ecology & Evolution* 27: 244–252.

Violle C, Navas M-L, Vile D, *et al.* 2007. Let the concept of trait be functional! *Oikos* 116: 882–892.

Wagner F, Rossi V, Stahl C, Bonal D, Hérault B. 2012. Water Availability Is the Main Climate Driver of Neotropical Tree Growth. *PLOS ONE* 7: e34074.

Zalamea P-C, Heuret P, Sarmiento C, *et al.* 2012. The Genus Cecropia: A Biological Clock to
Estimate the Age of Recently Disturbed Areas in the Neotropics. *PLoS ONE* 7: e42643.

Zalamea P-C, Sarmiento C, Stevenson PR, Rodríguez M, Nicolini E, Heuret P. 2013. Effect
of rainfall seasonality on the growth of Cecropia sciadophylla: intra-annual variation in leaf
production and node length. *Journal of Tropical Ecology* 29: 361–365.

Zalamea P-C, Stevenson PR, Madriñán S, Aubert P-M, Heuret P. 2008. Growth pattern and
age determination for Cecropia sciadophylla (Urticaceae). *American Journal of Botany* 95:
263–271.

1   **TABLES**

2   **Table 1.** List of measured growth and tree-level traits.

Trait	Abbreviation	Unit
Architectural growth traits		
Internode length	IN	mm
Internode length residuals		-
Phyllochron		day
Annual shoot length	AS	cm
Number of nodes per AS		-
Cumulated tree height		m
Tree-level traits		
Age after determination	Age	year
Tree height	Height	m
Diameter at breast height	DBH	cm
Number of trunk internodes	IN_{A1}	-
Number of A2 bearing branches	Br_{bear}	-
Number of A2 dead branches	Br_{dead}	-
Total number of A2 branches	Br_{tot}	-
Branching order	Order	-
Total number of inflorescences	Fl_{tot}	-
Total number of leaves	$Leaf_{tot}$	-
Total estimated crown area	A_{crown}	m²
First branching height	Br1stH	m
First flowering height	Fl1stH	m
First branching node rank	Br1stIN	-
First flowering node rank	Fl1stIN	-
First branching age	Br1stAge	year
First flowering age	Fl1stAge	year

3
4
5

1 **Table 2.** List of measured functional traits and the infered growth rate.

Trait	Abbreviation	Unit
Leaf content traits		
Residual water content	$H_2O_{res,leaf}$	%
$\delta^{13}C$ content	$\delta^{13}C_{leaf}$	‰
Carbon content	C_{leaf}	$g\ kg^{-1}$
Nitrogen content	N_{leaf}	$g\ kg^{-1}$
Carbon/Nitrogen ratio	CN_{leaf}	-
Phosphorus content	P_{leaf}	$g\ kg^{-1}$
Potassium content	K_{leaf}	$g\ kg^{-1}$
Chlorophyll content	Chl_{leaf}	$mg\ ml^{-1}$
Leaf morphological traits		
Leaf lifespan	LL	day
Petiole length	L_{pet}	cm
Petiole cross-sectional area	A_{pet}	mm²
Number of lobes	N_{lobe}	-
Leaf mass area	LMA	$g\ cm^{-2}$
Leaf thickness	Th_{leaf}	μm
Estimated leaf area	A_{leaf}	cm²
Wood trait		
Wood specific gravity	WSG	$g\ cm^{-3}$

2
3
4

Table 3. Linear mixed-effect models for measured architectural dimension traits between soil types and sites.

Abbreviation	Unit	Age	Soil	Direction	Counami FS	Counami WS	Sparouine FS	Sparouine WS	Range
Height	m	***	***	+	16.13 ± 1.92	13.31 ± 1.10	18.98 ± 0.65	17.27 ± 0.99	$10.05 - 21.99$
DBH	cm	***	**	+	15.02 ± 2.24	12.44 ± 2.33	19.30 ± 0.97	$18.24\ 1 \pm.52$	$6.21 - 30.49$
IN_{A1}	-	***			429.2 ± 87.0	457.9 ± 35.0	289.5 ± 10.1	277.9 ± 14.8	$221 - 765$
Br_{bear}	-		*	+	7.85 ± 1.75	4.32 ± 1.55	10.74 ± 1.79	11.11 ± 1.82	$0 - 19$
Br_{dead}	-	***			10.23 ± 6.02	7.47 ± 1.97	12.84 ± 2.35	10.58 ± 2.81	$0 - 33$
Br_{tot}	-				13.62 ± 4.07	6.79 ± 2.90	21.47 ± 3.80	24.16 ± 8.17	$1 - 75$
Order	-	***	*	+	2.92 ± 0.41	2.32 ± 0.26	3.16 ± 0.17	3.26 ± 0.20	$1 - 4$
Fl_{tot}	-				55.54 ± 17.11	46.42 ± 12.31	75.68 ± 24.95	121.9 ± 68.0	$0 - 657$
$Leaf_{tot}$	-				140.6 ± 37.6	75.74 ± 31.54	170.74 ± 31.35	200.3 ± 72.7	$25 - 674$
A_{crown}	m²				16.44 ± 6.11	6.893 ± 3.472	21.79 ± 4.17	26.78 ± 7.89	$1.656 - 74.10$
Br1stH	m		**	+	9.04 ± 0.80	8.20 ± 0.80	10.45 ± 0.50	9.05 ± 0.85	$5.27 - 14.56$
Fl1stH	m	*	***	+	11.29 ± 1.38	9.12 ± 0.89	18.67 ± 0.71	16.71 ± 0.94	$6.62 - 20.99$
Br1stIN	-				141.8 ± 22.6	178.7 ± 32.1	117.0 ± 6.8	113.7 ± 8.6	$81 - 358$
Fl1stIN	-	*			188.6 ± 26.3	198.6 ± 32.7	284.2 ± 7.9	264.1 ± 13.9	$101 - 412$
Br1stAge	year				4.272 ± 0.752	5.297 ± 1.053	3.358 ± 0.220	3.263 ± 0.275	$2.332 - 11.94$
Fl1stAge	year				5.709 ± 0.821	8.833 ± 0.986	8.307 ± 0.231	7.699 ± 0.406	$3.132 - 11.49$

Age and Soil refer to potential fixed effects. Direction refers to the direction of the potential relationship between soil and the tested trait: + increasing trait toward FS; - decreasing trait toward FS. Site is set as a random effect.***: $P < 0.001$; **: $P < 0.01$; *: $P < 0.05$. See Table 2 for definitions of abbreviations.

Table 4. Linear-mixed effect models for measured functional traits between soil types and sites.

Abbreviation	Unit	Age	Soil	Direction	Counami FS	Counami WS	Sparouine FS	Sparouine WS	Range
$H_2O_{res,leaf}$	%		*	-	9.921 ± 0.557	10.18 ± 0.26	7.823 ± 0.132	8.255 ± 0.124	7.38 – 11.50
$\delta^{13}C_{leaf}$	‰				-29.61 ± 0.38	-29.74 ± 0.50	-29.30 ± 0.19	-29.78 ± 0.31	-32.12 - -28.41
C_{leaf}	g kg^{-1}				46.99 ± 0.39	46.27 ± 0.41	47.12 ± 0.22	46.74 ± 0.84	39.11– 48.04
N_{leaf}	g kg^{-1}	***			2.531 ± 0.253	2.312 ± 0.170	2.649 ± 0.111	2.743 ± 0.177	1.630 – 3.200
CN_{leaf}	-		**		19.13 ± 1.87	20.52 ± 1.52	17.95 ± 0.85	17.16 ± 0.67	14.84 – 28.01
P_{leaf}	g kg^{-1}				1.687 ± 0.305	1.698 ± 0.205	1.322 ± 0.231	1.178 ± 0.059	0.588 – 2.42
K_{leaf}	g kg^{-1}		*	+	16.31 ± 2.98	14.56 ± 2.43	12.30 ± 2.32	9.099 ± 0.865	4.88 – 27.50
Chl_{leaf}	mg ml^{-1}				62.16 ± 9.73	53.50 ± 6.51	101.4 ± 20.6	110.7 ± 31.5	31.1 – 309.0
LL	day	*			117.3 ± 16.8	110.3 ± 10.8	79.96 ± 0.542	83.82 ± 0.626	58 - 181
L_{pet}	cm	*			37.19 ± 4.60	27.34 ± 3.32	42.79 ± 3.62	47.5 ± 2.70	14.2 – 62.0
A_{pet}	mm²	*			67.78 ± 12.36	47.90 ± 7.22	77.07 ± 12.41	81.35 ± 8.88	25.1 – 126.3
N_{lobe}	-				7.923 ± 0.268	7.353 ± 0.316	7.667 ± 0.267	8.000 ± 0.218	6 - 9
LMA	g m^{-2}	***			97.42 ± 10.41	96.43 ± 7.95	99.55 ± 3.74	89.06 ± 5.30	63.49 – 130.3
Th_{leaf}	μm	*			366.2 ± 23.7	397.8 ± 19.9	314.7 ± 17.6	302.9 ± 29.8	213 - 482
A_{leaf}	cm²	***			1860 ± 393	1094 ± 215	2043 ± 362	2587 ± 472	321 - 4916
WSG	g cm^{-3}				0.389 ± 0.029	0.352 ± 0.019	0.325 ± 0.018	0.319 ± 0.014	0.252 – 0.516

Age and Soil refer to potential fixed effects. Direction refers to the direction of the potential relationship between soil and the tested trait: + increasing trait toward FS; - decreasing trait toward FS. Site is set as a random effect.***: $P < 0.001$; **: $P < 0.01$; *: $P < 0.05$. See Table 2 for definitions of abbreviations.

FIGURE LEGENDS

Fig. 1. Architectural growth traits according to node rank and age (years). Main boxes

represent features according to soil types. Inboxes represent mean trajectories after clustering

longitudinal analyses (*kml*). The left column represents Counami trees, the right column

represents Sparouine trees. Distributions between soil types and kml-trajectories are

represented with Pearson chi-squared test. Red: ferralitic soils; black: white-sand soils. Blue:

trajectory A; green: trajectory B; orange: trajectory C. Thick lines: means; dashed lines:

confidence intervals at 95%.

Fig. 2. Cumulated number of trunk nodes with pairs of inflorescences and number of branches

per annual shot according to the age (year). (a) and (b) Means for inflorescences for Counami

and Sparouine respectively. (c) and (d) Means for branches for Counami and Sparouine

respectively. Red: ferralitic soils; black: white-sand soils. Thick lines: means, dashed lines:

confidence intervals at 95%.

Fig. 3. Principal component analysis (PCA) on functional traits for the two sites for 70 trees.

(a) Correlation circle of data with the histogram of inertia. (b) Individual factor map of data

according to soil types and sites. In (a), the colour gradient indicates the contribution of each

variable to the axis. See Table 3 for definitions of abbreviations. "Res" prefixes indicate

residuals after removing the ontogenetical effect. In (b), significant differences in coordinates

($P < 0.05$; ANOVA) between soil types and sites are indicated by letter according to the

considered axis. Red: ferralitic soils; black: white-sand soils; COU: Counami; SPA:

Sparouine.

SUPPORTING INFORMATION LEGENDS

Appendix S1. Pedological characterization: Materials and methods, and results.

Appendix S2. Materials and methods: residuals, year delineation and inference of age.

Appendix S3. Materials and methods: measurement of leaf and trunk functional traits.

Appendix S4. Autocorrelation coefficients for growth, branching, and flowering processes:

Materials and methods, and results.

Fig. S1. Morphological features of *Cecropia obtusa* Trécul (Urticaceae). (a) Focus on an

apex, ca: calyptra; pe: petiole; if: inflorescence. (b) Focus on a branch tier, br: branch; in:

internode, axis as the trunk are made of a linear succession of internodes; ins: inflorescence

scars, these are twice just above the axillary leaf; les: leaf scar. (c) Focus on an internode, in:

internode; ins: inflorescence scars; les: leaf scar; no: a node marled by the calyptra scar,

allowing for the delineation of internodes along an axis as the trunk.

Fig. S2. Box plots of mean annual rainfall (mm) from 1980 to 2016. (a) Counami, (b)

Sparouine.

Fig. S3. Tree heights (m) according to diameters at breast height (cm). Red: ferralitic soils;

black: white-sand soils. Cross: Sparouine; square: Counami.

Fig. S4. Other architectural growth traits according to age (years): number of nodes per

annual shoot and annual shoot length. Main boxes represent features according to soil types.

Inboxes represent mean trajectories after clustering longitudinal analyses (*kml*). The left

column represents Counami trees, the right column represents Sparouine trees. Distributions

between soil types and kml-trajectories are represented with Pearson chi-squared test. Red:

ferralitic soils; black: white-sand soils. Blue: trajectory A; green: trajectory B; orange:

trajectory C. Thick lines: means; dashed lines: confidence intervals at 95%.

(a) PCA 1: 28.5%, PCA 2: 18.5%

Variables: res.Chl$_{leaf}$, d^{13}C$_{leaf}$, res.LMA, res.L$_{pet}$, N$_{lobe}$, res.A$_{pet}$, res.A$_{leaf}$, C$_{leaf}$, res.C:N$_{leaf}$, res.N$_{leaf}$, WSG, res.Th$_{leaf}$, K$_{leaf}$, res.LL, P$_{leaf}$, res.H$_2$O$_{res,leaf}$

Contrib. (%): 10.0, 7.5, 5.0, 2.5

(b) PCA 1: 28.5%, PCA 2: 18.5%

SPA WS, SPA FS, COU WS, COU FS

	COU FS	COU WS	SPA FS	SPA WS
PCA 1	a	a	a	a
PCA 2	a	a	b	b