1
2
3
4
5 **Saccades and Handedness Interact to Affect Scene Memory**
6
7

10 Timothy M. Ellmore[1*], Bridget Mackin[1], and Kenneth Ng[1]

13 [1]Department of Psychology,
14 City University of New York, City College
15 New York, New York, United States of America
16
17

*Corresponding author

Timothy M. Ellmore, Ph.D.
Associate Professor
Department of Psychology
The City College of the City University of New York
E-mail: tellmore@ccny.cuny.edu
Phone: (212) 650-5714
Fax: (212) 650-5659
28
29

**Abstract**

Repetitive saccades benefit memory when executed before retrieval, with greatest effects for episodic memory in consistent-handers. Questions remain including how saccades affect scene memory, an important visual component of episodic memory. The present study tested how repetitive saccades affect working and recognition memory for novel scenes. Handedness direction (left-right) and degree (strong/consistent vs. mixed/inconsistent) was measured by raw and absolute laterality quotients respectively from an 8-question handedness inventory completed by 111 adults. Each then performed either 30 seconds of repetitive horizontal saccades or fixation before or after tasks of scene working memory and scene recognition. Regression with criterion variables of overall percent correct accuracy and d-prime sensitivity showed that when saccades were made before working memory, there was better overall accuracy as a function of increased direction but not degree of handedness. Subjects who made saccades before working memory also performed worse during subsequent recognition memory, while subjects who fixated or made saccades after the working memory task performed better. Saccades made before recognition resulted in recognition accuracy that was better (Cohen's d=0.3729), but not significantly different from fixation before recognition. The results demonstrate saccades and handedness interact to affect scene memory with larger effects on encoding than recognition. Saccades before scene encoding in working memory are detrimental to short- and long-term memory, especially for those who are not consistently right-handed, while saccade execution before scene recognition does not appear to benefit recognition accuracy. The findings are discussed with respect to theories of interhemispheric interaction and control of visuospatial attention.

**Introduction**

*Saccade Induced Retrieval Enhancement*

As few as thirty seconds of horizontal bilateral saccadic eye movements before testing

selectively enhances explicit memory, most notably episodic memory retrieval for laboratory and

everyday events (Christman et al. 2003). This has been termed saccade induced retrieval

enhancement (SIRE) by Lyle and colleagues (Lyle et al. 2008a). The benefits of SIRE appear to

be specific to retrieval. Early preliminary findings reported that eye movements before encoding

hurt subsequent memory performance (Christman & Butler 2005). Later experiments by others

showed that the benefits of horizontal eye movements are seen when they immediately precede

episodic memory retrieval but not when they precede encoding (Brunye et al. 2009). One

hypothesis is that horizontal eye movements enhance interhemispheric interaction, which is

associated with superior episodic memory (Christman & Propper 2001). Increased

interhemispheric interaction has also been related to decreased false memories in a semantic

associates paradigm (Christman et al. 2004).

Bilateral eye movements also appear to enhance some types of recognition memory.

Subjects who made bilateral eye movements were more likely to correctly recognize previously

presented words, but less likely to falsely recognize critical non-studied associates (Parker &

Dagnall 2007). Parker *et al* conducted multiple experiments in a study investigating effects of

bilateral saccadic eye movements on item, associative, and contextual information. In tests of

item recognition they found bilateral horizontal eye movements versus no eye movements

enhanced item recognition by increasing hit rate and decreasing false alarms; an additional

remember-know analysis showed eye movements increased "remember" responses (Parker et al.

2008). For associative recognition, they found bilateral eye movements increased correct

responses to intact pairs and decreased false alarms to rearranged pairs (Parker et al. 2008).

Bilateral eye movements also increased correct recall for both intrinsic (color) and extrinsic

(spatial location) context (Parker et al. 2008).

More recently, the horizontal SIRE effect has been extended beyond the visual domain in

a study that showed retrieval enhancement in the somatosensory system after alternating left-

right tactile stimulation (Nieuwenhuis et al. 2013). In what appears to be the only study of the

effects of horizontal saccadic eye movements on the retrieval of landmark shape and location

information, increased recognition sensitivity and decreased response times were reported in a

spatial memory test with effects only seen when eye movements preceded episodic memory

retrieval, but not when they preceded encoding (Brunye et al. 2009). Furthermore, these same

authors found that eye movements were only beneficial in an old-new recognition paradigm,

which purportedly involves more elaborate recollective processing and presumably demands a

high degree of right and left-hemisphere activity, compared to a forced-choice recognition test,

which is thought to be more dependent on familiarity.

Despite the significant clinical and applied implications for the effects of bilateral

saccades (Lyle & Jacobs 2010; Propper & Christman 2008; Stickgold 2002), there are only a few

studies employing neuroscientific methods to measure brain-based correlates of bilateral

saccades. Bilateral eye movements have been reported to have significant effects on

interhemispheric coherence in the gamma band as measured by EEG (Propper et al. 2007). But a

more recent EEG study found scant evidence that eye movements altered interhemispheric

coherence or that improvements in recall were correlated with changes in coherence (Samara et

al. 2011).

*SIRE and Handedness*

A growing literature documents that SIRE is modulated by handedness. A study of both

recall and recognition with strongly right-handed (SR) and non-strongly right handed (nSR)

subjects found that eye movements largely benefited the former while it was shown to be

somewhat detrimental to the latter (Lyle et al. 2008a). This finding was initially interpreted to

support the hemispheric interaction hypothesis. Better memory without eye movements is

thought to exist in nSR individuals because nSR handedness is thought to be a behavioral marker

for greater interhemispheric interaction. Consistent with this idea, middle-age nSR subjects

perform better on tasks like paired associate recall and source memory, which likely depend on

hemispheric interaction, compared to tasks like face recognition and forward digit span which

depend less on hemispheric interaction (Lyle et al. 2008b). For example one study (Lyle &

Orsborn 2011) required subjects to classify faces as famous or novel with face presentation

occurring in the left and right visual fields simultaneously (bilaterally) or in one field only

(unilaterally). Famous faces were classified more quickly and accurately during bilateral

presentation, reflecting that interhemispheric interaction facilitates famous face recognition, but

neither inconsistent handedness nor saccades increased the size of bilateral gain.

Strong right-handedness, which is argued to be associated with decreased

interhemispheric interaction, was associated with higher rates of false memories while bilateral

saccades were associated with fewer false memories. Other evidence suggests that mixed-

handers display better episodic memory in comparison with strong right-handers on assessments

of explicit word recall and recall of real world events, but when corrected scores are analyzed

handedness does not influence implicit word fragment completion (Propper et al. 2005). Mixed-

handedness and bilateral saccadic eye movements have also both been associated with an earlier

offset of childhood amnesia and support the idea that interhemispheric interaction exerts effects

on retrieval but not encoding of episodic memories. Mixed-handed subjects also demonstrate

greater autobiographical recollection on components of seeing hearing and emotion compared to

right-handed individuals, and 30 sec of bilateral eye movements induce greater levels of

autobiographical recollection across a range of components (Parker & Dagnall 2010).

*From Interhemispheric Interaction to Top Down Control*

A recent experiment in which subjects were shown arrays of lateralized letters and were

asked whether target letters matched either of two probe letters, saccades were reported to

enhance retrieval by increasing interaction among brain hemispheres (Lyle & Martin 2010).

Matching targets and probes were presented to either the same hemisphere or to separate

hemispheres. Interhemispheric interaction was required on the across-hemisphere trials and

intrahemispheric processing was required on the same hemisphere trials. Increased match

detection accuracy was found on the within-hemisphere trials as a function of pre-task eye-

movements suggesting saccades enhance intrahemispheric processing but not interhemispheric

interaction. The nSR subjects showed higher across-hemisphere accuracy suggesting that an

absence of strong right-handedness may reflect greater interhemispheric interaction. A recent

comparison of consistent and inconsistent left- and right-handers on associative memory tests

after saccade or no-saccade conditions showed that saccades enhanced retrieval for consistent-

handers only, but impaired retrieval for inconsistent-handers (Lyle et al. 2012). This important

study established that handedness consistency, regardless of left or right direction, is an

important factor to consider when studying memory.

Given the lack of convincing support for the idea that interhemispheric interaction

underlies the bilateral saccade effects, more recent work has focused on the idea that saccade

execution enhances cognition by altering attentional control. In one study, performance on the

well-established revised attentional network test (Fan et al. 2009) was assessed after either

repetitive bilateral saccades or central fixation (Edlin & Lyle 2013). Saccade execution increased

the executive function network, which encompasses attentional control, by decreasing response

times to target stimuli in presence of response-incongruent flankers. In this study, the saccade-

induced enhancement of attentional control occurred independently of handedness consistency.

This raises the possibility that there could be a larger role for top-down attention when memories

are more difficult to access, and recent results support that saccade execution has a greater

facilitative effect on retrieval when recall and recognition are more difficult (Lyle & Edlin 2015).

*Outstanding Questions and Current Objectives*

There remain several questions about the behavioral effects of repetitive saccades that

have not yet been explored. The first is how do repetitive saccades affect memory for complex

novel visual stimuli? The second involves how saccades affect stages of memory processing

other than retrieval. Most previous investigations have had subjects perform bilateral saccades

after encoding but before retrieval. It is not well understood how bilateral saccades affect the

encoding of complex novel visual stimuli, nor is it understood how these effects are modulated

by handedness. Understanding whether eye movements and handedness influence working

memory or long-term recognition memory or both is also relevant to a decades-long theoretical

debate concerning potential relationships between working memory and long-term memory.

"Buffer" accounts (e.g., Baddeley's multi-component model) argue for a central executive

coordinating separate phonological and visuo-spatial short-term stores that are separate from

long-term memory (Baddeley 2003), while the hierarchically arranged embedded process

account argues working memory is a subset of activated memory in the focus of attention and the

subset of long-term memory that is currently activated (Cowan 1999). To address these

questions, we conducted an exploratory study of the effects of repetitive bilateral saccades and

handedness on scene memory. We utilized a variant of the well-known Sternberg working

memory paradigm (Sternberg 1966) in which during a series of trials five pre-experimentally

novel scenes are presented and held online during a short 6 sec delay period. After a 10-minute

period of awake rest, we tested subjects' long-term recognition memory for scenes previously

maintained in short-term working memory. We have shown previously that subjects perform this

type of working memory task with scenes as stimuli and can recognize far above chance dozens

of scenes shown during the working memory task after a long-term retention interval (Ellmore et

al. 2015). Recent source analysis of scalp EEG acquired with this task has implicated a right

hemisphere parieto-occipital region that is active when maintaining scenes in working memory,

with temporal-spectral delay activity that correlates with subsequent probe memory retrieval

(Ellmore et al. 2017).

The specific objective of the present study was to test alternative hypotheses that making

repetitive saccades before or after encoding novel scenes in working memory would have either

beneficial or detrimental effects on subsequent memory as a function of handedness. A between-

subjects design was employed that allowed us to test 1) whether making repetitive saccades

*before* working memory encoding modulates subsequent working and long-term memory

retrieval as a function of handedness and 2) whether making repetitive saccades *after* working

memory encoding and before recognition modulates subsequent long-term recognition as a

function of handedness.

195
196     **Materials and Methods**
197
198     *Subjects*
199
A total of 111 subjects ($M$=20.74, $SD$=3.91, range 18 to 40; 68 females, 43 males) were recruited

through the Sona Systems online experimental scheduling system of the City College of New

York Psychology Department. Each provided written informed consent, completed a revised

Edinburgh Handedness Inventory (EHI) (Williams 2010), and completed study procedures

according to a protocol approved (approval number 2015-0550) by the Institutional Review

Board of the City College of New York Human Research Protection Program. At the end of the

study, each subject received extra course credit for a total of one hour of participation.

The target sample size of over 100 subjects was determined based on similar sample sizes

used in previous published studies of eye movements and memory as well as on our ability to

recruit as many participants as possible during the date range for study recruitment which began

September 16, 2015 and ended November 22, 2017. In addition, for this study all measures,

conditions, and data exclusions are reported herein.

212
213     *Experimental Design*
214
All subjects completed a working memory task, followed by a 10-minute period of quiet awake

rest, followed lastly by a test of recognition memory. The study utilized a between-subjects

design to manipulate eye movements by performance of two different eye tasks (saccade or

fixation) executed either before performance of the working memory task (encoding group) or

after the working memory task but before a test of recognition memory (recognition group).

There was a total of 65 subjects (*M*=20.86, *SD*=3.63, range 18 to 38; 41 females, 24 males) in the

encoding group and a total of 46 subjects (*M*=20.56, *SD*=4.30, range 18 to 40; 27 females, 19

males) in the recognition group. Within these two groups, subjects were assigned to perform

either the saccade or fixation task using a pre-determined randomization schedule to minimize

experimenter bias.

225
226    *Apparatus*
227
All subjects completed tasks inside a closed room to minimize auditory and visual distractions.

Tasks were programmed in SuperLab 5 (Cedrus Corporation). Tasks were displayed on a 28-inch

LED monitor with a refresh rate of 60 Hertz and a screen resolution of 1920 by 1080 pixels.

Participants sat in the dark 96 cm from the monitor on an adjustable stool and positioned their

head in an Ultra Precision Head Positioner™ (Arrington Research, Inc), a combined chin and

forehead rest, to help minimize head movements. The experimenter remained in the room for the

duration of the experiment, but was seated outside of subjects' view. A black curtain separated

the experimenter from the subject to minimize distraction. To ensure subjects complied with

instructions when required to make saccades or fixation, an infrared light emitter and a 220 Hz

eye-tracking    camera    (*ViewPoint    EyeTracker*®    by    Arrington    Research,    Inc,

[www.ArringtonResearch.com](www.ArringtonResearch.com)) was placed between the monitor and the subject to track the pupil

of one eye. The experimenter monitored the x and y eye position as measured by real-time

computation of pupil center-of-mass on a separate monitor only visible to the experimenter.

Subject behavioral responses were collected using a Cedrus RB-740 response pad placed on the

table in front of them.

243
244    *Tasks*
245

246    *Working Memory*
247
Subjects completed 40 trials of a variant of a Sternberg working memory task (Sternberg 1966)

used in previous experiments (Ellmore et al. 2015; Ellmore et al. 2017). Each trial consisted of a

fixed load of 5 scene stimuli. A total of 220 pre-experimentally novel scenes were shown across

the 40 trials of the working memory task. Each was a 24-bit color image of an outdoor scene

sampled from the SUN database (Xiao et al. 2010). Scene stimuli were 800 by 600 pixels and

were presented at the center of the monitor subtending a visual angle of 16 by 12 degrees. On

each trial, five scenes appeared sequentially for 2 sec each, followed by a 6 sec delay period

consisting of a white screen with a foveal crosshair, during which subjects were required to keep

these scenes in mind, and then a probe scene was displayed for 2 sec (Fig. 1a). Each trial was

separated by 5 sec of black screen. Half (20) of the trials contained a positive probe, which was

one of the five scenes presented before the delay period. The other 20 trials contained a negative

probe, a new scene not previously shown in the set of scenes presented before the delay period.

Positive probes were randomly selected from each trial's five encoding stimuli so that the serial

position of the probe was equally likely to come from encoding position 1, 2, 3, 4, or 5 across the

set of WM trials. Positive and negative probe trials were randomly distributed across the set of

40 working memory trials. Upon the presentation of the probe in each working memory trial,

subjects were instructed to press a green button on the response pad if the probe scene matched

one of the scenes presented in the set of 5 scenes shown before the 6 sec delay period; subjects

were instructed to press a red button if the probe scene did not match one of the scenes presented

in the set of 5 scenes shown before the delay. For each working memory probe presentation,

subjects were required to respond within the 2 seconds that the probe was on the screen. Before

performing the working memory task, subjects were given a 5-minute demonstration task with

different stimuli (animals) to ensure that they understood task instructions before the actual

working memory task began.

*Recognition Memory*

After completing the working memory task, subjects were allowed to disengage from the head

restraint and rest quietly in the experiment room for ten minutes. At the end of this ten-minute

retention interval, subjects completed a task of recognition memory (Fig. 1b) that included a total

of 200 scenes. One hundred of these scenes were randomly sampled from the sets of scenes

presented before the delay in each of the 40 previous working memory trials. A total of 8 out of

these 100 "old" stimuli were previously used as positive probes in the WM task. The other 100

scenes were completely new scenes sampled randomly from the SUN database. The 100 old and

100 new scenes were presented one at a time for 2 sec each in random order as individual trials.

For each scene presentation during the recognition test, subjects were required to respond within

the 2 seconds that the probe was on the screen. A 2 sec black screen separated each trial. For

each trial, subjects were instructed to press the green button on the response pad to indicate that

they recognized having previously viewed the scene; subjects were instructed to press the red

button to indicate that they did not recognize the scene.

*[insert Figure 1 here]*

*Eye Tasks*

The main experimental manipulation required subjects to perform one of two eye tasks (saccade

or fixation) either before the working memory task or before the recognition memory task. The

saccade task required subjects to make deliberate repetitive horizontal saccades for 30 sec. Each

horizontal saccade was cued by the appearance of a white disc on a black background. The

diameter of the white disk was 70 pixels (1.49 degrees of visual angle). Each dot appeared for

500 ms and alternated its position between the right and left sides of the screen for a total of 30

sec (Fig. 2a). The center-to-center distance of the disk as it alternated between the right and left

sides of the screen was 1285 pixels (27.3 degrees of visual angle). The fixation task required that

subjects fixate a single disc (same size as the white disc in the saccade task) that was stationary

at the center of the screen. The disc appeared on a black background for 30 sec and cycled

through 6 colors (red, blue, yellow, green, pink, and purple), with each color presented

individually for 500 ms (Fig. 2b). To keep subjects' attention on the different eye tasks, subjects

were instructed to count covertly the number of times the disc alternated positions between the

left and right screen positions; during the fixation task, subjects were instructed to count covertly

the number of times the disc changed colors. The number of disc alternations and color changes

was equal.

311  
312    *[insert Figure 2 here]*  
313  
314  
315  
316    *Analysis*  
317  
The analysis began with the computation of a laterality quotient (LQ) from the responses to the

eight    questions    of    the    EHI-R    (Williams    2010).    The    EHI-R

(https://doi.org/10.13140/RG.2.2.33298.25284) is similar to the original EHI published in 1971

(Oldfield 1971). Highly variable usage of modified versions of the original EHI are prominent in

the scientific literature, which may imperil efforts to produce replicable and convergent findings

(Edlin et al. 2015). Therefore, it is important to explain and scientifically justify differences in

any revision and note differences with the inventory used in many other labs (Lyle et al. 2008b).

Compared to the original EHI, the EHI-R discards three items ("broom", "opening box", and

"drawing") and adds a new one – "computer mouse" – making the total number of items eight. It

also uses the Likert format in an attempt to simplify the confusing instructions of the original

EHI. The items "opening box" and "broom" are removed based on statistical evidence including

factor analytic studies (Dragovic 2004; Williams 1986) that find a single Handedness factor

whose loadings on it of these two activities are low outliers. "Drawing" is removed because it

could be used as a substitute for writing since drawing and writing are very highly (about 0.9)

correlated and the inclusion of both adds little new information. The addition of "computer

mouse" is based on a suggestion by Dragovic (Dragovic 2004). The use of a computer mouse is

one major unimanual activity in today's world that was not widespread forty years ago when the

EHI was developed. Additionally, the EHI-R has been used in several recent studies (Eriksen et

al. 2018; Holcombe et al. 2017; Kielar et al. 2016; Rice & M. Gosser 2015). The EHI-R asks

subject to indicate which hand they use to complete various everyday activities (e.g., striking a

match, using a computer mouse). Participants indicated their handedness preference for each

item by checking off one of the following response options (with corresponding point values):

"always right" (50) "usually right" (25), "no preference" (0), "usually left" (-25), and "always

left (-50). The LQ was calculated by tallying the point values for all eight items and dividing the

total score by four, resulting in LQ scores that ranged from +100 (right handed) to -100 (left

handed). A total of 7 subjects scored as left-handed (LQ<0), one subject scored perfectly mixed

(LQ=0), 33 subjects scored non-strongly right-handed (LQ<80), and 78 subjects scored strongly-

right handed (LQ>=80).

The next step in the analysis was to compute for each subject performance expressed as

overall percentage correct and sensitivity as d-prime for the 40 trials of the working memory task

and separately for the 200 trials of the recognition memory task. The d-prime sensitivity measure

from signal detection theory accounts for effects of response bias (Stanislaw & Todorov 1999).

The proportions of total hits and false alarms were first calculated for each subject and each task.

A hit was counted when a previously presented stimulus (i.e., a positive probe in the working

memory task) was signaled by the subject pressing a button indicating, correctly, that the

stimulus had been previously seen (an old stimulus correctly classified as old). A false alarm was

counted when a (new) stimulus not previously presented (i.e., a negative probe in the working

memory task) was indicated by the subject pressing a button indicating, incorrectly, that the

stimulus had been previously presented (a new stimulus incorrectly classified as an old stimulus).

Then d-prime was computed as the difference in standardized normal deviates of hits minus false

alarms: *Z(hit rate) – Z(false alarm rate)*. Separate linear regression analyses were then performed

with the criterion variables (Y) percent correct accuracy and d-prime sensitivity. The predictor

variables (X) were raw LQ for direction of handedness and the absolute values of LQ for degree

of handedness. Regressions were computed to predict the criterion variables of accuracy or d-

prime by either direction or degree of handedness as a function of the eye-movement task

(fixation vs. saccade) and placement of the eye-movement task (before the working memory or

recognition memory task). Regression analyses were computed in GraphPad Prism version 7.0

(GraphPad Software, La Jolla California USA, www.graphpad.com). Statistical tests of the

regression slopes were also computed to assess differences in performance as a function of

laterality and type and placement of the eye-movement task. Statistical analysis of regression

intercepts (elevations) was used to assess differences in overall performance as a function of type

and placement of the eye-movement task. The raw data input to these analyses is provided in the

Supplemental File.

371
372 **Results**
373
374 *Working Memory Performance*
375
376 *Analysis of Regression Slopes*
377
Linear regression was used to quantify the relationship between working memory performance,

as measured by both percent correct accuracy and d-prime, and direction of laterality as a

function of the eye task. The statistical tests assessed whether the slope of the regression

equation differed significantly from zero. Similar tests were used to assess the relationship

between the dependent measures and the degree of laterality, indicating consistency of

handedness, using the absolute value of the laterality quotient.

When fixation occurred before the working memory task, working memory percent

correct accuracy increased non-significantly as a function of direction of laterality

(slope=0.03098 $\pm$ 0.03765 SE, $F_{(1,31)}$= 0.6774, p=0.4168, Fig 3a blue line and blue circles). The

d-prime measure also increased non-significantly as a function of direction of laterality

(slope=0.00168 $\pm$ 0.00337 SE, $F_{(1,31)}$=0.2473, p=0.6225, Fig 3b blue line and blue circles).

Working memory percent correct accuracy increased non-significantly as a function of degree of

laterality (slope=0.02432 $\pm$ 0.05178 SE, $F_{(1,31)}$= 0.2206, p=0.6419, Fig 3c blue line and blue

circles). The d-prime measure also increased non-significantly as a function of degree of

laterality (slope=0.001734 $\pm$ 0.00461 SE, $F_{(1,31)}$=0.141, p=0.7098, Fig 3d blue line and blue

circles).

When saccades occurred before the working memory task, working memory percent

correct accuracy increased significantly as a function of direction of laterality (slope=0.1029 $\pm$

0.04365 SE, $F_{(1,30)}$= 5.56, p=0.0251, Fig 3a red line and red triangles). The d-prime measure

increased non-significantly as a function of direction of laterality (slope=0.0029 $\pm$ 0.0032 SE,

$F_{(1,30)}$= 0.8206, p=0.3722, Fig 3b red line and red triangles). Working memory percent correct

accuracy increased non-significantly as a function of degree of laterality (slope=0.1212 ± 0.1455

SE, $F_{(1,30)}$= 0.694, p=0.4114, Fig 3c red line and red triangles). The d-prime measure also

increased non-significantly as a function of degree of laterality (slope=0.01137 ± 0.00984 SE,

$F_{(1,30)}$= 1.334, p=0.2571, Fig 3d red line and red triangles).

The subjects who performed fixation and saccades after the working memory task serve

as a control group, as one would expect no influence of these eye tasks on working memory

performance if they occur after the working memory task. Consistent with this prediction, when

fixation occurred after the working memory task, working memory percent correct accuracy

increased non-significantly as a function of direction of laterality (slope=0.02973 ± 0.02872 SE,

$F_{(1,21)}$=1.072, p=0.3123, Fig 3a grey line and grey boxes). The d-prime measure also increased

non-significantly as a function of direction of laterality (slope= 0.00116 ± 0.00268 SE,

$F_{(1,21)}$=0.1873, p=0.6696, Fig 3b grey line and grey boxes). Working memory percent correct

accuracy increased non-significantly as a function of degree of laterality (slope=0.06811 ±

0.05039 SE, $F_{(1,21)}$=1.827, p=0.1909, Fig 3c grey line and grey boxes). The d-prime measure also

increased non-significantly as a function of degree of laterality (slope= 0.00620 ± 0.00462 SE,

$F_{(1,21)}$=1.805, p=0.1935, Fig 3d grey line and grey boxes).

When saccades occurred after the working memory task, working memory percent

correct accuracy increased non-significantly as a function of direction of laterality

(slope=0.005429 ± 0.02251 SE, $F_{(1,21)}$=0.05818, p=0.8117, Fig 3a cyan line and cyan stars). The

d-prime measure decreased non-significantly as a function of direction of laterality (slope=-

0.00034 ± 0.0025 SE, $F_{(1,21)}$=0.0187, p=0.8926, Fig 3b cyan line and cyan stars). Working

memory percent correct accuracy increased non-significantly as a function of degree of laterality

(slope=0.0035 ± 0.03747 SE, $F_{(1,21)}$=0.0087, p=0.9265, Fig 3c cyan line and cyan stars). The d-

prime measure also increased non-significantly as a function of degree of laterality (slope=-

6.445e-005 ± 0.00422 SE, $F_{(1,21)}$=0.00023, p=0.9880, Fig 3d cyan line and cyan stars).

424
425 *Analysis of Regression Intercepts*
426
While analysis of regression slopes indicates change in performance as a function of direction or

degree of laterality and eye task timing, evaluation of the regression intercepts (i.e., elevations)

indicates whether the overall levels of performance differ.

When considering relationships of performance with direction of laterality, the four

intercepts of Fig 3a were significantly non-zero ($F_{(3,106)}$=6.164, p=0.0007) with best working

memory percent correct performance when saccades were made after the working memory task

(93.28% ± 1.844 SE, 95% C.I.: 89.45 to 97.12) and worst working memory percent correct

performance when saccades were made before the working memory task (77.79% ± 3.84, 95%

C.I.: 69.95 to 85.64%). The four intercepts of Fig 3b were also significantly non-zero

($F_{(3,106)}$=3.742, p=0.0133) with best working memory d-prime occurring when saccades were

made after the working memory task (3.179 ± 0.2082 SE, 95% C.I.: 2.746 to 3.612) and worst

working memory d-prime when saccades were made before the working memory task (2.442 ±

0.2819, 95% C.I.: 1.847 to 2.998).

When considering relationships of performance with degree of laterality, the four

intercepts of Fig 3c were significantly non-zero ($F_{(3,106)}$=5.699, p=0.0012) with best working

memory percent correct performance when saccades were made after the working memory task

(93.38% ± 3.07 SE, 95% C.I.: 87.00 to 99.77) and worst working memory percent correct

performance when saccades were performed before the working memory task (75.2% ± 12.80,

95% C.I.: 49.05 to 101.30%). The four intercepts of Fig 3d were also significantly non-zero

$(F_{(3,106)}=3.909$, p=0.0108) with best working memory d-prime occurring when saccades were

made after the working memory task (3.15 ± 0.3463 SE, 95% C.I.: 2.43 to 3.87) and worst

working memory d-prime when saccades were made before the working memory task (1.658 ±

0.866, 95% C.I.: -0.1106 to 3.427).

450
451    *[insert Figure 3 here]*
452
453
454
455    *Recognition Performance*
456
457    *Analysis of Regression Slopes*
458
Linear regression was also used to quantify the relationship between recognition memory, as

measured by both percent correct accuracy and d-prime, and direction of laterality as a function

of the eye task. The statistical test assessed whether the slope of the regression equation differed

significantly from zero. Similar tests were used to assess the relationship between the dependent

measures and the degree of laterality, indicating consistency of handedness, using the absolute

value of the laterality quotient.

When fixation occurred before the working memory task, later recognition percent

correct accuracy increased non-significantly as a function of direction of laterality

(slope=0.03746 ± 0.04449 SE, $F_{(1,31)}=0.7091$, p=0.4062, Fig 4a blue line and blue circles). The

d-prime measure also increased non-significantly as a function of direction of laterality

(slope=0.00208 ± 0.00281 SE, $F_{(1,31)}=1.351$, p=0.2542, Fig 4b blue line and blue circles).

Recognition memory percent correct accuracy increased non-significantly as a function of degree

of laterality (slope=0.03984 ± 0.06102 SE, $F_{(1,31)}= 0.4263$, p=0.5186, Fig 4c blue line and blue

circles). The d-prime measure also increased non-significantly as a function of degree of

laterality (slope=0.00344 ± 0.00383 SE, $F_{(1,31)}$=0.811, p=0.3747, Fig 4d blue line and blue

circles).

When saccades occurred before the working memory task, later recognition percent

correct accuracy increased non-significantly as a function of direction of laterality

(slope=0.06502 ± 0.04615 SE, $F_{(1,30)}$=1.985, p=0.1691, Fig 4a red line and red triangles). The d-

prime measure for later recognition also increased non-significantly as a function of direction of

laterality (slope=0.0032 ± 0.0027 SE, $F_{(1,30)}$= 1.351, p=0.2542, Fig 4b red line and red triangles).

Recognition memory percent correct accuracy also increased non-significantly as a function of

degree of laterality (slope=0.0629 ± 0.1472 SE, $F_{(1,30)}$= 0.1829, p=0.6720, Fig 4c red line and red

triangles). The d-prime measure for later recognition also increased non-significantly as a

function of degree of laterality (slope=0.0043 ± 0.00863 SE, $F_{(1,30)}$= 0.2443, p=0.6247, Fig 4d

red line and red triangles).

When fixation occurred after the working memory task and before the recognition

memory task, recognition percent correct accuracy increased significantly as a function of

direction of laterality (slope=0.08738 ± 0.03277 SE, $F_{(1,21)}$=7.111, p=0.0144, Fig 4a grey line and

grey boxes). The d-prime measure for recognition increased marginally significantly as a

function of direction of laterality (slope=0.00507 ± 0.00247 SE, $F_{(1,21)}$=4.23, p=0.0523, Fig 4b

grey line and grey boxes). Recognition memory percent correct accuracy increased non-

significantly as a function of degree of laterality (slope=0.1014 ± 0.0639 SE, $F_{(1,30)}$=2.517,

p=0.1276, Fig 4c grey line and grey triangles). The d-prime measure for recognition also

increased non-significantly as a function of degree of laterality (slope=0.00714 ± 0.00456 SE,

$F_{(1,30)}$=2.449, p=0.1326, Fig 4d grey line and grey triangles).

When saccades occurred after the working memory task and before the recognition

memory task, recognition percent correct accuracy increased non-significantly as a function of

laterality (slope=0.03398 ± 0.03797 SE, $F_{(1,21)}$=0.8009, p=0.3810, Fig 4a cyan line and cyan

stars). The d-prime measure for recognition increased non-significantly as a function of direction

of laterality (slope=0.0028 ± 0.00253 SE, $F_{(1,21)}$=1.274, p=0.2718, Fig 4b cyan line and cyan

stars). Recognition memory percent correct accuracy increased non-significantly as a function of

degree of laterality (slope=0.0679 ± 0.0626 SE, $F_{(1,30)}$=1.18, p=0.2897, Fig 4c cyan line and cyan

stars). The d-prime measure for recognition also increased non-significantly as a function of

degree of laterality (slope=1.115 ± 0.3462 SE, $F_{(1,30)}$=1.076, p=0.3113, Fig 4d cyan line and cyan

stars).

505
506    *Analysis of Regression Intercepts*
507
Analysis of regression intercepts was conducted to determine whether the overall level of

recognition performance differed as a function of the eye task timing. The four intercepts of Fig

4a representing recognition percent correct accuracy versus direction of laterality were

significantly non-zero ($F_{(3,106)}$=4.721, p=0.0039) with best recognition memory performance for

the condition of fixation occurring before the working memory task (70.08% ± 3.899 SE, 95%

C.I.: 62.12 to 78.03%, Fig 4a blue line and blue circles) and the lowest recognition memory

performance in the condition of saccades occurring before the working memory task (60.67% ±

4.06, 95% C.I.: 52.38 to 68.96%, Fig 4a red line and red triangles). Recognition performance

during the condition when saccades were made before the recognition memory task (70.06% ±

3.111, 95% C.I.: 63.59 to 76.53, Fig 4a cyan line and cyan stars) was greater than performance

during the condition of fixation before the recognition memory task (64.75% ± 2.819, 95% C.I.:

58.38 to 70.62%, Fig 4a grey line and grey boxes), but the difference was not significantly

different (two-tailed $t_{(44)}$=1.2648, p=0.2126, Cohen's d=0.3729). The level of recognition

performance in the condition when saccades were made before the recognition task was nearly

identical to performance after the condition of fixation before the working memory task (70.08%

± 3.899, 95% C.I.: 62.12 to 78.03%, Fig 4a blue line and blue circles). These virtually identical

levels of performance make the blue and cyan lines of Fig 4a appear to overlap.

The four intercepts of Fig 4b representing d-prime versus direction of laterality were

significantly non-zero ($F_{(3,106)}$=3.964, p=0.0101) with best recognition memory performance for

the condition of fixation occurring before the working memory task (1.327 ± 0.2467 SE, 95%

C.I.: 0.8235 to 1.83, Fig 4b blue line and blue circles) and the lowest recognition memory

performance in the condition of saccades occurring before the working memory task (0.8231 ±

0.2408, 95% C.I.: 0.3313 to 1.315, Fig 4b red line and red triangles).

The four intercepts of Fig 4c representing recognition percent correct accuracy versus

degree of laterality were significantly non-zero ($F_{(3,106)}$=4.863, p=0.0033) with best recognition

memory performance for the condition of fixation occurring before the working memory task

(69.76% ± 5.348 SE, 95% C.I.: 58.86 to 80.67, Fig 4c blue line and blue circles) and the lowest

recognition memory performance in the condition of saccades occurring before the working

memory task (60.22 ± 12.95, 95% C.I.: 33.78 to 86.66, Fig 4c red line and red triangles).

The four intercepts of Fig 4d representing recognition d-prime versus degree of laterality

were significantly non-zero ($F_{(3,106)}$=4.235, p=0.0072) with best recognition memory

performance for the condition of fixation occurring before the working memory task (1.205 ±

0.3354 SE, 95% C.I.: 0.5211 to 1.889, Fig 4d blue line and blue circles) and the lowest

recognition memory performance in the condition of saccades occurring before the working

memory task (0.6978 ± 0.7593, 95% C.I.: -0.853 to 2.249, Fig 4d red line and red triangles).

*[insert Figure 4 here]*
546
547
548     **Discussion**
549
Humans have a remarkable ability to remember a large number of scenes and other complex

visual stimuli even when given only a short period of time to encode them (Brown & Scott 1971;

Shepard 1967; Standing 1973; Standing et al. 1970). Scene memory is also an important

component of episodic autobiographical memory. Normal as well as disordered reminiscence

often includes vivid mental recollection of scenes where past events took place (Brewin 2014;

Burgess et al. 2002). In addition, recognition of a previously encountered scene in, for example

an old photograph, is often quick, highly accurate, and can trigger episodic memory retrieval.

Therefore, it is important to understand how scene memory is influenced by multifactorial traits

like handedness and by simple behavioral manipulations (e.g., bilateral saccades) shown

previously to influence episodic memory retrieval.

The present study was conducted to understand how both short- and long-term scene

memory is modulated by handedness and by bilateral horizontal saccades made before scene

encoding and recognition. Previous studies have shown that strongly right-handed and

consistently-handed benefit most from bilateral saccades (Chu et al. 2012; Lyle et al. 2012; Lyle

& Orsborn 2011; Prichard & Christman 2017; Propper et al. 2017), but these studies focused on

mostly verbal aspects of explicit or episodic memory retrieval and had subjects make the

saccades after encoding and before retrieval. The two novel aspects of the present study include,

first, the focus on visual scene memory and, second, having different groups of subjects make

saccades before encoding as well as before recognition.

There are two main findings of the present study. First, when saccades were made *before*

encoding scenes in working memory, performance as measured by regression slopes relating

overall percent correct with direction of handedness increased significantly positively (Fig 3a)

with right-handers (+LQ) performing better than left-handers (-LQ). Regression slopes exhibited

the same trend of increasing magnitude when a d-prime measure of sensitivity was related to raw

LQ scores (Fig 3b), but there was no significant relationship suggesting some effects of response

bias may have been present in the subjects when a measure of overall percent correct accuracy

was used. Regression slopes were mostly positive when percent correct and d-prime were

regressed against absolute LQ, a measure of degree or consistency of handedness, but these tests

also did not reach significance. Performance also did not correlate significantly as a function of

direction of handedness when either the saccade or fixation condition came *after* the scene

working memory task.

The second main result was that there was no apparent benefit of direction or degree of

handedness on making saccades before scene recognition. Recognition performance in the group

of subjects who made saccades after working memory and before recognition increased non-

significantly as a function of both direction and degree of handedness. Recognition performance

as measured by both overall percent correct and d-prime in the group of subjects who fixated

after working memory and before recognition did increase significantly as a function of direction

of handedness, with better performance as LQ increased indicating an advantage for right-

handed subjects. The relationships between recognition performance and degree of handedness

in those who fixated after working memory and before recognition were non-significantly

positive.  Finally, the subjects who made saccades before the working memory task performed

the poorest during a subsequent test of recognition memory suggesting a link between poor

encoding in working memory and poor subsequent long-term recognition memory.

Why would making saccades before encoding scenes in working memory result in better

performance for strongly right-handed (+LQ) but not necessarily for consistent-handed

individuals? There is some neuroimaging evidence suggesting that form-specific perceptual

aspects of scene encoding may be right lateralized in the brain (Brewer et al. 1998; Stevens et al.

2012), with form-abstract and language-specific aspects including the verbalizability of the

visual stimuli lateralized to the left (Golby et al. 2001; Stevens et al. 2012). Our recent source

analysis of working memory delay period EEG implicates right parieto-occipital focus of activity

that covaries positively as a function of performance. Due to limited sample size in that study,

it's not possible to draw conclusions about whether mixed- or left-handers have a more

symmetrical pattern of activity during scene encoding and memory maintenance that correlates

with worse performance. However, a tentative explanation is that strongly right-handed

individuals may have an advantage over mixed- and left-handers in controlling spatial attention

for scene encoding and memory. Right handers perform better than either mixed or left handers

on tasks requiring reproduction from memory of some aspects of geometric visual material,

while left and mixed handers do not differ from one another (Nebes & Briggs 1974). A

hemispheric advantage for directing spatial attention could protect these right-handers from

potentially destabilizing effects of repetitive saccades on visuospatial attention processes needed

to encode the perceptual aspects of scenes. While saccades may be beneficial to

mixed/inconsistent-handers before the retrieval of explicit episodic memories by improving

attentional control – a form of top-down inward control – they may be detrimental to control of

visuospatial attention directed outward to maintain a stable reference frame for encoding details

of complex visual information contained in scenes. Fitting with this idea, there is evidence for

hemispheric specialization at early levels of visual analysis to the right for processing low spatial

frequencies and to the left for processing high spatial frequencies, with supplementary activation

of right inferior parietal lobule reflecting attentional modulation (Peyrin et al. 2004).

A growing number of studies have demonstrated saccade induced benefits specifically on

recognition (Brunye et al. 2009; Lyle & Orsborn 2011; Parker & Dagnall 2007; Parker et al.

2008). In the present study, when bilateral saccades were made before scene recognition but not

before scene working memory, we found a small but not statistically significant increase in

performance relative to simple fixation. Performance was best for consistent right-handers,

which includes those who have an LQ near +100 in Fig 3 and Fig 4 because they indicate they

perform all tasks "always" with their right hand. A finding reported recently is that beneficial

effects of pre-task saccades occur in consistent-handed individuals (Lyle et al. 2012), those who

perform most actions consistently with either their right or their left hand. If this pattern was

present in our data, we might see an inverted U-shaped distribution of data points in Fig 4 with

lower baseline performance for both -LQ (consistent left-handers) and +LQ (consistent right-

handers) relative to mixed-handers. The ends of the inverted U-shaped might then be shifted up

and/or flattened relative to fixation following saccades. Benefits are not seen in Fig 4 for both

left and right consistent-handed individuals. Rather, it appears that only consistent right-handers

are better able to deal with the detrimental effects of bilateral saccades made before scene

working memory (as indicated by the significant slope of the red line in Fig 3a). Consistent right-

handers also have better recognition memory in general because when they fixate before the

recognition test they perform better (as indicated by the significant slope of the grey line in Fig

4a).

The result of the present study represents a novel amendment to the important findings of

previous studies showing better overall memory in inconsistent handers and beneficial effects of

saccades in consistent-handers. Those findings have replicated in more recent studies (Chu et al.

2012; Parker et al. 2017; Prichard & Christman 2017; Propper et al. 2017). Many of the tasks

used to assess episodic and/or autobiographical memories operate in the verbal domain. In the

present study, subjects might default to a familiarity-based and highly-visual implicit "know"

strategy rather than rely on a verbal episodic-like type of remembering. Adopting a familiarity-

based strategy in the present experiments is even more likely given the speeded choices subjects

were required to make within the 2 sec stimulus viewing windows. Eye movement effects have

been demonstrated to be more beneficial for remember/recollective responses or for tasks that

demand recollection for accurate responses rather than familiarity (Lyle et al. 2012; Parker et al.

2008). In addition, many recognition tasks shown to benefit from bilateral saccades rely on word

lists for memoranda (Parker & Dagnall 2007; Propper et al. 2017; Samara et al. 2011), content

that is much different from the novel scene stimuli used in the present study. However, it is also

worth noting that the beneficial effects of saccades do not manifest in implicit memory tasks

even when those tasks involve words (Christman et al. 2003) had participants study a short word

list and then tested them on a word-stem completion task, consisting of an equal mix of new and

previously seen words. The findings showed that none of the eye movement conditions enhanced

word fragment completion performance compared to the fixation condition. This was interpreted

to indicate that saccades do not enhance implicit forms of memory that are likely reflective of

intra-hemispheric activity (Christman et al. 2003).

One recent study of saccade related enhancement is notable for its use of visual-spatial

stimuli and an exploration of whether effects could be obtained when eye movements were made

before encoding. Brunye et al. (Brunye et al. 2009) used an encoding task that lasted a total of

only 120 seconds and comprised a small number of just four stimuli, all aerial maps that were

acquired from satellite pictures. The authors found increased recognition sensitivity and

decreased response times with effects only seen when eye movements preceded episodic

memory retrieval, but not when they preceded encoding. The effects were strongest for strongly

right-handed individuals. On the basis of the HERA model (Nyberg et al. 1996), Brunye et al

theorized that eye movements would not benefit encoding due to unilateral activity, but would be

beneficial with recognition tests demanding a high degree of right and left-hemisphere activity.

The present study has some limitations. One limitation is the few strongly left-handed

subjects in our sample. Although we found that when saccades occurred before the working

memory task the WM percent correct accuracy increased significantly as a function of direction

of laterality, this result could have been driven by the single very low performing left-hander

represented by the red triangle in the lower left corner of Figure 3a. The few left-handed subjects

and large number of strongly-right handed subjects in the sample rules out a traditional analysis

to assess group-level differences. Instead in the present study relative performance differences as

a function of LQ were characterized using regression slopes. Therefore, summaries of results

involving right-handers (+LQ) performing better than left-handers (-LQ) are not intended and

should not be taken to imply there were group-level differences between equal samples of left-

and right-handers. Another limitation is that the working memory task utilized a relatively high

load of 5 scene stimuli per trial and required a large number of scenes to be memorized and later

discriminated in the recognition test. The 200-trial recognition test took over thirteen minutes to

complete and it is possible that if there were any beneficial effects of preceding eye movements

they may have reduced across the lengthy time interval taken for subjects to complete the task.

This increased the overall difficulty of the memory tasks with performance during the working

memory task nowhere near ceiling. Prior studies involving saccade manipulation have used

relatively smaller numbers of items for encoding and have frequently required participants to

view stimuli passively during learning, rather than engage them in working memory. Both the

limited number of items and the nature of the encoding tasks used in previous research require a

lower degree of cognitive processing compared to the high-load working memory and

recognition tasks used here. Both tasks far exceeded the numbers of stimuli used in prior studies

and could have potentially decreased overall encoding and consolidation into long-term memory

given the short 10-minute interval separating working memory and recognition testing. For the

eye tasks, the requirement for subjects to count disc alternations and color changes was to ensure

subjects performed the tasks. However, an important difference to acknowledge is that counting

has not commonly been used in previous studies of saccade effects on memory. While our

sample included 111 participants, there was an unequal sample of strongly left-handed compared

to strongly-right handed subjects making the degree of handedness (i.e., consistency

comparisons) weighted unequally with strongly-right handed subjects. Finally, the study design

relied on between-subjects manipulation of conditions, which tends to increase variance and

decrease statistical power relative to a within-subject design.

700
701    **Conclusion**
702
We found evidence that performance of repetitive horizontal saccades before scene working and

recognition memory affects performance differently depending on the direction but not degree of

handedness. Making repetitive saccades before encoding scenes in working memory has

detrimental effects on both short- and long-term memory, with the most pronounced effects for

those who are not strongly right-handed. Moreover, saccade execution before scene recognition

does not appear to benefit recognition accuracy appreciably. Making repetitive saccades before

scene encoding may be detrimental because the eye movements destabilize subsequent

visuospatial processing. If working and long-term memory systems interact, then scenes that are

poorly encoded initially will not transfer to long-term memory. Right handers may have an

advantage in dealing with such disruption due to more lateralized mechanisms for scene

processing. These tentative explanations require confirmation by future experiments combining

both behavior and brain measurements.

**Acknowledgements**
717

718    The authors thank Iffat Noor and Chelsea Reichert for assistance with data collection. The

719    content is solely the responsibility of the authors and does not necessarily represent the official

720    views of the National Institutes of Health.

721
722
723    **Figure Legends**
724
**Figure 1. Example Scene Working and Recognition Memory Trials.** Each working memory

trial (a) consisted of an encoding phase of five scene stimuli, each presented for 2 sec, followed

by a 6 sec delay period of crosshair fixation and a 2 sec presentation of either a positive probe

(scene from the encoding set) or a negative probe (new scene). On each trial, there was a 50%

chance of a positive probe appearing. Each recognition memory trial (b) consisted of alternating

presentation of a scene stimuli for 2 sec followed by a black screen for 2 sec. Old and new scenes

were randomly intermixed. Subjects were instructed to press a green button if they had seen a

stimulus in any of the previous working memory trials and a red button if they had never seen the

stimulus before. In this example, image numbers from the set of scene stimuli are listed rather

than the actual scene images to illustrate how scenes during encoding could be presented as a

positive probe (a) or as old stimuli in the recognition task (b).

**Figure 2. The Saccade and Fixation Eye Tasks.** The saccade task (top left) required subjects to

alternate for 30 seconds looking toward the left and right as a white disc moved back and forth

across the screen. Example eye traces (top right) show saccadic movements with large periodic

deviations in the x position (horizontal) trace with a stable y position (vertical) trace. The

fixation task (bottom left) required subjects to maintain fixation on a center disc as it changed

color. Example eye-tracking traces (bottom right) show fixation, with minimal deviations in the

x (horizontal) and y position (vertical) traces.

**Figure 3. Scene Working Memory Performance as a Function of Laterality and Eye Task**

**Timing.** There was a significant relationship in working memory percent correct performance as

a function of direction of laterality when saccades were made before the working memory task

(non-zero slope of $0.1029 \pm 0.04365$ SE, $F_{(1,30)}= 5.56$, p=0.0251, red line and red triangles, panel

a). Overall mean performance among conditions differed (non-zero intercepts, $F_{(3,106)}=6.164$,

p=0.0007) with lowest working memory percent correct performance obtained when saccades

were made before the working memory task ($77.79\% \pm 3.84$, 95% C.I.: 69.95 to 85.64%) and

best performance when saccades were made after the working memory task ($93.28\% \pm 1.844$ SE,

95% C.I.: 89.45 to 97.12). Panel b shows the relationship of the sensitivity measured d-prime

with direction of laterality, while panels c and d show relationships between percent correct

accuracy and d-prime respectively with degree of laterality as expressed by the absolute value of

laterality quotients.

**Figure 4. Scene Recognition Memory Performance as a Function of Laterality and Eye**

**Task Timing.** There was a significant relationship in recognition percent correct performance as

a function of direction of laterality during the condition of fixation before the recognition

memory task (non-zero slope=0.08738 ± 0.03277 SE, $F_{(1,21)}$=7.111, p=0.0144, grey line and grey

boxes, panel a). Overall mean performance among the conditions differed ($F_{(3,106)}$=4.721,

p=0.0039) with best recognition memory performance during the condition of fixation before the

working memory task (70.08% ± 3.899 SE, 95% C.I.: 62.12 to 78.03, blue line and blue circles,

panel a) and lowest recognition memory performance during the condition when saccades were

made before the working memory task (60.67% ± 4.06, 95% C.I.: 52.38 to 68.96, red line and

red triangles, panel a). Panel b shows the relationship of the sensitivity measured d-prime with

direction of laterality, while panels c and d show relationships between percent correct accuracy

and d-prime respectively with degree of laterality as expressed by the absolute value of laterality

quotients.

771
772
773

774 **References**
775
Baddeley A. 2003. Working memory: looking back and looking forward. *Nature reviews*
*neuroscience* 4:829.

Brewer JB, Zhao Z, Desmond JE, Glover GH, and Gabrieli JD. 1998. Making memories: brain
activity that predicts how well visual experience will be remembered. *Science* 281:1185-
1187.

Brewin CR. 2014. Episodic memory, perceptual memory, and their interaction: foundations for a
theory of posttraumatic stress disorder. *Psychol Bull* 140:69-97. 10.1037/a0033722

Brown AL, and Scott MS. 1971. Recognition memory for pictures in preschool children. *Journal*
*of Experimental Child Psychology* 11:401-412.

Brunye TT, Mahoney CR, Augustyn JS, and Taylor HA. 2009. Horizontal saccadic eye
movements enhance the retrieval of landmark shape and location information. *Brain*
*Cogn* 70:279-288. 10.1016/j.bandc.2009.03.003

Burgess N, Maguire EA, and O'Keefe J. 2002. The human hippocampus and spatial and episodic
memory. *Neuron* 35:625-641.

Christman S, and Butler M. 2005. Bilateral eye movements impair the encoding and enhance the
retrieval of episodic memories. 46th annual meeting of the Psychonomic Society,
Toronto, Ontario.

Christman SD, Garvey KJ, Propper RE, and Phaneuf KA. 2003. Bilateral eye movements
enhance the retrieval of episodic memories. *Neuropsychology* 17:221-229.

Christman SD, and Propper RE. 2001. Superior episodic memory is associated with
interhemispheric processing. *Neuropsychology* 15:607.

Christman SD, Propper RE, and Dion A. 2004. Increased interhemispheric interaction is
associated with decreased false memories in a verbal converging semantic associates
paradigm. *Brain Cogn* 56:313-319. 10.1016/j.bandc.2004.08.005

Chu O, Abeare CA, and Bondy MA. 2012. Inconsistent vs consistent right-handers' performance
on an episodic memory task: evidence from the California Verbal Learning Test.
*Laterality* 17:306-317. 10.1080/1357650X.2011.568490

Cowan N. 1999. An embedded-processes model of working memory. *Models of working*
*memory: Mechanisms of active maintenance and executive control* 20:506.

Dragovic M. 2004. Towards an improved measure of the Edinburgh Handedness Inventory: A
one-factor congeneric measurement model using confirmatory factor analysis. *Laterality:*
*Asymmetries of Body, Brain and Cognition* 9:411-419.

Edlin JM, Leppanen ML, Fain RJ, Hackländer RP, Hanaver-Torrez SD, and Lyle KB. 2015. On
the use (and misuse?) of the Edinburgh Handedness Inventory. *Brain and cognition*
94:44-51.

Edlin JM, and Lyle KB. 2013. The effect of repetitive saccade execution on the attention
network test: enhancing executive function with a flick of the eyes. *Brain Cogn* 81:345-
351. 10.1016/j.bandc.2012.12.006

Ellmore TM, Feng A, Ng K, Dewan L, and Root JC. 2015. The Effects of Changing Attention
and Context in an Awake Offline Processing Period on Visual Long-Term Memory.
*Front Psychol* 6:1902. 10.3389/fpsyg.2015.01902

Ellmore TM, Ng K, and Reichert CP. 2017. Early and late components of EEG delay activity
correlate differently with scene working memory performance. *PLoS One* 12:e0186072.
10.1371/journal.pone.0186072

Eriksen A, Lorås H, Vorland Pedersen A, and Sigmundsson H. 2018. *Proximal–Distal Motor*
*Control in Skilled Drummers: The Effect on Tapping Frequency of Mechanically*
*Constraining Joints of the Arms in Skilled Drummers and Unskilled Controls*.

Fan J, Gu X, Guise KG, Liu X, Fossella J, Wang H, and Posner MI. 2009. Testing the behavioral
interaction and integration of attentional networks. *Brain Cogn* 70:209-220.
10.1016/j.bandc.2009.02.002

Golby AJ, Poldrack RA, Brewer JB, Spencer D, Desmond JE, Aron AP, and Gabrieli JD. 2001.
Material-specific lateralization in the medial temporal lobe and prefrontal cortex during
memory encoding. *Brain* 124:1841-1854.

Holcombe A, Nguyen E, and Goodbourn P. 2017. *Implied Reading Direction and Prioritization*
*of Letter Encoding*.

Kielar A, Deschamps T, Chu R, Jokel R, Khatamian Y, Chen JJ, and Meltzer J. 2016. *Identifying*
*Dysfunctional Cortex: Dissociable Effects of Stroke and Aging on Resting State*
*Dynamics in MEG and fMRI*.

Lyle KB, and Edlin JM. 2015. Why does saccade execution increase episodic memory retrieval?
A test of the top-down attentional control hypothesis. *Memory* 23:187-202.
10.1080/09658211.2013.877487

Lyle KB, Hanaver-Torrez SD, Hacklander RP, and Edlin JM. 2012. Consistency of handedness,
regardless of direction, predicts baseline memory accuracy and potential for memory
enhancement. *J Exp Psychol Learn Mem Cogn* 38:187-193. 10.1037/a0024831

Lyle KB, and Jacobs NE. 2010. Is saccade-induced retrieval enhancement a potential means of
improving eyewitness evidence? *Memory* 18:581-594. 10.1080/09658211.2010.493891

Lyle KB, Logan JM, and Roediger HL, 3rd. 2008a. Eye movements enhance memory for
individuals who are strongly right-handed and harm it for individuals who are not.
*Psychon Bull Rev* 15:515-520.

Lyle KB, and Martin JM. 2010. Bilateral saccades increase intrahemispheric processing but not
interhemispheric interaction: Implications for saccade-induced retrieval enhancement.
*Brain Cogn* 73:128-134. 10.1016/j.bandc.2010.04.004

Lyle KB, McCabe DP, and Roediger HL. 2008b. Handedness is related to memory via
hemispheric interaction: evidence from paired associate recall and source memory tasks.
*Neuropsychology* 22:523-530. 10.1037/0894-4105.22.4.523

Lyle KB, and Orsborn AE. 2011. Inconsistent handedness and saccade execution benefit face
memory without affecting interhemispheric interaction. *Memory* 19:613-624.
10.1080/09658211.2011.595418

Nebes RD, and Briggs GG. 1974. Handedness and the retention of visual material. *Cortex*
10:209-214.

Nieuwenhuis S, Elzinga BM, Ras PH, Berends F, Duijs P, Samara Z, and Slagter HA. 2013.
Bilateral saccadic eye movements and tactile stimulation, but not auditory stimulation,
enhance memory retrieval. *Brain Cogn* 81:52-56. 10.1016/j.bandc.2012.10.003

Nyberg L, Cabeza R, and Tulving E. 1996. PET studies of encoding and retrieval: The HERA
model. *Psychon Bull Rev* 3:135-148. 10.3758/BF03212412

Oldfield RC. 1971. The assessment and analysis of handedness: the Edinburgh inventory.
*Neuropsychologia* 9:97-113.

Parker A, and Dagnall N. 2007. Effects of bilateral eye movements on gist based false
recognition in the DRM paradigm. *Brain Cogn* 63:221-225. 10.1016/j.bandc.2006.08.005

Parker A, and Dagnall N. 2010. Effects of handedness and saccadic bilateral eye movements on
components of autobiographical recollection. *Brain Cogn* 73:93-101.
10.1016/j.bandc.2010.03.005

Parker A, Parkin A, and Dagnall N. 2017. Effects of handedness & saccadic bilateral eye
movements on the specificity of past autobiographical memory & episodic future
thinking. *Brain Cogn* 114:40-51. 10.1016/j.bandc.2017.03.006

Parker A, Relph S, and Dagnall N. 2008. Effects of bilateral eye movements on the retrieval of
item, associative, and contextual information. *Neuropsychology* 22:136-145.
10.1037/0894-4105.22.1.136

Peyrin C, Baciu M, Segebarth C, and Marendaz C. 2004. Cerebral regions and hemispheric
specialization for processing spatial frequencies during natural scene recognition. An
event-related fMRI study. *Neuroimage* 23:698-707. 10.1016/j.neuroimage.2004.06.020

Prichard EC, and Christman SD. 2017. Inconsistent-handed advantage in episodic memory
extends to paragraph-level materials. *Memory* 25:1063-1071.
10.1080/09658211.2016.1257725

Propper RE, and Christman SD. 2008. Interhemispheric Interaction and Saccadic Horizontal Eye
MovementsImplications for Episodic Memory, EMDR, and PTSD. *Journal of EMDR*
*Practice and Research* 2:269-281.

Propper RE, Christman SD, and Phaneuf KA. 2005. A mixed-handed advantage in episodic
memory: a possible role of interhemispheric interaction. *Mem Cognit* 33:751-757.

Propper RE, Patel N, Christman SD, and Carlei C. 2017. Superior episodic memory in
inconsistent-handers: a replication and extension using fNIRS. *Memory* 25:1390-1395.
10.1080/09658211.2017.1309051

Propper RE, Pierce J, Geisler MW, Christman SD, and Bellorado N. 2007. Effect of bilateral eye
movements on frontal interhemispheric gamma EEG coherence: implications for EMDR
therapy. *J Nerv Ment Dis* 195:785-788. 10.1097/NMD.0b013e318142cf73

Rice M, and M. Gosser S. 2015. *Efficiency of unimanual and bimanual reach in persons with*
*and without stroke*.

Samara Z, Elzinga BM, Slagter HA, and Nieuwenhuis S. 2011. Do Horizontal Saccadic Eye
Movements Increase Interhemispheric Coherence? Investigation of a Hypothesized
Neural Mechanism Underlying EMDR. *Front Psychiatry* 2:4. 10.3389/fpsyt.2011.00004

Shepard RN. 1967. Recognition memory for words, sentences, and pictures. *Journal of verbal*
*Learning and verbal Behavior* 6:156-163.

Standing L. 1973. Learning 10000 pictures. *The Quarterly journal of experimental psychology*
25:207-222.

Standing L, Conezio J, and Haber RN. 1970. Perception and memory for pictures: Single-trial
learning of 2500 visual stimuli. *Psychonomic Science* 19:73-74.

Stanislaw H, and Todorov N. 1999. Calculation of signal detection theory measures. *Behav Res*
*Methods Instrum Comput* 31:137-149.

Sternberg S. 1966. High-speed scanning in human memory. *Science* 153:652-654.

Stevens WD, Kahn I, Wig GS, and Schacter DL. 2012. Hemispheric asymmetry of visual scene
processing in the human brain: evidence from repetition priming and intrinsic activity.
*Cereb Cortex* 22:1935-1949. 10.1093/cercor/bhr273

Stickgold R. 2002. EMDR: a putative neurobiological mechanism of action. *J Clin Psychol*
58:61-75.

Williams S. 2010. A major revision of the Edinburgh Handedness Inventory. Colchester, Essex,
United Kingdom, https://doi.org/10.13140/RG.2.2.33298.25284.

Williams SM. 1986. Factor analysis of the Edinburgh Handedness Inventory. *Cortex: A Journal*
*Devoted to the Study of the Nervous System and Behavior* 22:325-326.

Xiao J, Hays J, Ehinger K, Oliva A, and Torralba A. 2010. SUN database: Large-scale scene
recognition from abbey to zoo. 2010 IEEE Conference on Computer Vision and Pattern
Recognition (CVPR). San Francisco, CA: IEEE. p 3485-3492.
917