Potential impacts of transport infrastructure and traffic on bird
conservation in Swedish Special Protection Areas

Jan Olof Helldin

Affiliation during the study: Calluna AB, Linköpings slott, 58228 Linköping, Sweden

Current affiliation: Swedish Biodiversity Centre, SLU, PO Box 7016, 75007 Uppsala,

Sweden

Email: j-o.helldin@slu.se

Tel: +46 70 6075322

Abstract

The ecological impacts of roads and railways extend into the surrounding landscape, leading

to habitat degradation and reduced animal densities within an area that is considerably larger

than the actual road or railway corridor. For birds, an extensive meta-analysis has pointed at

an average of 20% density reduction within 1 km from the infrastructure. I investigated to

what extent this density reduction could potentially compromise the habitat quality and

conservation value of Swedish Natura 2000 areas designated for the protection of birds

(Special Protection Areas; SPAs). A majority (63%) of Swedish SPAs are at least to some

extent found within this 1 km potential effect zone. The total overlap between SPA and effect

zone is 126,000 ha, or 4.2% of the country's SPA area. There are however large differences

among biogeographical regions. In the southern (continental) and coastal regions combined,

25.8% of the total SPA area fall within the effect zone, representing an estimated 4-7%

reduction in bird abundance within SPAs. The probability of overlap with effect zone is

higher for larger SPAs. However, the proportion of overlap is higher for smaller SPAs, and

accordingly smaller sites can be assumed to experience a greater impact from transport

infrastructure and traffic. The impacts on Natura 2000 sites are particularly concerning as this

network of protected areas is a cornerstone to maintain and restore biodiversity within EU. I

recommend putting a stronger emphasis in the management of Natura 2000 sites on the

potential threats to wildlife conservation caused by transport infrastructure and traffic. Special

attention should be paid in sites with a large overlap with the effect zone, and in sites hosting

particularly vulnerable taxa or habitats. Infrastructure owners and managers should do their

best to minimize and compensate for the negative impacts of roads and railways and related

traffic in SPAs and other protected areas.

Keywords

Bird conservation, Effect zone, Natura 2000, Railways, Roads, Special Protection Areas,

Sweden, Transport infrastructure

Introduction

Ecological impact of transport infrastructures

Infrastructure development is recognized as one of the significant drivers of the global

biodiversity loss, and with increasing traffic and expanding infrastructure networks

worldwide the pressure on biodiversity is expected to increase in the nearest decades (EEA

2011, 2012, OECD 2012). The impacts of transport infrastructures on wildlife are well

described (reviewed by, e.g., Forman et al. 2003, van der Ree et al. 2015), and include loss of

habitat, traffic casualties, creation of physical barriers, disturbance by noise, light and other

visual cues, spread of chemicals, dust and alien species, changes in hydrology and

microclimate, and accidental spill. Most of these impacts extend into the surrounding

landscape, leading to a degradation and fragmentation of habitats, and for some animal

species to restricted movements, increased mortality, and avoidance of a zone around the

infrastructure (Forman et al. 2003, EEA 2011, van der Ree et al. 2015).

Due to these impacts, the population densities of many animal species are reduced within a

distance from larger infrastructures (Rytwinski & Fahrig 2015). For example, the population

densities may be reduced on up to about 1 km distance for birds (Forman & Deblinger 2000,

Forman et al. 2002, Benítez-López et al. 2010) and anurans (Eigenbrod et al. 2009), and up to

5 km for mammals (Benítez-López et al. 2010). Not only large infrastructures but also minor

and unpaved roads may have a considerable impact on some wildlife species (e.g., van

Langevelde & Jaarsma 2009, Benítez-López et al. 2010, Shanley & Pyare 2011). Based on an

extensive meta-analysis, Benítez-López et al. (2010) showed that the mean bird and mammal

abundance in an effect zone around infrastructure is reduced by 20-30%, and with an

increasing reduction with proximity to the infrastructure (see Fig. 1).

Accordingly, in regions with dense infrastructure networks, large natural areas may be

situated within this effect zone and therefore impoverished in species sensitive to traffic and

transport infrastructures. For example, in the United States, the road effect zone covers 15-

22% of the total land area, and more than 60% of some particularly exposed biomes, such as

coastal regions and river basins (Forman 2000, Riiters & Wickham 2003). In Spain, a country

with intermediate road density with European standards, reduced bird densities due to

transportation infrastructure are expected in 49% of the country, and reduced mammal densities in as much as 96% of the country (Torres et al. 2016). Some habitats of particular importance to biodiversity in Europe, such as wetlands, semi-natural grasslands and temperate broad-leaved forest, may be disproportionately affected by roads because of the landscape structure (Helldin et al. 2013, Karlson & Mörtberg 2015, Torres et al. 2016). Disturbance (noise and visual cues) tend to be stronger and extend further from the road in open habitats as compared to forest (Forman & Deblinger 2000, Reijnen & Foppen 2006). There is a growing concern globally about impact of roads and traffic on wildlife populations in protected areas and other biodiversity hotspots (Forman & Deblinger 2000, Ament et al. 2008, Selva et al. 2011, Laurence & Balmford 2013, Bager et al. 2015, Gadd 2015, Seshadri & Ganesh 2015).

Integrity of the Natura 2000 network

The Natura 2000 network of protected areas is a key tool in the maintenance and restoration of biodiversity within the European Union (EU). The network consists of Special Protection Areas (SPAs) designated according to the Birds Directive, and Special Areas of Conservation (SACs) designated according to the Habitats Directive (EEC 1992, 2010). Under these directives, the network is supposed to provide protection for vulnerable wildlife and habitats. One important motivation for designating Natura 2000 sites, particularly in coastal and other lower-elevation areas, is the protection against negative impacts of urbanization and infrastructure development (EEA 2012). There are presently more than 26,400 sites in the Natura 2000 network, accounting for *ca* 18 % of EUs land territory.

The network is however biased toward highland areas, and lowland areas are underrepresented (Oldfield et al. 2004, Mariorano et al. 2007). In addition, the average size of the Natura 2000 sites is quite low, and particularly so in lowlands (Maiorano et al. 2007, Gaston et al. 2008, EEA 2012). As smaller sites are more susceptible to pressure from land use and human activities surrounding them, major concerns are expressed about the capacity of existing protected areas to maintain their biodiversity values (Chape et al. 2005, Gaston et al. 2008, Maiorano et al. 2008, Kati et al. 2015).

Though many assessments of the Natura 2000 network's effectiveness in protecting species have been reported in the last years, a vast majority of these relied on rather static population data, such as species' geographic distribution, species presence/absence, or habitat suitability (e.g., Araujo et al. 2007, Maiorano et al. 2007, 2015, Sánchez-Fernández et al. 2008, Iojă et al 2010, López-López et al. 2011, Gruber et al. 2012, Albuquerque et al. 2013, D'Amen et al.

2013, Lison et al. 2013, Rubio-Salcedo et al. 2013, Trochet & Schmeller 2013), and

accordingly were designed to assess ecological representativeness of the network rather than

tracking changes in population densities due to environmental impacts. In view of the many

negative population trajectories reported for both avian and non-avian species in the EU (EEA

2015), it appears necessary to analyze the ecological functionality of the Natura 2000 network

with regard to pressures both within and outside the designated areas, but few studies have

taken this course.

Frequency of transport infrastructure within Natura 2000 sites was investigated by Tsiafouli et

al. (2013), showing that as a European average, roads are present in 29% of the sites, with a

higher frequency in the countries in the south. A preceding study of Greek Natura 2000 sites,

by Votsi et al. (2012), showed that 85% of sites are bisected by roads. Insufficient functional

connectivity of the Natura 2000 network has been reported by Gurrutxaga et al. (2011) and

Opermanis et al. (2012), suggesting that dispersal barriers exist between sites, for example in

the form of large roads (Gurrutxaga et al. 2011). With regard to disturbance, the European

Environmental Agency estimated that almost 20% of Natura 2000 areas are presently

adversely affected by high levels of environmental noise, largely owing to major transport

infrastructures (EEA 2016). In an assessment of the impact on Natura 2000 sites of major

traffic arteries planned or under construction as part of EU's TEN-T framework, Byron &

Arnold (2008) estimated that 379 SPAs (8% of sites) and 953 SCIs (4% of sites) would be

adversely affected by these new traffic arteries, with potential effects also on Natura 2000

network coherence.

Aim of the study

Sweden is one of the European countries that are least fragmented by transport infrastructures

and built-up areas (EEA 2011), and the physical impact of infrastructure is generally not well

acknowledged in Swedish nature conservation. It is not until recently that national status

reports for biodiversity have described the threats from infrastructure and traffic on species

conservation (Bernes 2011, Naturvårdsverket 2015a), and the current national conservation

action plan contains few requirements to infrastructure managers (Naturvårdsverket 2015b).

Biodiversity is generally insufficiently described in impact assessments of transportation

infrastructure plans and projects (Wärnbeck 2013, Karlson et al. 2014). Few, if any,

management plans for protected areas address the full array of potential ecological impacts of

transport infrastructures on the areas' conservation status or management (Helldin & Tytor

2017). This ignorance is not unique for Sweden, but appear to be largely similar in most EU

member states (Tsiafouli et al. 2013, EEA 2015; but see, e.g., Selva et al. 2011, Votsi et al.

2012).

In order to illustrate and highlight the impacts of infrastructure on protected areas in

particular, I estimated the frequency and proportion of Swedish SPAs situated within the

potential effect zone for birds around existing larger transport infrastructures (roads and

railways), and hence can be expected not to reach their full conservation potential due to

infrastructure impacts. I included only SPAs in the estimation, i.e., areas designated

specifically for the protection of birds, because the effects of roads and railways on birds are

well described in literature and apparently can impact a majority of bird species (Reijnen &

Foppen 2006, Benítez-López et al. 2010, Rytwinski & Fahrig 2015). I assessed the

conservation value of SPAs that is lost due to infrastructure in terms of reduced predicted bird

abundance. Because of the large geographic variation over the country in density of the

infrastructure network and proportion of area within SPA, I separated the analyses between

biogeographical regions. I tentatively explored the association between the infrastructure

impact on an individual SPA and its size and dominating habitats. In this paper, I present the

results of these analyses and propose improvements for the management of Swedish SPAs.

Methods

A map of the effect zone for birds around the existing larger Swedish transport infrastructure

was produced. I assumed a standardized effect distance of 1 km from the infrastructure,

following the result from a meta-analysis presented by Benítez-López et al. (2010). To my

knowledge, this is the most comprehensive analysis of infrastructure impacts on bird

densities, including 49 bird datasets and 201 bird species. Most studies in the meta-analysis

were conducted in biomes that occur in Sweden, i.e., taiga, temperate broadleaf forest or

alpine/tundra (39 of the 49 datasets), and on species that occur in Sweden (105 of the 201

species), and I therefore judged the results relevant in a Swedish perspective. The results have

also previously been applied to assess the impacts of the road network on birds in Sweden

(Karlson & Mörtberg 2015) and Europe (Torres et al. 2016). I consider the assumption of a 1

km effect distance to be conservative because i) individual studies in the meta-analysis

indicated reduced bird populations on larger distances, and ii) impacts on longer distances

may not necessarily result in direct population declines, but yet be of ecological significance.

Infrastructure data were obtained from Open Street Map (http://openstreetmapdata.com),

using only the following road classes: primary road, secondary road, tertiary road, motorway,

trunk road, railway (thus excluding minor roads for which ecological effects are less well

known).

The effect zone map was overlaid with all Swedish SPAs to calculate the area and proportion

of each site situated within the effect zone. Shape of areas and habitat distribution were

obtained from European Environment Agency's Natura 2000 database

(http://www.eea.europa.eu/data-and-maps/data/natura-2000-eunis-database). SPAs were

separated depending on biome (based on a combination of the Natura 2000 database and

Global Biomes data from the CIESIN; http://sedac.ciesin.columbia.edu/data/set/nagdc-

population-landscape-climate-estimates-v3/maps?facets=theme:climate) and proximity to

coast (data on coastline obtained from Open Street Map; http://openstreetmapdata.com) on the

following terms:

• *continental region:* sites with >50% of the area within EU continental region,

• *mixed-forest region:* sites with >50% of the area within EU boreal region and in CIESIN

broadleaf and mixed-forest region,

• *boreal region:* rest of sites within EU boreal region but with no part within EU alpine

region, or

• *alpine region:* sites with at least some part of the area within EU alpine region;

in combination with

• *coastal:* mainland sites with at least some part within 20 km from coast of mainland

Sweden (including mainland islands Öland and Gotland), or

• *inland:* the rest of mainland sites.

The alpine region in Sweden is only inland, i.e., no alpine coastal sites exist. Because only

two continental sites are inland, all continental sites were pooled in one region. In addition, an

off-coastal region was formed including all sites with no contact with mainland Sweden

irrespective of terrestrial biome. Hereby a total of seven biogeographical regions were

obtained (see Table 1 and Fig. 2).

To assess the bird conservation value of SPAs that is lost due to infrastructure, I assumed an

average of 20% (C.I. 12-33%) reduction of bird abundance within the 1 km effect distance

from infrastructure, as indicated by the results presented by Benítez-López et al. (2010; see

Fig. 1). GIS analyses were conducted using ArcGIS version 10.2 and QGIS version 2.6.

To explore how the degree of impact on an SPA is associated with its size and dominating

habitat, two different analyses where performed for each region. For the probability of

overlap with effect zone (response variable either 0 or 1), a generalized linear model with a

logit link function (logistic regression) was used, and for the proportion of overlap with the
effect zone (i.e., a response variable between 0 and 1), a beta-regression model (Ferrari &
Cribari-Neto 2004) was used. In both types of models explanatory variables were 1) SPA size,
2) proportion forest habitat, 3) proportion wetlands, and 4) proportion agricultural land and
grasslands. SPA size was log-transformed to improve normality and all variables were
standardized to make parameter estimates comparable. Model selection was based on AIC
and the final models were the ones with a combination of explanatory variables resulting in
the lowest AIC. Statistical analyses were conducted using program R version 3.4.2.

Results
The overlay of SPAs and effect zone of larger transport infrastructures showed that 339 of
Sweden´s 538 SPAs (63%) have at least some part, and 123 (23%) have most of their area,
within the effect zone (Table 1). In terms of area, a total of ca 126,000 ha, or 4.2% of the total
SPA area in the country, lies within the effect zone.

National level figures on impacted area however gives a crude picture, as the results pointed
at large differences among the biogeographical regions (Table 1). Alpine and marine areas
have a number of large SPAs and hold most of Sweden´s total SPA area, but have sparse
networks of large (terrestrial) transport infrastructures, and accordingly they reduce the
national average. Continental, mixed-forest coastal and boreal coastal regions, however, are
comparably more impacted; in these three regions combined a total of 46,046 ha, or 25.8% of
the total SPA area, fall within the effect zone. In terms of bird conservation value, present
infrastructure impacts are estimated to cause a 4-7% reduction in bird abundance in SPAs in
the three most impacted regions (continental, mixed-forest coastal and boreal coastal), and an
average of 1% reduction when all SPAs in the country are taken in consideration (Table 1).

The overlap models with the lowest AIC included SPA size or habitat composition in all
regions except the continental (Table 2). As the most emergent pattern, larger SPAs appear to
have a higher probability of overlap with effect zone (in three regions) but a lower proportion
of overlap with effect zone (in all regions except continental). Another pattern, however less
emergent, is that the overlap with effect zone is larger in SPAs with more agricultural land
and grasslands (in four regions), and smaller in SPAs with more forest habitat (in two
regions).

Discussion

Discussion

The results indicate that a significant proportion of Swedish SPAs, both in terms of area and

number of sites, lies within a potential effect zone for birds around present larger transport

infrastructures and therefore can be assumed not to reach their full potential as bird habitat.

The reduction due to transport infrastructure impacts may not be dramatic when seen on the

country as a whole, with only around 4% of the total SPA area affected, corresponding to a ca

1% reduction in predicted bird abundance within SPAs. However, for more urbanized parts of

the country, with a denser infrastructure network, the predicted impact and reduction is nearly

an order of magnitude larger and may well be one of the main factors determining bird

abundance in protected areas. This is the case in the southern (continental) and coastal regions

of the country, where the urbanization and landscape fragmentation is in level with that of

most western and central European countries (EEA 2011).

At the level of individual SPAs, smaller sites tend to have a higher proportion of overlap with

the effect zone and accordingly can be assumed to experience a greater impact than larger

sites. This pattern emerge in all biogeographical regions, except the continental where most

SPAs are small and indeed impacted to a large degree. In effect, the larger impact on smaller

sites amplifies the bias against area protection in the lowlands, i.e., the southern and coastal

regions. This is in line with general concerns previously expressed about the small size of

many protected areas in Europe and about the impacts from transport infrastructure, traffic

and other urban development in the landscape surrounding them (Shafer 1995, Gaston et al.

2008, Maiorano et al. 2008, Kati et al. 2015). However, as indicated by Helldin & Tytor

(2017), these concerns are not well expressed in the management plans for Swedish SPAs,

and appear particularly underestimated in the regions where the impacts are the largest.

The function of predicted bird abundance and distance to infrastructure described by Benítez-

López et al. (2010) have previously been applied twice to assess the ecological impact of an

infrastructure network on a larger geographical level. Karlson & Mörtberg (2015) presented

an assessment of the impacts of roads on habitats of high diversity value in Sweden

(irrespective of their protection status), concluding that natural grasslands and southern

broadleaved forest are likely to be particularly impacted; on national level, 13-19% and 16-

24% of the total areas of these habitats are found within predicted effect zone for birds. Torres

et al. (2016) estimated a 19.0% (CI = 9.6–25.6%) reduction in national bird numbers due to

transport infrastructure for Spain, with all land in view and not only protected areas. They too

concluded that some habitats (most notably farmland and maritime wetlands) might be

disproportionally affected by transport infrastructure. In relation to these previous

assessments, the present study is unique in that it points out the impacts specifically on

protected areas, i.e., areas where nature conservation should be a top priority.

The assessment was aimed to give a general picture, and was therefore simplified in several

respects. A fixed-width effect zone is less realistic, as the actual effect depend on the local

context, such as the habitat distribution, topography, species and ecological processes

involved (Forman & Dubliner 2000, Ritter's & Wickham 2003, Bilging & Duping-Giroux

2006) or the road characteristics (e.g., Reined & Fop pen 2006, Rytwinski & Fahrig 2015).

Also the stronger reduction in bird densities near the infrastructure within the effect zone (Fig.

1) provides an opportunity to assess the decline in bird abundance within the effect zone in

individual sites in greater detail than here conducted (e.g., Torres et al. 2016).

Furthermore, the analysis of impacts of different habitats within SPAs was rather coarse,

since the Natura 2000 database does not provide habitat maps. Therefore, I could not explore

to what degree EU priority habitats (habitat types of community interest; EEC 1992) are

distributed disproportionally within the effect zone.

Implications for management of protected areas

The present study underlines the concern about the impact of transport infrastructures on

wildlife in protected areas in general (Forman & Deblinger 2000, Ament et al. 2008, Selva et

al. 2011, Tsiafouli et al. 2013), and in the smaller areas in particular (Shafer 1995, Maiorano

et al. 2008). Following article 4 of the Birds Directive, EU Member States must take

appropriate steps to avoid habitat deterioration and significant species disturbance within

SPAs (EEC 2010). Accordingly, a stronger emphasis on keeping natural areas free from the

impacts of heavy traffic and new roads and railways has been proposed for conservation and

transport planning (Selva et al. 2011, 2015, Laurance & Balmford 2013, IENE 2015).

Management plans for Natura 2000 sites should better acknowledge the threats to wildlife

conservation caused by both present transport infrastructure and new development projects in

and near sites (Cortina & Boggia 2014, Helldin & Tytor 2017). Assessments of the

effectiveness of individual Natura 2000 sites in maintaining biodiversity should include

monitoring of population density and demography of species of special conservation concern

(Gaston et al. 2006). Managers of Natura 2000 sites have the opportunity to conduct more

detailed assessments of predicted impacts from transport infrastructure based on habitat maps,

species occurrences, and local road characteristics, to serve as a basis for priorities in

conservation planning and action. Special attention to road effects should thus be paid in

protected areas with a large overlap with the effect zone, in areas hosting particularly

vulnerable taxa, and in areas with pronounced impacts on EU priority habitats.

Conservation authorities should secure that infrastructure owners and managers do their best

to minimize the negative impact of nearby roads and railways and related traffic. Technical

mitigation of impacts of transport infrastructure on birds could include preventing bird—

vehicle collision (e.g., with flight diverters), planning the timing of infrastructure maintenance

and construction work to avoid particularly sensitive periods, providing crossing structures

(safe over-/underpasses), and reducing noise and visual impacts through walls, berms or

adapted paving. Traffic calming, speed reduction or road closing (permanent or temporary)

would also provide reductions in road mortality, disturbance and barrier effects. Finally,

compensatory measures such as habitat improvement or additional area protection could

reinforce remnant populations and restore vital ecological processes.

Acknowledgements

I am grateful to Vadym Sokol for assisting in GIS data retrieval and GIS analyses, and to

Victor Johansson for assisting in the statistical analyses. I am also grateful to Lars Nilsson for

providing helpful comments on earlier drafts of this paper. I thank Ana Benítez-López and

coworkers for kind permission to reproduce their published figure. The study was financed by

the Swedish Transport Administration as part of the research program TRIEKOL (triekol.se).

References

Albuquerque FS, Assunção-Albuquerque MJT, Cayuela L, Zamora R, Benito BM (2013)
European Bird distribution is ''well'' represented by Special Protected Areas: Mission
accomplished? Biological Conservation 159: 45–50.

Ament R, Clevenger AP, Yu O, Hardy A (2008) An Assessment of Road Impacts on Wildlife
Populations in U.S. National Parks. Environmental Management 42: 480–496.

Araújo MB, Lobo JM, Moreno JC (2007) The effectiveness of Iberian protected areas in
conserving terrestrial biodiversity. Conservation Biology 21: 1423–1432.

Bager A, Borghi CE, Secco H (2015) The influence of economics, politics and environment
on road ecology in South America. Pp 407-413 in: van der Ree R, Smith DJ, Grilo C
(Eds.). Handbook of Road Ecology. John Wiley & Sons Ltd, UK.

Benítez-López A, Alkemade R, Verweij PA (2010) The impacts of roads and other
infrastructure on mammal and bird populations: A meta-analysis. Biological
Conservation 143: 1307–1316.

Bernes C (2011) Biodiversity in Sweden. Monitor 22, Swedish Environmental Protection
Agency, Stockholm.

Biglin K, Dupigny-Giroux L-A (2006) Mapping the road-effect zone to assess impacts of
proposed road segments. Journal of Conservation Planning 2: 1-16.

Byron H, Arnold L (2008) TEN-T and Natura 2000: the way forward. An assessment of the
potential impact of the TEN-T Priority Projects on Natura 2000. Report from BirdLife
International. http://www.birdlife.org/eu/pdfs/TEN_T_report2008_final.pdf

Chape S, Harrison J, Spalding M, Lysenko I (2005) Measuring the extent and effectiveness of
protected areas as an indicator for meeting global biodiversity targets. Phil. Trans. R. Soc.
B 360: 443-455.

Cortina C, Boggia A (2014) Development of policies for Natura 2000 sites: A multi-criteria
approach to support decision makers. Journal of Environmental Management 141: 138-
145.

D'Amen M, Bombi P, Campanaro A, Zapponi L, Bologna MA, Mason F (2013) Protected
areas and insect conservation: questioning the effectiveness of Natura 2000 network for
saproxylic beetles in Italy. Animal Conservation 16(4): 370-378.

EEA (2011) Landscape fragmentation in Europe; Joint EEA-FOEN report. EEA report No
2/2011. European Environment Agency, Copenhagen.

EEA (2012) Protected areas in Europe – an overview. EEA report No 5/2012. European
Environment Agency, Copenhagen.

EEA (2015) State of nature in the EU; Results from reporting under the nature directives
2007–2012. EEA report No 2/2015. European Environment Agency, Copenhagen.

EEA (2016) Quiet areas in Europe; The environment unaffected by noise pollution. EEA
report No 14/2016. European Environment Agency, Copenhagen.

EEC (1992) Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural
habitats and of wild fauna and flora. Official Journal of the European Communities No L
206/7-50.

EEC (2010) Directive 2009/147/EC of the European Parliament and of the Council of 30
November 2009 on the conservation of wild birds. Official Journal of the European
Union No L 20/7-25.

Eigenbrod F, Hecnar SJ, Fahrig L (2009) Quantifying the Road-Effect Zone: Threshold
Effects of a Motorway on Anuran Populations in Ontario, Canada. Ecology and Society
14(1): 24.

Ferrari S, Cribari-Neto F (2004) Beta Regression for Modelling Rates and Proportions.
Journal of Applied Statistics 31(7): 799-815.

Forman RTT (2000) Estimate of the area affected ecologically by the road system in the
United States. Conservation Biology 14(1): 31-35.

Forman RTT, Deblinger RD (2000) The ecological road-effect zone of a Massachusetts
(U.S.A.) suburban highway. Conservation Biology 14: 36-46.

Forman RTT, Reineking B, Hersperger AM (2002) Road traffic and nearby grassland bird
patterns in a suburbanizing landscape. Environmental Management 29(6): 782-800.

Forman RTT, Sperling D, Bissonette JA, Clevenger AP, Cutshall CD, Dale VH, Fahrig L,
France R, Goldman CR, Haenue K, Jones JA, Swanson FJ, Turrentine T, Winter TC
(2003) Road ecology – Science and solutions. Island Press (Washington, USA).

Gadd ME (2015) Expected effects of a road across the Serengeti. Pg 455-464 in: van der Ree
R, Smith DJ, Grilo C (Eds). Handbook of Road Ecology. John Wiley & Sons Ltd, UK.

Gaston KJ, Jackson SF, Nagy A, Cantú-Salazar L, Johnson M (2008) Protected Areas in
Europe; Principle and Practice. Ann. N.Y. Acad. Sci. 1134: 97–119.

Gruber B, Evans D, Henle K, Bauch B, Schmeller D, Dziock F, Henry PY, Lengyel S,
Margules C, Dormann C (2012) "Mind the gap!" – How well does Natura 2000 cover
species of European interest? Nature Conservation 3: 45–62.

Gurrutxaga M, Rubio L, Saurac S (2011) Key connectors in protected forest area networks
and the impact of highways: A transnational case study from the Cantabrian Range to the
Western Alps (SW Europe). Landscape and Urban Planning 101: 310–320.

Helldin J-O, Tytor S (2017) Hur uppmärksammas ekologiska effekter av vägar och järnvägar
inom naturvården? – innehållsanalys av bevarandeplaner för Natura 2000-områden.
Report from Calluna AB.
http://www.calluna.se/userfiles/files/5_Innehållsanalys_SPA_bevarandeplaner_2017(1).p
df (In Swedish)

Helldin J-O, Collinder P, Bengtsson D, Karlberg Å, Askling J (2013) Assessment of traffic
noise impact in important bird sites in Sweden – a practical method for the regional scale.
Oecologia Australis 17(1): 70-84.

Iojă CI, Pâtroescu M, Rozylowicz L, Popescu VD, Verghelet M¸ Zotta MI, Felciuc M (2010)
The efficacy of Romania's protected areas network in conserving biodiversity. Biological
Conservation 143: 2468–2476.

Karlson M, Mörtberg U (2015) A spatial ecological assessment of fragmentation and
disturbance effects of the Swedish road network. Landscape and Urban Planning 134:
53–65.

Karlson M, Mörtberg U, Balfors B (2014) Road ecology in environmental impact assessment.
Environmental Impact Assessment Review 48: 10–19.

Kati V, Hovardas T, Dieterich M, Ibisch PL, Mihok B, Selva N (2015) The challenge of
implementing the European network of protected areas Natura 2000. Conservation
Biology 29(1): 260–270.

Laurance WF, Balmford A (2013) A global map for road building. Nature 495: 308-309.

Lison F, Palazon JA, Calvo JF (2013) Effectiveness of the Natura 2000 Network for the
conservation of cave-dwelling bats in a Mediterranean region. Animal Conservation
16(5): 528-537.

López-López P, Maiorano L, Falcucci A, Barba E, Boitani L (2011) Hotspots of species
richness, threat and endemism for terrestrial vertebrates in SW Europe. Acta Oecologica
37(5): 399-412.

Maiorano L, Falcucci A, Garton EO, Boitani L (2007) Contribution of the Natura 2000
Network to Biodiversity Conservation in Italy. Conservation Biology 21(6): 1433–1444.

Maiorano L, Falcucci A, Boitani L (2008) Size-dependent resistance of protected areas to
land-use change. Proc. R. Soc. B 275: 1297–1304.

Naturvårdsverket (2015a) Målövergripande analys av miljömålen; Underlag till fördjupad
utvärdering av miljömålen 2015. Swedish Environmental Protection Agency, Report No
6692. (In Swedish)

Naturvårdsverket (2015b) Styr med sikte på miljömålen – Naturvårdsverkets fördjupade utvärdering av miljömålen 2015. Swedish Environmental Protection Agency, Report No 6666. (In Swedish)

OECD (2012) OECD Environmental Outlook to 2050; The Consequences of Inaction. OECD Publishing, Paris. http://dx.doi.org/10.1787/9789264122246-en

Oldfield TEE, Smith RJ, Harrop SR, Leader-Williams N (2004) A gap analysis of terrestrial protected areas in England and its implications for conservation policy. Biological Conservation 120: 307–313.

Opermanis O, MacSharry B, Aunins A, Sipkova Z (2012) Connectedness and connectivity of the Natura 2000 network of protected areas across country borders in the European Union. Biological Conservation 153: 227-238.

Reijnen R, Foppen R (2006) Impact of road traffic on breeding bird populations. Pp. 255-274 in: Davenport, Davenport J (Eds) The Ecology of Transportation: Managing Mobility for the Environment. Springer, Dordrecht, the Netherlands.

Riitters KH, Wickham JD (2003) How far to the nearest road? Front. Ecol. Environ. 1(3): 125–129.

Rubio-Salcedo M, Martínez I, Carreño F, Escudero A (2013) Poor Effectiveness of the Natura 2000 Network Protecting Mediterranean Lichen Species. Journal for Nature Conservation 21(1): 1–9.

Rytwinski T, Fahrig L (2015) The Impacts of Roads and Traffic on Terrestrial Animal Populations. Pp 237-246 in: van der Ree R, Smith DJ, Grilo C (Eds.). Handbook of Road Ecology. John Wiley & Sons Ltd, UK.

Sánchez-Fernández D, Bilton DT, Abellán P, Ribera I, Velasco J, Millán A (2008) Are the endemic water beetles of the Iberian Peninsula and the Balearic Islands effectively protected? Biological Conservation 141(6): 1612-1627.

Selva N, Kreft S, Kati V, Schluck M, Jonsson B-G, Mihok B, Okarma H, Ibisch PL (2011) Roadless and Low-Traffic Areas as Conservation Targets in Europe. Environmental Management 48: 865–877.

Seshradi KS, Ganesh T (2015) Road ecology in south India: issues and mitigation opportunities. Pg 425-429 in: van der Ree R, Smith DJ, Grilo C (Eds). Handbook of Road Ecology. John Wiley & Sons Ltd, UK.

Shanley CS, Pyare S (2011) Evaluating the road-effect zone on wildlife distribution in a rural landscape. Ecosphere 2(2): 16.

Shafer CL (1995) Values and Shortcomings of Small Reserves: Dealing with the smallest habitat fragments when some of them are all that is left. BioScience 45(2): 80–88.

Torres A, Jaeger JAG, Alonsoa JC (2016) Assessing large-scale wildlife responses to human infrastructure development. PNAS 113(30): 8472-8477. https://doi.org/10.1073/pnas.1522488113

Trochet A, Schmeller DS (2013) Effectiveness of the Natura 2000 network to cover threatened species. Nature Conservation 4: 35–53.

Tsiafouli MA, Apostolopoulou E, Mazaris AD, Kallimanis AS, Drakou EG, Pantis JD (2013) Human Activities in Natura 2000 Sites: A Highly Diversified Conservation Network. Environmental Management 51: 1025–1033.

van der Ree R, Smith DJ, Grilo C (2015). Handbook of Road Ecology. John Wiley & Sons
Ltd, UK.

van Langevelde F, Jaarsma CF (2009) Modeling the effect of traffic calming on local animal
population persistence. Ecology and Society 14(2): 39.

Votsi N-E, Mazaris AD, Kallimanis AS, Zomeni MS, Vogiatzakis IN, Sgardelis SP, Pantis JD
(2012) Road effects on habitat richness of the Greek Natura 2000 network. Nature
Conservation 1: 53–71.

Wärnbäck A (2013). Beaktande av biologisk mångfald och ekosystemtjänster i
miljökonsekvensbeskrivningar och miljöbedömningar. Annex 5 to SOU 2013:68,
Synliggöra värdet av ekosystemtjänster – Åtgärder för välfärd genom biologisk mångfald
och ekosystemtjänster.
http://www.regeringen.se/49bba7/contentassets/ba53cd9f18b74f348eb0ff31e8280d60/syn
liggora-vardet-av-ekosystemtjanster-sou-201368 (In Swedish)

Tables

Table 1. Number, area and proportion of Swedish SPAs within an assumed effect zone of 1
km from larger transport infrastructure, and predicted total reduction in bird abundance (with
95% confidence interval) due to the effects. Results are given for the entire country and
separated by biogeographical region.

Region	Total no. of SPAs	Total area of SPA (ha)	No. of SPAs affected to		Total area of SPA in effect zone (ha)	Proportion of total SPA in effect zone (%)	Reduction in bird abundance (% with C.I.)
			>0%	>50%			
Continental	41	53,331	41	18	19,202	36.0	7.2 (4.3-11.9)
Mixed-forest coastal	94	103,064	71	24	20,865	20.2	4.0 (2.4-6.7)
Mixed-forest inland	161	258,804	119	43	34,622	13.4	2.7 (1.6-4.4)
Boreal coastal	22	21,712	15	8	5,968	27.5	5.5 (3.3-9.1)
Boreal inland	130	162,924	63	30	13,122	8.1	1.6 (1.0-2.7)
Alpine (only inland)	26	1,984,005	16	0	28,242	1.4	0.3 (0.2-0.5)
Off-coastal	64	415,308	14	0	3,913	1.0	0.2 (0.1-0.3)
All Sweden	*538*	*2,999,149*	*339*	*123*	*125,946*	*4.2*	*0.8 (0.5-1.4)*

Table 2. Variables associated with effect zone overlap with Swedish SPAs divided by biogeographical region. Values given are mean estimates of coefficients of logistic regressions and beta-regressions with standard errors (SE) and probabilities (P). Values are only given for variables that were included in the final model.

Region	Log(size) Estimate (SE)	P	Forested area Estimate (SE)	P	Wetland Estimate (SE)	P	Agri & grassland Estimate (SE)	P
Logistic regression (probability of overlap)								
Continental †	-	-	-	-	-	-	-	-
Mixed-forest coastal	0.45(0.27)	0.094						
Mixed-forest inland					0.49(0.21)	0.019	0.91(0.38)	0.016
Boreal coastal			-1.06(0.57)	0.065				
Boreal inland			-2.61(0.75)	<0.001	-2.06(0.70)	<0.001		
Alpine (only inland)	1.31(0.64)	0.039						
Off-coastal	1.07(0.40)	0.008						
Beta-regression (proportion of overlap)								
Continental ‡								
Mixed-forest coastal	-0.56(0.15)	<0.001					0.44(0.15)	0.002
Mixed-forest inland	-0.64(0.12)	<0.001					0.47(0.11)	<0.001
Boreal coastal	-0.85(0.34)	0.013						
Boreal inland	-0.56(0.16)	<0.001	-0.48(0.16)	0.004			0.43(0.16)	0.009
Alpine (only inland)	-0.78(0.11)	<0.001			-0.38(0.17)	0.022		
Off-coastal	-1.09(0.24)	<0.001					0.40(0.19)	0.036

† Could not be analyzed as all sites had value 1, i.e., overlap to some degree with the effect zone.

‡ No variable contributed to the final model; i.e., the model including only the intercept had the lowest AIC.

Figures

Figure 1. Mean species abundance (MSA) of birds as a function of distance to infrastructure (logistic regression). Open dots represent the pooled results of a meta-analysis per distance interval ±SE. Solid black line denotes the estimated curve for the decline of MSA in proximity to infrastructure; dashed lines are the 95% upper and lower limits of the confidence bands of the curve. Figure from Benítez-López et al. (2010).

Figure 2. Special Protection Areas (SPAs) in Sweden divided by biogeographical region; see text for further explanation.