DoubletFinder: Doublet detection in single-cell RNA sequencing data using artificial nearest neighbors

Christopher S. McGinnis[1], Lyndsay M. Murrow[1] and Zev J. Gartner[1,2,3,4]*
[1]Department of Pharmaceutical Chemistry, University of California, San Francisco
[2]Chan Zuckerbeg Biohub, University of California, San Francisco
[3]Center for Cellular Construction, University of California, San Francisco, San Francisco, CA 94158, USA
[4]Lead Contact
*Correspondence: zev.gartner@ucsf.edu

SUMMARY

Single-cell RNA sequencing (scRNA-seq) using droplet microfluidics occasionally produces transcriptome data representing more than one cell. These technical artifacts are caused by cell doublets formed during cell capture and occur at a frequency proportional to the total number of sequenced cells. The presence of doublets can lead to spurious biological conclusions, which justifies the practice of sequencing fewer cells to limit doublet formation rates. Here, we present a computational doublet detection tool – DoubletFinder – that identifies doublets based solely on gene expression features. DoubletFinder infers the putative gene expression profile of real doublets by generating artificial doublets from existing scRNA-seq data. Neighborhood detection in gene expression space then identifies sequenced cells with increased probability of being doublets based on their proximity to artificial doublets. DoubletFinder robustly identifies doublets across scRNA-seq datasets with variable numbers of cells and sequencing depth, and predicts false-negative and false-positive doublets defined using conventional barcoding approaches. We anticipate that DoubletFinder will aid in scRNA-seq data analysis and will increase the throughput and accuracy of scRNA-seq experiments.

INTRODUCTION

Since its introduction nearly a decade ago, scRNA-seq has been used to elucidate previously unknown cell types and reconstruct developmental dynamics among heterogeneous cell populations (Human Cell Atlas Consortium, 2017). At first, scRNA-seq workflows were

limited to tens to hundreds of cells which hindered data interpretation due to batch effects and

low statistical power (Stegle et al., 2016). Today, sequencing thousands to hundreds of

thousands of cells is routine due to the advent of droplet microfluidics and nanowell-based

sequencing strategies (Macosko et al., 2015; Klein et al., 2015; Zheng et al., 2017; Gierahn et

al., 2017; Takara Bio USA, 2018). These techniques rely on a Poisson loading strategy to

compartmentalize individual cells and mRNA capture beads before cell lysis, mRNA capture,

and transcript barcoding via reverse transcription. Since cells are captured randomly, the

proportion of droplets containing >1 cell – known as doublets – scales linearly across an

experimentally-relevant range of input cell concentrations (10X Genomics, 2017), justifying the

practice of limiting the number of sequenced cells to minimize doublet formation rates.

The confounding effects of doublets in scRNA-seq data are well-appreciated (Ilicic et al.,

2016). However, genomic and cellular barcoding techniques for identifying doublets have only

recently been developed (Stoeckius et al., 2017; Kang et al., 2018; Gehring et al., 2018; Guo et

al., 2018; Rosenberg et al., 2018). In one such strategy, distinct samples receive unique

oligonucleotide barcodes delivered by conjugation to antibodies targeting broadly expressed

cell-surface antigens. When the barcoded pools are combined and sequenced, doublets can be

identified according to the co-occurrence of orthogonal cell 'hashtags' (Stoeckius et al., 2017).

In a second strategy, doublets in a pooled population of cells from different individuals are

identified by a computational pipeline, Demuxlet, which facilitates doublet inference based on

the co-occurrence of mutually-exclusive SNP profiles (Kang et al., 2018).

By detecting doublets, both Demuxlet and Cell Hashing minimize technical artifacts while

enabling users to "superload" droplet microfluidics devices for increased scRNA-seq throughput.

However, both methods have limitations. First, neither method can identify doublets formed from

identically-barcoded cells. Second, neither method is universally applicable across experimental

systems, since Demuxlet requires genetically distinct samples and Cell Hashing requires unique antibody-oligonucleotide conjugate panels for the cell types and species of interest. Third, neither method can be used to analyze existing scRNA-seq datasets. For these reasons, computational methods for defining doublets based on gene expression patterns alone are highly desirable.

Here, we present DoubletFinder, a computational doublet detection tool that relies solely on gene expression data. Beginning with the observation that doublets cluster separately from singlets in high-dimensional gene expression space (Stoeckius et al., 2017; Kang et al., 2018), we reasoned that real doublets would cluster together with synthetic doublets formed by averaging the expression data of two real cells. By merging artificial doublets with existing scRNA-seq data, we can distinguish doublets from singlets according to the proportion of artificial nearest neighbors (pANN) for each real cell in gene expression space. Thresholding the resulting pANN distribution to match the expected number of doublets provides an accurate metric for doublet prediction that can be applied to any scRNA-seq dataset.

RESULTS

DoubletFinder predicts doublets more accurately than nUMIs: Existing strategies for identifying doublets using gene expression features primarily rely on two sources of information. First, since the total number of captured mRNA molecules is expected to be greater for doublets than singlets, doublets are commonly excluded by thresholding cells with high numbers of unique molecular identifiers (nUMIs; Islam et al., 2014; Ziegenhain et al., 2017). While intuitively appealing, technical variability in mRNA capture efficiency and biological variability in mRNA content limits the utility of nUMI-based doublet predictions (Stoeckius et al., 2017). In a second strategy, doublets are removed from scRNA-seq data by identifying groups of cells exhibiting

the co-expression of genes with non-overlapping expression patterns *in vivo* (Rosenberg et al.,

2018). This strategy cannot be applied to biological systems where such marker genes are

unknown, undetected, or unavailable. Moreover, such a strategy could theoretically lead to the

erroneous removal of new cell types or developmental states with intermediate expression

profiles (Fig. 1A). Given these shortcomings, new methods for predicting doublets using gene

expression features alone would greatly benefit the single-cell genomics field.

DoubletFinder predicts doublets in a fashion agnostic to nUMIs, marker gene expression,

genetic background or exogenous barcodes and can be split into four distinct steps: (1) Generate

artificial doublets, (2) Merge real and artificial data and reduce dimensionality with principal

component analysis (PCA), (3) Define the nearest neighbors for every real cell in PC space, and

(4) Compute and threshold the proportion of artificial nearest neighbors (pANN; Fig. 1B). We

tested the efficacy of DoubletFinder against scRNA-seq datasets where doublets are empirically-

defined: The publically-available Cell Hashing and Demuxlet datasets comprised of 15,178 and

35,524 peripheral blood mononuclear cells (PBMCs), respectively. Demuxlet PBMCs were

derived from 8 genetically-distinct human sources while Cell Hashing PBMCs were barcoded

with 8 distinct antibody-oligonucleotide conjugate panels. Using optimized input parameters

(Supplementary Materials, Fig. S1), we tested whether pANN outperforms nUMIs as a doublet-

prediction feature by using receiver operating curve (ROC) analysis to compare logistic

regression models trained on the Cell Hashing data using pANN alone, nUMIs alone, or both

(Fig. 1C). ROC analysis demonstrates that pANN predicts doublets more accurately than nUMIs.

Moreover, the model trained with both features performed nearly indistinguishably to the pANN-

alone model, suggesting that DoubletFinder captures all of the doublet-specific information

inherent to nUMIs.

<u>DoubletFinder predicts Cell Hashing doublets</u>: To make specific doublet predictions for each

cell, DoubletFinder rank-orders cells by their pANN values and thresholds this list according to

the number of expected doublets. To test the robustness of pANN thresholding, the number of

expected doublets was determined using two different strategies (Fig. 1D). First, since 8 samples

were multiplexed in the Cell Hashing and Demuxlet studies, we reasoned that 1/8 of the true

doublets were undetected because they were formed from genetically-identical or identically-

barcoded cells. Thus, we thresholded pANN according to the number of detected doublets with

an assumed 12.5% false negative rate (β). Second, since the doublet formation rate can be

accurately estimated by applying Poisson statistics to the number of cells loaded into the droplet

microfluidics device (10X Genomics, 2017), we thresholded pANN according to this rate. For

standard scRNA-seq experiments where doublets are not empirically-defined, pANN can only

be thresholded using the Poisson strategy.

Depending on the threshold used, DoubletFinder predicted 2680 or 2155 doublets when

applied to the full Cell Hashing dataset. Single-cell gene expression data was visualized using

t-stochastic neighborhood embedding (t-SNE; van der Maaten and Hinton, 2008) and cells were

colored according to their real and predicted doublet status (Fig. 1E). Visual comparison of

doublets in t-SNE space illustrates that DoubletFinder predictions closely track Cell Hashing

results. This result is further supported by the observation that the frequency of DoubletFinder

predictions is highly enriched in Cell Hashing-defined doublet groups relative to singlet groups

(Fig. 1F), regardless of the thresholding strategy used.

<u>DoubletFinder predicts Cell Hashing false-negatives</u>: DoubletFinder predictions exhibit a less

'speckled' appearance in t-SNE space relative to the Cell Hashing results (Fig. 2A, insets).

Considering that doublets often cluster separately from singlets in gene expression space

(Stoeckius et al., 2017; Kang et al., 2018), we reasoned that cells called as singlets via Cell

Hashing, that nonetheless co-cluster with high-confidence doublets, are actually false-negatives

derived from identically-barcoded cells. Two main predictions follow from this line of reasoning.

First, if the putative false negatives are truly doublets, then they should exhibit gene expression

patterns associated with distinct cell types. In line with this prediction, Cell Hashing-defined

doublets and singlets in the highlighted region express marker genes for both B cells and NK

cells (Fig. 2B) – hematopoietic cell types that do not share a common progenitor in peripheral

blood. Second, since false negative cells would be associated with the combined barcodes of

two cells, the nUMI counts for the most abundant barcode should be significantly higher in false

negatives than high-confidence doublets. Moreover, since false negatives would not be

associated with high levels of multiple barcodes, the second most abundant barcode should be

similar to high-confidence singlets and significantly lower relative to high-confidence doublets.

Statistical analysis supports these predictions (Wilcoxon rank sum test, $p < 10^{-13}$; Fig. 2C).

Collectively, these results demonstrate that DoubletFinder robustly recapitulates doublet

assignments and accurately predicts Cell Hashing false-negatives.

<u>DoubletFinder predicts Demuxlet doublets and identifies putative false-positives</u>: To test whether

DoubletFinder performance is sensitive to changes in the number of sequenced cells and

sequencing depth, we applied DoubletFinder to the Demuxlet dataset. In addition to having more

cells than the Cell Hashing data, the average number of UMIs (2408 vs 676) and genes (837 vs

376) per cell is also greater in the Demuxlet data. In line with our previous results, visual

comparison of real and predicted doublets using t-SNE illustrates that DoubletFinder

successfully identifies all doublet-enriched regions in gene expression space (Fig. 2D).

As with our Cell Hashing comparison, there were a number of regions in gene expression

space where DoubletFinder predictions differed from Demuxlet classifications. Specifically, there

were many DoubletFinder-defined doublets called as singlets by Demuxlet that give doublet-

enriched clusters the 'speckled' appearance discussed above. Moreover, in contrast to the Cell

Hashing comparison, there was a subset of cells classified as doublets by Demuxlet and singlets

using DoubletFinder (Fig. 2D, insets). Interestingly, the majority of these discordant calls are

scattered amongst high-confidence singlet clusters in gene expression space. This observation

can be explained by two alterative models. In one model, these discordant calls are caused by

homotypic doublets – i.e., doublets formed from cells of the same type – which presumably have

a similar transcriptional profile to singlets and, thus, would be more difficult for DoubletFinder to

detect relative to heterotypic doublets. Alternatively, the discordant calls are due to false-positive

Demuxlet classifications.

If these cells were in fact homotypic doublets left undetected by DoubletFinder, then one

would expect that DoubletFinder was insensitive to homotypic doublets throughout the Demuxlet

dataset. To test this possibility, we tracked the cell types comprising each artificial doublet and

deconvolved pANN values into homotypic and heterotypic components. Visualization of cells

with majority homotypic or heterotypic nearest neighbors highlights a region of homotypic

doublets formed from $CD4^+$ T-cells (Fig. 2E; Supplementary Materials, Fig. S2), which suggests

that DoubletFinder has the sensitivity to detected certain classes of homotypic doublets.

Moreover, the scattering of discordant doublet classifications amongst high-confidence singlet

clusters in gene expression space is also evident in Demuxlet classifications of the Cell Hashing

data (Supplementary Materials, Fig. S2). Cell Hashing is sensitive to homotypic doublets, which

suggests that the discordant calls are not a consequence of homotypic doublets missed by

DoubletFinder. Finally, if the putative false-positive calls are homotypic doublets, one would

expect the number of RNA UMIs to approximate levels observed for high-confidence doublets.

Interestingly, the RNA nUMI distribution for putative false-positives is nearly indistinguishable to

high-confidence singlets (Wilcoxon rank sum test, p = 0.34), while high-confidence doublets and

putative false-negatives are both significantly enriched for RNA nUMIs (Wilcoxon rank sum test,

$p < 10^{-15}$; Fig. 2F). While it is difficult to definitively ascertain the ground truth for these discordant

calls, these results collectively demonstrate that DoubletFinder is robust across a range of cell

numbers and sequencing depths and prospectively predicts Demuxlet false-positives.

DISCUSSION

High-throughput scRNA-seq suffers from the formation of doublets due to the inherent

nature of Poisson cell loading. Doublets can lead to spurious conclusions during analysis when

left unidentified because the resulting artefactual expression data may be interpreted as

previously-undescribed cell types, developmental intermediates, or disease states. As a result,

it has become common practice to minimize the doublet formation rate by minimizing the ratio

of sequenced cells to mRNA capture beads. Although recent advances in direct doublet

detection methodologies have proven to be effective, they are not universally or retroactively

applicable. For this reason, complementary techniques for predicting doublets based only on

gene expression data have the potential to further increase scRNA-seq throughput while

removing technical artifacts.

Towards this goal, DoubletFinder accurately identifies doublets in scRNA-seq data by

integrating artificial doublets into real data and computing the pANN for every real cell. We have

shown that DoubletFinder distinguishes real doublets from singlets better than nUMIs in the Cell

Hashing dataset. Moreover, we demonstrate that DoubletFinder accurately predicts doublets for

two independent PBMC scRNA-seq datasets of different sizes and sequencing depths. As these

are the only publically available data with empirically-defined doublets, it is unclear whether

DoubletFinder will require further optimization for scRNA-seq datasets describing different

tissues or biological systems. We have also shown that DoubletFinder identifies false-negative

and putative false-positive doublet classifications present in these datasets, which supports the

use of DoubletFinder in concert with Cell Hashing, Demuxlet, and other barcoding approaches.

Finally, we demonstrate that DoubletFinder performs robustly with pANN thresholding strategies

that differed by >5000 cells (Fig. 1D). This suggests that DoubletFinder can be applied in

experimental contexts where doublet formation rates differ significantly from industry estimates

– e.g., clumpy single-cell suspensions or especially cohesive cell types. Collectively,

DoubletFinder represents a fast, easy-to-use doublet detection strategy that will aid the single-

cell genomics community in data analysis and enable high-throughput scRNA-seq technologies

to be utilized to their fullest potential.

MATERIALS & METHODS

<u>DoubletFinder Overview</u>: Artificial doublets were generated from raw UMI count matrices via

random sampling of cell expression profiles without replacement before pre-processing using

the 'Seurat' R package, as described previously (Butler et al., 2018). Notably, no sources of

variation were regressed out of the merged data before PCA, and the top 10 PCs – chosen via

inflection point estimation on the corresponding elbow plot – were used to define the Euclidean

distance matrix using the 'dist' R function.

<u>ROC Analysis</u>: ROC analysis-based model comparisons were performed using the 'ROCR'

(Sing et al., 2005) and 'pROC' (Robin et al., 2011) R packages. Briefly, logistic regression

models were defined on a training set comprising half the total data using the 'glm' R function

with the link argument set to 'logit'. These models were then used to create a vector describing

each cell's doublet probability with the 'predict' R function. ROC analysis was then performed by

calculating the sensitivity and specificity of doublet predictions based on the aforementioned

probability vector at varying probability thresholds. The AUC was then calculated for the resulting

curve, and AUC was used as a proxy for doublet detection model performance.

Statistical Analysis: Statistically-significant differences between UMI levels were defined using

the Wilcoxon rank sum test implemented with the 'pairwisewilcox.test' R function. Multiple

comparison correction was performed using the Benjamini-Hochberg procedure.

Data Availability: Cell Hashing (GEO: GSE108313) and Demuxlet (GEO: GSE96583) UMI count

matrices were downloaded from the Gene Expression Omnibus. DoubletFinder is implemented

as a fast, easy-to-use R package that interfaces with Seurat version 2.0 and higher.

DoubletFinder can be downloaded from GitHub (github.com/chris-mcginnis-ucsf/DoubletFinder).

ACKNOWLEDGMENTS

We thank Matt Thomson (California Institute of Technology) for helpful discussion, as well as

Jimmie Ye (UCSF), Marlon Stoeckius and Shiwei Zheng (New York Genome Center) for

providing data access.

AUTHOR CONTRIBUTIONS

C.S.M., L.M.M., and Z.J.G conceptualized the method and wrote the manuscript. C.S.M. wrote

the software and performed all bioinformatics analyses.

DECLARATION OF INTERESTS

The authors declare no conflict of interest.

FUNDING

This research was supported in part by grants from the Department of Defense Breast Cancer Research Program (W81XWH-10-1-1023 and W81XWH-13-1-0221), the NIH Common Fund (DP2 HD080351-01), the NSF (MCB-1330864), and the UCSF Center for Cellular Construction (DBI-1548297), an NSF Science and Technology Center. Z.J.G is a Chan-Zuckerberg Biohub Investigator. L.M.M is a Damon Runyon Fellow supported by the Damon Runyon Cancer Research Foundation (DRG-2239-15).

REFERENCES

1. The HCA Consortium. The Human Cell Atlas White Paper. 2017. Retrieved from: humancellatlas.org/files/HCA_WhitePaper_18Oct2017.pdf
2. Stegle O, Teichmann SA, Marioni JC. Computational and analytical challenges in single-cell transcriptomics. Nat Rev Genet. 2015; 16(3):133-45.
3. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. 2015. Cell; 161(5):1202-1214.
4. Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, et al. Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. 2015. Cell; 161(5):1187-1201.
5. Zheng GX, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel digital transcriptional profiling of single cells. Nat Commun. 2017; 8:14049.
6. Gierahn TM, Wadsworth MH 2nd, Hughes TK, Bryson BD, Butler A, Satija R, *et al*. Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput. Nat Methods. 2017; 14(4):395-398.
7. Takara Bio USA. Full-length, single-cell RNA-seq with the SMARTer™ ICELL8® Single-Cell System (Application note 633878). Mountain View, CA: Takara Bio. 2018. Available from: clontech.com/US/Products/cDNA_Synthesis_and_Library_Construction/NGS_Automation/Single-Cell_Automation/Single-Cell_Automation_Overview?sitex=10020:22372:US
8. 10X Genomics. ChromiumTM Single Cell 3' Reagent Kits v2 User Guide. Pleasanton, CA: 10X Genomics. 2017.

9. Ilicic T, Kim JK, Kolodziejczyk AA, Bagger FO, McCarthy DJ, Marioni JC, Teichmann SA. Classification of low quality cells from single-cell RNA-seq data. Genome Biol. 2016 17:29.

10. Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Smibert P, Satija R. Cell "hashing" with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. 2017. Preprint. bioRxiv doi: 10.1101/237693.

11. Kang HM, Subramaniam M, Targ S, Nguyen M, Maliskova L, McCarthy E. Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. Nat Biotechnol. 2018; 36(1):89-94.

12. Gehring J, Park JH, Chen S, Thomson M, Pachter L. Highly Multiplexed Single-Cell RNA-seq for Defining Cell Population and Transcriptional Spaces. 2018. Preprint: bioRxiv doi: 10.1101/315333**.**

13. Guo C, Biddy BA, Kamimoto K, Kong W, Morris SA. CellTag Indexing: a genetic barcode-based multiplexing tool for single- cell technologies. 2018. Preprint: bioRxiv doi: 10.1101/335547.

14. Rosenberg AB, Roco CM, Muscat RA, Kuchina A, Sample P, Yao Z, et al. Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding. Science. 2018; 360(6385):176-182.

15. Islam S, Zeisel A, Joost S, La Manno G, Zajac P, Kasper M, Lönnerberg P, Linnarsson S. Quantitative single-cell RNA-seq with unique molecular identifiers. Nat Methods. 2014; 11(2):163-6.

16. Ziegenhain C, Vieth B, Parekh S, Reinius B, Guillaumet-Adkins A, Smets M, et al. Comparative Analysis of Single-Cell RNA Sequencing Methods. Mol Cell. 2017. 65(4):631-643.e4.

17. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol. 2018; doi: 10.1038/nbt.4096.

18. van der Maaten L, Hinton G. Visualizing Data using t-SNE. J. Mach. Learn. Res. 2008; 9: 2579–2605.

19. Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier performance in R. Bioinformatics. 2005; 21(20): 3940-1.

20. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Müller M. pROC: an open-source package for R and S+ to analyze and compare ROC curves. BMC Bioinformatics, 2011; 12: 77.

327
328
329
330 **FIGURE LEGENDS**
331

Figure 1: DoubletFinder robustly predicts Cell Hashing doublets and outperforms nUMI
thresholding. (A) Schematic describing importance of doublet detection. Developmental
intermediates (light red) can express genes associated with both progenitor (grey) and
differentiated (dark red) cell types. Doublets formed from progenitor and mature cells may mimic
the expression profile of intermediate cell states, and thereby confound analysis. (B) Schematic
of DoubletFinder workflow. After artificial doublet (red outline) generation, the proportion of
artificial nearest neighbors is defined for every real cell (examples highlighted yellow). These
results are thresholded to define doublet predictions. (C) Density plot of pANN values with red
dotted lines denoting expected doublet thresholds. Histogram is colored according to whether
the cells were called as singlets (grey) or doublets by one (light red) or both thresholding
strategies (dark red). (D) t-SNE visualization of real and predicted Cell Hashing doublets (red)
and singlets (grey), DoubletFinder predictions, and nUMI thresholding predictions. nUMI
predictions deviate significantly from Cell Hashing results (red dashed boxes). (E) Heat map
showing the proportion of DoubletFinder and nUMI-predicted doublets present in Cell Hashing
singlet (S) and doublet (D) groups. (F) ROC analysis of logistic regression models trained using
nUMIs alone (dotted red), pANN alone (solid blue) and both nUMIs and pANN (dashed orange)
as features.

Figure 2: DoubletFinder detects false-negative and false-positive doublet predictions in
Cell Hashing and Demuxlet datasets. (A) t-SNE visualization of real and predicted Cell
Hashing doublets (red) and singlets (grey) highlighting a doublet-enriched, 'speckled' region.
(B) Violin plots describing the distribution of marker gene expression in high-confidence doublets
(red), putative false-negatives (blue), and singlet B-cells (teal) and NK cells (orange). (C)
Barcode UMI box plots for the 1st and 2nd most abundant barcodes in high-confidence singlets
(black) and doublets (red), as well as false-negative singlets (blue). (D) t-SNE visualization of
real and predicted Demuxlet doublets (red) and singlets (grey) highlighting putative false-
positives. (E) Violin plots describing the contributions of homotypic and heterotypic doublets to
each DoubletFinder-defined doublet's nearest neighborhood. t-SNE visualization of putative
homotypic and heterotypic doublet clusters in gene expression space. (F) RNA UMI box plots
for high- confidence singlets (black) and doublets (red) as well as discordant doublet predictions
between Demuxlet and DoubletFinder (orange).