Running head: LISSAMPHIBIAN ORIGIN AND OSSIFICATION SEQUENCES

Title:

What do ossification sequences tell us about the origin of extant amphibians?

Michel Laurin[1],*, Océane Lapauze[1], David Marjanović[2]

[1] *CR2P (Centre de Recherche sur la Paléodiversité et les Paléoenvironments; UMR 7207),*

CNRS/MNHN/UPMC–Sorbonne Universités, Muséum national d'Histoire naturelle,

Département Histoire de la Terre, 57 rue Cuvier, F-75231 Paris cedex 05, France; [2] *Museum*

für Naturkunde (Leibniz Institute for Evolutionary and Biodiversity Research),

Invalidenstraße 43, D-10115 Berlin, Germany, david.marjanovic@gmx.at

*Correspondence to be sent to: Muséum national d'Histoire naturelle, Département Histoire

de la Terre, 57 rue Cuvier, F-75231 Paris cedex 05, France; michel.laurin@mnhn.fr

ABSTRACT—The controversial origin of extant amphibians has been studied using several sources of data and methods, including phylogenetic analyses of morphological data, molecular dating, stratigraphic data, and integration of ossification sequence data, but a consensus has failed to emerge. We have compiled five datasets to assess the relative support for six competing hypotheses about the origin of extant amphibians: a monophyletic origin among temnospondyls, a monophyletic origin among lepospondyls, a diphyletic origin among both temnospondyls and lepospondyls, a diphyletic origin among temnospondyls alone, and two variants of a triphyletic origin, in which anurans and urodeles come from different temnospondyl taxa while caecilians come from lepospondyls and are either closer to anurans and urodeles or to amniotes. Our datasets comprise ossification sequences of up to 107 terminal taxa and up to eight cranial bones, and up to 65 terminal taxa and up to seven appendicular bones, respectively. Among extinct taxa, only two or three temnospondyl can be analyzed simultaneously for cranial data, but this is not an insuperable problem because each of the six tested hypotheses implies a different position of temnospondyls and caecilians relative to other sampled taxa. For appendicular data, more extinct taxa can be analyzed, including some lepospondyls and the finned tetrapodomorph *Eusthenopteron*, in addition to temnospondyls. The data are analyzed through maximum likelihood, and the AICc (corrected Akaike Information Criterion) weights of the six hypotheses allow us to assess their relative support. By an unexpectedly large margin, our analyses of the cranial data support a monophyletic origin among lepospondyls; a monophyletic origin among temnospondyls, the current near-consensus, is a distant second. All other hypotheses are exceedingly unlikely according to our data. Surprisingly, analysis of the appendicular data supports triphyly of extant amphibians within a clade that unites lepospondyls and temnospondyls, contrary to all molecular and recent paleontological phylogenies, but this conclusion is not very robust.

Keywords: macroevolution; paleontology; evo-devo; ossification sequences; Lissamphibia;

Tetrapoda; phylogeny

Introduction

Paleontologists have been studying the origin of the extant amphibian clades for more than a

century. Early studies generally proposed an origin of at least some extant amphibians from

temnospondyls. Cope (1888) initially suggested that batrachians (anurans and urodeles)

derived from temnospondyls (a large clade of limbed vertebrates known from the Early

Carboniferous to the Early Cretaceous) because he believed that the batrachian vertebral

centrum was an intercentrum, the dominant central element of temnospondyls. Later, Watson

(1940) argued that anurans were derived from temnospondyls because of similarities (mostly

in the palate) between the temnospondyl "*Miobatrachus*" (now considered a junior synonym

of *Amphibamus*) and anurans. Monophyly of extant amphibians (Lissamphibia) was proposed

by Parsons and Williams (1962, 1963), an idea that was accepted more quickly by

herpetologists than by paleontologists. Lissamphibian monophyly was supported by (among a

few other character states) the widespread occurrence of pedicellate, bicuspid teeth. The

subsequent discovery of such teeth in the amphibamid temnospondyl *Doleserpeton* (Bolt

1969) reinforced the widespread acceptance of an origin of Lissamphibia from within

temnospondyls (e.g., Schoch and Milner 2004). Recently, this hypothesis, referred to as the

temnospondyl hypothesis or TH for short (Fig. 1c), has been supported by several

phylogenetic analyses based on phenotypic data matrices (e.g. Ruta and Coates 2007;

Sigurdsen and Green 2011; Maddin et al. 2012; Pardo et al. 2017a, b: fig. S6; Mann et al.

2019).

Dissenting opinions about the origin of extant amphibians have been expressed for

several decades (see Schoch and Milner 2004 for a historical review). These were initially

formulated especially for the urodeles and caecilians, which are less similar to temnospondyls

and lack a tympanic middle ear (which is present in most anurans and often inferred for at

least some temnospondyls but absent in lepospondyls). Thus, Steen (1938) highlighted

similarities in the palate (broad cultriform process of the parasphenoid) and cheek (loss of

several bones) between lysorophian lepospondyls and urodeles. Carroll and Currie (1975) and

Carroll and Holmes (1980) argued that the exant amphibians had three distinct origins among

early stegocephalians; while they accepted an origin of anurans among temnospondyls, they

suggested that urodeles and caecilians originated from two distinct groups of lepospondyls

(*Rhynchonkos* for caecilians, Hapsidopareiidae for urodeles). Later, based mostly on

developmental similarities between the temnospondyl *Apateon* and urodeles, Carroll (2001,

2007) and Fröbisch et al. (2007) proposed another hypothesis involving a triphyletic origin of

lissamphibians, with an origin of anurans and urodeles from two distinct temnospondyl

groups, while the caecilians would remain in the lepospondyl clade. This is what we call the

polyphyly hypothesis (PH). We have tested two versions. One (called PH1; Fig. 1e) was

cautiously suggested by Fröbisch et al. (2007); it agrees with the paleontological consensus in

placing all or most lepospondyls closer to Amniota than to Temnospondyli (Fig. 1b;

Sigurdsen and Green 2011; Pardo et al. 2017a, b: fig. S6; Marjanović and Laurin 2019; Clack

et al. 2019; Mann et al. 2019). The other (PH2; Fig. 1f) is modified to make Lissamphibia

monophyletic with respect to Amniota, a fact we consider demonstrated beyond reasonable

doubt by multiple phylogenetic analyses of molecular data (Fig. 1a; Irisarri et al. 2017; Feng

et al. 2017; and references cited therein); this comes at the expense of contradicting the

paleontological consensus, which was not yet established when Milner (1993: 16–18, fig. 5B)

argued for something like the PH2 as one of two more or less equal possibilities. Anderson

(2007) and Anderson et al. (2008) found lissamphibian diphyly, specifically a monophyletic,

exclusive Batrachia among the temnospondyls while keeping the caecilians among the

lepospondyls (DH1; Fig. 1g). Pardo et al. (2017b: fig. 2, S7) presented a similar hypothesis,

with batrachians and caecilians having separate origins within the temnospondyls (DH2; Fig.

1h). Further, a monophyletic origin of all extant amphibians among lepospondyls has also

been proposed (Laurin 1998; Pawley 2006: appendix 16; Marjanović and Laurin 2009, 2013,

2019). This will be referred to below as the lepospondyl hypothesis (LH; Fig. 1d).

Phylogenetic analyses of molecular data cannot distinguish the TH, the PH2, the DH2

or the LH from each other by topology (Fig. 1) because all of these imply lissamphibian

monophyly with respect to amniotes. Several other types of data and methods have, however,

been used to try to discriminate between the various hypotheses on the origin of extant

amphibians. In addition to classical phylogenetic analyses of morphological data matrices,

these include the use of molecular dating (Zhang et al. 2005; Marjanović and Laurin 2007;

Pardo et al. 2017b) and stratigraphic data (Marjanović and Laurin 2008) to compare the

inferred divergence dates between the three main extant amphibian clades on the basis of

molecular data with predictions based on the fossil record under the TH and the LH on the

one side and the PH and the DH on the other. However, developmental data, in the form of

ossification sequences, have been the second-most frequently used (after classical

morphological data) to argue for particular phylogenetic hypotheses. These data include

mainly cranial (e.g. Schoch 2002, 2006; Schoch and Carroll 2003; Schoch and Milner 2004;

Anderson 2007; Carroll 2007; Germain and Laurin 2009) and autopodial ossification

sequences (e.g. Fröbisch et al. 2007, 2015). Ossification sequences of other parts of the

skeleton, like the vertebrae, shoulder girdle and scales, are also documented in a few

Paleozoic stegocephalians (e.g. Carroll et al. 1999; Witzmann and Schoch 2006; Anderson

2007; Carroll 2007; Olori 2013), not to mention finned tetrapodomorphs (Cloutier 2009), but

these have played a minor role in the controversy about the origin of extant amphibians, and

recently, Danto et al. (2019) concluded that vertebral ossification sequences varied too

quickly and could not be used to assess the origin of lissamphibians. This study relies on both

cranial and appendicular ossification sequences and compares their implications for tetrapod

phylogeny.

METHODS

Ossification sequence data

From all the literature we could access, we compiled the most extensive database on

ossification sequences for osteichthyans that exists to date. The most useful sources for extant

taxa included compilations: Harrington et al. (2013) for amphibians, Weisbecker and

Mitgutsch (2010) for anurans, Hugi et al. (2012) for squamates, Maxwell et al. (2010) for

birds, and Koyabu et al. (2014) and Weisbecker (2011) for mammals. The cranial and

appendicular sequences of Permian temnospondyls (the stereospondylomorphs

Sclerocephalus and *Archegosaurus*, the non-branchiosaurid "branchiosaur" *Micromelerpeton*

and the branchiosaurids "*Melanerpeton*" *humbergense*, *Apateon caducus* and *A. pedestris*)

were assembled from several references cited in the Appendix; note that the two *Apateon*

species are each represented by two different sequences scored after populations from two

separate paleo-lakes (Erdesbach and Obermoschel) in which both species occur. Appendicular

ossification sequences of the lepospondyls *Microbrachis* and *Hyloplesion* are incorporated

from Olori (2013), that for the finned tetrapodomorph *Eusthenopteron* was combined from

Cote et al. (2002) and Leblanc and Cloutier (2005).

All sources of our sequence data can be found in the Appendix. The sequences

themselves and the phylogenetic trees corresponding to the tested hypotheses are included in

the supplementary material. The sequences were not used to generate the tree topology or the

branch lengths (which represent evolutionary time); the tree is compiled from published

sources (provided below) which did not use any ossification sequences in their phylogenetic

analyses.

The software we used to compute AICc weights, the CoMET module (Lee et al. 2006)

for Mesquite 3.6 (Maddison and Maddison 2018), cannot handle missing data. This

unfortunately meant we had to discard much information. In order to keep as many taxa as

possible in the analysis, we first compiled a matrix (not shown) of 244 taxa and 213

characters. All of these characters are positions of skeletal elements (cranial, appendicular,

axial and others) in ossification sequences, standardized between 0 and 1 following Germain

and Laurin (2009), as explained below. Of these, we kept characters that were scored in the

Paleozoic taxa in our initial database, and extant taxa that were scored for the same sets of

characters. This resulted in two initial datasets, one of cranial and one of appendicular

sequences (it was not possible to include both sets of sequences together because this would

have left too few taxa in the matrix). In the end, we were left with three overlapping cranial

datasets. Dataset 1 contains 107 taxa (104 extant, *Apateon* spp. from Erdesbach, and

Sclerocephalus) and only six characters. Dataset 2 (see Table 1) has 105 taxa (103 extant, plus

the two species of *Apateon* scored from Erdesbach) and seven characters (nasal, parietal,

squamosal, premaxilla, maxilla, pterygoid, and exoccipital); The third cranial dataset (dataset

5) includes 84 taxa (81 extant, *Apateon* spp. from Erdesbach, and *Sclerocephalus*) and eight

cranial characters (the frontal bone is added). For the appendicular characters, in addition to

dataset 3 which contains seven characters (humerus, radius, ulna, ilium, femur, tibia and

fibula) and 62 taxa (54 extant, *Apateon* spp. from Obermoschel, *Sclerocephalus*,

Archegosaurus, *Micromelerpeton*, *Hyloplesion*, *Microbrachis* and *Eusthenopteron*), another

(dataset 4) includes only four characters (radius, ulna, ilium, and femur), but it features 65

sequences, the additional data being *Apateon* spp. from Erdesbach and "*Melanerpeton*"

humbergense. See Table 1 for a list of these datasets and the SM for the datasets themselves.

The data loss in these various datasets is not as severe as it may first seem, because

most of the characters that have collectively been excluded from these analyses had less than

10% scored cells (sometimes less than 1%), and most of them could not be scored for any

temnospondyl or lepospondyl, so they could not have helped resolve the main question

examined in this study.

The order in which the eight cranial bones ossify varies substantially in our sample of

taxa, but based on simple (not phylogenetically-weighted) average position, the frontal

appears first, followed closely by the premaxilla, parietal, and maxilla (in close succession),

and then by the squamosal, exoccipital, pterygoid, and last by the nasal. However, all of these

bones ossify first (among these seven bones; not necessarily in the whole skeleton) in at least

one of the included taxa. Among the appendicular bones, there is more variability; all ossify

first in at least one of the 62 sampled taxa, and three (radius, ulna and ilium) ossify last in at

least one taxon.

Of the eight cranial characters, *Sclerocephalus* cannot presently be scored for the

squamosal. Because of the potential importance of *Sclerocephalus* as a stem-caecilian

according to the DH2 (Fig. 1h) and as one of only three sampled extinct taxa with any known

cranial ossification sequence, we ran variants of the analyses of cranial data with

Sclerocephalus and six characters (dataset 1), and without *Sclerocephalus* and with seven

characters (dataset 2; see Table 1).

Due to the homology problems between the skull bones of tetrapods and

actinopterygians and missing data, we had to omit all actinopterygians from our analyses. As

cranial ossification sequences remain unknown for extant finned sarcopterygians (except

perhaps lungfish, whose skull bones seem mostly impossible to homologize), our analyses of

those data are restricted to limbed vertebrates. However, for appendicular data, we were able

to include the Devonian tristichopterid *Eusthenopteron foordi*.

Unfortunately, the only cranial ossification sequence available for any supposed

lepospondyl, that of the aïstopod *Phlegethontia longissima*, is documented from only three

ossification stages (Anderson et al. 2003; Anderson 2007). This poses a problem for our

analysis method, which assumes that character evolution can be modeled as Brownian

motion; this assumption is decreasingly realistic as the number of character states (sequence

positions) decreases, because the resulting distribution deviates increasingly from that of a

continuous character. Furthermore, some recent anatomical restudies and phylogenetic

analyses suggest that aïstopods are not lepospondyls, but early-branching stem-

stegocephalians (Pardo et al. 2017a, 2018; Mann et al. 2019; Clack et al. 2019).

The low taxon sample is more limiting for this analysis than the low character sample.

However, as explained below, the absence of lepospondyl sequences in our cranial dataset

does not preclude testing the six hypotheses (TH, PH1, PH2, DH1, DH2, LH; see above or

Figure 1 for the explanation of these abbreviations) because each of these six hypotheses

makes different predictions about where temnospondyls and caecilians fit relative to other

taxa. Thus, in the absence of lepospondyls in our dataset, the tests of these hypotheses are

somewhat indirect and inference-based, but they remain possible. Our tests based on

appendicular data include two lepospondyls (*Hyloplesion longicostatum* and *Microbrachis*

pelikani), but the absence of caecilians in that dataset proves more limiting than the absence

of lepospondyls in the cranial dataset because the TH, DH1 and DH2 become

indistinguishable (Fig. 1 c, g, h). However, the presence of lepospondyls allows us to test two

variants of the TH/DH distinguished by the monophyly (e.g. Ruta and Coates 2007) or

polyphyly (e.g. Schoch 2019) of "branchiosaurs" (the temnospondyls *Apateon*,

"*Melanerpeton*" and *Micromelerpeton*).

Sensitivity analysis for sequence polymorphism

Given the potential impact of infraspecific variability in ossification sequence on inferred

nodal sequences and heterochrony (Olori 2013; Sheil et al. 2014), we compiled two consensus

212 sequences for *Apateon caducus* and *A. pedestris* each, representing two localities where both

213 species occur, the paleo-lakes of Erdesbach (Schoch 2004) and Obermoschel (Werneburg

214 2018). Based on dataset 4 (see Table 1), we incorporated these into a global and two separate

215 analyses (one analysis per locality) to determine the impact of the observed variability.

216 Incorporating the sequences from Erdesbach reduced the number of characters from seven to

217 only four because the software used cannot handle missing data (see below), but this

218 information loss is compensated by the great increase in number of sequences from extinct

219 taxa (eleven instead of two, when counting the sequences of *Apateon* from both localities

220 separately) and the fact that this includes some lepospondyls (see below). It would have been

221 even better to perform a sensitivity analysis incorporating variability for all taxa for which

222 such information was available, but given the scope and nature of our study, this would have

223 been exceedingly time-consuming and is best left for the future.

224 *Standardization of the data*

225 Given that various taxa differ in their numbers of bones and that the resolution of the

226 sequences is also variable between taxa, these data needed to be standardized to make

227 comparisons and computations meaningful, as suggested by Germain and Laurin (2009).

228 Note that we performed this standardization on the complete dataset of characters, before

229 filtering for data completeness. This complete dataset (not shown) includes 213 cranial,

230 appendicular and other characters, but no taxon is scored for all characters, given that the

231 original (complete) matrix has much missing data. For instance, the most completely scored

232 taxon, *Amia calva*, still has 57.4% missing data (more than half), which indicates that 92

233 characters were scored for this taxon, including several ties (the resolution was 41 positions,

234 so they varied by increments of 0.025 or 2.5% of the recorded ontogeny). We did not re-

235 standardize after filtering characters out because we believe that the initial standardization

236 better reflects the relative position of events in development than a standardization based on

only seven events in ontogeny would. Because of this, in the reduced dataset of seven

characters used in the calculations, for some taxa, no character has a score of 0 or 1 because

the first or last events in the ontogenetic sequence were filtered out. Thus, we used the

position in the sequence (from first to last, in the complete dataset) and standardized this

relative sequence position between 0 and 1 using the simple formula given by Germain and

Laurin (2009). The standardized sequence position (X_s) is:

$X_s = (X_i - X_{min})/(X_{max} - X_{min})$,

where:

X_i is the position of a given bone in the sequence

X_{min} is the lowest position in the sequence (generally denoted 0 or 1)

X_{max} is the highest position in the sequence (for instance, if there are 20 bones, X_{min} is 1 and

the sequence is completely resolved, $X_{max} = 20$).

This yields a standardized scale that varies between 0 and 1 for each taxon, in which 0 and 1

are the positions of the first and last events in the sequence, respectively. For instance, for

Ambystoma maculatum (an extant urodele), in the original dataset, the first events (tied) were

the ossification of premaxilla, vomer, dentary and coronoid (standardized position: 0); the last

event was the articular (standardized position: 1), and there is a resolution of 12 positions

(hence, increments of 0.0909 or 1/11). However, in the final dataset of 7 charcters, the

articular is absent; hence, the first bone in the sequence is the premaxilla, at a standardized

position of 0, and the last is the nasal, as a standardized position of 0.8181 because all events

in position 1 (articular) and 0.9091 (stapes) have been filtered out.

We also experimented with using size (skull length) or developmental stage as standards, but

this led to lower sequence resolution because body size is not available for all sequence

positions and for all taxa (results not shown), so we worked only with sequences standardized

using sequence position. Given that our data filtering procedure retains little data (only six,

seven or eight characters for the cranial dataset, and four or seven characters for the

postcranial dataset), it is important to use the method that discards the least amount of data,

and this was achieved by using sequence position. We do not imply that standardizing by size

is not recommended in general. On the contrary, if good body size data were available for all

taxa and all developmental stages, this should be a better strategy, and only having access to

absolute time should be even better. However, practical limitations of data availability prevent

us from using these methods now.

Our ossification sequence data (reduced dataset of four to eight characters) of extant

and extinct taxa, and the phylogenetic trees we used, are available in the supplement to this

paper.

Analysis methods

To discriminate between the six hypotheses about the origin of extant amphibians, two

methods are available: direct phylogenetic analysis of the sequence data, and comparisons of

the tree length (number of steps in regular parsimony, squared length in squared-change

parsimony, likelihood, or similar measures) of various trees selected a priori to represent these

hypotheses (in these trees, only the position of caecilians and extinct taxa, here temnopondyls

and lepospondyls, varies). We used both approaches but expected the second to perform much

better because relatively few data are available, and thus, phylogenetic analysis of such data is

unlikely to provide a well-resolved tree.

For the first approach, we first transformed the standardized sequence positions back

into discrete characters using formulae in a spreadsheet and scaled the characters so that the

highest state in all would be 9. This ensures that each character has an equal weight in the

analysis, regardless of its variability in the ossification sequence. The characters were ordered

to reflect the assumed evolutionary model (ontogenetic timing is a quantitative character that

was discretized) and because for such characters, ordering yields better results (Rineau et al.

2015, 2017; see discussion in Marjanović & Laurin 2019). The resulting data matrices (one

for cranial and another for appendicular characters, both with seven characters each) were

analysed using parsimony in PAUP* 4.0a165 (Swofford 2019). We used the TBR (tree

bisection-reconnection) branch swapping algorithm and performed a search with 50 random

addition replicates (or several such searches, for the cranial data) while holding two trees at

each step and with a maximal number of trees set at one million. For cranial data, the main

search lasted about 100 hours on a MacBook Pro Retina with a 2.5 GHz iCore 7 quadri-core

processor and 16 GB RAM. The exact search time cannot be reported because PAUP*

crashed after saving the trees to a file for one of the longest runs (several analyses were made,

over several days), but before the log could be saved. The analysis of the seven appendicular

characters was much faster (27 minutes and a half), presumably because that matrix has fewer

taxa (62 instead of 105).

For the second approach (comparison of fit of various trees selected a priori to reflect

previously published hypotheses), we used the CoMET module (Lee et al. 2006) for Mesquite

3.6 (Maddison and Maddison 2018) to test the relative fit of the data on trees representing the

six hypotheses. CoMET calculates the likelihood and the AIC (Akaike Information Criterion)

of nine evolutionary models given continuous data and a tree. Note that our data only

represent an approximation of continuous data; if standardization had been performed on

developmental time or body size, the data would actually have been continuous.

Standardization was carried out using sequence position because of data limitation problems,

so the data actually follow a decimalized meristic scale. However, the difference between

these situations decreases as the number of sequence positions increases, and our global scale

includes up to 41 positions (and an average of 10.9 positions), so our data should approximate

a continuous distribution sufficiently well for our analyses to be valid. This consideration

prevents us from adding the highly apomorphic aïstopod *Phlegethontia*, for which only three

cranial ossification stages are known (Anderson et al. 2003; Anderson 2007); moreover, five

of the seven bones included in our analyses appear in the last two of these stages, and two of

the relevant bones (parietal and exoccipital) are not present as separate ossifications, which

would create additional missing data. In that case, the very low number of stages would create

strong departures from the assumption of continuous data. This would probably create

statistical artifacts, and the uncertainty about the position of *Phlegethontia* (Pardo et al.

2017a, 2018; Marjanović and Laurin 2019; Clack et al. 2019) would complicate interpretation

of the results.

The nine models evaluated by CoMET are obtained by modifying the branch lengths

of the reference tree. Thus, branches can be set to 0 (for internal branches only, to yield a non-

phylogenetic model), to 1 (equal or speciational model), left unchanged from their original

length (gradual evolution), or set free and evaluated from the data (free model). This can be

applied to internal and/or external branches, and various combinations of these yield nine

models (Lee et al. 2006: fig. 1). Among these nine models two have been frequently discussed

in the literature and are especially relevant: gradual evolution, in which branch lengths (here

representing evolutionary time) have not been changed, and a speciational model, in which all

branches are set to the same length, and which has some similarities with Eldredge and

Gould's (1972) punctuated equilibria model (though a model with one internal branch

stemming from each node set to 0 and the other set to 1 would be even closer to the original

formulation of that model). In this study, we assessed the fit of six of the nine models covered

by CoMET; the other three (the punctuated versions of distance [original branch length],

equal and free) in which the one of each pair of daughter-lineages has a branch length of zero,

could not be assessed due to problems in the current version of CoMET and possibly the size

of our dataset.

Provided that the same evolutionary model is optimal for all compared phylogenetic

hypotheses (this condition is met, as shown below), the AIC weights of the various trees

under that model can be used to assess the support for each tree. In such comparisons, the

topology is part of the evolutionary model, and the data are the sequences. These comparisons

can show not only which tree is best supported, but how many times more probable the best

tree is compared to the alternatives. This quantification is another reason to prefer this

approach over a phylogenetic analysis (performed below, but with the poor results that we

anticipated), which can at best yield a set of trees showing where the extinct taxa most

parsimoniously fit (if we had dozens of characters, this might be feasible). Comparisons with

other hypotheses through direct phylogenetic analysis are not possible. Given the small

sample size (which here is the number of characters), we computed the corrected AIC (AICc)

and the AICc weights using the formulae given by Anderson and Burnham (2002) and

Wagenmakers and Farrell (2004).

Our tests make sense only in the presence of a phylogenetic signal in the data. In

addition to the test of evolutionary model in CoMET evoked above (which tests non-

phylogenetic as well as phylogenetic models), we performed a test based on squared-change

parsimony (Maddison 1991) and random taxon reshuffling (Laurin 2004). For this test, we

compared the length of the LH (lepospondyl hypothesis; Fig. 1d) reference tree (with and

without *Sclerocephalus*) to a population of 10,000 random trees produced by taxon

reshuffling.

It could be argued that using other methods (in addition to the method outlined above)

would have facilitated comparisons with previous studies. However, the two main alternative

methods, event-pair cracking with Parsimov (Jeffery et al. 2005) and Parsimov-based genetic

inference (PGI; Harrison and Larsson 2008), have drawbacks that decided us against using

them. Our objections against event-pair cracking with Parsimov were detailed by Germain

and Laurin (2009) but can be summarized briefly as including the unnecessary decomposition

of sequences into event pairs and the fact that the method cannot incorporate absolute timing

information (in the form of time, developmental stage or body size, for instance) or branch

length information. More importantly, the simulations performed by Germain and Laurin

(2009) showed that event-pair cracking with Parsimov yields more artefactual change and has

lower power to detect real sequence shifts. That method is also problematic when trying to

infer ancestral sequences and can lead to impossible ancestral reconstructions (e.g. A occurs

before B, B occurs before C, and C occurs before A), as had been documented previously.

This would create problems when trying to compare the fit of the data on various

phylogenetic hypotheses. The performance of Parsimov-based genetic inference (PGI;

Harrison and Larsson 2008) has not been assessed by simulations, but it rests on an edit cost

function that is contrary to our working hypothesis (that the timing of developmental events

can be modeled with a bounded Brownian motion model, which is assumed by continuous

analysis). More specifically, Harrison and Larsson (2008: 380) stated that their function

attempts to minimize the number of sequence changes, regardless of the magnitude of these

changes. We believe that disregarding the size of changes is unrealistic, as shown by the fact

that Poe's (2006) analyses of thirteen empirical datasets rejected that model (which he called

UC, for unconstrained change) in favor of the model we accept (AJ for adjacent states, which

favors small changes over large ones). Furthermore, analyses of ossification sequence data

using techniques for continuous data as done here (see above) have been performed by an

increasingly large number of studies (e.g., Skawiński and Borczyk 2017; Spiekman and

Werneburg 2017; Werneburg and Geiger 2017, just to mention papers published in 2017), so

the issue of ease of comparisons of our results with other studies is not as serious as it would

have been only a few years ago, and it should be decreasingly so in the future.

Reference phylogenies

We built a reference timetree that attempts to capture the established consensus (Fig. 2). The

tree was compiled in Mesquite versions up to 3.6 (Maddison and Maddison 2018) and time-

calibrated using the Stratigraphic Tools module for Mesquite (Josse et al. 2006). For

consistency and to avoid the effects of gaps in the fossil record, we used molecular divergence

dates whenever possible. The tree had to be time-scaled because many of the evolutionary

models that we fit on the tree in the first series of tests (to determine which evolutionary

model can be used to compare the fit of the hypotheses) use branch lengths to assess model

fit. Note that our procedure requires estimating divergence times between all taxa (geological

ages of all nodes). When taxa are pruned, branch lengths are adjusted automatically. The main

sources we used for topology and divergence times (and hence branch lengths) are as follows:

The phylogeny of lissamphibians follows the work of Jetz and Pyron (2018).

However, several other sources have been used for the temporal calibration of the tree:

Germain and Laurin (2009) was used for the urodeles, whereas Feng et al. (2017),

supplemented by Bossuyt and Roelants (2009) and Pyron (2014), was used for the anurans as

well as more rootward nodes (Batrachia, Lissamphibia, Tetrapoda; also Amniota). Marjanović

and Laurin (2014) was used for the Ranidae, Ceratophryidae and Hylidae.

The sediments that have preserved the temnospondyls *Apateon* and *Sclerocephalus* are

not easy to correlate with each other or with the global chronostratigraphic scale. Combining

stratigraphic information from Schoch (2014a), Schneider et al. (2015) and Werneburg

(2018), we have placed all three sampled species (*A. pedestris*, *A. caducus*, *S. haeuseri*) at the

Sakmarian/Artinskian stage boundary (Permian; 290.1 Ma ago); combining stratigraphic

information from Schneider et al. (2015) with the phylogeny in Schoch (2014a), we have

tentatively placed the divergence between the two *Apateon* species (which are not sister-

groups: Schoch 2014a) at the Kasimovian/Gzhelian stage boundary (Carboniferous; 303.7 Ma

ago). The age of the last common ancestor of *Apateon* and *Sclerocephalus* depends strongly

on temnospondyl phylogeny, which remains unresolved (Pardo et al. 2017b; Marjanović and

Laurin 2019; and numerous references in both); as a compromise between the various options,

we have provisionally placed it at the boundary between the Early and the Late Carboniferous

(Serpukhovian/Bashkirian, 323.2 Ma ago) where applicable.

For the birds, Pons et al. (2005) was used for the Laridae, Wang et al. (2013) for the

Phasianidae and Gonzales et al. (2009) for the Anatidae. The temporal calibration was taken

from Prum et al. (2015) as recommended by Berv and Field (2017); gaps were filled in using

the database www.birdtree.org.

Several papers, mainly Tarver et al. (2016), were used for the phylogeny and

divergence times of mammals. For the Muridae, three references were used: Lecompte et al.

(2008), Zhuang et al. (2015), and Lu et al. (2017) for the position of two taxa: *Mesocricetus*

auratus and *Peromyscus melanophrys*. Other species were placed following the work of

Meredith et al. (2011), which also gives divergence times. We caution, however, that all

available molecular dates for Paleogene and earlier mammal nodes are controversial and may

be overestimates (Berv and Field 2017).

Three references were also used to integrate squamates in the phylogenetic tree and for

the calibration of divergence times: Brandley et al. (2005), Rabosky et al. (2014), Reeder

(2003). Sterli et al. (2013) was used for turtles.

For turtles, there is now a near-consensus that they are diapsids, a hypothesis that is

not necessarily incompatible with an origin among "parareptiles" (Laurin and Piñeiro 2017).

Thus, following most recent molecular phylogenetic analyses (e.g., Hugall et al. 2007; Irisarri

et al. 2017), we have inserted them as the sister-group of Archosauria.

We disagree with several of the calibration dates in Irisarri et al. (2017), which often

appear unreasonably old. For instance, they place the divergence between caecilians and

batrachians and the divergence between anurans and urodeles in the Early Carboniferous,

around 330 and 320 Ma, respectively, but our thorough analyses of the fossil record, with due

consideration of its incompleteness, suggest significantly more recent dates, in the Permian

(Marjanović and Laurin 2007, 2008, 2014). This is not surprising because some of the dating

constraints used by Irisarri et al. (2017: table S8) are wrong. For instance, they enforced a

minimal divergence age between cryptodiran and pleurodiran turtles of 210 Ma (Late

Triassic), but all analyses of the last fifteen years (e.g. Sterli et al. 2013, 2018) strongly

suggest that the oldest known turtles that fit within this dichotomy date from the Late Jurassic,

less than 165 Ma. The divergence between humans and armadillos (boreotherian and

xenarthran placentals) was constrained to the middle of the Cretaceous (95.3–113 Ma), based

on outdated literature that assigned a wide variety of stem-eutherians to highly nested

positions in the placental crown; there are currently no clear placentals known from any

Cretaceous sediments even as young as 66 Ma (see e.g. Wible et al. 2009), barely half the age

of the older end of the constraint range. Conversely, the divergence between diapsids (hence

sauropsids) and synapsids had a minimal age constraint of 288 Ma (Early Permian), which is

much too young given the presence of sauropsids (and presumed synapsids) in Joggins, in

sediments that have recently been dated (Carpenter 2015) around 317–319 Ma (early Late

Carboniferous). Thus, we have not used divergence dates from that source.

To discriminate among the hypotheses on lissamphibian origins, we inserted the

temnospondyl *Apateon* in the tree where each predicts that it should be (Fig. 1c–h). Thus,

according to the TH (temnospondyl hypothesis; Fig. 1c), *Apateon* lies on the lissamphibian

stem. Under the LH (lepospondyl hypothesis; Fig. 1d), *Apateon* lies on the tetrapod stem.

Under both versions of the DH (diphyly hypothesis; Fig. 1g, h), *Apateon* lies on the

batrachian stem. Under both versions of the PH (polyphyly hypothesis; Fig. 1e, f), *Apateon*

lies on the caudate stem. Within the DH and the PH, both versions of each differ in the

position of Gymnophiona. Thus, despite the absence of any lepospondyl in our cranial

ossification sequence dataset, our taxonomic sample allows us to test all these competing

hypotheses. The appendicular datasets allow more direct tests of some of these hypotheses

because they include two lepospondyl taxa, which were likewise placed in trees representing

the tested hypotheses (Fig. 1).

Sclerocephalus is the sister-group of *Apateon* under the LH (Fig. 1d), immediately

rootward of it (on the lissamphibian stem) under the TH (Fig. 1c) and likewise (but on the

batrachian stem) under the DH1 (Fig. 1g), on the caecilian stem under the DH2 (Fig. 1h) and

the sister-group of Batrachia (including *Apateon*) under both versions of the PH (Fig. 1e, f).

"*Melanerpeton*" *humbergense* (appendicular data only) is the sister-group of *Apateon*

in all trees, except under the hypothesis of branchiosaur paraphyly; *Eusthenopteron*

(appendicular data only) forms the outgroup in all trees.

The lepospondyls *Microbrachis* and *Hyloplesion*, from both of which only

appendicular data are available, form an exclusive clade (Marjanović and Laurin 2019; Clack

2019). This clade is the sister-group of Lissamphibia (represented only by Batrachia) under

the LH (because caecilians are lacking from the appendicular datasets), of Amniota under the

TH and both versions of the DH (these three cannot be distinguished due to the absence of

caecilians) as well as under the PH1, and of Temnospondyli (including Batrachia) under the

PH2.

The temnospondyl *Micromelerpeton*, from which likewise only appendicular data are

available, forms the sister-group of *Apateon* under the LH. The uncertainty over its

phylogenetic position within Dissorophoidea (as the sister-group to the rest, including anurans

and urodeles: e.g. Schoch 2019; as the sister-group of *Apateon* + "*Melanerpeton*"

humbergense: e.g. Ruta & Coates 2007; Marjanović and Laurin 2019) generates two versions

of the TH/DH1/DH2 tree for the appendicular dataset. We tested both of these versions

against that dataset, for a total of five trees.

To ensure that our analyses were not biased in favor of a given hypothesis, and in case

that a continuous evolutionary model were favored, we initially adjusted the branch lengths

such that the sum of branch lengths was equal between the compared topologies and that the

root was approximately at the same age (in this case in the Tournaisian, the first stage of the

Carboniferous). This was done for the trees used to compare the hypotheses using the cranial

dataset because if a model incorporating (variable) branch length information had been

selected, and if the trees representing the various hypotheses had not all had the same total

length (the sum of all branch lengths), the resulting distortions in branch lengths created

around the extinct taxa (whose height compared to extant taxa is specified by their geological

age) would have introduced another variable influencing the AICc. But given that the selected

model ignores branch lengths, this precaution turned out to be superfluous. We have therefore

not made these time-consuming adjustments to the additional trees we generated later to

analyze the appendicular data.

RESULTS

In the phylogenetic analysis of cranial data, a single tree island of 22,077 trees of 438 steps

was found, only once, so there might be more trees of that length and perhaps even shorter

trees. Initially, an island of 22,075 trees was found; we swapped on each of these in a

subsequent run, which only recovered two additional trees. Given that slightly longer trees did

not differ much from those that we obtained, the low quality of the results (poor congruence

with the established consensus about the monophyly of major clades such as squamates, birds,

mammals and turtles) and the fact that about four full days of computer time had been spent

on analysis of the cranial data, we did not pursue that search further. As expected, the strict

consensus tree is poorly resolved (Fig. 3). For the appendicular matrix, 22,757 trees of 164

steps were found. Their strict consensus (Fig. 4) deviates even more from the established

consensus than the tree obtained from cranial data.

This visual assessment of phylogenetic signal through an examination of the

consensus trees (Figs. 3, 4) is congruent with the test based on squared-change parsimony and

random taxon reshuffling (Laurin 2004). Indeed, the latter indicates that the phylogenetic

signal in the cranial data is fairly strong, with a probability of less than 0.0001 that the

observed covariation between the data and the tree reflects a random distribution (none of the

10,000 random trees generated were as short as the reference tree), but it is weaker, with a

probability of 0.0017, for the appendicular data.

The speciational model of evolution, in which all branch lengths are equal, has

overwhelming support among cranial data, whether or not the Permian temnospondyl

Sclerocephalus (Table 2) or the squamosal (Table 3) are included (including *Sclerocephalus*

adds a second temnospondyl genus, but given that the timing of ossification of the squamosal

is unknown in *Sclerocephalus*, including it requires excluding the squamosal from the

analysis); the five other examined models have AICc weights $< 10^{-11}$. For the appendicular

data, the speciational model also has the most support, but that support is not as strong and

varies depending on which dataset is analyzed (seven characters or four) and under which

phylogenetic hypothesis. In three of the four tests performed, support for the second-best

model, the non-phylogenetic/equal model, varied between 5% and 19% (Table 4).

528   Two main conclusions can be drawn from these tests (Tables 2–4). First, given that

529 both of the best-supported models imply equal branch lengths, actual time represented by

530 branches can be ignored, so we compare support of the six competing topologies using only

531 the best-supported model (speciational). This simplifies the discussion, because it means that

532 the original branch lengths are irrelevant (under that model, all branch lengths are equal);

533 unfortunately, the branch length (evolutionary time) data were needed to reach this

534 conclusion. Thus, the only remaining variable is the topology. Second, model fitting, along

535 with the test based on squared-change parsimony and random taxon reshuffling, indicates that

536 the phylogenetic signal in the cranial data is strong, but that it is noticeably weaker in the

537 appendicular data (this is shown mostly by the non-negligible support for the non-

538 phylogenetic/equal model). Thus, comparisons of the fit of the various phylogenetic

539 hypotheses for the cranial data should be more reliable than for the appendicular data.

540 However, given that for several Paleozoic taxa (most importantly both of the sampled

541 lepospondyls), comparisons can be performed only for the appendicular data, these were

542 performed as well.

543   Using the speciational model, the AICc weights of the six compared topologies

544 indicate that there is strong support in the cranial data for the LH (lepospondyl hypothesis),

545 with an AICc weight of 0.9885 when *Sclerocephalus* is included (Table 5) and 0.8848 when

546 the squamosal is included instead (Table 6). Of the other topologies, the TH (temnospondyl

547 hypothesis) was by far the best supported, with an AICc weight of 0.01144 (with

548 *Sclerocephalus*) or 0.1056 (with the squamosal), which is 86.44 or 8.38 times less than for the

549 LH. Both versions of the DH (diphyly hypothesis) and of the PH (polyphyly hypothesis) have

550 negligible support (AICc weights < 0.01 when the squamosal is included, < 0.0001 when

551 *Sclerocephalus* is included). The least support is found for the PH2 when *Sclerocephalus* is

552 included, and for the DH1 when the squamosal is included. In both cases, the recently

proposed DH2 (Pardo et al. 2017b) fares second-worst by a small margin. Notably, the DH1

contradicts the modern consensus on lissamphibian monophyly (Fig. 1g), while the PH2 and

the DH2 fulfill this constraint from the molecular but not the paleontological point of view,

having lissamphibian monophyly with respect to amniotes but not with respect to

temnospondyls (Fig. 1f, h).

A slightly different dataset is used (only 84 taxa, but eight cranial characters, the

additional one being the frontal, and *Apateon* sequences for both species from Erdesbach

rather than Obermoschel) provides even stronger support for the LH, with an AICc weight of

0.9935 (Table 7). The next best-supported topology, which simultaneously represents the TH,

DH1 and DH2, has an AICc weight of only 0.0065.

The appendicular data are available in far more Paleozoic taxa than the cranial data;

these include *Sclerocephalus haeuseri*, *Archegosaurus decheni*, and the non-branchiosaurid

"branchiosaur" *Micromelerpeton credneri* among temnospondyls, the lepospondyls

Hyloplesion longicaudatum and *Microbrachis pelikani*, and the tristichopterid finned stem-

tetrapodomorph *Eusthenopteron foordi,* in addition to the same two species of *Apateon* as for

the cranial datasets, *A. caducus* and *A. pedestris*. Analysis of these postcranial data (seven

characters: humerus, radius, ulna, ilium, femur, tibia and fibula) yields surprising results, with

the PH2 having the most support, with an AICc weight of 0.7978 when using the dataset of

seven bones (Table 8). The TH, DH1 and DH2 with "branchiosaur" monophyly are

collectively (they cannot be distinguished with that taxonomic sample) the second-best

hypotheses with that dataset, with an AICc weight of only 0.1874. The least-supported

hypothesis with these data is the TH with "branchiosaur" polyphyly.

Using the other postcranial dataset with only four bones (radius, ulna, ilium, and

femur) but with more taxa (notably the branchiosaurid temnospondyl "*Melanerpeton*"

humbergense) shows that infraspecific variation in the postcranial ossification sequences of

Apateon do not significantly impact our assessment of the support for various hypotheses.

Whether both sequences of *Apateon* (from the Erdesbach and Obermoschel localities) are

included (treated as if they were distinct taxa, such as subspecies), or whether either one of

these is used in isolation, the PH2 retains the highest support, with AICc weights of 0.62 to

0.65. The LH is a distant second, at 0.20–0.23, but still well ahead of the TH/DH and the PH1,

which all receive AICc weights between 0.03 and 0.06 (Table 9).

DISCUSSION

Phylogenetic signal

In his discussion of previous phylogenetic conclusions from ossification sequences (e.g.

Schoch and Carroll 2003), Anderson (2007) noted that ossification sequences seemed to

abound in symplesiomorphies and in autapomorphies of terminal taxa, while potential

synapomorphies were scarce. This pessimism was seemingly confirmed by Schoch (2006) in

a paper that was published after Anderson's (2007) book chapter had gone to press: not only

were many similarities in the cranial ossification sequences across Osteichthyes found to be

symplesiomorphies, but a phylogenetic analysis of cranial ossification sequences did not

recover Mammalia, Sauropsida, Amniota or Lissamphibia as monophyletic. Along with these

results, Schoch (2006) dismissed another: the position of the temnospondyl *Apateon caducus*

(the only included extinct taxon) outside the tetrapod crown-group, i.e. the lepospondyl

hypothesis on lissamphibian origins (LH).

While ossification sequences alone may not provide enough data for a phylogenetic

analysis, as shown by our results (Fig. 3, 4), our datasets (with much larger taxon samples

than in Schoch 2006) fit some tree topologies much better than others. Both the tests using

CoMET and squared-change parsimony with random taxon reshuffling overwhelmingly

support the presence of a strong phylogenetic signal in the cranial data; the null hypothesis of

the absence of a phylogenetic signal can be rejected in both cases, given that it has a

probability of $< 10^{-97}$ for the cranial and $< 10^{-4}$ for the appendicular dataset. We conclude that

the cranial dataset contains a strong phylogenetic signal, and are therefore cautiously

optimistic about future contributions of ossification sequences to phylogenetics. We are less

optimistic about the appendicular sequence data, which both tests suggest contains less

phylogenetic signal.

The sizable effect on nodal estimates and inferred heterochronies of infraspecific

variation found by Sheil et al. (2014) in lissamphibians could raise doubts about the

robustness of our findings. We have been able to incorporate infraspecific variability in only

two terminal taxa (*Apateon caducus* and *A. pedestris*), but *Apateon* has played a prominent

role in discussions about the significance of cranial ossification sequences on the origins of

extant amphibians (Schoch and Carroll 2003; Schoch 2006; Germain and Laurin 2009). Thus,

incorporation of infraspecific variability in *Apateon* is presumably much more important than

in extant taxa, even though variability in the latter would obviously add to the analysis and

should be tackled in the future. The variability in *Apateon* should be exempt from two sources

of artefactual variability in ossification sequences discussed by Sheil et al. (2014), namely the

way in which the specimens were collected (there can be no lab-raised specimens in long-

extinct taxa) and the fixing method used (in this case, fossilization under quite consistent

taphonomic conditions). The finding that whether we used the *Apateon* sequences from

Erdesbach, Obermoschel, or both, we find very similar results (Table 9), is reassuring. In this

case, infraspecific variation has negligible impact. However, future studies should attempt to

assess the effect of more generalized incorporation of infraspecific variability (in a greater

proportion of the OTUs).

Of course, these results do not preclude functional or developmental constraints from

applying to the same data. This phenomenon has been documented, among other taxa, in

urodeles, whose development has often been compared with that of temnospondyls (e.g.

Schoch 2006; Schoch and Carroll 2003; Fröbisch et al. 2007, 2015; Germain and Laurin

2009). For instance, Vorobyeva and Hinchliffe (1996) documented the larval functional

constraints linked to early forelimb use that may cause an early development of manual digits

1 and 2, compared with other tetrapods, as briefly discussed below. However, in the case of

our seven cranial characters, there is no evidence of functional constraints. This is a little-

investigated topic, but all these bones apparently form a single developmental module of the

urodele skull (Laurin 2014). For the appendicular data, functional constraints might explain

the more subdued phylogenetic signal, but this will have to be determined by additional

research.

The finding that the postcranial characters that we analyzed contain relatively little

phylogenetic signal may raise doubts about the claims that have been made about the

phylogenetic implications of other such data. Specifically, Carroll et al. (1999) stated that the

neural arches ossify before the centra in frogs and temnospondyls, but not in salamanders,

caecilians or lepospondyls. When it was found that the centra do ossify first in a few

cryptobranchoid salamanders, Carroll (2007: 30) took this as "strong evidence that the most

primitive crown-group salamanders had a sequence of vertebral development that is common

to frogs and labyrinthodonts (but distinct from that of lepospondyls)". In fact, apart from tail

regeneration in *Hyloplesion* and *Microbrachis* (where the centra ossify before the neural

arches: Olori 2015; Fröbisch et al. 2015; van der Vos et al. 2017), only one incompletely

ossified vertebral column (referred to *Utaherpeton*) is known of any putative lepospondyl. "In

this specimen, […] five neural arches […] have ossified behind the most posterior centrum."

(Carroll and Chorn 1995: 40–41) Carroll's (2007: 85) claim that "the centra always ossified

prior to the arches" in lepospondyls is therefore rather puzzling.

Fröbisch et al. (2007, 2015) pointed out that the first two digital rays (digits,

metapodials and distal carpals/tarsals) ossify before the others ("preaxial polarity") in

salamanders and the temnospondyls *Apateon*, *Micromelerpeton* and *Sclerocephalus*, while the

fourth ossifies first ("postaxial polarity") in amniotes, frogs and "probably" (Fröbisch et al.

2015: 233, 234) the lepospondyls *Microbrachis* and *Hyloplesion*. This latter inference,

however, is based only on a delay in the ossification of the fifth ray that is shared specifically

with sauropsid amniotes (Olori 2015). Ossification sequences (however partial) of the other

four rays in any lepospondyl are currently limited to the tarsus of *Batropetes*, which clearly

shows preaxial polarity (Glienke 2015: fig. 6O–S; Marjanović and Laurin 2019), and that of

the putative (but see Clack et al. 2019) lepospondyl *Sauropleura*, in which likewise the

second distal tarsal ossified before all others (Marjanović and Laurin 2019). Outside of

temno- and lepospondyls, Marjanović and Laurin (2013, 2019) presented evidence that

preaxial polarity is plesiomorphic, widespread and dependent on the use of the still

developing limbs for locomotion, which would explain why it was independently lost in

amniotes and frogs and reduced (the third ray ossifies first) in direct-developing salamanders.

It may be relevant here that the PH2 (Fig. 1f), favored by our appendicular data, groups

exactly those sampled taxa in a clade that are known to have preaxial polarity in limb

development. To sum up, neither our own analyses nor the previous works that we cited

above demonstrated conclusively that ossification sequences of postcranial elements provide

reliable clues about the origin of extant amphibians.

In contrast, we are reasonably confident about our results on the cranial ossification

sequences. Given the phylogenetic signal we have found in our cranial datasets, we think that

ossification sequence data should eventually be added to phenotypic datasets for analyses of

tetrapod phylogeny. Indeed, an analysis of amniote phylogeny using data from organogenesis

sequences (coded using event-pairing in Parsimov) already exists (Werneburg and Sánchez-

Villagra 2009). The usefulness of such data for phylogenetic inference was further tested,

with encouraging results, by Laurin and Germain (2011), and the present analysis adds

additional support for it.

Indirect support for the lepospondyl hypothesis from temnospondyls

The strong support for the lepospondyl hypothesis that we have found in cranial data is

surprising because cranial ossification sequence data, especially those of the Permo-

Carboniferous temnospondyl *Apateon*, have often been claimed to contradict the LH

(lepospondyl hypothesis, Fig. 1d). Similarities between *Apateon* and extant urodeles, in

particular the supposedly "primitive" hynobiid *Ranodon*, have often been emphasized

(Schoch and Carroll 2003; Schoch and Milner 2004; Carroll 2007; Schoch 2014b). However,

other studies have already raised doubts about some of these claims (e.g. Schoch 2006;

Anderson 2007; Germain and Laurin 2009). Schoch (2006) and Anderson (2007) concluded

that most characters shared between *Apateon* and urodeles were plesiomorphies. Germain and

Laurin (2009) also demonstrated that, far from being very similar to the ancestral urodele

morphotype (contra Schoch and Carroll 2003 or Carroll 2007), the cranial ossification

sequence of *Apateon* was statistically significantly different from that of the hypothetical last

common ancestor of all urodeles (as suspected by Anderson 2007). However, these earlier

studies did not clearly show which of the various hypotheses on lissamphibian origins the

ossification sequences of *Apateon* spp. – or the newly available partial sequence (Werneburg

2018) of the phylogenetically distant temnospondyl *Sclerocephalus* – supported most. This is

what we have attempted to do here.

Unfortunately, the development of lepospondyls is too poorly documented to be

incorporated into the cranial analyses, but we included two lepospondyls in analyses of

appendicular data. These analyses weakly favor a polyphyletic origin of extant amphibians,

with both temno- and lepospondyls in the amphibian clade, a hypothesis that has not been

advocated seriously for decades (Milner 1993: fig. 5B) as far as we know. However, given the

moderate phylogenetic signal in these data, we view these results with skepticism. Olori

(2011), using event-pairing with Parsimov (Jeffery et al. 2005) and PGi (Harrison and Larsson

2008), analyzed lepospondyl postcranial ossification sequences and concluded that support for

the three hypotheses that she tested (TH/DH with two different positions for

Micromelerpeton, and LH) did not differ significantly. By contrast, our analyses of the

postcranial data indicate a stronger support for polyphyly (PH2) than for the TH/DH, which is

only a distant second (Table 8) or third (behind PH2 and LH; Table 9) depending on the

analyses. Olori (2011) performed no statistical test of phylogenetic signal of her data, though

a related test (performing phylogenetic analyses on the data) yielded trees (Olori, 2011: fig.

5.5–5.7) that are largely incongruent with the established consensus, in which most large taxa

(Mammalia, Testudines, Lissamphibia, etc.) are para- or polyphyletic. Olori's (2011) results,

like ours, support the conclusion that the phylogenetic signal in postcranial ossification

sequence data is low.

Given the current limitations in the availability of developmental data in Paleozoic

stegocephalians, we hope to have demonstrated that cranial ossification sequences of

amniotes, lissamphibians and temnospondyls provide support for the LH that is independent

of the phylogenetic analyses of Laurin (1998), Pawley (2006: appendix 16) or Marjanović and

Laurin (2009, 2018). This independence is important because the cranial ossification sequence

data cannot rival the morphological data in terms of data availability, simply because growth

sequences of extinct taxa are rare (Sánchez-Villagra 2012), but having a fairly independent

722 line of evidence to investigate a major evolutionary problem is clearly advantageous. We

723 hope that the modest methodological progress made in this study will stimulate the search for

724 fossilized ontogenies (Cloutier 2009; Sánchez-Villagra 2010).

725 FUNDING

726 This work was supported by the Centre National de la Recherche Scientifique and the French

727 Ministry of Research (unnumbered recurring grants to the CR2P, for ML).

728 ACKNOWLEDGEMENTS

729 Jennifer Olori, two anonymous reviewers and the editor Robert Asher made helpful comments

730 that improved this paper. D. M. would further like to thank Ralf Werneburg for an electronic

731 reprint of his 2018 paper, and Daniel Field for discussion of molecular divergence times and

732 the fossil record.

REFERENCES

Anderson J.S. 2007. Incorporating ontogeny into the matrix: a phylogenetic evaluation of developmental evidence for the origin of modern amphibians. Pages 182–227 in Anderson J.S., Sues H.-D., editors. Major transitions in vertebrate evolution. Bloomington: Indiana University Press.

Anderson D.R., Burnham K.P. 2002. Avoiding pitfalls when using information-theoretic methods. J. Wildl. Manag. 66:912–918.

Anderson J.S., Carroll R.L., Rowe T.B. 2003. New information on *Lethiscus stocki* (Tetrapoda: Lepospondyli: Aistopoda) from high-resolution computed tomography and a phylogenetic analysis of Aistopoda. Can. J. Earth Sci. 40:1071–1083.

Anderson J.S., Reisz R.R., Scott D., Fröbisch N.B., Sumida S.S. 2008. A stem batrachian from the Early Permian of Texas and the origin of frogs and salamanders. Nature 453:515–518.

Berv J.S., Field D.J. 2017 (printed 2018). Genomic signature of an avian Lilliput Effect across the K-Pg Extinction. Syst. Biol. 67:1–13.

Bolt J.R. 1969. Lissamphibian origins: possible protolissamphibian from the Lower Permian of Oklahoma. Science 166:888–891.

Bossuyt F., Roelants K. 2009. Frogs and toads (Anura). Pages 357–364 in Hedges S.B., Kumar S., editors. The Timetree of Life. New York: Oxford University Press.

Brandley M.C., Schmitz A., Reeder T.W. 2005. Partitioned Bayesian analyses, partition choice, and the phylogenetic relationships of scincid lizards. Syst. Biol. 54:373–390.

Carpenter D.K., Falcon-Lang H.J., Benton M.J., Grey M. 2015. Early Pennsylvanian (Langsettian) fish assemblages from the Joggins Formation, Canada, and their implications for palaeoecology and palaeogeography. Palaeontology 58:661–690.

Carroll R.L. 2001. The origin and early radiation of terrestrial vertebrates. J. Paleontol. 75:1202–1213.

Carroll R.L. 2007. The Palaeozoic ancestry of salamanders, frogs and caecilians. Zool. J. Linn. Soc. 150 (suppl. 1):1–140.

Carroll R.L., Chorn J. 1995. Vertebral development in the oldest microsaur and the problem of "lepospondyl" relationships. J. Vert. Paleont. 15:37–56.

Carroll R.L., Currie P.J. 1975. Microsaurs as possible apodan ancestors. Zool. J. Linn. Soc. 57:229–247.

Carroll R.L., Holmes R. 1980. The skull and jaw musculature as guides to the ancestry of salamanders. Zool. J. Linn. Soc. 68:1–40.

Carroll R.L., Kuntz A., Albright K. 1999. Vertebral development and amphibian evolution. Evol. Dev. 1:36–48.

Cope E.D. 1888. On the intercentrum of the terrestrial Vertebrata. Trans. Am. Phil. Soc. 16:243–253.

Cloutier R. 2009 (printed 2010). The fossil record of fish ontogenies: Insights into developmental patterns and processes. Semin. Cell Dev. Biol. 21:400–413.

Danto M., Witzmann F., Kamenz S., Fröbisch N. 2019. How informative is vertebral development for the origin of lissamphibians? J. Zool. (Lond.) 307:292–305.

Eldredge N., Gould S.J. 1972. Punctuated equilibria: an alternative to phyletic gradualism. Pages 82–115 in Schopf T.J.M., editor. Models in Paleobiology. San Francisco: Freeman, Cooper & Company.

Feng Y.-J., Blackburn D.C., Liang D., Hillis D.M., Wake D.B., Cannatella D.C., Zhang P. 2017. Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary. Proc. Natl. Acad. Sci. USA E5864–E5870.

Fröbisch N.B., Carroll R.L., Schoch R.R. 2007. Limb ossification in the Paleozoic

branchiosaurid *Apateon* (Temnospondyli) and the early evolution of preaxial dominance

in tetrapod limb development. Evol. Dev. 9:69–75.

Fröbisch N.B., Bickelmann C., Olori J.C., Witzmann F. 2015. Deep-time evolution of

regeneration and preaxial polarity in tetrapod limb development. Nature 527:231–234.

Germain D., Laurin M. 2009. Evolution of ossification sequences in salamanders and urodele

origins assessed through event-pairing and new methods. Evol. Dev. 11:170–190.

Glienke S. 2015. Two new species of the genus *Batropetes* (Tetrapoda, Lepospondyli) from

the Central European Rotliegend (basal Permian) in Germany. J. Vert. Paleont.

35:e918041.

Gonzalez J., Düttmann H., Wink M. 2009. Phylogenetic relationships based on two

mitochondrial genes and hybridization patterns in Anatidae. J. Zool. 279:310–318.

Harrington S.M., Harrison L.B., Sheil C.A. 2013. Ossification sequence heterochrony among

amphibians. Evol. Dev. 15:344–364.

Harrison L.B., Larsson H.C.E. 2008. Estimating evolution of temporal sequence changes: a

practical approach to inferring ancestral developmental sequences and sequence

heterochrony. Syst. Biol. 57:378–387.

Hugall A.F., Foster R., Lee M.S.Y. 2007. Calibration choice, rate smoothing, and the pattern

of tetrapod diversification according to the long nuclear gene RAG-1. Syst. Biol.

56:543–563.

Hugi J., Hutchinson M.N., Koyabu D., Sánchez-Villagra M.R. 2012. Heterochronic shifts in

the ossification sequences of surface- and subsurface-dwelling skinks are correlated with

the degree of limb reduction. Zoology 115:188–198.

Irisarri I., Baurain D., Brinkmann H., Delsuc F., Sire J.-Y., Kupfer A., Petersen J., Jarek M.,

Meyer A., Vences M., Philippe H. 2017. Phylotranscriptomic consolidation of the jawed

vertebrate timetree. Nature Ecol. Evol. 1:1370–1378.

Jeffery J.E., Bininda-Emonds O.R.P., Coates M.I., Richardson M.K. 2005. A new technique

for identifying sequence heterochrony. Syst. Biol. 54:230–240.

Jetz W., Pyron R.A. 2018. The interplay of past diversification and evolutionary isolation

with present imperilment across the amphibian tree of life. Nat. Ecol. Evol. 2:850–858.

Josse S., Moreau T., Laurin M. 2006. Stratigraphic tools for Mesquite, version 1.0.

http://mesquiteproject.org/packages/stratigraphicTools/

Koyabu D., Werneburg I., Morimoto N., Zollikofer C.P.E., Forasiepi A.M., Endo H., Kimura

J., Ohdachi S.D., Son N.T., Sánchez-Villagra M.R. 2014. Mammalian skull

heterochrony reveals modular evolution and a link between cranial development and

brain size. Nat. Commun. 5:3625.

Laurin M. 1998. The importance of global parsimony and historical bias in understanding

tetrapod evolution. Part I. Systematics, middle ear evolution, and jaw suspension. Ann.

Sci. Nat., Zool., 13^e Sér. 19:1–42.

Laurin M. 2004. The evolution of body size, Cope's rule and the origin of amniotes. Syst.

Biol. 53:594–622.

Laurin M. 2014. Assessment of modularity in the urodele skull: an exploratory analysis using

ossification sequence data. J. Exp. Zool. B (Mol. Dev. Evol.) 322:567–585.

Laurin M., Germain D. 2011. Developmental characters in phylogenetic inference and their

absolute timing information. Syst. Biol. 60:630–644.

Laurin M., Piñeiro G. 2017. A reassessment of the taxonomic position of mesosaurs, and a

surprising phylogeny of early amniotes. Front. Earth Sci. 5:88.

Lecompte E., Aplin K., Denys C., Catzeflis F., Chades M., Chevret P. 2008. Phylogeny and

biogeography of African Murinae based on mitochondrial and nuclear gene sequences,

with a new tribal classification of the subfamily. BMC Evol. Biol. 8:199.

Lee C., Blay S., Mooers A.Ø., Singh A., Oakley T.H. 2006. CoMET: A Mesquite package for

comparing models of continuous character evolution on phylogenies. Evol.

Bioinformatics Online 2:193–196.

Lu T., Zhu M., Yi C., Si C., Yang C., Chen H. 2017. Complete mitochondrial genome of the

gray red-backed vole (*Myodes rufocanus*) and a complete estimate of the phylogenetic

relationships in Cricetidae. Mitochondrial DNA Part A 28:62-64.

Maddin H.C., Jenkins F.A. Jr., Anderson J.S. 2012. The braincase of *Eocaecilia micropodia*

(Lissamphibia, Gymnophiona) and the origin of caecilians. PLOS ONE 7:e50743.

Maddison W.P. 1991. Squared-change parsimony reconstructions of ancestral states for

continuous-valued characters on a phylogenetic tree. Syst. Zool. 40:304–314.

Maddison W.P., Maddison D.R. 2018. Mesquite: a modular system for evolutionary analysis.

Version 3.6. http://mesquite.wikispaces.com

Mann A., Pardo J.D., Maddin H.C. 2019. *Infernovenator steenae*, a new serpentine

recumbirostran from the 'Mazon Creek' *Lagertätte* [sic] further clarifies lysorophian

origins. Zool. J. Linn. Soc. online early (12 pp.).

Marjanović D., Laurin M. 2007. Fossils, molecules, divergence times, and the origin of

lissamphibians. Syst. Biol. 56:369–388.

Marjanović D., Laurin M. 2008. Assessing confidence intervals for stratigraphic ranges of

higher taxa: The case of Lissamphibia. Acta Palaeont. Pol. 53:413–432.

Marjanović D., Laurin M. 2009. The origin(s) of modern amphibians: a commentary. Evol.

Biol. 36:336–338.

Marjanović D., Laurin M. 2013. The origin(s) of extant amphibians: a review with emphasis

on the "lepospondyl hypothesis". Geodiversitas 35:207–272.

Marjanović D., Laurin M. 2014. An updated paleontological timetree of lissamphibians, with

comments on the anatomy of Jurassic crown-group salamanders (Urodela). Hist. Biol.

26:535–550.

Marjanović D., Laurin M. 2019. Phylogeny of Paleozoic limbed vertebrates reassessed through

revision and expansion of the largest published relevant data matrix. PeerJ 6:e5565.

Maxwell E.E., Harrison L.B., Larsson H.C.E. 2010. Assessing the phylogenetic utility of

sequence heterochrony: evolution of avian ossification sequences as a case study.

Zoology 113:57–66.

Meredith R.W., Janečka J.E., Gatesy J., Ryder O.A., Fisher C.A., Teeling E.C., Goodbla A.,

Eizirik E., Simão T.L.L., Stadler T., Rabosky D.L., Honeycutt R.L., Flynn J.J., Ingram

C.M., Steiner C., Williams T.L., Robinson T.J., Burk-Herrick A., Westerman M.,

Ayoub N.A., Springer M.S., Murphy W.J. 2011. Impacts of the Cretaceous terrestrial

revolution and KPg extinction on mammal diversification. Science 334:521–524.

Milner A.R. 1993. The Paleozoic relatives of lissamphibians. Herpetol. Monogr. 7:8–27.

Olori J.C. 2011. The evolution of skeletal development in early tetrapods: anatomy and

ontogeny of microsaurs (Lepospondyli) [doctoral thesis]. Austin: University of Texas at

Austin. http://hdl.handle.net/2152/ETD-UT-2011-05-3535

Olori J.C. 2013. Ontogenetic sequence reconstruction and sequence polymorphism in extinct

taxa: an example using early tetrapods (Tetrapoda: Lepospondyli). Paleobiology

39:400–428.

Olori J.C. 2015. Skeletal morphogenesis of *Microbrachis* and *Hyloplesion* (Tetrapoda:

Lepospondyli), and implications for the developmental patterns of extinct, early

tetrapods. PLOS ONE 10:e0128333.

Pardo J.D., Szostakiwskyj M., Ahlberg P.E., Anderson J.S. 2017a. Hidden morphological

diversity among early tetrapods. Nature 546:642–645.

Pardo J.D., Small B.J., Huttenlocker A.K. 2017b. Stem caecilian from the Triassic of

Colorado sheds light on the origins of Lissamphibia. Proc. Natl. Acad. Sci. U.S.A.

114:E5389–E5395.

Parsons T.S., Williams E.E. 1962. The teeth of Amphibia and their relation to amphibian

phylogeny. J. Morph. 110:375–389.

Parsons T.S., Williams E.E. 1963. The relationships of the modern Amphibia: A re-

examination. Q. Rev. Biol. 38:26–53.

Pawley K. 2006. The postcranial skeleton of temnospondyls (Tetrapoda: Temnospondyli)

[doctoral thesis]. Melbourne: La Trobe University. http://hdl.handle.net/1959.9/405644

Poe S. 2006. Test of von Baer's law of the conservation of early development. Evolution

60:2239–2245.

Pons J.-M., Hassanin A., Crochet P.-A. 2005. Phylogenetic relationships within the Laridae

(Charadriiformes: Aves) inferred from mitochondrial markers. Mol. Phyl. Evol. 37:686–

699.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon A.M., Lemmon A.R.

2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA

sequencing. Nature 526:569–573.

Pyron R.A. 2014. Biogeographic analysis reveals ancient continental vicariance and recent

oceanic dispersal in amphibians. Syst. Biol. 63:779–797.

Rabosky D.L., Donnellan S.C., Grundler M., Lovette I.J. 2014. Analysis and visualization of

complex macroevolutionary dynamics: an example from Australian scincid lizards. Syst.

Biol. 63:610–627.

Reeder T.W. 2003. A phylogeny of the Australian *Sphenomorphus* group (Scincidae:

Squamata) and the phylogenetic placement of the crocodile skinks (*Tribolonotus*):

Bayesian approaches to assessing congruence and obtaining confidence in maximum

likelihood inferred relationships. Mol. Phyl. Evol. 27:384–397.

Rineau V., Grand A., Zaragüeta R., Laurin M. 2015. Experimental systematics: sensitivity of

cladistic methods to polarization and character ordering schemes. Contr. Zool. 84:129–

148.

Rineau V., Zaragüeta i Bagils R., Laurin M. 2018. Impact of errors on cladistic inference:

simulation-based comparison between parsimony and three-taxon analysis. Contr. Zool.

87:25–40.

Ruta M., Coates M.I. 2007. Dates, nodes and character conflict: addressing the lissamphibian

origin problem. J. Syst. Palaeontol. 5:69–122.

Sánchez M. 2012. Embryos in Deep Time: The Rock Record of Biological Development. U.

of California Press, Berkeley.

Sánchez-Villagra M.R. 2010. Contributions on fossilised ontogenies: The rock record of

vertebrate development. Semin. Cell Dev. Biol. 21:399.

Schneider J.W., Werneburg R., Rößler R., Voigt S., Scholze F. 2015. Example for the

description of basins in the CPT Nonmarine-Marine Correlation Chart – Thuringian

Forest Basin, East Germany. Permophiles 61:29–35.

Schoch R.R. 2002. The formation of the skull in Paleozoic and extant amphibians.

Paleobiology 28:378–396.

Schoch R.R. 2004. Skeleton formation in the Branchiosauridae: a case study in comparing

ontogenetic trajectories. J. Vert. Paleont. 24:309–319.

Schoch R.R. 2006. Skull ontogeny: developmental patterns of fishes conserved across major

tetrapod clades. Evol. Dev. 8:524–536.

Schoch R.R. 2014a. First evidence of the branchiosaurid temnospondyl *Leptorophus* in the

Early Permian of the Saar-Nahe Basin (SW Germany). N. Jb. Geol. Paläont. Abh.

272:225–236.

Schoch R.R. 2014b. Amphibian skull evolution: the developmental and functional context of

simplification, bone loss and heterotopy. J. Exp. Zool. (Mol. Dev. Evol.) 322B:619–630.

Schoch R.R. 2019. The putative lissamphibian stem-group: phylogeny and evolution of the

dissorophoid temnospondyls. J. Paleont. 93:37–156.

Schoch R.R., Carroll R.L. 2003. Ontogenetic evidence for the Paleozoic ancestry of

salamanders. Evol. Dev. 5:314–324.

Schoch R.R., Milner A.R. 2004. Structure and implications of theories on the origin of

lissamphibians. Pages 345–377 in Arratia G., Wilson M.V.H., Cloutier R., editors.

Recent Advances in the Origin and Early Radiation of Vertebrates. Munich: Dr.

Friedrich Pfeil.

Sheil C.A., Jorgensen M., Tulenko F., Harrington S. 2014. Variation in timing of ossification

affects inferred heterochrony of cranial bones in Lissamphibia. Evol. Dev. 16:292–305.

Sigurdsen T., Green D.M. 2011. The origin of modern amphibians: a re-evaluation. Zool. J.

Linn. Soc. 162:457–469.

Skawiński T., Borczyk B. 2017. Evolution of developmental sequences in lepidosaurs. PeerJ

5:e3262.

Spiekman S.N., Werneburg I. 2017. Patterns in the bony skull development of marsupials:

high variation in onset of ossification and conserved regions of bone contact. Sci. Rep.

7:43197.

Steen M.C. 1938. On the fossil Amphibia from the Gas Coal of Nýřany and other deposits in

Czechoslovakia. Proc. Zool. Soc. Lond. 108:205–283.

Sterli J., Pol D., Laurin M. 2013. Incorporating phylogenetic uncertainty on phylogeny-based

paleontological dating and the timing of turtle diversification. Cladistics 29:233–246.

Sterli J., de la Fuente M.S., Rougier G.W. 2018. New remains of *Condorchelys antiqua*

(Testudinata) from the Early-Middle Jurassic of Patagonia: anatomy, phylogeny, and

paedomorphosis in the early evolution of turtles. J. Vert. Paleont. 38:e1480112.

Swofford D.L. 2019. PAUP*: Phylogenetic Analysis Using Parsimony (*and other methods),

version 4.0a165. Sinauer Associates.

Tarver J.E., dos Reis M., Mirarab S., Moran R.J., Parker S., O'Reilly J.E., King B.L.,

O'Connell M.J., Asher R.J., Warnow T., Peterson K.J., Donoghue P.C.J., Pisani D.

2016. The interrelationships of placental mammals and the limits of phylogenetic

inference. Genome Biol. Evol. 8:330–344.

van der Vos W., Witzmann F., Fröbisch N.B. 2017. Tail regeneration in the Paleozoic

tetrapod *Microbrachis pelikani* and comparison with extant salamanders and squamates.

J. Zool. 304:34–44.

Vorobyeva E.I., Hinchliffe J.R. 1996. Developmental pattern and morphology of

Salamendrella keyserlingii limbs (Amphibia, Hynobiidae) including some

evolutionary aspects. Russ. J. Herpetol. 1:68–81.

Wagenmakers E.-J., Farrell S. 2004. AIC model selection using Akaike weights. Psychon.

Bull. Rev. 11:192–196.

Wang N., Kimball R.T., Braun E.L., Liang B., Zhang Z. 2013. Assessing phylogenetic

relationships among Galliformes: a multigene phylogeny with expanded taxon sampling

in Phasianidae. PLOS ONE 8:1–12.

Watson D.M.S. 1940. The origin of frogs. Trans. R. Soc. Edinburgh 60:195–231.

Weisbecker V. 2011. Monotreme ossification sequences and the riddle of mammalian skeletal

development. Evolution 65:1323–1335.

Weisbecker V., Mitgutsch C. 2010. A large-scale survey of heterochrony in anuran cranial

ossification patterns. J. Zool. Syst. Evol. Research 48:332–347.

Werneburg R. 2018 (for 2017). Earliest 'nursery ground' of temnospondyl amphibians in the

Permian. Semana 32:3–42.

Werneburg I., Geiger M. 2017. Ontogeny of domestic dogs and the developmental

foundations of carnivoran domestication. J. Mammal. Evol. 24:323–343.

Werneburg I., Sánchez-Villagra M.R. 2009. Timing of organogenesis support basal position

of turtles in the amniote tree of life. BMC Evol. Biol. 9:82.

Witzmann F., Schoch R.R. 2006. Skeletal development of the temnospondyl

Acanthostomatops vorax from the Lower Permian Döhlen Basin of Saxony. Trans. R.

Soc. Edinburgh: Earth Sci. 96:365–385.

Zhang P., Zhou H., Chen Y.-Q., Liu Y.-F., Qu L.-H. 2005. Mitogenomic perspectives on the

origin and phylogeny of living amphibians. Syst. Biol. 54:391–400.

Zhuang L., Bluteau G., Trueb B. 2015. Phylogenetic analysis of receptor FgfrL1 shows

divergence of the C-terminal end in rodents. Comp. Biochem. Physiol. B 186:43–50.

.

FIGURE LEGENDS

FIGURE 1. Hypotheses on the relationships of the extant amphibian clades since the late 20th

century. The names of terminal taxa sampled here for cranial characters are in boldface, those

sampled for appendicular characters are underlined; the names of larger clades are placed

toward the right end of a branch if they have minimum-clade (node-based) definitions, to the

left if they have maximum-clade (branch-based) definitions. Names in parentheses would,

given that phylogenetic hypothesis, not be used, but replaced by synonyms. Among terminal

taxa, "*Melanerpeton*" *humbergense*, sampled for appendicular characters, is not shown, but is

always the sister-group of *Apateon*; *Microbrachis*, likewise sampled for appendicular

characters, is not shown either, but is always the sister-group of *Hyloplesion*; *Eusthenopteron*

is not shown in c)–h), where it forms the outgroup (b)). For complications involving the

dissorophoid temnospondyl *Micromelerpeton*, see the text. The first two trees (a, b) show the

current consensus; the other trees (c–h) show the various tested paleontological hypotheses.

Abbreviations: D., Dissorophoidea; S., Stereospondylomorpha. a) Consensus of the latest and

largest phylogenetic analyses of molecular data (Irisarri et al. 2017; Feng et al. 2017; Jetz and

Pyron 2018); all named clades are therefore extant. Note the monophyly of the extant

amphibians (Lissamphibia, marked with a light gray dot) with respect to Amniota. b)

Consensus of all analyses of Paleozoic limbed vertebrates (latest and largest: Pawley 2006;

Sigurdsen and Green 2011; Pardo et al. 2017a, b: fig. S6; Marjanović and Laurin 2019; Clack

et al. 2019), omitting the extant amphibian clades. Note the monophyly of "lepospondyls" +

amniotes (marked with a dark gray dot). c) TH: "temnospondyl hypothesis" (most recently

found by Sigurdsen and Green 2011; Maddin et al. 2012; Pardo et al. 2017a, b: fig. S6; argued

for by Schoch and Milner 2004, Schoch 2014b and others). Lissamphibia nested among

dissorophoid temnospondyls. Compatible with both a) and b) (gray dots). d) LH:

"lepospondyl hypothesis" (found most recently by Pawley 2006; Marjanović and Laurin

2009, 2018). Lissamphibia nested among "lepospondyls"; consequently, temnospondyls are

not crown-group tetrapods. Compatible with both a) and b) (gray dots). e) PH1: "polyphyly

hypothesis", first variant (argued for by Carroll 2001, 2007; Schoch and Carroll 2003; very

cautiously Fröbisch et al. 2007). Urodela as dissorophoid temnospondyls close to *Apateon*,

Anura as a separate clade of dissorophoid temnospondyls, Gymnophiona as "lepospondyls".

Compatible with b) (dark gray dot) but not with a) (light gray circle). f) PH2: "polyphyly

hypothesis", second variant (argued for, as one of two options, by Milner 1993). Like PH1,

but with restored monophyly of extant amphibians with respect to amniotes (light gray dot;

see a)) at the expense of compatibility with the paleontological consensus concerning the

position of temnospondyls, lepospondyls, and amniotes (dark gray circle; see b)). g) DH1:

"diphyly hypothesis", first variant (found by Anderson 2007; Anderson et al. 2008). Batrachia

as dissorophoid temnospondyls, Gymnophiona as "lepospondyls". Compatible with b) (dark

gray dot) but not with a) (light gray circle). h) DH2: "diphyly hypothesis", second variant

(found by Pardo et al. 2017b in an analysis that included only temnospondyls and

lissamphibians: fig. 2, S7). Batrachia as dissorophoid temnospondyls, Gymnophiona as

stereospondylomorph temnospondyls . Compatible with both a) and b).

FIGURE 2. Reference phylogeny used for some of the analyses, illustrating the LH

(lepospondyl hypothesis) of lissamphibian origins. The tree was time-calibrated, but analyses

showed that branch lengths are irrelevant, given that the best model is speciational (Tables 2–

4).

FIGURE 3. Strict consensus of the most parsimonious trees obtained by analyzing cranial

dataset 2, which is comprised of 105 taxa and seven characters (see Table 1). Note that

several higher taxa whose monophyly is well-established appear to be para- or polyphyletic

here, which strongly suggests that these data are insufficient to reliably estimate a phylogeny,

but there is clearly a phylogenetic signal because the taxa are not randomly scattered over the

tree. The majority-rule consensus (not shown, but available in SM 1) is more resolved but not

necessarily better because much of the additional resolution contradicts the established

consensus.

FIGURE 4. Strict consensus of the most parsimonious trees obtained by analyzing appendicular

dataset 3, which is comprised of 62 taxa and seven characters (see Table 1). The phylogenetic

signal in these data seems to be lower than in the cranial data. As for the cranial data, the

majority-rule consensus (not shown, but available in SM 1) is more resolved but not

necessarily better because much of the additional resolution contradicts the established

consensus.

1053

1054

Figure 1

Figure 2

Character 1: Taxa
Parsimony
reconstruction (
Unordered) [Steps: 23]

- actinopterygians
- temnospondyls
- lepospondyls
- gymnophionans
- urodeles
- anurans
- turtles
- squamates
- archosaurs
- mammals

Strict, cranial analysis 6, 438 steps

Figure 3

Figure 4

1055   TABLE 1. List of datasets used in this paper. All are subsets of our global compilation that

1056   were selected to meet the requirement of the method used (missing data cannot be handled).

1057   The temnospondyl species *Apateon caducus* and *A. pedestris* are included in all datasets, but

1058   scored after populations from two different paleo-lakes in which both species occur.

Dataset number	1	2	3	4	5
Type of characters	cranial	cranial	appendicular	appendicular	cranial
Number of characters	6	7	7	4	8
Number of taxa	107	105	62	65	84
Sclerocephalus	yes	no	yes	yes	yes
Source of data for *Apateon*	Erdesbach	Erdesbach	Obermoschel	Erdesbach and Obermoschel	Erdesbach
Additional Paleozoic taxa	None	None	*Archegosaurus, Micromelerpeton, Hyloplesion, Microbrachis, Eusthenopteron*	*Archegosaurus, Micromelerpeton,* "*Melanerpeton*" *humbergense, Hyloplesion, Microbrachis, Eusthenopteron*	None
Table in which it is used	2, 5	3, 6	4, 8	4, 9	7

1059

1060

TABLE 2. Support (AICc and AICc weights) for six evolutionary models given our reference

tree (LH) and dataset 1 (see Table 1), which comprises six cranial characters (nasal, parietal,

squamosal, maxilla, pterygoid, and exoccipital) scored in 107 taxa, including the

temnospondyl *Sclerocephalus*. This was performed on the tree representing the LH

(lepospondyl hypothesis), but doing this on other trees leads to similar results. Numbers

presented with four significant digits; best values in boldface. "Distance" refers to keeping the

original branch lengths (which represent evolutionary time), "equal" sets all branch lengths

(internal and terminal) to 1, "free" infers them from the data. Abbreviations: k, number of

estimable parameters; l, likelihood; wi, weight; Δ_i, difference of AICc from that of the Pure-

Phylogenetic / Equal model.

	AIC	l	k	AICc	Δ_i AICc	wi(AICc)
Pure-Phylogenetic / Distance	−584.4	293.2	**1**	−583.4	641.2	5.85 E−140
Pure-Phylogenetic / Equal (speciational)	**−1225.6**	**613.8**	**1**	**−1224.6**	**0**	**1.000**
Pure-Phylogenetic / Free	2.000 E10	−1.000 E10	486	2.000 E10	2.000 E10	< E−165
Non-Phylogenetic / Distance	−473.6	237.8	**1**	−472.6	752.0	4.97 E−164
Non-Phylogenetic / Equal	−959.9	481.0	**1**	−958.9	265.7	2.02 E−58
Non-Phylogenetic / Free	2.000 E10	−1.000 E10	244	2.000 E10	2.000 E10	< E−165

1071

1072   TABLE 3. Support (AICc and AICc weights) for six evolutionary models given our reference

1073   tree (LH) and dataset 2 (see Table 1), which comprises seven cranial characters (nasal,

1074   parietal, squamosal, premaxilla, maxilla, pterygoid, and exoccipital) and 105 taxa, excluding

1075   *Sclerocephalus*. Abbreviations and boldface as in Table 2.

Evolutionary model	AIC	L	k	AICc	Δ_i AICc	wi(AICc)
Pure-Phylogenetic / Distance	−715.9	359.0	**1**	−714.9	683.5	< E−26
Pure-Phylogenetic / Equal	**−1399.5**	**700.7**	**1**	**−1398.5**	**0**	**1.000**
Pure-Phylogenetic / Free	2.000 E10	−1.000 E10	306	2.000 E10	2.000 E10	0
Non-Phylogenetic / Distance	−580.6	291.3	**1**	−579.6	818.8	< E−26
Non-Phylogenetic / Equal	−1106.0	554.0	**1**	−1105.0	293.5	2.278 E−98
Non-Phylogenetic / Free	2.000 E10	−1.000 E10	244	2.000 E10	2.000 E10	< E−26

1076

1077 TABLE 4. AICc weights showing relative support for six evolutionary models given various

1078 appendicular datasets (3 and 4; see Table 1) and various hypotheses. Because of the number

1079 of analyses presented below, only the AICc weights are presented (best values in boldface).

1080 Abbreviations: DH, diphyly hypothesis (both versions); LH, lepospondyl hypothesis; TH,

1081 temnospondyl hypothesis.

Evolutionary model	7 characters, LH	7 characters, LH	4 characters, LH	4 characters, TH/DH
Pure-Phylogenetic / Distance	5.1857 E−149	2.340 E−70	1.227 E−52	2.646 E−52
Pure-Phylogenetic / Equal	**1**	**0.9335**	**0.94459**	**0.8139**
Pure-Phylogenetic / Free	< E−179	1.598 E−277	4.012 E−158	3.002 E−155
Non-Phylogenetic / Distance	7.515 E−179	4.843 E−52	2.162 E-42	7.262 E−42
Non-Phylogenetic / Equal	2.14914 E−64	6.648 E−02	5.541 E−02	0.1861
Non-Phylogenetic / Free	< E−179	< E−179	< E−179	< E−179

1082

1083

TABLE 5. Support (AIC and AICc weights) for the six topologies, reflecting the six

hypotheses about the origin of extant amphibians, under the speciational model (called Pure-

Phylogenetic / Equal in Tables 2–4), with dataset 1 (see Table 1), which includes six cranial

characters (nasal, parietal, squamosal, maxilla, pterygoid, and exoccipital) and 107 taxa

(including, among Paleozoic taxa, *Apateon* and *Sclerocephalus*). Abbreviations and boldface

as in Table 2, except Δ_i: difference of AICc from that of the LH. Hypotheses from top to

bottom: LH: monophyletic origin from lepospondyls; TH: monophyletic origin among

temnospondyls; DH1: diphyletic origin, caecilians from lepospondyls and batrachians from

temnospondyls, as in Anderson et al. (2008); DH2: diphyletic origin (batrachians and

caecilians from different temnospondyls: Pardo et al. 2017b); PH1: triphyletic (polyphyletic)

origin with anurans and urodeles from different temnospondyls, caecilians from lepospondyls,

and lepospondyls closer to Amniota than to Batrachia (Fröbisch et al. 2007); PH2: triphyletic

(polyphyletic) origin as above, but with lepospondyls and caecilians closer to temnospondyls

than to amniotes (Milner 1993), reflecting the well-established lissamphibian monophyly

among extant taxa (e.g. Irisarri et al. 2017; Feng et al. 2017).

Hypothesis	AIC	L	AICc	Δ_i AICc	wi(AICc)
TH	−1217	609.4	−1215	8.919	0.01144
LH	**−1226**	**613.8**	**−1224**	**0**	**0.9885**
DH1	−1204	602.9	−1202	21.90	1.738 E−05
DH2	−1195	598.3	−1193	31.01	1.827 E−07
PH1	−1194	597.9	−1192	31.86	1.196 E−07
PH2	−1193	597.4	−1191	32.89	7.143 E−08

1099

1100

1101   TABLE 6. Support (AIC and AICc weights) for the six topologies, reflecting the six

1102   hypotheses about the origin of extant amphibians, for dataset 2 (see Table 1), which includes

1103   seven cranial characters (nasal, parietal, squamosal, premaxilla, maxilla, pterygoid, and

1104   exoccipital) and 105 taxa, excluding *Sclerocephalus* (among Paleozoic taxa, only *Apateon* is

1105   present). Abbreviations, boldface and hypotheses as in Tables 2 and 5.

Hypothesis	AIC	L	AICc	Δ_i AICc	wi(AICc)
TH	−1395	698.6	−1394	4.251	0.1056
LH	**−1399**	**700.7**	**−1398**	**0**	**0.8848**
DH1	−1384	693.1	−1383	15.203	4.42 E−4
DH2	−1385	693.6	−1384	14.315	6.89 E−4
PH1	−1387	694.5	−1386	12.404	1.792 E−3
PH2	−1390	695.8	−1388	9.792	6.615 E−3

1106

1107

1108   TABLE 7. Support for the various hypotheses about amphibian origins for dataset 5 (see Table

1109   1), which includes eight cranial characters (frontal added) and 84 taxa, with *Apateon*

1110   sequences from Erdesbach (in addition to *Sclerocephalus* among Paleozoic taxa).

1111   Abbreviations, boldface and hypotheses as in Tables 2 and 5. Because of the taxon sample,

1112   only three topologies can be tested.

Hypothesis	AIC	L	AICc	Δ_i AICc	wi(AICc)
LH	**−1296**	**649.0**	**−1294**	**0**	**0.9935**
TH, DH1, DH2	−1286	644.0	−1284	10.061	6.493 E−3
PH	−1274	638.0	−1272	22.038	1.628 E−5

1113

1114

TABLE 8. Support (AICc weights) for the various hypotheses about amphibian origins

according to dataset 3 (see Table 1), which features seven appendicular characters (humerus,

radius, ulna, ilium, femur, tibia and fibula) and 62 taxa, including several Paleozoic taxa (the

temnospondyls *Archegosaurus decheni* and *Micromelerpeton credneri*, the lepospondyls

Hyloplesion longicaudatum and *Microbrachis pelikani*, and the tristichopterid *Eusthenopteron*

foordi) in addition to *Apateon* (two species, *A. caducus* and *A. pedestris*) and *Sclerocephalus*

haeuseri. The *Apateon* sequences come from Obermoschel. Abbreviations, boldface and

hypotheses as in Table 5, except that the TH and both variants of the DH become

indistinguishable, but the phylogenetic position of the "branchiosaur" *Micromelerpeton* can

be tested.

Hypothesis	AIC	l	AICc	Δ_i AICc	wi(AICc)
LH	−885.0	443.5	−884.2	11.808	2.177 E−3
TH, DH (branchiosaur monophyly)	−881.1	441.6	−880.3	2.897	0.1874
TH, DH (branchiosaur polyphyly)	−886.4	444.2	−885.6	15.754	3.027 E−4
PH1	−888.5	445.3	−887.7	8.341	0.01232
PH2	**−896.9**	**449.4**	**−896.1**	**0.000**	**0.7978**

1125

1126

TABLE 9. Effect of the intraspecific variability in ossification sequences of *Apateon* on the

support (AICc weight; best values in boldface) for the various hypotheses about amphibian

origins. The dataset (number 4; Table 1) includes only four appendicular bones (radius, ulna,

ilium, and femur) and 63 to 65 taxa but it allows testing the impact of infraspecific variability

in ossification sequences in *Apateon*, which are documented in two localities (Erdesbach and

Obermoschel). Because of the number of tests presented (15: five topologies x three sets of

sequences), only the AICc weights are given. In all tests, the following Paleozoic taxa are

present: *Sclerocephalus haeuseri*, *Archegosaurus decheni*, "*Melanerpeton*" *humbergense*,

Micromelerpeton credneri, *Apateon* (two species, *A. caducus* and *A. pedestris*) among

temnospondyls, *Hyloplesion longicaudatum* and *Microbrachis pelikani* among lepospondyls,

and the tristichopterid *Eusthenopteron foordi*. For abbreviations of the hypotheses, see Table

5.

Hypothesis	Erdesbach and Obermoschel	Erdesbach	Obermoschel
LH	0.21407	0.20169	0.22657
TH, DH (branchiosaur monophyly)	0.05492	0.05265	0.05532
TH, DH (branchiosaur polyphyly)	0.03713	0.04285	0.03342
PH1	0.05653	0.05491	0.05638
PH2	**0.63735**	**0.64790**	**0.62832**

1139

Appendix 1: Sources of data for ossification sequences.

Empty cells indicate that these data are unavailable. Three methods were examined, and we used the one for which most data were available (position in the ossification sequence, last column).

Taxa	Standardization method (data type used)		
	Ontogenetic stages	Snout-vent length (mm)	Ossification sequence position
Actinopterygii			
Amia calva		Grande and Bemis 1998	Grande and Bemis 1998
Clarias gariepinus		Adriaens and Verraes 1998	Adriaens and Verraes 1998
Danio rerio		Cubbage and Mabee 1996	Cubbage and Mabee 1996
Oryzias latipes	Langille and Hall 1987		
Tristichopteridae			
Eusthenopteron foordi		Cote et al. 2002; Leblanc and Cloutier 2005	Cote et al. 2002; Leblanc and Cloutier 2005
Temnospondyli			
Archegosaurus decheni		Witzmann 2006	Witzmann 2006

Apateon caducus (Erdesbach)	Schoch 2004	Schoch 2004	Schoch 2004
Apateon caducus (Obermoschel)		Werneburg 2018	Werneburg 2018
Apateon pedestris (Erdesbach)	Schoch 2004		Schoch 2004
Apateon pedestris (Obermoschel)		Werneburg 2018	Werneburg 2018
"*Melanerpeton*" *humbergense*	Schoch 2004		Schoch 2004
Micromelerpeton credneri		Boy 1995; Lillich and Schoch 2007; Witzmann and Pfretzschner 2009; Schoch 2009	Boy 1995; Lillich and Schoch 2007; Witzmann and Pfretzschner 2009; Schoch 2009
Sclerocephalus haeuseri	Lohmann and Sachs 2001; Schoch 2003; Schoch and Witzmann 2009; Werneburg 2018	Lohmann and Sachs 2001; Schoch 2003; Schoch and Witzmann 2009; Werneburg 2018	Lohmann and Sachs 2001; Schoch 2003; Schoch and Witzmann 2009; Werneburg 2018
Lepospondyli			
Hyloplesion longicaudatum		Olori 2013	Olori 2013
Microbrachis pelikani		Olori 2013	Olori 2013

Gymnophiona

Gegeneophis ramaswamii	Müller et al. 2005		Harrington et al. 2013
Hypogeophis rostratus	Müller 2006		Harrington et al. 2013

Urodela

Aneides lugubris		Wake et al. 1983	Wake et al. 1983
Ambystoma macrodactylum			Harrington et al. 2013
Ambystoma maculatum	Moore 1989		Harrington et al. 2013
Ambystoma mexicanum		Laurin and Germain 2011	Harrington et al. 2013
Ambystoma talpoideum	Reilly 1987	Reilly 1987	Reilly 1987
Ambystoma texanum		Laurin and Germain 2011	Harrington et al. 2013
Ambystoma tigrinum			Harrington et al. 2013
Amphiuma means			Harrington et al. 2013
Andrias japonicus			Harrington et al. 2013
Bolitoglossa subpalmata			Ehmcke and Clemen 2000
Dicamptodon tenebrosus			Harrington et al. 2013
Eurycea bislineata			Harrington et al. 2013
Gyrinophilus porphyriticus			Harrington et al. 2013
Hemidactylium scutatum			Harrington et al. 2013
Lissotriton vulgaris		Laurin and Germain 2011	Harrington et al. 2013

Necturus maculosus			Harrington et al. 2013
Notophthalmus viridescens	Reilly 1986	Reilly 1986	Harrington et al. 2013
Onychodactylus japonicus			Harrington et al. 2013
Pleurodeles waltl			Harrington et al. 2013
Ranodon sibiricus			Harrington et al. 2013
Salamandra salamandra			Harrington et al. 2013
Salamandrella keyserlingii			Harrington et al. 2013
Siren intermedia	Reilly and Altig 1996	Reilly and Altig 1996	Reilly and Altig 1996
Triturus karelinii			Harrington et al. 2013

Anura

Alytes obstetricans			Yeh 2002
Ascaphus truei			Harrington et al. 2013
Anaxyrus boreas			Gaudin 1978
Bombina orientalis			Harrington et al. 2013
Bufo bufo			Harrington et al. 2013
Cornufer guentheri			Harrington et al. 2013
Ceratophrys cornuta			Harrington et al. 2013
Chacophrys pierotti			Harrington et al. 2013
Crinia signifera			Harrington et al. 2013
Dendrobates auratus	de Sá and Hill 1998	de Sá and Hill 1998	Harrington et al. 2013
Discoglossus sardus			Pugener and Maglia 1997
Eleutherodactylus coqui			Harrington et al. 2013

Eleutherodactylus nubicola			Harrington et al. 2013
Epidalea calamita			Harrington et al. 2013
Epipedobates tricolor	de Sá and Hill 1998	de Sá and Hill 1998	Harrington et al. 2013
Fejervarya cancrivora			Harrington et al. 2013
Hamptophryne boliviana			Harrington et al. 2013
Hyla versicolor			Harrington et al. 2013
Hylorina sylvatica			Harrington et al. 2013
Hymenochirus boettgeri			de Sá and Swart 1999
Hypsiboas lanciformis	de Sá 1988	de Sá 1988	de Sá 1988
Kassina senegalensis			Harrington et al. 2013
Leptodactylus chaquensis			Harrington et al. 2013
Osteopilus septentrionalis			Trueb 1966
Palaeobatrachus sp.			Harrington et al. 2013
Pelobates cultripes			Harrington et al. 2013
Philautus silus			Harrington et al. 2013
Phyllomedusa vaillanti			Harrington et al. 2013
Pipa myersi			Yeh 2002
Pipa pipa		Trueb et al. 2000	Harrington et al. 2013
Pseudacris regilla			Harrington et al. 2013
Pseudacris triseriata			Harrington et al. 2013
Pseudis platensis			Harrington et al. 2013
Pseudophryne bibronii			Harrington et al. 2013
Pyxicephalus adspersus			Harrington et al. 2013
Rana (Amerana) aurora			Harrington et al. 2013

Rana (Amerana) cascadae			Harrington et al. 2013
Rana (Amerana) pretiosa			Harrington et al. 2013
Rana (Rana) temporaria			Harrington et al. 2013
Rana (Pantherana) pipiens			Kemp and Hoyt 1969
Rhinophrynus dorsalis			Harrington et al. 2013
Shomronella jordanica			Harrington et al. 2013
Smilisca baudini			Harrington et al. 2013
Spea bombifrons	Wiens 1989	Wiens 1989	Wiens 1989
Spea multiplicata			Harrington et al. 2013
Triprion petasatus			Harrington et al. 2013
Uperoleia laevigata			Harrington et al. 2013
Xenopus laevis			Harrington et al. 2013

Mammalia

Bradypus variegatus			Hautier et al. 2011
Cavia porcellus			Hautier et al. 2013
Choloepus didactylus			Hautier et al. 2011
Cryptotis parva			Koyabu et al. 2011
Cyclopes didactylus			Hautier et al. 2011
Dasypus novemcinctus			Hautier et al. 2011
Dasyurus viverrinus			Hautier et al. 2013
Didelphis albiventris		de Oliveira et al. 1998	de Oliveira et al. 1998
Echinops telfairi			Werneburg et al. 2013
Elephantulus rozeti			Hautier et al. 2013
Eremitalpa granti			Hautier et al. 2013

Erinaceus amurensis		Koyabu et al. 2011
Felis silvestris		Sánchez-Villagra et al. 2008
Homo sapiens		Hautier et al. 2013
Heterohyrax brucei		Hautier et al. 2013
Loxodonta africana		Hautier et al. 2012
Macropus eugenii		Hautier et al. 2013
Macroscelides proboscideus		Hautier et al. 2013
Manis javanica		Hautier et al. 2013
Meriones unguiculatus	Yukawa et al. 1999	Yukawa et al. 1999
Mesocricetus auratus		Hautier et al. 2013
Mogera wogura		Koyabu et al. 2011
Monodelphis domestica		Hautier et al. 2013
Mus musculus		Hautier et al. 2013
Ornithorhynchus anatinus		Weisbecker 2011
Orycteropus afer		Hautier et al. 2013
Perameles nasuta		Hautier et al. 2013
Peromyscus melanophrys		Hautier et al. 2013
Procavia capensis		Hautier et al. 2013
Rattus norvegicus		Hautier et al. 2013
Rhabdomys pumilio		Hautier et al. 2013
Rousettus amplexicaudatus		Hautier et al. 2013
Sus scrofa		Hautier et al. 2013
Tachyglossus aculeatus		Weisbecker 2011

Talpa spp.			Sánchez-Villagra et al. 2008
Tenrec ecaudatus			Werneburg et al. 2013
Tamandua tetradactyla			Hautier et al. 2011
Tarsius spectrum			Hautier et al. 2013
Trichosurus vulpecula	Weisbecker et al. 2008		Hautier et al. 2013
Tupaia javanica			Hautier et al. 2013

Squamata

Lacerta vivipara			Hautier et al. 2013
Lerista bougainvillii		Hugi et al. 2012	Hugi et al. 2012
Liopholis whitii		Hugi et al. 2012	Hugi et al. 2012
Hemiergis peronii		Hugi et al. 2012	Hugi et al. 2012
Saiphos equalis		Hugi et al. 2012	Hugi et al. 2012

Crocodylia

Alligator mississipiensis	Rieppel 1993a		Rieppel 1993a

Aves

Anas platyrhynchos			Maxwell et al. 2010
Cairina moschata			Maxwell et al. 2010
Coturnix coturnix			Maxwell et al. 2010
Coturnix coturnix (N&T)			Maxwell et al. 2010
Dromaius novaehollandiae			Maxwell et al. 2010
Dromaius novaehollandiae (YPM)			Maxwell et al. 2010

Gallus gallus			Maxwell et al. 2010
Gallus gallus (S&W)			Maxwell et al. 2010
Larus argentatus			Maxwell et al. 2010
Larus canus			Maxwell et al. 2010
Larus ridibundus			Maxwell et al. 2010
Meleagris gallopavo			Maxwell et al. 2010
Phalacrocorax auritus			Maxwell et al. 2010
Somateria mollissima			Maxwell et al. 2010
Stercorarius skua			Maxwell et al. 2010
Sterna hirundo			Maxwell et al. 2010
Struthio camelus			Maxwell et al. 2010

Testudines

Apalone spinifera			Sánchez-Villagra et al. 2008
Chelydra serpentina	Rieppel 1993b	Rieppel 1990, 1993b	Rieppel 1993b
Macrochelys temminckii			Sánchez-Villagra et al. 2008
Pelodiscus sinensis			Sánchez-Villagra et al. 2008

1145

1146

Appendix references

Adriaens D., Verraes W. 1998. Ontogeny of the osteocranium in the African Catfish, *Clarias gariepinus* Burchell (1822) (Siluriformes: Clariidae): ossification sequence as a response to functional demands. J. Morph. 235:183–237.

Boy J. 1995. Über die Micromelerpetontidae (Amphibia: Temnospondyli). 1. Morphologie und Paläoökologie des *Micromelerpeton credneri* (Unter-Perm; SW-Deutschland). Paläont. Z. 69:429–457.

Cote S., Carroll R., Cloutier R., Bar-Sagi L. 2002. Vertebral development in the Devonian sarcopterygian fish *Eusthenopteron foordi* and the polarity of vertebral evolution in non-amniote tetrapods. J. Vertebr. Paleontol. 22:487–502.

Cubbage C.C., Mabee P.M. 1996. Development of the cranium of paired fins in the Zebrafish *Danio rerio* (Ostariophysi, Cyprinidae). J. Morph. 229:121–160.

Ehmcke J., Clemen G. 2000. The structure and development of the skull of Costa Rican plethodontid salamanders (Amphibia: Urodela). Ann. Anat. 182: 537–547.

Gaudin A.J. 1978. The sequence of cranial ossification in the California toad, *Bufo boreas* (Amphibia, Anura, Bufonidae). J. Herpetol. 12:309–318.

Grande L., Bemis W.E. 1998. A comprehensive phylogenetic study of amiid fishes (Amiidae) based on comparative skeletal anatomy. An empirical search for interconnected patterns of natural history. J. Vertebr. Paleontol. 18(1, suppl.; Soc. of Vert. Paleont. Memoir 4):688 pages.

Hautier L., Weisbecker V., Goswami A., Knight F., Kardjilov N., Asher R. J. 2011. Skeletal ossification and sequence heterochrony in xenarthran evolution. Evol. Dev. 13:460–476.

Hautier L., Stansfield F. J., Allen W. R., Asher R. J. 2012. Skeletal development in the African elephant and ossification timing in placental mammals. Proc. R. Soc. B 279:2188–2195.

Hautier L., Bennett N.C., Viljoen H., Howard L., Milinkovitch M.C., Tzika A.C., Goswami

A., Asher R.J. 2013. Patterns of ossification in southern versus northern placental

mammals: mammal skeletogenesis. Evolution 67:1994–2010.

Harrington S.M., Harrison L.B., Sheil C.A. 2013. Ossification sequence heterochrony among

amphibians. Evol. Dev. 15:344–364.

Hugi J., Hutchinson M.N., Koyabu D., Sánchez-Villagra M.R. 2012. Heterochronic shifts in

the ossification sequences of surface- and subsurface-dwelling skinks are correlated with

the degree of limb reduction. Zoology 115:188–198.

Kemp N. E. and Hoyt J. 1969. Sequence of ossification in the skeleton of growing and

metamorphosing tadpoles of *Rana pipiens*. J. Morphol. 129:415–444.

Koyabu D., Endo H., Mitgutsch C., Suwa G., Catania K. C., Zollikofer C. P. E., Oda S.,

Koyasu K., Ando M., Sánchez-Villagra M. R. 2011. Heterochrony and developmental

modularity of cranial osteogenesis in lipotyphlan mammals. EvoDevo 2:1–18.

Laurin M., Germain D. 2011. Developmental characters in phylogenetic inference and their

absolute timing information. Syst. Biol. 60:630–644.

Langille R.M., Hall B.K. 1987. Development of the head skeleton of the Japanese medaka,

Oryzias latipes (Teleostei). J. Morphol. 193:135–158.

Leblanc J., Cloutier R. 2005. Developmental modularity and saltatory ontogeny in the Late

Devonian osteolepiform *Eusthenopteron foordi*. Pp. 32–84 in Leblanc J: Précisions sur

l'anatomie de l'ostéolépiforme *Eusthenopteron foordi* du Dévonien supérieur de

Miguasha, Québec. Mémoire de maîtrise (~ M.Sc. thesis), Université du Québec à

Rimouski. Available at http://semaphore.uqar.ca/283/1/Joel_Leblanc_aout2005.pdf

Lillich R., Schoch R. 2007. Finally grown up—the significance of adult *Micromelerpeton*

[abstract]. J. Vert. Paleontol. 27(3, suppl.):106A.

Lohmann U., Sachs S. 2001. Observations on the postcranial morphology, ontogeny and

palaeobiology of *Sclerocephalus haeuseri* (Amphibia: Actinodontidae) from the Lower

Permian of Southwest Germany. Mem. Queensland Mus. 46:771–781.

Maxwell E.E., Harrison L.B., Larsson H.C.E. 2010. Assessing the phylogenetic utility of

sequence heterochrony: evolution of avian ossification sequences as a case study.

Zoology 113:57–66.

Müller H., Oommen O.V., Bartsch P. 2005. Skeletal development of the direct-developing

caecilian *Gegeneophis ramaswamii* (Amphibia: Gymnophiona: Caeciliidae).

Zoomorphology 124:171–188.

Müller H. 2006. Ontogeny of the skull, lower jaw, and hyobranchial skeleton of *Hypogeophis*

rostratus (Amphibia: Gymnophiona: Caeciliidae) revisited. J. Morph. 267:968–986.

de Oliveira C.A., Nogueira J.C., Mahecha G.A.B. 1998. Sequential order of appearance of

ossification centers in the opossum Didelphis albiventris (Didelphidae) skeleton during

development in the Marsupium. Ann. Anat. 180:113–121. [Lack of italics in the

original.]

Olori, J.C. 2013. Ontogenetic sequence reconstruction and sequence polymorphism in extinct

taxa: an example using early tetrapods (Tetrapoda: Lepospondyli). Paleobiology

39:400–428.

Púgener L.A., Maglia A.M. 1997. Osteology and skeletal development of *Discoglossus*

sardus (Anura: Discoglossidae). J. Morphol. 233:267–286.

Reilly S.M. 1986. Ontogeny of cranial ossification in the eastern newt, *Notophthalmus*

viridescens (Caudata: Salamandridae), and its relationship to metamorphosis and

neoteny. J. Morphol. 188:315–326.

Reilly S.M. 1987. Ontogeny of the hyobranchial apparatus in the salamanders *Ambystoma*

talpoideum (Ambystomatidae) and *Notophthalmus viridescens* (Salamandridae): the

ecological morphology of two neotenic strategies. J. Morphol. 191:205–214.

Reilly S.M., Altig R. 1996. Cranial ontogeny in *Siren intermedia* (Amphibia: Sirenidae):

Paedomorphic, metamorphic, and novel patterns of heterochrony. Copeia 1996:29–41.

Rieppel O. 1990. The structure and development of the jaw adductor musculature in the turtle

Chelydra serpentina. Zool. J. Linn. Soc. 98:27–62.

Rieppel O. 1993a. Studies on skeleton formation in reptiles. V. Patterns of ossification in the

skeleton of *Alligator mississippiensis* Daudin (Reptilia, Crocodylia). Zool. J. Linn. Soc.

109:301–325.

Rieppel O. 1993b. Studies on skeleton formation in reptiles: Patterns of ossification in the

skeleton of *Chelydra serpentina* (Reptilia, Testudines). J. Zool. (Lond.) 231:487–509.

Sánchez-Villagra M., Goswami A.,Weisbecker V., Mock O., Kuratani S. 2008. Conserved

relative timing of cranial ossification patterns in early mammalian evolution. Evol.

Dev. 10:519–530.

de Sá R.O. 1988. Chondrocranium and ossification sequence of *Hyla lanciformis*. J. Morphol.

195:345–355.

de Sá R.O., Swart C.C. 1999. Development of the suprarostral plate of pipoid frogs. J.

Morphol. 240:143–153.

Schoch R.R. 2004. Skeleton formation in the Branchiosauridae: a case study in comparing

ontogenetic trajectories. J. Vert. Paleontol. 24:309–319.

Schoch R.R. 2003. Early larval ontogeny of the Permo-Carboniferous temnospondyl

Sclerocephalus. Palaeontology 46:1055–1072.

Schoch R.R. 2009. Evolution of life cycles in early amphibians. Annu. Rev. Earth Planet. Sci.

37:125–162.

Schoch R.R., Witzmann F. 2009. Osteology and relationships of the temnospondyl genus

Sclerocephalus. Zool. J. Linn. Soc. 157:135–168.

Trueb L. 1966. Morphology and development of the skull in the frog *Hyla septentrionalis*.

Copeia 1966:562–573.

Trueb L., Pugener L.A., Maglia A.M. 2000. Ontogeny of the bizarre: An osteological

description of *Pipa pipa* (Anura: Pipidae), with an account of skeletal development in

the species. J. Morphol. 243:75–104.

Wake T.A., Wake D.B., Wake M.H. 1983. The ossification sequence of *Aneides lugubris*,

with comments on heterochrony. J. Herpetol. 17:10–22.

Weisbecker V., Goswami A., Wroe S., Sánchez-Villagra M.R. 2008. Ossification

heterochrony in the therian postcranial skeleton and the marsupial–placental dichotomy.

Evolution 62:2027–2041.

Weisbecker V. 2011. Monotreme ossification sequences and the riddle of mammalian skeletal

development. Evolution 65:1323–1335.

Werneburg R. 2018 (for 2017). Earliest 'nursery ground' of temnospondyl amphibians in the

Permian. Semana 32:3–42.

Werneburg I., Tzika A.C., Hautier L., Asher R.J., Milinkovitch M.C., Sánchez-Villagra M.R.

2013. Development and embryonic staging in non-model organisms: the case of an

afrotherian mammal: Development and embryonic staging. J. Anat. 222:2–18.

Wiens J.J. 1989. Ontogeny of the skeleton of *Spea bombifrons* (Anura: Pelobatidae). J.

Morphol. 202:29–51.

Witzmann F. 2006. Developmental patterns and ossification sequence in the Permo-

Carboniferous temnospondyl *Archegosaurus decheni* (Saar-Nahe Basin, Germany). J.

Vert. Paleontol. 26:7–17.

Witzmann F., Pfretzschner H.-U. 2003. Larval ontogeny of *Micromelerpeton credneri*

(Temnospondyli, Dissorophoidea). J. Vert. Paleontol. 23:750–768.

Yeh J. 2002. The evolution of development: two portraits of skull ossification in pipoid frogs.

Evolution 56:2484–2498.

Yukawa M., Hayashi N., Takagi K., Mochizuki K. 1999. The normal development of

Mongolian Gerbil foetuses and, in particular, the timing and sequence of the appearance

of ossification centres. Anat. Histol. Embryol. 28:319–324.

1275

1276

1277   SUPPLEMENTARY MATERIAL

1278   Data matrix in NEXUS format for Mesquite.

1279