Running head:    LISSAMPHIBIAN ORIGIN AND OSSIFICATION SEQUENCES

Title:

What do ossification sequences tell us about the origin of extant amphibians?

Michel Laurin[1,]*, Océane Lapauze[1], David Marjanović[2]

[1] *CR2P (Centre de Recherche sur la Paléodiversité et les Paléoenvironments; UMR 7207),*

*CNRS/MNHN/UPMC–Sorbonne Universités, Muséum national d'Histoire naturelle,*

*Département Histoire de la Terre, 57 rue Cuvier, F-75231 Paris cedex 05, France;* [2] *Museum*

*für Naturkunde (Leibniz Institute for Evolutionary and Biodiversity Research),*

*Invalidenstraße 43, D-10115 Berlin, Germany, david.marjanovic@gmx.at*

*\*Correspondence to be sent to: Muséum national d'Histoire naturelle, Département Histoire*

*de la Terre, 57 rue Cuvier, F-75231 Paris cedex 05, France; michel.laurin@mnhn.fr*

ABSTRACT—The controversial origin of extant amphibians has been studied using several

sources of data and methods, including phylogenetic analyses of morphological data,

molecular dating, stratigraphic data, and comparisons of ossification sequences. A consensus

has failed to emerge, perhaps because the diversity of methods used hampers comparisons.

We have compiled a dataset of ossification sequences of 102 terminal taxa and seven cranial

bones to assess the relative support for six competing hypotheses about the origin of extant

amphibians: a monophyletic origin among temnospondyls, a monophyletic origin among

lepospondyls, a diphyletic origin among both temnospondyls and lepospondyls, a diphyletic

origin among temnospondyls alone, and two variants of a triphyletic origin, in which anurans

and urodeles come from different temnospondyl taxa while caecilians come from

lepospondyls and are either closer to anurans and urodeles or to amniotes. The data are

analyzed through maximum likelihood, and the AICc (corrected Akaike Information

Criterion) weights of the six hypotheses allow us to assess their relative support. By an

unexpectedly large margin, our analyses support a monophyletic origin among lepospondyls;

a monophyletic origin among temnospondyls, the current near-consensus, is a distant second.

All other hypotheses are exceedingly unlikely according to our data.

**Keywords:** macroevolution; paleontology; evo-devo; ossification sequences; Lissamphibia;

Tetrapoda; phylogeny

## Introduction

Paleontologists have been studying the origin of the extant amphibian clades for more than a

century. Early studies generally proposed an origin of at least some extant amphibians from

temnospondyls. Cope (1888) initially suggested that batrachians (anurans and urodeles)

derived from temnospondyls (a large clade of limbed vertebrates known from the Early

Carboniferous to the Early Cretaceous) because he believed that the batrachian vertebral

centrum was an intercentrum, the dominant central element of temnospondyls. Later, Watson

(1940) argued that anurans were derived from temnospondyls because of similarities (mostly

in the palate) between the temnospondyl "*Miobatrachus*" (now considered a junior synonym

of *Amphibamus*) and anurans. Monophyly of extant amphibians (Lissamphibia) was proposed

by Parsons and Williams (1962, 1963), an idea that was accepted more quickly by

herpetologists than by paleontologists. Lissamphibian monophyly was supported by (among a

few other character states) the widespread occurrence of pedicellate, bicuspid teeth. The

subsequent discovery of such teeth in the amphibamid temnospondyl *Doleserpeton* (Bolt

1969) reinforced the widespread (though not universal) acceptance of an origin of

Lissamphibia from within temnospondyls (e.g., Schoch and Milner 2004). Recently, this

hypothesis, referred to as the temnospondyl hypothesis or TH for short (Fig. 1c), has been

supported by several phylogenetic analyses based on phenotypic data matrices (e.g. Ruta and

Coates 2007; Sigurdsen and Green 2011; Maddin et al. 2012; Pardo et al. 2017a, b: fig. S6).

Dissenting opinions about the origin of extant amphibians have been expressed for

several decades (see Schoch and Milner 2004 for a historical review). These were initially

formulated especially for the urodeles and caecilians, which are less similar to temnospondyls

and lack a tympanic middle ear (which is present in most anurans and often inferred for at

least some temnospondyls but absent in lepospondyls). Thus, Steen (1938) highlighted

similarities in the palate (broad cultriform process of the parasphenoid) and cheek (loss of

several bones) between lysorophian lepospondyls and urodeles. Carroll and Currie (1975) and

Carroll and Holmes (1980) argued that the exant amphibians had three distinct origins among

early stegocephalians; while they accepted an origin of anurans among temnospondyls, they

suggested that urodeles and caecilians originated from two distinct groups of lepospondyls

(*Rhynchonkos* for caecilians, Hapsidopareiidae for urodeles). Later, based mostly on

developmental similarities between the temnospondyl *Apateon* and urodeles, Carroll (2001,

2007) and Fröbisch et al. (2007) even proposed a second hypothesis involving a triphyletic

origin of lissamphibians, with an origin of anurans and urodeles from two distinct

temnospondyl groups, while the caecilians would remain in the lepospondyl clade. This is

what we call the polyphyly hypothesis (PH). We have tested two versions: the one (called

PH1; Fig. 1e) cautiously suggested by Fröbisch et al. (2007), which agrees with the

paleontological consensus in placing all or most lepospondyls closer to Amniota than to

Temnospondyli (Fig. 1b; Sigurdsen and Green 2011; Pardo et al. 2017a, b: fig. S6;

Marjanović and Laurin 2018), and a version modified (PH2; Fig. 1f) to make Lissamphibia

monophyletic with respect to Amniota, a fact we consider demonstrated beyond reasonable

doubt by multiple molecular phylogenetic analyses (Fig. 1a; Pyron 2014; Irisarri et al. 2017;

Feng et al. 2017; and references cited therein). Anderson (2007) and Anderson et al. (2008)

found lissamphibian diphyly, specifically a monophyletic Batrachia among the

temnospondyls while keeping the caecilians among the lepospondyls (DH1; Fig. 1g). Pardo et

al. (2017b: fig. 2, S7) presented a similar hypothesis, with batrachians and caecilians having

separate origins within the temnospondyls (DH2; Fig. 1h). Further, a monophyletic origin of

all extant amphibians among lepospondyls has also been proposed (Laurin 1998; Pawley

2006: appendix 16; Marjanović and Laurin 2009, 2013, 2018). This will be referred to below

as the lepospondyl hypothesis (LH; Fig. 1d).

Phylogenetic analyses of molecular data cannot distinguish the TH, the PH2, the DH2

or the LH from each other by topology (Fig. 1). Several other types of data and methods have,

however, been used to try to discriminate between the various hypotheses on the origin of

extant amphibians. In addition to classical phylogenetic analyses of morphological data

matrices, these include the use of molecular (Zhang et al. 2005; Marjanović and Laurin 2007;

Pardo et al. 2017b) and stratigraphic data (Marjanović and Laurin 2008) to compare the

inferred divergence dates between the three main extant amphibian clades with predictions

based on the TH and the LH on the one side and the PH and the DH on the other. However,

developmental data, in the form of ossification sequences, have been the second-most

frequently used (after classical morphological data) to argue for particular phylogenetic

hypotheses. These data include mainly cranial (e.g. Schoch 2002, 2006; Schoch and Carroll

2003; Schoch and Milner 2004; Anderson 2007; Carroll 2007; Germain and Laurin 2009) and

autopodial ossification sequences (e.g. Fröbisch et al. 2007, 2015). Ossification sequences of

other parts of the skeleton, like the vertebrae, shoulder girdle and scales, are also documented

in a few Paleozoic stegocephalians (e.g. Carroll et al. 1999; Witzmann and Schoch 2006;

Anderson 2007; Carroll 2007; Olori 2013), but these have played a minor role in the

controversy about the origin of extant amphibians. This study will emphasize cranial

ossification sequences because these are available for the greatest number of extant taxa.

METHODS

*Ossification sequence data*

From all the literature we could get access to, we compiled the most extensive

database on ossification sequences for osteichthyans that exists to date. The most useful

sources for extant taxa included compilations: Harrington et al. (2013) for amphibians, Hugi

et al. (2012) for squamates, Maxwell et al. (2010) for birds, and Weisbecker (2011) for

mammals. For the temnospondyl *Apateon* (*A. caducus* and *A. pedestris*), data come from

Schoch (2004); the partial cranial sequence for the temnospondyl *Sclerocephalus* (*S.*

*haeuseri*) comes from Werneburg (2018). The other sources of sequence data can be found in

SM (Supplementary Material) 1. The data themselves and the phylogenetic trees

corresponding to the tested hypotheses are included in SM 2.

The software we used to compute AICc weights (see below) cannot handle missing

data. This unfortunately meant we had to discard much information. In order to keep as many

taxa as possible in the analysis, we first compiled a matrix (not shown) of 153 taxa and 216

characters. Of these, we first kept characters that were scored in at least 75% of the taxa in our

initial database, and taxa that were scored for at least six characters out of the seven that are

scored in more than 90% of the taxa. All seven of these characters are cranial; postcranial

characters are systematically scored in fewer than 25% of the taxa, so we could not use them

here. In the end, we were left with 102 taxa and only seven characters (all cranial: nasal,

parietal, squamosal, premaxilla, maxilla, pterygoid, and exoccipital). This data loss is not as

severe as it may first seem because many excluded characters (most of them postcranial) had

less than 10% scored cells (sometimes less than 1%), and most of them could not be scored

for the temnospondyls *Apateon* and *Sclerocephalus*, so they could not have helped resolve the

main question examined in this study. This reduced dataset used in our analyses is included in

SM 2.

Of these seven characters, *Sclerocephalus* cannot presently be scored for the

squamosal. Because of the potential importance of *Sclerocephalus* as a stem-caecilian

according to the DH2 (Fig. 1h) and as one of only three sampled extinct taxa, we ran all

analyses twice: once with *Sclerocephalus* and six characters, once without *Sclerocephalus* and

with seven characters. The first set of analyses is presented in Tables 1 and 3; the second set

gave very similar results and is presented in Tables 2 and 4.

Due to the scarcity of clear homologies between the skull bones of tetrapods and

actinopterygians, we had to omit all actinopterygians from our analyses. As cranial

ossification sequences remain unknown for finned sarcopterygians (except perhaps lungfish,

whose skull bones seem mostly impossible to homologize), our analyses are restricted to

limbed vertebrates.

Unfortunately, no cranial ossification sequence is available for any lepospondyl or

possible lepospondyl except the highly apomorphic aïstopod *Phlegethontia*, for which three

cranial ossification stages are known (Anderson et al. 2003; Anderson 2007). A recent

anatomical restudy and phylogenetic analysis suggests that aïstopods are not lepospondyls

(Pardo et al., 2017a); the resulting uncertainty about the position of the aïstopods (among

lepospondyls or on the tetrapod stem) prevents us from using the developmental sequence of

*Phlegethontia* here. The low taxon sample is more limiting for this analysis than the low

character sample. However, as explained below, the absence of lepospondyl sequences in our

dataset does not preclude testing the six hypotheses (TH, PH1, PH2, DH1, DH2, LH).

Given that various taxa differ in the numbers of bones and that the resolution of the

sequences is also variable between taxa, these data needed to be standardized to make

comparisons and computations meaningful, as suggested by Germain and Laurin (2009). For

this, we used the position in the sequence (from first to last) and standardized this relative

sequence position between 0 and 1 using the simple formula given by Germain and Laurin

(2009). The standardized sequence position ($X_s$) is:

$X_s = (X_i - X_{min})/(X_{max} - X_{min})$,

where:

$X_i$ is the position of a given bone in the sequence

$X_{min}$ is the lowest position in the sequence (generally denoted 0 or 1)

$X_{max}$ is the highest position in the sequence (for instance, if there are 20 bones, $X_{min}$ is 1 and

the sequence is completely resolved, $X_{max} = 20$).

We also experimented with using size (skull length) or developmental stage as standards, but

this led to lower resolution (results not shown), so we worked only with sequences

standardized using sequence position.

Our ossification sequence data of extant and extinct taxa and the phylogenetic trees we

used are available online in the supplements to this paper on the bioRxiv web site.

*Analytical methods*

To discriminate between the six hypotheses about the origin of extant amphibians, two

methods are available: direct phylogenetic analysis of the sequence data, and comparisons of

the tree length (number of steps in regular parsimony, squared length in squared-change

parsimony, or similar measures) of various trees selected a priori to represent these

hypotheses. We selected the latter approach because relatively few data are available, and

thus, phylogenetic analysis of such data is unlikely to provide a well-resolved tree.

We used the CoMET module (Lee et al. 2006) for Mesquite 3.2 (Maddison and

Maddison 2017) to test the relative fit of the data on trees representing the six hypotheses.

CoMET calculates the likelihood and the AIC (Akaike Information Criterion) of nine

evolutionary models given continuous data and a tree. The nine models are obtained by

modifying the branch lengths of the reference tree. Thus, branches can be set to 0 (for internal

branches only, to yield a non-phylogenetic model), to 1 (equal or speciational model), left

unchanged from their original length (gradual evolution), or set free and evaluated from the

data (free model). This can be applied to internal and/or external branches, and various

combinations of these yield nine models (Lee et al. 2006: fig. 1). Among these nine models

two have been frequently discussed in the literature and are especially relevant: gradual

evolution, in which branch lengths (here representing evolutionary time) have not been

changed, and a speciational model, in which all branches are set to the same length, and which

has some similarities with Gould and Eldredge's (1972) punctuated equilibria model (though

a model with one internal branch stemming from each note set to 0 and the other set to 1

would be even closer to the original formulation of that model). In this study, we assessed the

fit of six of the nine models covered by CoMET; the other three (the punctuated version of

distance, equal and free) could not be assessed due to problems in the current version of

CoMET and possibly the size of our dataset (the corresponding author T. Oakley, pers.

comm., has not yet been able to look into them).

Provided that the same evolutionary model is optimal for all compared hypotheses

(this condition is met, as shown below), the AIC weights of the various trees under the same

model can be used to assess the support for each tree. This can show not only which tree is

best supported, but how many times more probable the best tree is compared to the

alternatives – another reason to prefer this approach over a phylogenetic analysis. Given the

small sample size (which here is the number of characters), we computed the corrected AIC

(AICc) and the AICc weights using the formulae given by Anderson and Burnham (2002) and

Wagenmakers and Farrell (2004).

Our tests make sense only in the presence of a phylogenetic signal in the data. In

addition to the test of evolutionary model in CoMET evoked above (which tests non-

phylogenetic as well as phylogenetic models), we performed a test based on squared-change

parsimony (Maddison 1991) and random taxon reshuffling (Laurin 2004). For this test, we

compared the length of the LH reference tree (with and without *Sclerocephalus*) to a

population of 10,000 random trees produced by taxon reshuffling.

*Reference phylogenies*

We built a reference timetree that attempts to capture the most consensual relationships (Fig.

2). The tree was compiled in Mesquite (Maddison and Maddison 2017) and time-calibrated

using the Stratigraphic Tools (Josse et al. 2006). For consistency and to avoid the effects of

gaps in the fossil record, we used molecular divergence dates whenever possible. The main

sources we used for topology and divergence times are as follows:

The phylogeny of lissamphibians follows the work of Pyron (2014). However, several

data bases have been used for the temporal calibration of the tree: Germain and Laurin (2009)

was used for the urodeles, whereas Feng et al. (2017), supplemented by Bossuyt and Roelants

(2009) and Pyron (2014), was used for the anurans as well as more rootward nodes

(Batrachia, Lissamphibia, Tetrapoda; also Amniota). Marjanović and Laurin (2014) was used

for the Ranidae, Ceratophryidae, and Hylidae.

The sediments that have preserved the temnospondyls *Apateon* and *Sclerocephalus* are

not easy to correlate with each other or with the global chronostratigraphic scale. Combining

stratigraphic information from Schoch (2014a), Schneider et al. (2015) and Werneburg

(2018), we have placed all three sampled species (*A. pedestris*, *A. caducus*, *S. haeuseri*) at the

Sakmarian/Artinskian stage boundary (Permian; 290.1 Ma ago); combining stratigraphic

information from Schneider et al. (2015) with the phylogeny in Schoch (2014a), we have

tentatively placed the divergence between the two *Apateon* species (which are not sister-

groups: Schoch 2014a) at the Kasimovian/Gzhelian stage boundary (Carboniferous; 303.7 Ma

ago). The age of the last common ancestor of *Apateon* and *Sclerocephalus* depends strongly

on temnospondyl phylogeny, which remains unresolved (Pardo et al. 2017b; Marjanović and

Laurin 2018; and numerous references in both); as a compromise between the different

options, we have provisionally placed it at the boundary between the Early and the Late

Carboniferous (Serpukhovian/Bashkirian, 323.2 Ma ago) where applicable.

For the birds, Pons et al. (2005) was used for the Laridae, Wang et al. (2013) for the

Phasianidae and Gonzales et al. (2009) for the Anatidae. The temporal calibration was taken

from Prum et al. (2015) as recommended by Berv and Field (2017); gaps were filled in using

the database http://www.birdtree.org.

Several papers, mainly Tarver et al. (2016), were used for the phylogeny and

divergence times of mammals. For the Muridae, two references were used: Lecompte et al.

(2008) and the website "Mammals species of the World" (http://vertebrates.si.edu) for the

position of two taxa: *Mesocricetus auratus* and *Peromyscus melanophrys.* Other species were

placed following the work of Meredith et al. (2011), which also gives divergence times. We

caution, however, that all available molecular dates for Paleogene and earlier mammal nodes

may be overestimates (Berv and Field 2017).

Three references were also used to integrate squamates in the phylogenetic tree and for

the calibration of divergence times: Brandley et al. (2005), Rabosky et al. (2014), Reeder

(2003). Sterli et al. (2013) was used for turtles.

For turtles, there is now a near-consensus that they are diapsids, a hypothesis that is

not necessarily incompatible with an origin among "parareptiles" (Laurin and Piñeiro 2017).

Thus, following most recent molecular phylogenies (e.g., Hugall et al. 2007; Irisarri et al.

2017), we have inserted them as the sister-group of Archosauria.

Given that we disagree with several of the calibration dates in Irisarri et al. (2017), we

have not used divergence dates from that source.

To discriminate between the hypotheses on lissamphibian origins, we inserted the

temnospondyl *Apateon* in the tree where each predicts that it should be (Fig. 1c–h). Thus,

according to the TH (Fig. 1c), *Apateon* lies on the lissamphibian stem (Ruta and Coates 2007;

Anderson 2007). Under the LH (Fig. 1d), *Apateon* lies on the tetrapod stem (Laurin 1998).

Under both versions of the DH (Fig. 1g, h), *Apateon* lies on the batrachian stem (Pardo et al.

2017b: fig. 2, S7). Under both versions of the PH (Fig. 1e, f), *Apateon* lies on the caudate

stem (Carroll 2007; Fröbisch et al. 2007). Within the DH and the PH, both versions of each

differ in the position of Gymnophiona. Thus, despite the absence of any lepospondyl in our

ossification sequence dataset, our taxonomic sample allows us to test all competing

hypotheses.

*Sclerocephalus* is the sister-group of *Apateon* under the LH (Fig. 1d), immediately

rootward of it (on the lissamphibian stem) under the TH (Fig. 1c) and likewise (but on the

batrachian stem) under the DH1 (Fig. 1g), on the caecilian stem under the DH2 (Fig. 1h) and

the sister-group of Batrachia (including *Apateon*) under both versions of the PH (Fig. 1e, f).

To ensure that our analyses were not biased in favor of a given hypothesis, in case that

a continuous evolutionary model were favored, we adjusted the branch lengths such that the

sum of branch lengths was equal between the compared topologies.

RESULTS

The phylogenetic speciational model (in which all branch lengths are equal) had

overwhelming support whether or not *Sclerocephalus* (Table 1) or the squamosal (Table 2)

were included; the five other examined models had AICc weights $< 10^{-11}$. Thus, from here on,

we compare support of the six competing topologies using only this model. This simplifies

the discussion, because it means that the original branch lengths are irrelevant (under that

model, all branch lengths are equal). Thus, the only variable is the topology. This, along with

the test based on squared-change parsimony and random taxon reshuffling indicates that the

phylogenetic signal in these data is strong; the latter test (Laurin, 2004) indicates that the

probability that the observed covariation between the data and the tree reflects a random

distribution is less than 0.0001 (none of the 10,000 random trees generated were as short as

the reference tree).

Using the speciational model, the AICc weights of the six compared topologies

indicate that there is strong support for the LH, with an AICc weight of 0.987 when

*Sclerocephalus* is included (Table 3) and 0.895 when the squamosal is included instead (Table

4). Of the other topologies, the TH was by far the best supported, with an AICc weight of

0.013 (with *Sclerocephalus*) or 0.094 (with the squamosal), which is 75.11 or 9.54 times less

than for the LH. Both versions of the DH and of the PH have negligible support (AICc

weights < 0.01 when the squamosal is included, < 0.0001 when *Sclerocephalus* is included).

The least support is found for the PH2 when *Sclerocephalus* is included, and for the DH1

when the squamosal is included. In both cases, the recently proposed DH2 (Pardo et al.

2017b) fares second-worst by a small margin. Notably, the DH1 contradicts the modern

consensus on lissamphibian monophyly (Fig. 1g), while the PH2 and the DH2 fulfill this

constraint from the molecular but not the paleontological point of view, having lissamphibian

monophyly with respect to amniotes but not with respect to temnospondyls (Fig. 1f, h).

DISCUSSION

*Phylogenetic signal*

In his discussion of previous attempts to draw phylogenetic conclusions from ossification

sequences (e.g. Schoch and Carroll 2003), Anderson (2007) noted that ossification sequences

seemed to abound in symplesiomorphies and in autapomorphies of terminal taxa, while

potential synapomorphies were scarce. This pessimism was seemingly confirmed by Schoch

(2006) in a paper that was published after Anderson's (2007) book chapter had gone to press:

not only were many similarities in the cranial ossification sequences across Osteichthyes

found to be symplesiomorphies, but a phylogenetic analysis of cranial ossification sequences

did not recover Mammalia, Sauropsida, Amniota or Lissamphibia as monophyletic. Along

with these untenable results, Schoch (2006) dismissed another: the position of the

temnospondyl *Apateon caducus* (the only included extinct taxon) outside the tetrapod crown-

group, i.e. the lepospondyl hypothesis on lissamphibian origins (LH).

While ossification sequences alone may not provide enough data for a phylogenetic

analysis, our datasets (with much larger taxon samples than in Schoch 2006) fit some tree

topologies much better than others. Both the tests using CoMET and squared-change

parsimony with random taxon reshuffling overwhelmingly support the presence of a

phylogenetic signal in these data; the null hypothesis of the absence of a phylogenetic signal

can be rejected in both cases, given that it has a probability of $< 10^{-97}$ in the first case, and $<$

$10^{-4}$ in the second case. We conclude that it contains a strong phylogenetic signal, and are

therefore cautiously optimistic about future contributions of ossification sequences to

phylogenetics.

*Suggestions for future extensions and uses of our dataset*

The postcranial ossification sequences of the lepospondyls *Microbrachis* and *Hyloplesion*

were described by Olori (2013). While they feature lower resolution than most sequences

used here, a study comparable to ours but focusing on postcranial elements could be carried

out in the future, if only with fewer taxa (as long as missing data remain a problem). This

would be worth doing because postcranial ossification sequences, like cranial ones, have been

used as arguments in the discussion on lissamphibian origins.

Specifically, Carroll et al. (1999) stated that the neural arches ossify before the centra

in frogs and temnospondyls, but not in salamanders, caecilians or lepospondyls. When it was

found that the centra do ossify first in a few cryptobranchoid salamanders, Carroll (2007: 30)

took this as "strong evidence that the most primitive crown-group salamanders had a sequence

of vertebral development that is common to frogs and labyrinthodonts (but distinct from that

of lepospondyls)". In fact, apart from tail regeneration in *Hyloplesion* and *Microbrachis*

(where the centra ossify before the neural arches: Olori 2015; Fröbisch et al. 2015; van der

Vos et al. 2017), only one incompletely ossified vertebral column (referred to *Utaherpeton*) is

known of any putative lepospondyl. "In this specimen, […] five neural arches […] have

ossified behind the most posterior centrum." (Carroll and Chorn 1995: 40–41) Carroll's

(2007: 85) claim that "the centra always ossified prior to the arches" in lepospondyls is

therefore rather puzzling.

Fröbisch et al. (2007, 2015) pointed out that the first two digital rays (digits,

metapodials and distal carpals/tarsals) ossify before the others ("preaxial polarity") in

salamanders and the temnospondyls *Apateon*, *Micromelerpeton* and *Sclerocephalus*, while the

fourth ossifies first ("postaxial polarity") in amniotes, frogs and "probably" (Fröbisch et al.

2015: 233, 234) the lepospondyls *Microbrachis* and *Hyloplesion*. This latter inference,

however, is based only on a delay in the ossification of the fifth ray that is shared specifically

with sauropsid amniotes (Olori 2015). Ossification sequences (however partial) of the other

four rays in any lepospondyl are currently limited to the tarsus of *Batropetes*, which clearly

shows preaxial polarity (Glienke 2015: fig. 6O–S; Marjanović and Laurin 2018), and that of

the putative lepospondyl *Sauropleura*, in which likewise the second distal tarsal ossified

before all others (Marjanović and Laurin 2018). Outside of temno- and lepospondyls,

Marjanović and Laurin (2013, 2018) presented evidence that preaxial polarity is

plesiomorphic, widespread and dependent on the use of the still developing limbs for

locomotion, which would explain why it was independently lost in amniotes and frogs and

reduced (the third ray ossifies first) in direct-developing salamanders.

Given the phylogenetic signal we have found in cranial ossification sequences, we

think that ossification sequence data should eventually be added to phenotypic datasets for

analyses of tetrapod phylogeny. Indeed, an analysis of amniote phylogeny using data from

organogenesis sequences (coded using event-pairing in Parsimov) already exists (Werneburg

and Sánchez-Villagra 2009). The usefulness of such data for phylogenetic inference was

further tested, with encouraging results, by Laurin and Germain (2011), and the present

analysis adds additional support for it.

*On the origin of lissamphibians, or the preservation of favoured lepospondyls in the struggle*

*for life*

The strong support for the lepospondyl hypothesis that we have found is surprising because

cranial ossification sequence data, especially those of the Permo-Carboniferous temnospondyl

*Apateon*, have often been claimed to contradict the LH; especially, similarities between

*Apateon* and extant urodeles, in particular the supposedly "primitive" hynobiid *Ranodon*, have

often been emphasized (Schoch and Carroll 2003; Schoch and Milner 2004; Carroll 2007;

Schoch 2014b). However, other studies have already raised doubts about some of these claims

(e.g. Schoch 2006; Anderson 2007; Germain and Laurin 2009). Schoch (2006) and Anderson

(2007) concluded that most characters shared between *Apateon* and urodeles were

plesiomorphies. Germain and Laurin (2009) also demonstrated that, far from being very

similar to the ancestral urodele morphotype (contra Schoch and Carroll 2003 or Carroll 2007),

the cranial ossification sequence of *Apateon* was statistically significantly different from that

of the hypothetical last common ancestor of all urodeles (as suspected by Anderson 2007).

However, these earlier studies did not clearly show which of the various hypotheses on

lissamphibian origins the ossification sequences of *Apateon* – or the newly available partial

sequence (Werneburg 2018) of the phylogenetically distant temnospondyl *Sclerocephalus* –

supported most. This is what we have attempted to do here.

We conclude that cranial ossification sequences of amniotes, lissamphibians and

temnospondyls provide support for the LH that is independent of the phylogenetic analyses of

Laurin (1998), Pawley (2006: appendix 16) or Marjanović and Laurin (2009, 2018).

FUNDING

This work was supported by the Centre National de la Recherche Scientifique; and the French

Ministry of Research (unnumbered recurring grants to the CR2P, for ML).

ACKNOWLEDGEMENTS

D. M. would like to thank Ralf Werneburg for an electronic reprint of his 2018 paper, and

Daniel Field for discussion of molecular divergence times and the fossil record.

REFERENCES

Anderson J.S. 2007. Incorporating ontogeny into the matrix: a phylogenetic evaluation of

developmental evidence for the origin of modern amphibians. Pages 182–227 in

Anderson J.S., Sues H.-D., editors. Major transitions in vertebrate evolution.

Bloomington: Indiana University Press.

Anderson J.S., Carroll R.L., Rowe T.B. 2003. New information on *Lethiscus stocki*

(Tetrapoda: Lepospondyli: Aistopoda) from high-resolution computed tomography and

a phylogenetic analysis of Aistopoda. Can. J. Earth Sci. 40:1071–1083.

Anderson D.R., Burnham K.P. 2002. Avoiding pitfalls when using information-theoretic

methods. J. Wildl. Manag. 66:912–918.

Anderson J.S., Reisz R.R., Scott D., Fröbisch N.B., Sumida S.S. 2008. A stem batrachian

from the Early Permian of Texas and the origin of frogs and salamanders. Nature

453:515–518.

Berv J.S., Field D.J. 2017 (printed 2018). Genomic signature of an avian Lilliput Effect across

the K-Pg Extinction. Syst. Biol. 67:1–13.

Bolt J.R. 1969. Lissamphibian origins: possible protolissamphibian from the Lower Permian

of Oklahoma. Science 166:888–891.

Bossuyt F., Roelants K. 2009. Frogs and toads (Anura). Pages 357–364 in Hedges S.B.,

Kumar S., editors. The Timetree of Life. New York: Oxford University Press.

Brandley M.C., Schmitz A., Reeder T.W. 2005. Partitioned Bayesian analyses, partition

choice, and the phylogenetic relationships of scincid lizards. Syst. Biol. 54:373-390.

Carroll R.L. 2001. The origin and early radiation of terrestrial vertebrates. J. Paleontol.

75:1202–1213.

Carroll R.L. 2007. The Palaeozoic ancestry of salamanders, frogs and caecilians. Zool. J.

Linn. Soc. 150 (suppl. 1):1–140.

Carroll R.L., Chorn J. 1995. Vertebral development in the oldest microsaur and the problem

of "lepospondyl" relationships. J. Vert. Paleont. 15:37–56.

Carroll R.L., Currie P.J. 1975. Microsaurs as possible apodan ancestors. Zool. J. Linn. Soc.

57:229–247.

Carroll R.L., Holmes R. 1980. The skull and jaw musculature as guides to the ancestry of

salamanders. Zool. J. Linn. Soc. 68:1–40.

Carroll R.L., Kuntz A., Albright K. 1999. Vertebral development and amphibian evolution.

Evol. Dev. 1:36–48.

Cope E.D. 1888. On the intercentrum of the terrestrial Vertebrata. Trans. Am. Phil. Soc.

16:243–253.

Eldredge N., Gould S.J. 1972. Punctuated equilibria: an alternative to phyletic gradualism.

Pages 82–115 in Schopf T.J.M., editor. Models in Paleobiology. San Francisco:

Freeman, Cooper & Company.

Feng Y.-J., Blackburn D.C., Liang D., Hillis D.M., Wake D.B., Cannatella D.C., Zhang P.

2017. Phylogenomics reveals rapid, simultaneous diversification of three major clades

of Gondwanan frogs at the Cretaceous–Paleogene boundary. Proc. Natl. Acad. Sci. USA

E5864–E5870.

Fröbisch N.B., Carroll R.L., Schoch R.R. 2007. Limb ossification in the Paleozoic

branchiosaurid *Apateon* (Temnospondyli) and the early evolution of preaxial dominance

in tetrapod limb development. Evol. Dev. 9:69–75.

Fröbisch N.B., Bickelmann C., Olori J.C., Witzmann F. 2015. Deep-time evolution of

regeneration and preaxial polarity in tetrapod limb development. Nature 527:231–234.

Germain D., Laurin M. 2009. Evolution of ossification sequences in salamanders and urodele

origins assessed through event-pairing and new methods. Evol. Dev. 11:170–190.

Glienke S. 2015. Two new species of the genus *Batropetes* (Tetrapoda, Lepospondyli) from

the Central European Rotliegend (basal Permian) in Germany. J. Vert. Paleont.

35:e918041.

Gonzalez J., Düttmann H., Wink M. 2009. Phylogenetic relationships based on two

mitochondrial genes and hybridization patterns in Anatidae. J. Zool. 279:310–318.

Harrington S.M., Harrison L.B., Sheil C.A. 2013. Ossification sequence heterochrony among

amphibians. Evol. Dev. 15:344–364.

Hugall A.F., Foster R., Lee M.S.Y. 2007. Calibration choice, rate smoothing, and the pattern

of tetrapod diversification according to the long nuclear gene RAG-1. Syst. Biol.

56:543–563.

Hugi J., Hutchinson M.N., Koyabu D., Sánchez-Villagra M.R. 2012. Heterochronic shifts in

the ossification sequences of surface- and subsurface-dwelling skinks are correlated with

the degree of limb reduction. Zoology 115:188–198.

Irisarri I., Baurain D., Brinkmann H., Delsuc F., Sire J.-Y., Kupfer A., Petersen J., Jarek M.,

Meyer A., Vences M., Philippe H. 2017. Phylotranscriptomic consolidation of the jawed

vertebrate timetree. Nature Ecol. Evol. 1:1370–1378.

Josse S., Moreau T., Laurin M. 2006. Stratigraphic tools for Mesquite, version 1.0.

http://mesquiteproject.org/packages/stratigraphicTools/

Laurin M. 1998. The importance of global parsimony and historical bias in understanding

tetrapod evolution. Part I. Systematics, middle ear evolution, and jaw suspension. Ann.

Sci. Nat., Zool., 13$^e$ Sér. 19:1–42.

Laurin M. 2004. The evolution of body size, Cope's rule and the origin of amniotes. Syst.

Biol. 53:594-622.

Laurin M., Germain D. 2011. Developmental characters in phylogenetic inference and their

absolute timing information. Syst. Biol. 60:630–644.

Laurin M., Piñeiro G. 2017. A reassessment of the taxonomic position of mesosaurs, and a

surprising phylogeny of early amniotes. Front. Earth Sci. 5:88.

Lecompte E., Aplin K., Denys C., Catzeflis F., Chades M., Chevret P. 2008. Phylogeny and

biogeography of African Murinae based on mitochondrial and nuclear gene sequences,

with a new tribal classification of the subfamily. BMC Evol. Biol. 8:199.

Lee C., Blay S., Mooers A.Ø., Singh A., Oakley T.H. 2006. CoMET: A Mesquite package for

comparing models of continuous character evolution on phylogenies. Evol.

Bioinformatics Online 2:193–196.

Maddin H.C., Jenkins F.A. Jr., Anderson J.S. 2012. The braincase of *Eocaecilia micropodia*

(Lissamphibia, Gymnophiona) and the origin of caecilians. PLOS ONE 7:e50743.

Maddison W.P. 1991. Squared-change parsimony reconstructions of ancestral states for

continuous-valued characters on a phylogenetic tree. Syst. Zool. 40:304–314.

Maddison W.P., Maddison D.R. 2017. Mesquite: a modular system for evolutionary analysis.

Version 3.2. http://mesquite.wikispaces.com

Marjanović D., Laurin M. 2008. Assessing confidence intervals for stratigraphic ranges of

higher taxa: The case of Lissamphibia. Acta Palaeont. Pol. 53:413–432.

Marjanović D., Laurin M. 2009. The origin(s) of modern amphibians: a commentary. Evol.

Biol. 36:336–338.

Marjanović D., Laurin M. 2013. The origin(s) of extant amphibians: a review with emphasis

on the "lepospondyl hypothesis". Geodiversitas 35:207–272.

Marjanović D., Laurin M. 2014. An updated paleontological timetree of lissamphibians, with

comments on the anatomy of Jurassic crown-group salamanders (Urodela). Hist. Biol.

26:535–550.

Marjanović D., Laurin M. 2018. Reproducibility in phylogenetics: reevaluation of the largest

published morphological data matrix for phylogenetic analysis of Paleozoic limbed

vertebrates. PeerJ Preprints 6:e1596v3.

Maxwell E.E., Harrison L.B., Larsson H.C.E. 2010. Assessing the phylogenetic utility of

sequence heterochrony: evolution of avian ossification sequences as a case study.

Zoology 113:57–66.

Meredith R.W., Janečka J.E., Gatesy J., Ryder O.A., Fisher C.A., Teeling E.C., Goodbla A.,

Eizirik E., Simão T.L.L., Stadler T., Rabosky D.L., Honeycutt R.L., Flynn J.J., Ingram

C.M., Steiner C., Williams T.L., Robinson T.J., Burk-Herrick A., Westerman M.,

Ayoub N.A., Springer M.S., Murphy W.J. 2011. Impacts of the Cretaceous terrestrial

revolution and KPg extinction on mammal diversification. Science 334:521–524.

Olori J.C. 2013. Ontogenetic sequence reconstruction and sequence polymorphism in extinct

taxa: an example using early tetrapods (Tetrapoda: Lepospondyli). Paleobiology

39:400–428.

Olori J.C. 2015. Skeletal morphogenesis of *Microbrachis* and *Hyloplesion* (Tetrapoda:

Lepospondyli), and implications for the developmental patterns of extinct, early

tetrapods. PLOS ONE 10:e0128333.

Pardo J.D., Szostakiwskyj M., Ahlberg P.E., Anderson J.S. 2017a. Hidden morphological

diversity among early tetrapods. Nature 546:642–645.

Pardo J.D., Small B.J., Huttenlocker A.K. 2017b. Stem caecilian from the Triassic of

Colorado sheds light on the origins of Lissamphibia. Proc. Natl. Acad. Sci. U.S.A.

114:E5389–E5395.

Parsons T.S., Williams E.E. 1962. The teeth of Amphibia and their relation to amphibian

phylogeny. J. Morph. 110:375–389.

Parsons T.S., Williams E.E. 1963. The relationships of the modern Amphibia: A re-

examination. Q. Rev. Biol. 38:26–53.

Pawley K. 2006. The postcranial skeleton of temnospondyls (Tetrapoda: Temnospondyli)

[doctoral thesis]. Melbourne: La Trobe University. http://hdl.handle.net/1959.9/405644

Pons J.-M., Hassanin A., Crochet P.-A. 2005. Phylogenetic relationships within the Laridae

(Charadriiformes: Aves) inferred from mitochondrial markers. Mol. Phyl. Evol. 37:686–

699.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon A.M., Lemmon A.R.

2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA

sequencing. Nature 526:569–573.

Pyron R.A. 2014. Biogeographic analysis reveals ancient continental vicariance and recent

oceanic dispersal in amphibians. Syst. Biol. 63:779–797.

Rabosky D.L., Donnellan S.C., Grundler M., Lovette I.J. 2014. Analysis and visualization of

complex macroevolutionary dynamics: an example from Australian scincid lizards. Syst.

Biol. 63:610-627.

Reeder T.W. 2003. A phylogeny of the Australian *Sphenomorphus* group (Scincidae:

Squamata) and the phylogenetic placement of the crocodile skinks (*Tribolonotus*):

Bayesian approaches to assessing congruence and obtaining confidence in maximum

likelihood inferred relationships. Mol. Phyl. Evol. 27:384-397.

Ruta M., Coates M.I. 2007. Dates, nodes and character conflict: addressing the lissamphibian

origin problem. J. Syst. Palaeontol. 5:69–122.

Schneider J.W., Werneburg R., Rößler R., Voigt S., Scholze F. 2015. Example for the

description of basins in the CPT Nonmarine-Marine Correlation Chart – Thuringian

Forest Basin, East Germany. Permophiles 61:29–35.

Schoch R.R. 2002. The formation of the skull in Paleozoic and extant amphibians.

Paleobiology 28:378–396.

Schoch R.R. 2004. Skeleton formation in the Branchiosauridae: a case study in comparing

ontogenetic trajectories. J. Vert. Paleont. 24:309–319.

Schoch R.R. 2006. Skull ontogeny: developmental patterns of fishes conserved across major

tetrapod clades. Evol. Dev. 8:524–536.

Schoch R.R. 2014a. First evidence of the branchiosaurid temnospondyl *Leptorophus* in the

Early Permian of the Saar-Nahe Basin (SW Germany). N. Jb. Geol. Paläont. Abh.

272:225–236.

Schoch R.R. 2014b. Amphibian skull evolution: the developmental and functional context of

simplification, bone loss and heterotopy. J. Exp. Zool. (Mol. Dev. Evol.) 322B:619–630.

Schoch R.R., Carroll R.L. 2003. Ontogenetic evidence for the Paleozoic ancestry of

salamanders. Evol. Dev. 5:314–324.

Schoch R.R., Milner A.R. 2004. Structure and implications of theories on the origin of

lissamphibians. Pages 345–377 in Arratia G., Wilson M.V.H., Cloutier R., editors.

Recent Advances in the Origin and Early Radiation of Vertebrates. Munich: Dr.

Friedrich Pfeil.

Sigurdsen T., Green D.M. 2011. The origin of modern amphibians: a re-evaluation. Zool. J.

Linn. Soc. 162:457–469.

Steen M.C. 1938. On the fossil Amphibia from the Gas Coal of Nýřany and other deposits in

Czechoslovakia. Proc. Zool. Soc. Lond. 108:205–283.

Sterli J., Pol D., Laurin M. 2013. Incorporating phylogenetic uncertainty on phylogeny-based

paleontological dating and the timing of turtle diversification. Cladistics 29:233–246.

Tarver J.E., dos Reis M., Mirarab S., Moran R.J., Parker S., O'Reilly J.E., King B.L.,

O'Connell M.J., Asher R.J., Warnow T., Peterson K.J., Donoghue P.C.J., Pisani D.

2016. The interrelationships of placental mammals and the limits of phylogenetic

inference. Genome Biol. Evol. 8:330–344.

van der Vos W., Witzmann F., Fröbisch N.B. 2017. Tail regeneration in the Paleozoic

tetrapod *Microbrachis pelikani* and comparison with extant salamanders and squamates.

J. Zool. 304:34–44

Wagenmakers E.-J., Farrell S. 2004. AIC model selection using Akaike weights. Psychon.

Bull. Rev. 11:192–196.

Wang N., Kimball R.T., Braun E.L., Liang B., Zhang Z. 2013. Assessing phylogenetic

relationships among Galliformes: a multigene phylogeny with expanded taxon sampling

in Phasianidae. PLOS ONE 8:1–12.

Watson D.M.S. 1940. The origin of frogs. Trans. R. Soc. Edinburgh 60:195–231.

Weisbecker V. 2011. Monotreme ossification sequences and the riddle of mammalian skeletal

development. Evolution 65:1323–1335.

Werneburg R. 2018 (for 2017). Earliest 'nursery ground' of temnospondyl amphibians in the

Permian. Semana 32:3–42.

Werneburg I., Sánchez-Villagra M.R. 2009. Timing of organogenesis support basal position

of turtles in the amniote tree of life. BMC Evol. Biol. 9:82.

Witzmann F., Schoch R.R. 2006. Skeletal development of the temnospondyl

*Acanthostomatops vorax* from the Lower Permian Döhlen Basin of Saxony. Trans. R.

Soc. Edinburgh: Earth Sci. 96:365–385.

Zhang P., Zhou H., Chen Y.-Q., Liu Y.-F., Qu L.-H. 2005. Mitogenomic perspectives on the

origin and phylogeny of living amphibians. Syst. Biol. 54:391–400.

.

FIGURE LEGENDS

FIGURE 1. Hypotheses on the relationships of the extant amphibian clades since the mid-20[th]

century. The names of terminal taxa sampled here are in boldface; the names of larger clades

are placed toward the right end of a branch if they have minimum-clade (node-based)

definitions, to the left if they have maximum-clade (branch-based) definitions. The first two

trees (a, b) show the current consensus; the other trees (c–h) show the various tested

paleontological hypotheses. a) Consensus of the latest and largest phylogenetic analyses of

molecular data (Pyron 2014; Irisarri et al. 2017; Feng et al. 2017); all named clades are

therefore extant. Note the monophyly of the extant amphibians (Lissamphibia, marked with a

light gray dot) with respect to Amniota. b) Consensus of all analyses of Paleozoic limbed

vertebrates (latest and largest: Pawley 2006; Sigurdsen and Green 2011; Pardo et al. 2017a, b:

fig. S6; Marjanović and Laurin 2018), omitting the extant amphibian clades. Note the

monophyly of "lepospondyls" + amniotes (marked with a dark gray dot). c) TH: Lissamphibia

nested among temnospondyls, close to *Apateon* (most recently found by Sigurdsen and Green

2011; Maddin et al. 2012; Pardo et al. 2017a, b: fig. S6; argued for by Schoch and Milner

2004, Schoch 2014b and others). Compatible with both a and b (light gray dots). d) LH:

Lissamphibia nested among "lepospondyls" (found most recently by Pawley 2006;

Marjanović and Laurin 2009, 2018); consequently, the temnospondyls *Apateon* and

*Sclerocephalus* are not crown-group tetrapods. Compatible with both a) and b) (light gray

dots). e) PH1: Urodela as temnospondyls close to *Apateon*, Anura as a separate clade of

temnospondyls, Gymnophiona as "lepospondyls" (found by Carroll 2007; argued for by

Carroll 2001; Schoch and Carroll 2003; very cautiously Fröbisch et al. 2007). Compatible

with b) (dark gray dot) but not with a) (light gray circle). f) PH2: like PH1, but with

monophyletic extant amphibians with respect to amniotes (a) restored (light gray dot) at the

expense of compatibility with the paleontological consensus concerning the position of

temnospondyls, lepospondyls, and amniotes (b) (dark gray circle). g) DH1: Batrachia as

temnospondyls close to *Apateon*, Gymnophiona as "lepospondyls" (found by Anderson 2007;

Anderson et al. 2008). Compatible with b) (dark gray dot) but not with a) (light gray circle).

h) DH2: Batrachia as temnospondyls close to *Apateon*, Gymnophiona as temnospondyls

distantly related to *Sclerocephalus* (found by Pardo et al. 2017b in an analysis that included

only temnospondyls and lissamphibians: fig. 2, S7). Compatible with both a) and b).

FIGURE 2. Reference phylogeny used for some of the analyses, illustrating the LH. The tree

was time-calibrated, but analyses showed that branch lengths are irrelevant, given that the best

model is speciational (Table 1).

TABLE 1. AICc and AICc weights of six evolutionary models given our data (102 taxa,

including the temnospondyl *Sclerocephalus*; six characters) and reference tree. This was

performed on the tree representing the LH (lepospondyl hypothesis), but doing this on other

trees leads to similar results. Numbers presented with four significant digits; best values in

boldface. Abbreviations: k, number of estimable parameters; l, likelihood; wi, weight; $\Delta_i$,

difference of AICc from that of the Pure-Phylogenetic / Equal model.

| | AIC | l | k | AICc | $\Delta_i$ AICc | wi(AICc) |
|---|---|---|---|---|---|---|
| Pure-Phylogenetic / Distance | −569.0 | 285.5 | **1** | −567.5 | 780.4 | 3.461 E−170 |
| Pure-Phylogenetic / Equal | **−1173** | **587.5** | **1** | **−1171** | **0** | **1.000** |
| Pure-Phylogenetic / Free | 2.000 E10 | −1.000 E10 | 306 | 2.000 E10 | 2.000 E10 | 0 |
| Non-Phylogenetic / Distance | −439.8 | 220.9 | **1** | −438.3 | 909.6 | 3.026 E−198 |
| Non-Phylogenetic / | −899.5 | 450.7 | **1** | −898.0 | 450.0 | 1.959 E−98 |

Equal

Non-

Phylogenetic /     2.000

Free        E10   −1.000 E10   154     2.000 E10     2.000 E10         0

618 TABLE 2. As Table 1, but with 101 taxa, excluding *Sclerocephalus*, and six characters.

619 Abbreviations and boldface as in Table 1.

| | AIC | l | k | AICc | $\Delta_i$ AICc | wi(AICc) |
|---|---|---|---|---|---|---|
| Pure-Phylogenetic / Distance | −718.3 | 360.2 | **1** | −717.1 | 630.8 | 1.046 E−137 |
| Pure-Phylogenetic / Equal | **−1350** | **676.1** | **1** | **−1349** | **0** | **1.000** |
| Pure-Phylogenetic / Free | 2.000 E10 | −1.000 E10 | 306 | 2.000 E10 | 2.000 E10 | 0 |
| Non-Phylogenetic / Distance | −549.5 | 275.7 | **1** | −548.3 | 799.6 | 2.282 E−174 |
| Non-Phylogenetic / Equal | −899.5 | 450.7 | **1** | −898.3 | 449.7 | 2.278 E−98 |
| Non-Phylogenetic / Free | −649.8 | 478.9 | 154 | −1293 | 55.11 | 1.078 E−12 |

TABLE 3. AIC and AICc weights of the six topologies, reflecting the six hypotheses about the

origin of extant amphibians, for the speciational model (called Pure-Phylogenetic / Equal in

Tables 1 and 2), given a sample with 102 taxa (including *Sclerocephalus*) and six characters.

Abbreviations and boldface as in Table 1, except $\Delta_i$: difference of AICc from that of the LH.

Hypotheses from top to bottom: TH: monophyletic origin among temnospondyls; PH1:

triphyletic origin (anurans and urodeles from different temnospondyls, caecilians from

lepospondyls); PH2: triphyletic origin from temnospondyls; DH1: diphyletic origin (anurans

and urodeles from temnospondyls, caecilians from lepospondyls); DH2: diphyletic origin

from temnospondyls; LH: monophyletic origin from lepospondyls; see text for details and

references.

| Hypothesis | AIC | l | AICc | $\Delta_i$ AICc | wi(AICc) |
|---|---|---|---|---|---|
| TH | −1164 | 583.2 | −1163 | 8.638 | 0.01314 |
| PH1 | −1148 | 574.8 | −1146 | 25.37 | 3.062 E−06 |
| PH2 | −1147 | 574.4 | −1145 | 26.18 | 2.034 E−06 |
| DH1 | −1154 | 578.1 | −1153 | 18.65 | 8.799 E−05 |
| DH2 | −1147 | 574.5 | −1146 | 25.94 | 2.302 E−06 |
| LH | **−1173** | **587.5** | **−1171** | **0** | **0.9868** |

633 TABLE 4. As Table 3, but for a sample with 101 taxa, excluding *Sclerocephalus*, and seven

634 characters. Abbreviations, boldface and hypotheses as in Table 3.

| Hypothesis | AIC | l | AICc | $\Delta_i$ AICc | wi(AICc) |
|---|---|---|---|---|---|
| TH | −1346 | 673.8 | −1345 | 4.511 | 0.09377 |
| PH1 | −1338 | 670.2 | −1337 | 11.83 | 0.002409 |
| PH2 | −1341 | 671.4 | −1340 | 9.419 | 0.008059 |
| DH1 | −1335 | 668.6 | −1334 | 14.92 | 5.157 E−04 |
| DH2 | −1336 | 669.0 | −1335 | 14.24 | 7.224 E−04 |
| **LH** | **−1350** | **676.1** | **−1349** | **0** | **0.8945** |

636     SUPPLEMENTARY MATERIAL

637     ONLINE SUPPLEMENTARY DOCUMENT 1. Sources of the ossification sequence data.

638     ONLINE SUPPLEMENTARY DOCUMENT 2. Data matrix in NEXUS format for Mesquite.