Title: Antibody responses to the Zika virus polyprotein in pregnant and non-pregnant

macaques

Authors:

Emma L. Mohr[a]†, **Anna S. Heffron[b]†**, David Baker[b], Amelia K. Haj[b], Connor R.

Buechler[b], Adam Bailey[b], Dawn M. Dudley[b], Christina M. Newman[b], Mariel S. Mohns[b],

Michelle Koenig[b], Meghan E. Breitbach[b], Mustafa Rasheed[b], Laurel M. Stewart[b], Jens

Eickhoff[c], Richard S. Pinapati[d], Erica Beckman[d], Hanying Li[d], Jigar Patel[d], John C. Tan[d],

David H. O'Connor[b]#

Affiliations:

[a]Department of Pediatrics, School of Medicine and Public Health, University of

Wisconsin-Madison

[b]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison

[c]Department of Biostatistics & Medical Informatics, University of Wisconsin-Madison

[d]Roche Sequencing Solutions, Madison

Running title: IgG responses to the ZIKV polyprotein in NHPs

#Address correspondence to David H. O'Connor, dhoconno@wisc.edu

†E.L.M and A.S.H. contributed equally to this work.

ABSTRACT

The kinetics and specificity of antibody responses against Zika virus (ZIKV) are not well-

characterized. This is due, in part, to the antigenic similarity between ZIKV and closely

related Dengue virus (DENV) serotypes. Since these and other similar viruses co-

circulate and are spread by the same mosquito species, it is difficult to disentangle

ZIKV-specific antibody responses from responses to closely-related arboviruses in

humans. Here we use high-density peptide microarrays to differentiate ZIKV from DENV

antibody reactivity in pregnant and non-pregnant macaque monkeys with known

exposure histories. We independently confirm a ZIKV-specific IgG antibody response

targeting ZIKV nonstructural protein 2B (NS2B) that was recently reported in ZIKV-

infected people and show that antibody reactivity in pregnant animals can be detectable

as late as 113 days post-infection (dpi). We also show, however, these responses wane

over time, and in one case the response was elicited following DENV infection in a

previously ZIKV-exposed animal. The generally weak and short-lived antibody

responses to this NS2B epitope, the only reliably ZIKV-specific epitope identified to date

to our knowledge, suggest epidemiologic studies assessing seroprevalence specifically

to ZIKV using linear epitope-based strategies may be prone to false negative or

otherwise confounded results. These results additionally indicate some antibodies

produced in response to ZIKV infection are in circulation for only a short period of time.

IMPORTANCE

Infection with ZIKV during pregnancy can lead to congenital ZIKV infection and

associated birth defects. The vulnerability of communities to future ZIKV outbreaks will

depend, in part, on the prevalence of immunity, which is thought to be mediated

principally by antibodies. We currently lack diagnostic assays to differentiate ZIKV-

specific antibodies from antibodies directed against closely related DENV and we do not

know how long anti-ZIKV responses persist. Here we profile antibodies recognizing

linear epitopes throughout the entire ZIKV and DENV polyproteins and show that while

ZIKV-specific antibody responses can be detected, these responses are generally weak

and ephemeral. This may complicate efforts to identify people previously infected with

ZIKV and to determine ZIKV seroprevalence, and it raises questions about the utility of

linear epitope-based technologies in defining ZIKV infection history.

INTRODUCTION

Zika virus (ZIKV) serologic assays suffer from cross-reactivity with closely related and

co-endemic Dengue virus (DENV), due to the high level (average ~55%) of amino acid

sequence identity and structural similarity (1-4) of ZIKV and DENV. Serologic assays

have been developed to detect past ZIKV infection, reporting sensitivity varying from

37% to 97% and specificity varying from 20% to 90% (5-8). Most of these assays have

focused on antibodies to the ZIKV envelope protein or nonstructural protein 1 (NS1) (7,

9-11). A recent publication by Mishra et al. employed a high-density peptide microarray

to identify antibodies to linear ZIKV epitopes that do not cross-react with other

flaviviruses (12). This group identified an IgG immunoreactive peptide sequence in ZIKV

nonstructural protein 2B (NS2B) which induces little or no antibody binding in serum

from ZIKV-naive people or people with history of infection with other flaviviruses (12).

This suggests profiling antibody binding against the entire viral proteome may identify

unexpected epitopes with useful properties, such as cross-reactivity with multiple similar

viruses or specificity to a single virus.

Because ZIKV, DENV, and other flaviviruses are similar in structure and clinical

syndrome, are spread by the same mosquito vectors (13), and are co-endemic (14, 15),

it is difficult to identify people who have unequivocally been exposed to ZIKV only,

making it difficult to profile ZIKV-specific antibody responses. In contrast, macaques

raised in indoor colonies can be infected specifically with ZIKV or DENV. Macaque

models of ZIKV infection provide a close approximation of human ZIKV infection in

regards to natural history (16-19), tissue tropism (16, 17, 20, 21), and transmission (17,

22-25). Importantly, macaques infected with ZIKV during pregnancy provide insight into

the pathogenesis of congenital ZIKV infection (17, 18, 21, 22, 26, 27). Since the strain,

dose, and timing of ZIKV infection is exactly known, the kinetics and specificity of

humoral immune responses can be profiled in macaques with better resolution than is

possible in cross-sectional human studies.

Here, we used high-density peptide microarrays to map macaque IgG epitopes in full-

length ZIKV and DENV polyproteins. This peptide microarray technology allows for one

serum sample to be assayed against six million unique 16-residue peptides or, as was

done in this study, for 12 samples to each be assayed against 392,000 peptides on a

single chip (Roche Sequencing Solutions, Madison, WI). This technology has been

used in proteome-wide epitope mapping (28), profiling of antibody responses in

autoimmune disease (29), profiling venom toxin epitopes (30), determining functions of

cellular enzymes (31), *de novo* binding sequence discovery (32), epitope validation

following phage display screening (33), and screening for tick-borne disease

seroprevalence (34). We previously used this tool to examine antibody responses in

simian pegivirus (SPgV) and simian immunodeficiency virus (SIV) infections (35). As

mentioned above, a recent publication explored use of this technology in profiling

human antibody responses against flaviviruses (12). This linear peptide microarray

technology has advantages over previous assays through its capability to screen for

reactivity to a large number of pathogens while simultaneously mapping reactive

peptides precisely, and it has the distinct advantage of allowing detection of unexpected

epitopes due to its capacity to assay the entirety of a virus's proteome.

RESULTS

Ultradense peptide microarray identifies SIV-specific IgG epitopes

To assess the performance of the recently-developed peptide microarray platform we

sought to test its performance against a well-characterized immune response. We

synthesized 16-mer peptides overlapping by 12 amino acids representing the

SIVmac239 envelope protein (env) and analyzed serum from two cynomolgus

macaques for SIV-specific IgG reactivity before and approximately 125 days after

SIVmac239 infection (Table 1). We previously plotted data from these samples (35);

here we show the same data using an updated, improved data processing pipeline that

is used throughout this manuscript. Post-infection samples exhibited reactivity as high

as 1,000 times the reactivity in pre-infection samples (Figure 1). Regions of higher-fold

increases in reactivity corresponded to previously defined variable domains of env

116 which are known antibody targets, the variable loop regions, as well as others

117 corresponding to known epitopes (36). Taken together, these results demonstrate the

118 peptide microarray platform is capable of high resolution virus-specific IgG epitope

119 identification.

121 **Identification of ZIKV polyprotein epitopes**

122 Next we sought to determine reactivity to the ZIKV polyprotein following ZIKV infection.

123 We assessed reactivity against ZIKV polyproteins represented by 16-mer peptides

124 (overlapping by 15 amino acids) of serum from animals with recent ZIKV-FP (animals

125 B1, D1, and D2) or ZIKV-MR766 (animal C1) infections (Table 1). Animals in both

126 groups demonstrated neutralizing antibody responses at 28 dpi, measured by 90%

127 plaque reduction neutralization tests ($PRNT_{90}$) as described previously (16, 23, 37).

128 Peptides were defined as reactive if the normalized signal intensity was greater after

129 infection with a cognate strain (e.g., increased signal intensity against ZIKV peptides in

130 an animal infected with ZIKV) than it was before infection. Peptides were defined as

131 cross-reactive if the normalized signal intensity was greater after infection with a

132 noncognate strain (e.g., increased signal intensity against ZIKV peptides in an animal

133 infected with DENV). In regions of interest, such as the epitope in NS2B in which a

134 ZIKV-specific epitope was identified, statistical significance of the change in signal

135 intensity versus no change was calculated using a two-tailed log-ratio t-test. Given the

136 small offset between adjacent peptides, true epitopes were expected to induce antibody

137 binding to multiple peptides with overlapping sequences; thus regions were only

considered epitopic when there was a statistically significant increase in post-infection

reactivity relative to pre-infection reactivity in three or more consecutive peptides.

Reactivity was most commonly observed in three regions: envelope protein, NS2B, and

nonstructural protein 3 (NS3). Antibody responses against the envelope protein were

seen in four out of four animals at 28 dpi; responses against nonstructural protein NS3

(NS3) were seen in three out of four (Figure 2). All four animals exhibited an antibody

response to the ZIKV NS2B epitope similar to that documented in humans (12). Area

under the curve (AUC) values and corresponding receiver operating characteristic

(ROC) curves were calculated (supplemental figure S1), identifying ten peptides at

positions 1427-1436 in the polyprotein (for a total of 25 amino acids, sequence

RAGDITWEKDAEVTGNSPRLDVALD) as the best-performing epitope, hereafter

referred to as $NS2B_{1427-1451}RD25$ (AUC of 0.9375, 95% confidence interval of 0.89065

to 0.9375). This epitope is slightly larger in size than that found by Mishra et al. (nine

12-mer peptides, for a total of 20 amino acids, sequence DITWEKDAEVTGNSPRLDVA)

(12). The four animals exhibited similar responses against an Asian ZIKV strain, ZIKV-

FP (Figure 2), as against an African strain and an American strain (ZIKV-MR766 and

ZIKV-PR respectively, supplemental figure S2). Animals infected with either of the ZIKV-

FP or ZIKV-MR766 strains showed similar reactivity against the ZIKV polyprotein

(supplemental figure S3).

Cross-reactivity of ZIKV-convalescent serum with DENV polyproteins

DENV polyproteins share an average of 55% sequence identity with ZIKV polyproteins

(1, 2), and the DENV NS2B region corresponding to the immunoreactive ZIKV

$NS2B_{1427-1451}RD25$ shares an average of 41% identity (Figure 3 A). To examine cross-

reactivity, we analyzed the antibody responses of the ZIKV-convalescent animals to

DENV polyproteins represented on the array (Figure 3 B-E). Cross-reactivity was

apparent, with antibodies from the ZIKV-convalescent but DENV-naive animals

recognizing regions of DENV polyproteins. All four animals showed only minimal

reactivity against DENV envelope protein and DENV NS3. Animals exhibited

comparable cross-reactivities to all four DENV serotypes, with the highest single

instance of cross-reactivity observed against a DENV-3 epitope in the NS3 region. No

significant cross-reactivity against the DENV NS2B protein was observed.

Differentiating DENV from ZIKV serologic responses

Previous assays have struggled to distinguish DENV serologic responses from ZIKV

serologic responses (5-8). We investigated whether reactivity against the peptide

microarray and specifically against the ZIKV $NS2B_{1427-1451}RD25$ epitope could

distinguish a recent DENV infection in an animal with previous ZIKV exposure. Three

animals (cohort F) had been challenged twice with ZIKV-FP 12 months prior and 9.5

months prior (16); we infected these animals with DENV-2 and collected serum samples

prior to DENV-2 infection and 28 days after. These samples were analyzed against 16-

mer peptides, with amino acid overlap of 12, representing ZIKV-FP (Figure 4).

These animals showed reactivity against all four DENV polyproteins in regions

representing the DENV envelope protein, DENV NS3, and others (supplemental figure

S4) and cross-reactivity against corresponding regions of the ZIKV polyprotein (Figure

4). One animal (F3) showed reactivity to the $NS2B_{1427-1451}RD25$ epitope, though this

reactivity was slightly outside the area of typical $NS2B_{1427-1451}RD25$ reactivity (peptides

1421-1429 rather than 1427-1436).

Anti-ZIKV antibody response during pregnancy

Given the importance of determining an accurate ZIKV infection history in pregnancy,

we sought to characterize the gestational anti-ZIKV antibody response. We used the

peptide microarray to evaluate serum samples from six pregnant animals. Animals were

inoculated with ZIKV at 35-47 days post-conception (gestational date, gd) (Table 2).

Two animals (G1 and G2) had no history of flavivirus exposure and were inoculated with

ZIKV-FP at 36-38 gd. Three more flavivirus-naive animals (H1, H2, and H3) were

infected with ZIKV-PR at 45-47 gd. One animal (I1) had a history of exposure to DENV-

3 nine months prior to inoculation with a barcoded clone of ZIKV-PR at 35 dpi (38).

Serum samples collected approximately one week post-infection and at two to six week

intervals thereafter were analyzed against flavivirus polyproteins represented on the

peptide array (Figure 5). All six pregnant animals exhibited anti-$NS2B_{1427-1451}RD25$

reactivity by the early convalescent phase (21-29 dpi), though time of initial appearance

and duration of the response varied between animals.

In G1, elevated baseline intensity in the ZIKV $NS2B_{1427-1451}RD25$ region was present

prior to inoculation with ZIKV-FP. This animal experienced one of the highest levels of

anti-ZIKV IgG reactivity in this study (magnitude of fluorescence intensity reaching

6338.7 AU, or 13.1 on the log base 2 scale) in the acute phase (7 dpi). Reactivity

subsequently decreased but remained above pre-infection levels through 78 dpi. The

anti-$NS2B_{1427-1451}RD25$ intensity level fell below the pre-infection level by 127 dpi, after

G1 had given birth. In G2, reactivity was not appreciable until the early convalescent

phase (21 dpi) but remained above pre-infection levels until the latest time point

analyzed (113 dpi).

Both H1 and H2 showed reactivity against $NS2B_{1427-1451}RD25$ during acute infection (7

dpi) and all three animals in cohort H showed reactivity by the early convalescent phase

(21 dpi). This reactivity decreased to or below pre-infection levels for H1 and H3 but not

H2 by 70 dpi.

I1, an animal nine months post-DENV-3 infection, showed a pattern of reactivity similar

to the other animals, with anti-$NS2B_{1427-1451}RD25$ IgG reactivity first appearing at 8 dpi

and peaking at 29 dpi at a level 9 times pre-infection reactivity (Figure 5 F). I1 was the

only animal in this study for which reactivity remained close to peak reactivity by 78 dpi.

All pregnant animals' cross-reactivity against DENV polyproteins mirrored that in non-

pregnant animals (supplemental figure S6). All animals in which fluorescence intensity

at the $NS2B_{1427-1451}RD25$ epitope was higher at pre-infection time points than at post-

infection time points had this effect only for the $NS2B_{1427-1451}RD25$ epitope and not for

other immunoreactive regions of the ZIKV polyprotein (supplemental figure S5).

Kinetics of anti-ZIKV IgG responses

The duration of antibody responses following ZIKV infection has not yet been

established. Our data provide preliminary insight into the strength and duration of anti-

ZIKV antibody responses.

Kinetics of IgG binding at the $NS2B_{1427-1451}RD25$ epitope were followed during the

course of infection using fold change from pre-infection levels. Non-pregnant animals

(B1, C1, D1, and D2) exhibited a characteristic rise and fall in reactivity, with greatest

fold increase consistently noted at early convalescence, 28 dpi (supplemental figure S7

A). At 28 dpi B1 and C1 exhibited a 5.7-fold reactivity increase above pre-infection

levels; by 67 dpi these levels had fallen to 3.1-fold and 5.5-fold respectively,

representing a 46% decrease and a 4% decrease in anti-$NS2B_{1427-1451}RD25$ reactivity in

39 days. Animals D1 and D2 showed reactivity increased 2.4-fold and 1.2-fold above

pre-infection levels at 28 dpi. Likewise, samples from D1 and D2 in late convalescence

(621 and 582 dpi, respectively) showed a decline in IgG reactivity (to 0.92-fold and 0.82-

fold, respectively).

Kinetics in pregnant animals were tracked more extensively (supplemental figure S7 B).

One animal, G1, exhibited unusually high pre-infection fluorescence intensity (to a

normalized level of 10.2; see Figure 5) at the $NS2B_{1427-1451}RD25$ epitope prior to

infection; G1's reactivity peaked from 7 to 21 dpi at a 2.3-fold increase over pre-infection

levels. Reactivity decreased by 78 dpi and fell 4.6-fold below pre-infection levels by 127

dpi. Animal G2 first demonstrated reactivity at 21 dpi, reaching peak reactivity at 78 dpi

and falling thereafter, with levels still above pre-infection through the latest time

examined (113 dpi). Two animals in cohort H, H1 and H3, showed ZIKV-specific

reactivity at 7 dpi, and H3 reached peak reactivity at this time. H1 and H2 exhibited

peak reactivity at 21 dpi, and reactivity in animals H1 and H3 had returned to or fallen

below pre-infection levels by 70 dpi. Animal I1, with history of previous DENV-3

infection, first showed $NS2B_{1427-1451}RD25$ reactivity at 8 dpi with peak reactivity at 29

dpi, with reactivity remaining close to peak reactivity through the last day examined, 78

dpi.

DISCUSSION

We describe the anti-ZIKV IgG response in non-pregnant and pregnant rhesus

macaques and compare this to the anti-DENV response. Using a recently developed

high-density peptide microarray we show macaques infected with ZIKV produce IgG

antibodies reactive against the ZIKV polyprotein, including a conserved response

against an epitope in ZIKV NS2B, $NS2B_{1427-1451}RD25$, that is apparent regardless of the

ZIKV strain used for infection. We establish cross-reactivity between anti-ZIKV and anti-

DENV antibodies for the ZIKV and DENV polyproteins. Additionally we show the anti-

$NS2B_{1427-1451}RD25$ IgG response in ZIKV, an acute infection, is low in magnitude and

brief in duration compared to the antibody reactivity induced by SIV, a chronic infection.

As seen in previous assays (2, 12, 39-41), we observed antibody cross-reactivity

between ZIKV- and DENV-immune serum, though reactivity to the ZIKV $NS2B_{1427-}$

$_{1451}$RD25 epitope was observed in all cases of recent ZIKV infection (12 out of 12 ZIKV-

infected animals) and in only one out of three instances of recent DENV infection.

Inoculation with African, Asian, and American ZIKV strains induced some variability in

responses, though reactivity to the NS2B epitope was conserved throughout. All ZIKV-

infected animals produced anti-ZIKV IgG against the $NS2B_{1427-1451}$RD25 epitope, though

reactivity was sometimes only slightly above an animal's pre-infection baseline (as in

animal D2) or was measurable for only a short duration (as in animal H3). The lack of

uniformity of the anti-ZIKV IgG response is especially relevant since all ZIKV-exposed

animals could be followed, in contrast to work in humans where there may be a

selection bias for individuals with symptomatic ZIKV, thought to account for only

approximately half of ZIKV infections (12, 42). Though anti-$NS2B_{1427-1451}$RD25 reactivity

was consistently present in early convalescence, it was observed to decay in all but one

case during the time period assessed. These findings corroborate findings in humans, in

which antibody reactivity in symptomatic patients was strongly associated with a

detectable NS2B response in the early convalescent phase (96%), but less likely six

months post-infection (44%) (12). These results demonstrate the need for investigation

into the longevity and kinetics of the anti-ZIKV antibody response. Given that this is the

only linear epitope identified to date as being reliably ZIKV-specific (12), and given that

antibodies to this epitope appear to drop to undetectable levels in at least 50% of

individuals by six months post-infection, these results also call into question how useful

current serological methods may be in differentiating past ZIKV exposure from exposure

to other flaviviruses. Additionally, one animal (F3) showed reactivity against the ZIKV

$NS2B_{1427-1451}RD25$ epitope 28 days after DENV infection, possibly as an effect of having

been infected with ZIKV 12 and 9.5 months prior. Thus, while the specificity of this

epitope is important, its utility in guiding development of diagnostics will likely be limited.

This array detected anti-$NS2B_{1427-1451}RD25$ reactivity in all pregnant macaques exposed

to ZIKV. Production of IgG antibodies has relevance to both mother and fetus, since

maternal IgG crosses the placenta freely and maternal donation of IgG increases

throughout pregnancy (43). It can be assumed the anti-ZIKV IgG produced by these

animals also reached their fetuses, but whether these anti-$NS2B_{1427-1451}RD25$

antibodies provide protection or whether they contribute to the pathology of ZIKV

disease and ZIKV congenital effects is currently unknown. We did not note any variation

in outcomes with variation in antibody responses, but it is possible variation in antibody

responses during pregnancy could help explain differences in outcomes following

gestational ZIKV infection. Several pregnant animals also showed post-infection

reactivity against the $NS2B_{1427-1451}RD25$ epitope that was lower than their pre-infection

reactivity. It is possible these animals were exposed to some substance similar to the

$NS2B_{1427-1451}RD25$ epitope that caused them to produce antibodies against it before

exposure to ZIKV, or that there is some genetic characteristic or some aspect of

pregnancy that causes increased likelihood of producing anti-$NS2B_{1427-1451}RD25$

antibodies prior to ZIKV exposure, though we have not found this epitope present in any

other pathogen. These findings merit further and more thorough investigation than this

current study can provide.

The peptide array technology used in this study has several limitations. Utility of the

assay will depend upon the magnitude of the antibody response induced, which varies

from pathogen to pathogen. In this study the magnitude of antibody reactivity induced

by ZIKV infection was markedly lower (165 times lower at peak reactivity) than that

induced by SIV infection. Our study defined positive responses as an increase from an

animal's pre-infection reactivity; human patients cannot give pre-infection samples and

must rely upon on controls determined from humans having no known history of

flavivirus exposure (12). We simulated the performance of this assay without the

availability of pre-infection samples by defining a positivity threshold using the average

reactivity of macaque pre-infection samples + 1 standard deviation; application of this

control caused some animals with lower reactivity (D1 and D2) to appear to lose their

post-infection positivity. Applying a control using the average of pre-infection samples +

2 standard deviations resulted in even more false negatives. Our control was limited by

small sample size, but these results raise the possibility of lower sensitivity when

definitive pre-infection samples from the same individual are not available.

The peptide array approach is also limited due to its detection only of continuous linear

epitopes. Many documented ZIKV epitopes are conformational discontinuous epitopes

(44-49). Other epitope discovery methods will likely remain superior in defining

discontinuous epitopes, but peptide microarrays are useful in identifying

immunoreactive regions not previously considered as potential epitopes. Past work from

our laboratory, including results from some of the animals analyzed here, has shown

anti-ZIKV neutralizing antibody titers measured by $PRNT_{90}$ do not drop off but remain

elevated as late as 64 dpi (16). The discordance between levels of anti-$NS2B_{1427\text{-}}$

$_{1451}$RD25 IgG and neutralizing antibody titers may indicate the anti-$NS2B_{1427\text{-}1451}$RD25

antibodies do not play a role in protection against future infections, and thus their

production is discontinued in favor of antibodies which can neutralize ZIKV virions.

Despite these limitations, this technology could be useful as a screening tool for

antibody reactivity to ZIKV and many other pathogens given its enormous throughput.

Little is known about the pervasiveness of ZIKV in Africa, though the virus' presence

there was established in 1947. To this end, our group has designed a peptide

microarray capable of simultaneously assessing twelve samples' antibody responses

against every arbovirus strain known to cause human infection in Africa. We plan to use

this array to screen pooled samples for evidence of human antibody reactivity to ZIKV

and to the hundreds of other viruses represented. This assay could also be used to

compare IgM and IgG responses, a currently untapped potential of this technology.

In summary, this work in macaques demonstrates the capacity of a recently developed

peptide microarray to differentiate anti-ZIKV and anti-DENV IgG antibody responses, a

distinction which has previously been difficult. Our work shows the anti-$NS2B_{1427\text{-}}$

$_{1451}$RD25 IgG response is characterized by relatively low-magnitude antibody responses

prone to relatively rapid decay, mirroring the events seen in humans. The peptide

microarray design used shows particular promise in evaluating serological reactivity

specific to ZIKV while simultaneously evaluating for responses against a large number

of other similar or dissimilar pathogens. In the future this technology could be expanded

for use in screening populations for exposure to ZIKV, DENV, and a multitude of other

pathogens.

MATERIALS AND METHODS

Macaque study design

Animal demographics, inoculation strain, dose, and route, serum sample collection

timelines, and array design used for each animal are described in Table 1. Gestational

details and outcomes for pregnant animals are described in Table 2. Additional details

on the study histories of the animals in this study can be found at

https://go.wisc.edu/b726s1.

Ethics

All monkeys were cared for by the staff at the Wisconsin National Primate Research

Center (WNPRC) in accordance with the regulations and guidelines outlined in the

Animal Welfare Act and the Guide for the Care and Use of Laboratory Animals and the

recommendations of the Weatherall report (https://royalsociety.org/topics-

policy/publications/2006/weatherall-report/). Per WNPRC standard operating

procedures for animals assigned to protocols involving the experimental inoculation of

an infectious pathogen, environmental enhancement included constant visual, auditory,

and olfactory contact with conspecifics, the provision of feeding devices which inspire

foraging behavior, the provision and rotation of novel manipulanda (e.g., Kong toys,

nylabones, etc.), and enclosure furniture (i.e., perches, shelves). Per Animal Welfare

Regulations (Title 9, Chapter 1, Subchapter A, Parts 1–4, Section 3.80 Primary

enclosures) the animals were housed in a nonhuman primate Group 3 enclosure with at

least 4.3 square feet of floor space and at least 30 inches of height. This study was

approved by the University of Wisconsin-Madison Graduate School Institutional Animal

Care and Use Committee (animal protocol numbers G005401 and G005443).

Virus stocks

SIVmac239 (GenBank accession: M33262) stock was prepared by WNPRC Virology

Services. Vero cells were transfected with the SIVmac239 plasmid. Infectious

supernatant was then transferred onto rhesus PBMC. The stock was not passaged

before collecting and freezing. ZIKV strain H/PF/2013 (GenBank accession: KJ776791)

was obtained from Xavier de Lamballerie (European Virus Archive, Marseille, France)

and passage history is described in Dudley et al. (16). ZIKV strain MR766, ZIKV strain

PRVABC59, and DENV-2 strain New Guinea C were generously provided by Brandy

Russell (CDC, Ft. Collins, CO). ZIKV strain MR766 passage history has been described

in Aliota et al. (37). A molecularly-barcoded version of ZIKV strain PRVABC59 (Zika

virus/H.sapiens-tc/PUR/2015/PRVABC59; GenBank accession: KU501215) was

constructed as described in Aliota et al. (38). DENV-2 strain New Guinea C (GenBank

accession: FJ390389), originally isolated from a human in New Guinea, underwent 17

rounds of amplification on cells and/or suckling mice followed by a single round of

amplification on C6/36 cells; virus stocks were prepared by inoculation onto a confluent

monolayer of C6/36 mosquito cells. DENV-3 strain Sleman/78 was obtained from the

NIH; virus stocks were prepared by a single passage on C6/36 cells.

Multiple sequence alignment

Full-length ZIKV and DENV polyprotein sequences were extracted from the National

Center for Biotechnology Information (NCBI) database into Geneious Pro 9.1.8

(Biomatters, Ltd., Auckland, New Zealand). These sequences included the ZIKV and

DENV polyprotein sequences described above (see "Virus stocks"), as well as DENV-1

strain VR-1254 (Genbank accession: EU848545) and DENV-4 strain VID-V2055

(Genbank accession: KF955510). These amino acid sequences were aligned using the

Geneious alignment algorithm as implemented in Geneious Pro 9.1.8 using default

parameters (global alignment with free end gaps, cost matrix: Blosum62).

Peptide array design and synthesis

Viral protein sequences were selected and submitted to Roche Sequencing Solutions

(Madison, WI) for development into a peptide microarray as part of an early access

program. Sequences included three ZIKV polyproteins (an Asian strain, ZIKV-FP,

Genbank accession: KJ776791.2; an African strain, ZIKV-MR766, Genbank accession:

KU720415.1; and an American strain, ZIKV-PR, Genbank accession: KU501215.1), four

DENV polyproteins (DENV-1, Genbank accession: EU848545.1; DENV-2, Genbank

accession: KM204118.1; DENV-3, Genbank accession: M93130.1; and DENV-4,

Genbank accession: KF955510.1), and one SIVmac239 env protein (Genbank

accession: AAA47637.1), which were used for analyses in this study. Proteins were tiled

as non-redundant 16 amino acid peptides, overlapping by 12 or 15 amino acids. The

array designs are publicly available at https://go.wisc.edu/b726s1.

The peptide sequences were synthesized *in situ* with a Roche Sequencing Solutions

Maskless Array Synthesizer (MAS) by light-directed solid-phase peptide synthesis using

an amino-functionalized support (Geiner Bio-One) coupled with a 6-aminohexanoic acid

linker and amino acid derivatives carrying a photosensitive 2-(2-nitrophenyl)

propyloxycarbonyl (NPPOC) protection group (Orgentis Chemicals). Unique peptides

were printed in random positions on the array to minimize impact of positional bias.

Peptide array sample binding

Macaque serum samples were diluted 1:100 in binding buffer (0.01M Tris-Cl, pH 7.4,

1% alkali-soluble casein, 0.05% Tween-20). Diluted sample aliquots and binding buffer-

only negative controls were bound to arrays overnight for 16-20 h at 4°C. After binding,

the arrays were washed 3x in wash buffer (1x TBS, 0.05% Tween-20), 10 min per wash.

Primary sample binding was detected via 8F1-biotin mouse anti-primate IgG (NIH

Nonhuman Primate Reagent Resource) secondary antibody. The secondary antibody

was diluted 1:10,000 (final concentration 0.1 ng/µl) in secondary binding buffer (1x TBS,

1% alkali-soluble casein, 0.05% Tween-20) and incubated with arrays for 3 h at room

temperature, then washed 3x in wash buffer (10 min per wash) and 30 sec in reagent-

grade water. The secondary antibody was labeled with Cy5-Streptavidin (GE

Healthcare; 5 ng/µl in 0.5x TBS, 1% alkali-soluble casein, 0.05% Tween-20) for 1 h at

room temperature, then the array was washed 2x for 1 min in 1x TBS, and washed once

for 30 sec in reagent-grade water. Fluorescent signal of the secondary antibody was

detected by scanning at 635 nm at 2 μm resolution and 25% gain, using an MS200

microarray scanner (Roche NimbleGen).

Peptide array data processing

The datafiles and analysis code for figures is available from https://go.wisc.edu/b726s1.

Fluorescence signal intensities were normalized by background subtraction and quantile

normalization across samples. All figures use the log base 2 of the resulting intensity

values. For each sample, each unique peptide was assayed and processed once; then

results from peptides redundant on the chip (i.e. were present in more than one protein

represented) were restored to each protein.

Acknowledgments

We would like to thank Adam Ericsen for helpful discussions.

Grant support: National Institutes of Health grants R01AI116382 and R24OD017850

(ASH, AKH, CRB, AB, DMD, CMN, MK, MEB, MR, LMS, DHO).

Conflict of interest

This manuscript describes the use of a platform provided on an early-access basis by

Roche Sequencing Solutions. While scientists from Roche were involved in the

experimental design and data analysis, the manuscript was prepared independently

from Roche and did not require pre-approval from Roche prior to submission. RSP, EB,

HL, JP, and JCT are employed by Roche Sequencing Solutions.

Supplemental materials

Supplemental figure S1: ROC curves for best-performing epitope and multidimensional scaling (MDS) plot for different time points.

Supplemental figure S2: Reactivity of B1, C1, D1, and D2 against ZIKV-MR766 and ZIKV-PR.

Supplemental figure S3: Reactivity of B1, C1, D1, and D2 against the complete polyprotein of ZIKV-FP.

Supplemental figure S4: Reactivity of cohort F against the DENV-2 polyprotein.

Supplemental figure S5: Reactivity of cohort G, cohort H, and cohort I against the ZIKV-FP polyprotein.

Supplemental figure S6: Reactivity of cohort G, cohort H, and cohort I against the DENV-2 polyprotein.

Supplemental figure S7: Kinetics of IgG antibody reactivity against the ZIKV NS2B$_{1427\text{-}1451}$RD25 epitope

References

1. Chang, H. H., Huber, R. G., Bond, P. J., Grad, Y. H., Camerini, D., Maurer-Stroh, S., & Lipsitch, M. (2017). Systematic analysis of protein identity between Zika virus and other arthropod-borne viruses. *Bull World Health Organ, 95*(7), 517-525I. doi:10.2471/BLT.16.182105

2. Xu, X., Vaughan, K., Weiskopf, D., Grifoni, A., Diamond, M. S., Sette, A., & Peters, B. (2016). Identifying Candidate Targets of Immune Responses in Zika

Virus Based on Homology to Epitopes in Other Flavivirus Species. *PLoS Curr, 8*.

doi:10.1371/currents.outbreaks.9aa2e1fb61b0f632f58a098773008c4b

3. Xu, X., Song, H., Qi, J., Liu, Y., Wang, H., Su, C., Shi, Y., & Gao, G. F. (2016).

Contribution of intertwined loop to membrane association revealed by Zika virus

full-length NS1 structure. *EMBO J, 35*(20), 2170-2178.

doi:10.15252/embj.201695290

4. Harrison, S. C. (2016). Immunogenic cross-talk between dengue and Zika

viruses. *Nat Immunol, 17*(9), 1010-1012. doi:10.1038/ni.3539

5. Safronetz, D., Sloan, A., Stein, D. R., Mendoza, E., Barairo, N., Ranadheera, C.,

Scharikow, L., Holloway, K., Robinson, A., Traykova-Andonova, M., Makowski,

K., Dimitrova, K., Giles, E., Hiebert, J., Mogk, R., Beddome, S., & Drebot, M.

(2017). Evaluation of 5 Commercially Available Zika Virus Immunoassays. *Emerg*

Infect Dis, 23(9), 1577-1580. doi:10.3201/eid2309.162043

6. Wong, S. J., Furuya, A., Zou, J., Xie, X., Dupuis, A. P., 2nd, Kramer, L. D., & Shi,

P. Y. (2017). A Multiplex Microsphere Immunoassay for Zika Virus Diagnosis.

EBioMedicine, 16, 136-140. doi:10.1016/j.ebiom.2017.01.008

7. Granger, D., Hilgart, H., Misner, L., Christensen, J., Bistodeau, S., Palm, J.,

Strain, A. K., Konstantinovski, M., Liu, D., Tran, A., & Theel, E. S. (2017).

Serologic Testing for Zika Virus: Comparison of Three Zika Virus IgM-Screening

Enzyme-Linked Immunosorbent Assays and Initial Laboratory Experiences. *J*

Clin Microbiol, 55(7), 2127-2136. doi:10.1128/JCM.00580-17

8. Goncalves, A., Peeling, R. W., Chu, M. C., Gubler, D. J., de Silva, A. M., Harris,

E., Murtagh, M., Chua, A., Rodriguez, W., Kelly, C., & Wilder-Smith, A. (2018).

Innovative and New Approaches to Laboratory Diagnosis of Zika and Dengue: A

Meeting Report. *J Infect Dis, 217*(7), 1060-1068. doi:10.1093/infdis/jix678

9. Lee, A. J., Bhattacharya, R., Scheuermann, R. H., & Pickett, B. E. (2017).

Identification of diagnostic peptide regions that distinguish Zika virus from related

mosquito-borne Flaviviruses. *PLoS One, 12*(5), e0178199.

doi:10.1371/journal.pone.0178199

10. Balmaseda, A., Stettler, K., Medialdea-Carrera, R., Collado, D., Jin, X.,

Zambrana, J. V., Jaconi, S., Cameroni, E., Saborio, S., Rovida, F., Percivalle, E.,

Ijaz, S., Dicks, S., Ushiro-Lumb, I., Barzon, L., Siqueira, P., Brown, D. W. G.,

Baldanti, F., Tedder, R., Zambon, M., de Filippis, A. M. B., Harris, E., & Corti, D.

(2017). Antibody-based assay discriminates Zika virus infection from other

flaviviruses. *Proc Natl Acad Sci U S A, 114*(31), 8384-8389.

doi:10.1073/pnas.1704984114

11. L'Huillier, A. G., Hamid-Allie, A., Kristjanson, E., Papageorgiou, L., Hung, S.,

Wong, C. F., Stein, D. R., Olsha, R., Goneau, L. W., Dimitrova, K., Drebot, M.,

Safronetz, D., & Gubbay, J. B. (2017). Evaluation of Euroimmun Anti-Zika Virus

IgM and IgG Enzyme-Linked Immunosorbent Assays for Zika Virus Serologic

Testing. *J Clin Microbiol, 55*(8), 2462-2471. doi:10.1128/JCM.00442-17

12. Mishra, N., Caciula, A., Price, A., Thakkar, R., Ng, J., Chauhan, L. V., Jain, K.,

Che, X., Espinosa, D. A., Montoya Cruz, M., Balmaseda, A., Sullivan, E. H.,

Patel, J. J., Jarman, R. G., Rakeman, J. L., Egan, C. T., Reusken, C.,

Koopmans, M. P. G., Harris, E., Tokarz, R., Briese, T., & Lipkin, W. I. (2018).

Diagnosis of Zika Virus Infection by Peptide Array and Enzyme-Linked

Immunosorbent Assay. *MBio, 9*(2). doi:10.1128/mBio.00095-18

13. Paixao, E. S., Teixeira, M. G., & Rodrigues, L. C. (2018). Zika, chikungunya and

dengue: the causes and threats of new and re-emerging arboviral diseases. *BMJ*

Glob Health, 3(Suppl 1), e000530. doi:10.1136/bmjgh-2017-000530

14. Carrillo-Hernandez, M. Y., Ruiz-Saenz, J., Villamizar, L. J., Gomez-Rangel, S. Y.,

& Martinez-Gutierrez, M. (2018). Co-circulation and simultaneous co-infection of

dengue, chikungunya, and zika viruses in patients with febrile syndrome at the

Colombian-Venezuelan border. *BMC Infect Dis, 18*(1), 61. doi:10.1186/s12879-

018-2976-1

15. Azeredo, E. L., Dos Santos, F. B., Barbosa, L. S., Souza, T. M. A., Badolato-

Correa, J., Sanchez-Arcila, J. C., Nunes, P. C. G., de-Oliveira-Pinto, L. M., de

Filippis, A. M., Dal Fabbro, M., Hoscher Romanholi, I., & Venancio da Cunha, R.

(2018). Clinical and Laboratory Profile of Zika and Dengue Infected Patients:

Lessons Learned From the Co-circulation of Dengue, Zika and Chikungunya in

Brazil. *PLoS Curr, 10*.

doi:10.1371/currents.outbreaks.0bf6aeb4d30824de63c4d5d745b217f5

16. Dudley, D. M., Aliota, M. T., Mohr, E. L., Weiler, A. M., Lehrer-Brey, G.,

Weisgrau, K. L., Mohns, M. S., Breitbach, M. E., Rasheed, M. N., Newman, C.

M., Gellerup, D. D., Moncla, L. H., Post, J., Schultz-Darken, N., Schotzko, M. L.,

Hayes, J. M., Eudailey, J. A., Moody, M. A., Permar, S. R., O'Connor, S. L.,

Rakasz, E. G., Simmons, H. A., Capuano, S., Golos, T. G., Osorio, J. E.,

Friedrich, T. C., & O'Connor, D. H. (2016). A rhesus macaque model of Asian-

lineage Zika virus infection. *Nat Commun, 7*, 12204. doi:10.1038/ncomms12204

17. Nguyen, S. M., Antony, K. M., Dudley, D. M., Kohn, S., Simmons, H. A., Wolfe,

B., Salamat, M. S., Teixeira, L. B. C., Wiepz, G. J., Thoong, T. H., Aliota, M. T.,

Weiler, A. M., Barry, G. L., Weisgrau, K. L., Vosler, L. J., Mohns, M. S.,

Breitbach, M. E., Stewart, L. M., Rasheed, M. N., Newman, C. M., Graham, M.

E., Wieben, O. E., Turski, P. A., Johnson, K. M., Post, J., Hayes, J. M., Schultz-

Darken, N., Schotzko, M. L., Eudailey, J. A., Permar, S. R., Rakasz, E. G., Mohr,

E. L., Capuano, S., 3rd, Tarantal, A. F., Osorio, J. E., O'Connor, S. L., Friedrich,

T. C., O'Connor, D. H., & Golos, T. G. (2017). Highly efficient maternal-fetal Zika

virus transmission in pregnant rhesus macaques. *PLoS Pathog, 13*(5),

e1006378. doi:10.1371/journal.ppat.1006378

18. Martinot, A. J., Abbink, P., Afacan, O., Prohl, A. K., Bronson, R., Hecht, J. L.,

Borducchi, E. N., Larocca, R. A., Peterson, R. L., Rinaldi, W., Ferguson, M.,

Didier, P. J., Weiss, D., Lewis, M. G., De La Barrera, R. A., Yang, E., Warfield, S.

K., & Barouch, D. H. (2018). Fetal Neuropathology in Zika Virus-Infected

Pregnant Female Rhesus Monkeys. *Cell, 173*(5), 1111-1122 e1110.

doi:10.1016/j.cell.2018.03.019

19. Osuna, C. E., Lim, S. Y., Deleage, C., Griffin, B. D., Stein, D., Schroeder, L. T.,

Omange, R. W., Best, K., Luo, M., Hraber, P. T., Andersen-Elyard, H., Ojeda, E.

F., Huang, S., Vanlandingham, D. L., Higgs, S., Perelson, A. S., Estes, J. D.,

Safronetz, D., Lewis, M. G., & Whitney, J. B. (2016). Zika viral dynamics and

shedding in rhesus and cynomolgus macaques. *Nat Med, 22*(12), 1448-1455.

doi:10.1038/nm.4206

20. Hirsch, A. J., Smith, J. L., Haese, N. N., Broeckel, R. M., Parkins, C. J.,

Kreklywich, C., DeFilippis, V. R., Denton, M., Smith, P. P., Messer, W. B., Colgin,

L. M., Ducore, R. M., Grigsby, P. L., Hennebold, J. D., Swanson, T., Legasse, A.

W., Axthelm, M. K., MacAllister, R., Wiley, C. A., Nelson, J. A., & Streblow, D. N.

(2017). Zika Virus infection of rhesus macaques leads to viral persistence in

multiple tissues. *PLoS Pathog, 13*(3), e1006219.

doi:10.1371/journal.ppat.1006219

21. Hirsch, A. J., Roberts, V. H. J., Grigsby, P. L., Haese, N., Schabel, M. C., Wang,

X., Lo, J. O., Liu, Z., Kroenke, C. D., Smith, J. L., Kelleher, M., Broeckel, R.,

Kreklywich, C. N., Parkins, C. J., Denton, M., Smith, P., DeFilippis, V., Messer,

W., Nelson, J. A., Hennebold, J. D., Grafe, M., Colgin, L., Lewis, A., Ducore, R.,

Swanson, T., Legasse, A. W., Axthelm, M. K., MacAllister, R., Moses, A. V.,

Morgan, T. K., Frias, A. E., & Streblow, D. N. (2018). Zika virus infection in

pregnant rhesus macaques causes placental dysfunction and immunopathology.

Nat Commun, 9(1), 263. doi:10.1038/s41467-017-02499-9

22. Mohr, E. L., Block, L. N., Newman, C. M., Stewart, L. M., Koenig, M., Semler, M.,

Breitbach, M. E., Teixeira, L. B. C., Zeng, X., Weiler, A. M., Barry, G. L., Thoong,

T. H., Wiepz, G. J., Dudley, D. M., Simmons, H. A., Mejia, A., Morgan, T. K.,

Salamat, M. S., Kohn, S., Antony, K. M., Aliota, M. T., Mohns, M. S., Hayes, J.

M., Schultz-Darken, N., Schotzko, M. L., Peterson, E., Capuano, S., 3rd, Osorio,

J. E., O'Connor, S. L., Friedrich, T. C., O'Connor, D. H., & Golos, T. G. (2018).

Ocular and uteroplacental pathology in a macaque pregnancy with congenital Zika virus infection. *PLoS One, 13*(1), e0190617. doi:10.1371/journal.pone.0190617

23. Newman, C. M., Dudley, D. M., Aliota, M. T., Weiler, A. M., Barry, G. L., Mohns, M. S., Breitbach, M. E., Stewart, L. M., Buechler, C. R., Graham, M. E., Post, J., Schultz-Darken, N., Peterson, E., Newton, W., Mohr, E. L., Capuano, S., 3rd, O'Connor, D. H., & Friedrich, T. C. (2017). Oropharyngeal mucosal transmission of Zika virus in rhesus macaques. *Nat Commun, 8*(1), 169. doi:10.1038/s41467-017-00246-8

24. Dudley, D. M., Newman, C. M., Lalli, J., Stewart, L. M., Koenig, M. R., Weiler, A. M., Semler, M. R., Barry, G. L., Zarbock, K. R., Mohns, M. S., Breitbach, M. E., Schultz-Darken, N., Peterson, E., Newton, W., Mohr, E. L., Capuano, S., III, Osorio, J. E., O'Connor, S. L., O'Connor, D. H., Friedrich, T. C., & Aliota, M. T. (2017). Infection via mosquito bite alters Zika virus tissue tropism and replication kinetics in rhesus macaques. *Nat Commun, 8*(1), 2096. doi:10.1038/s41467-017-02222-8

25. Haddow, A. D., Nalca, A., Rossi, F. D., Miller, L. J., Wiley, M. R., Perez-Sautu, U., Washington, S. C., Norris, S. L., Wollen-Roberts, S. E., Shamblin, J. D., Kimmel, A. E., Bloomfield, H. A., Valdez, S. M., Sprague, T. R., Principe, L. M., Bellanca, S. A., Cinkovich, S. S., Lugo-Roman, L., Cazares, L. H., Pratt, W. D., Palacios, G. F., Bavari, S., Pitt, M. L., & Nasar, F. (2017). High Infection Rates for Adult Macaques after Intravaginal or Intrarectal Inoculation with Zika Virus. *Emerg Infect Dis, 23*(8), 1274-1281. doi:10.3201/eid2308.170036

26. Adams Waldorf, K. M., Stencel-Baerenwald, J. E., Kapur, R. P., Studholme, C.,

Boldenow, E., Vornhagen, J., Baldessari, A., Dighe, M. K., Thiel, J., Merillat, S.,

Armistead, B., Tisoncik-Go, J., Green, R. R., Davis, M. A., Dewey, E. C.,

Fairgrieve, M. R., Gatenby, J. C., Richards, T., Garden, G. A., Diamond, M. S.,

Juul, S. E., Grant, R. F., Kuller, L., Shaw, D. W., Ogle, J., Gough, G. M., Lee, W.,

English, C., Hevner, R. F., Dobyns, W. B., Gale, M., Jr., & Rajagopal, L. (2016).

Fetal brain lesions after subcutaneous inoculation of Zika virus in a pregnant

nonhuman primate. *Nat Med, 22*(11), 1256-1259. doi:10.1038/nm.4193

27. Adams Waldorf, K. M., Nelson, B. R., Stencel-Baerenwald, J. E., Studholme, C.,

Kapur, R. P., Armistead, B., Walker, C. L., Merillat, S., Vornhagen, J., Tisoncik-

Go, J., Baldessari, A., Coleman, M., Dighe, M. K., Shaw, D. W. W., Roby, J. A.,

Santana-Ufret, V., Boldenow, E., Li, J., Gao, X., Davis, M. A., Swanstrom, J. A.,

Jensen, K., Widman, D. G., Baric, R. S., Medwid, J. T., Hanley, K. A., Ogle, J.,

Gough, G. M., Lee, W., English, C., Durning, W. M., Thiel, J., Gatenby, C.,

Dewey, E. C., Fairgrieve, M. R., Hodge, R. D., Grant, R. F., Kuller, L., Dobyns,

W. B., Hevner, R. F., Gale, M., Jr., & Rajagopal, L. (2018). Congenital Zika virus

infection as a silent pathology with loss of neurogenic output in the fetal brain.

Nat Med, 24(3), 368-374. doi:10.1038/nm.4485

28. Forsstrom, B., Axnas, B. B., Stengele, K. P., Buhler, J., Albert, T. J., Richmond,

T. A., Hu, F. J., Nilsson, P., Hudson, E. P., Rockberg, J., & Uhlen, M. (2014).

Proteome-wide epitope mapping of antibodies using ultra-dense peptide arrays.

Mol Cell Proteomics, 13(6), 1585-1597. doi:10.1074/mcp.M113.033308

29. Zandian, A., Forsstrom, B., Haggmark-Manberg, A., Schwenk, J. M., Uhlen, M.,

Nilsson, P., & Ayoglu, B. (2017). Whole-Proteome Peptide Microarrays for

Profiling Autoantibody Repertoires within Multiple Sclerosis and Narcolepsy. *J*

Proteome Res, 16(3), 1300-1314. doi:10.1021/acs.jproteome.6b00916

30. Engmark, M., Andersen, M. R., Laustsen, A. H., Patel, J., Sullivan, E., de Masi,

F., Hansen, C. S., Kringelum, J. V., Lomonte, B., Gutierrez, J. M., & Lund, O.

(2016). High-throughput immuno-profiling of mamba (Dendroaspis) venom toxin

epitopes using high-density peptide microarrays. *Sci Rep, 6*, 36629.

doi:10.1038/srep36629

31. Steffen, W., Ko, F. C., Patel, J., Lyamichev, V., Albert, T. J., Benz, J., Rudolph,

M. G., Bergmann, F., Streidl, T., Kratzsch, P., Boenitz-Dulat, M., Oelschlaegel,

T., & Schraeml, M. (2017). Discovery of a microbial transglutaminase enabling

highly site-specific labeling of proteins. *J Biol Chem, 292*(38), 15622-15635.

doi:10.1074/jbc.M117.797811

32. Lyamichev, V. I., Goodrich, L. E., Sullivan, E. H., Bannen, R. M., Benz, J., Albert,

T. J., & Patel, J. J. (2017). Stepwise Evolution Improves Identification of Diverse

Peptides Binding to a Protein Target. *Sci Rep, 7*(1), 12116. doi:10.1038/s41598-

017-12440-1

33. Christiansen, A., Kringelum, J. V., Hansen, C. S., Bogh, K. L., Sullivan, E., Patel,

J., Rigby, N. M., Eiwegger, T., Szepfalusi, Z., de Masi, F., Nielsen, M., Lund, O.,

& Dufva, M. (2015). High-throughput sequencing enhanced phage display

enables the identification of patient-specific epitope motifs in serum. *Sci Rep, 5*,

12913. doi:10.1038/srep12913

34. Tokarz, R., Mishra, N., Tagliafierro, T., Sameroff, S., Caciula, A., Chauhan, L.,

Patel, J., Sullivan, E., Gucwa, A., Fallon, B., Golightly, M., Molins, C., Schriefer,

M., Marques, A., Briese, T., & Lipkin, W. I. (2018). A multiplex serologic platform

for diagnosis of tick-borne diseases. *Sci Rep, 8*(1), 3158. doi:10.1038/s41598-

018-21349-2

35. Bailey, A. L., Buechler, C. R., Matson, D. R., Peterson, E. J., Brunner, K. G.,

Mohns, M. S., Breitbach, M., Stewart, L. M., Ericsen, A. J., Newman, C. M.,

Koenig, M. R., Mohr, E., Tan, J., Capuano, S., 3rd, Simmons, H. A., Yang, D. T.,

& O'Connor, D. H. (2017). Pegivirus avoids immune recognition but does not

attenuate acute-phase disease in a macaque model of HIV infection. *PLoS*

Pathog, 13(10), e1006692. doi:10.1371/journal.ppat.1006692

36. Pahar, B., Kenway-Lynch, C. S., Marx, P., Srivastav, S. K., LaBranche, C.,

Montefiori, D. C., & Das, A. (2016). Breadth and magnitude of antigen-specific

antibody responses in the control of plasma viremia in simian immunodeficiency

virus infected macaques. *Virol J, 13*(1), 200. doi:10.1186/s12985-016-0652-x

37. Aliota, M. T., Dudley, D. M., Newman, C. M., Mohr, E. L., Gellerup, D. D.,

Breitbach, M. E., Buechler, C. R., Rasheed, M. N., Mohns, M. S., Weiler, A. M.,

Barry, G. L., Weisgrau, K. L., Eudailey, J. A., Rakasz, E. G., Vosler, L. J., Post,

J., Capuano, S., 3rd, Golos, T. G., Permar, S. R., Osorio, J. E., Friedrich, T. C.,

O'Connor, S. L., & O'Connor, D. H. (2016). Heterologous Protection against

Asian Zika Virus Challenge in Rhesus Macaques. *PLoS Negl Trop Dis, 10*(12),

e0005168. doi:10.1371/journal.pntd.0005168

38. Aliota, M. T., Dudley, D. M., Newman, C. M., Weger-Lucarelli, J., Stewart, L. M.,

Koenig, M. R., Breitbach, M. E., Weiler, A. M., Semler, M. R., Barry, G. L.,

Zarbock, K. R., Haj, A. K., Moriarty, R. V., Mohns, M. S., Mohr, E. L., Venturi, V.,

Schultz-Darken, N., Peterson, E., Newton, W., Schotzko, M. L., Simmons, H. A.,

Mejia, A., Hayes, J. M., Capuano, S., 3rd, Davenport, M. P., Friedrich, T. C.,

Ebel, G. D., O'Connor, S. L., & O'Connor, D. H. (2018). Molecularly barcoded

Zika virus libraries to probe in vivo evolutionary dynamics. *PLoS Pathog, 14*(3),

e1006964. doi:10.1371/journal.ppat.1006964

39. Lindsey, N. P., Staples, J. E., Powell, K., Rabe, I. B., Fischer, M., Powers, A. M.,

Kosoy, O. I., Mossel, E. C., Munoz-Jordan, J. L., Beltran, M., Hancock, W. T.,

Toews, K. E., Ellis, E. M., Ellis, B. R., Panella, A. J., Basile, A. J., Calvert, A. E.,

Laven, J., Goodman, C. H., Gould, C. V., Martin, S. W., Thomas, J. D.,

Villanueva, J., Mataia, M. L., Sciulli, R., Gose, R., Whelen, A. C., & Hills, S. L.

(2018). Ability To Serologically Confirm Recent Zika Virus Infection in Areas with

Varying Past Incidence of Dengue Virus Infection in the United States and U.S.

Territories in 2016. *J Clin Microbiol, 56*(1). doi:10.1128/JCM.01115-17

40. Tsai, W. Y., Youn, H. H., Brites, C., Tsai, J. J., Tyson, J., Pedroso, C., Drexler, J.

F., Stone, M., Simmons, G., Busch, M. P., Lanteri, M., Stramer, S. L.,

Balmaseda, A., Harris, E., & Wang, W. K. (2017). Distinguishing Secondary

Dengue Virus Infection From Zika Virus Infection With Previous Dengue by a

Combination of 3 Simple Serological Tests. *Clin Infect Dis, 65*(11), 1829-1836.

doi:10.1093/cid/cix672

41. Ronnberg, B., Gustafsson, A., Vapalahti, O., Emmerich, P., Lundkvist, A.,

Schmidt-Chanasit, J., & Blomberg, J. (2017). Compensating for cross-reactions

using avidity and computation in a suspension multiplex immunoassay for

serotyping of Zika versus other flavivirus infections. *Med Microbiol Immunol,*

206(5), 383-401. doi:10.1007/s00430-017-0517-y

42. Subissi, L., Daudens-Vaysse, E., Cassadou, S., Ledrans, M., Bompard, P.,

Gustave, J., Aubry, M., Cao-Lormeau, V. M., & Mallet, H. P. (2017). Revising

rates of asymptomatic Zika virus infection based on sentinel surveillance data

from French Overseas Territories. *Int J Infect Dis, 65*, 116-118.

doi:10.1016/j.ijid.2017.10.009

43. Palmeira, P., Quinello, C., Silveira-Lessa, A. L., Zago, C. A., & Carneiro-

Sampaio, M. (2012). IgG placental transfer in healthy and pathological

pregnancies. *Clin Dev Immunol, 2012*, 985646. doi:10.1155/2012/985646

44. Dai, L., Song, J., Lu, X., Deng, Y. Q., Musyoki, A. M., Cheng, H., Zhang, Y.,

Yuan, Y., Song, H., Haywood, J., Xiao, H., Yan, J., Shi, Y., Qin, C. F., Qi, J., &

Gao, G. F. (2016). Structures of the Zika Virus Envelope Protein and Its Complex

with a Flavivirus Broadly Protective Antibody. *Cell Host Microbe, 19*(5), 696-704.

doi:10.1016/j.chom.2016.04.013

45. Barba-Spaeth, G., Dejnirattisai, W., Rouvinski, A., Vaney, M. C., Medits, I.,

Sharma, A., Simon-Loriere, E., Sakuntabhai, A., Cao-Lormeau, V. M., Haouz, A.,

England, P., Stiasny, K., Mongkolsapaya, J., Heinz, F. X., Screaton, G. R., &

Rey, F. A. (2016). Structural basis of potent Zika-dengue virus antibody cross-

neutralization. *Nature, 536*(7614), 48-53. doi:10.1038/nature18938

46. Wang, Q., Yang, H., Liu, X., Dai, L., Ma, T., Qi, J., Wong, G., Peng, R., Liu, S.,

Li, J., Li, S., Song, J., Liu, J., He, J., Yuan, H., Xiong, Y., Liao, Y., Li, J., Yang, J.,

Tong, Z., Griffin, B. D., Bi, Y., Liang, M., Xu, X., Qin, C., Cheng, G., Zhang, X.,

Wang, P., Qiu, X., Kobinger, G., Shi, Y., Yan, J., & Gao, G. F. (2016). Molecular

determinants of human neutralizing antibodies isolated from a patient infected

with Zika virus. *Sci Transl Med, 8*(369), 369ra179.

doi:10.1126/scitranslmed.aai8336

47. Zhang, S., Kostyuchenko, V. A., Ng, T. S., Lim, X. N., Ooi, J. S., Lambert, S.,

Tan, T. Y., Widman, D. G., Shi, J., Baric, R. S., & Lok, S. M. (2016).

Neutralization mechanism of a highly potent antibody against Zika virus. *Nat*

Commun, 7, 13679. doi:10.1038/ncomms13679

48. Hasan, S. S., Miller, A., Sapparapu, G., Fernandez, E., Klose, T., Long, F.,

Fokine, A., Porta, J. C., Jiang, W., Diamond, M. S., Crowe, J. E., Jr., Kuhn, R. J.,

& Rossmann, M. G. (2017). A human antibody against Zika virus crosslinks the E

protein to prevent infection. *Nat Commun, 8*, 14722. doi:10.1038/ncomms14722

49. Robbiani, D. F., Bozzacco, L., Keeffe, J. R., Khouri, R., Olsen, P. C., Gazumyan,

A., Schaefer-Babajew, D., Avila-Rios, S., Nogueira, L., Patel, R., Azzopardi, S.

A., Uhl, L. F. K., Saeed, M., Sevilla-Reyes, E. E., Agudelo, M., Yao, K. H.,

Golijanin, J., Gristick, H. B., Lee, Y. E., Hurley, A., Caskey, M., Pai, J., Oliveira,

T., Wunder, E. A., Jr., Sacramento, G., Nery, N., Jr., Orge, C., Costa, F., Reis, M.

G., Thomas, N. M., Eisenreich, T., Weinberger, D. M., de Almeida, A. R. P.,

West, A. P., Jr., Rice, C. M., Bjorkman, P. J., Reyes-Teran, G., Ko, A. I.,

MacDonald, M. R., & Nussenzweig, M. C. (2017). Recurrent Potent Human

Neutralizing Antibodies to Zika Virus in Brazil and Mexico. *Cell, 169*(4), 597-609

e511. doi:10.1016/j.cell.2017.04.024

Figures and Tables

Figure 1: SIV env epitopes identified by a high-density peptide microarray overlap

known epitopes in variable loop regions. Serum from two cynomolgus macaques

(animals A1 and A2) before and after infection with SIVmac239 was evaluated for

reactivity to overlapping peptides representing the SIVmac239 env protein sequence on

a linear peptide microarray. SIV env variable loop regions (36) are highlighted in grey.

Figure 2: Reactivity of ZIKV-convalescent animals against the ZIKV-FP polyprotein.

Serum from rhesus macaques infected with ZIKV-FP (animals B1, D1, and D2) and

ZIKV-MR766 (animal C1) was assayed for IgG recognition of the ZIKV-FP polyprotein.

Reactivity at pre-infection and 28 dpi is shown. The $NS2B_{1427-1451}$RD25 epitope is

highlighted in grey. Using the mean signal intensity across this highlighted region, the

change of signal intensity from 0 dpi to 28 dpi was statistically significant versus no

change, by a two-tailed log-ratio t-test (p-value = 0.038 < 0.05, df = 3).

Figure 3: Reactivity of ZIKV-convalescent animals against polyproteins of the four

DENV serotypes. DENV serotypes share an average of 41% sequence similarity with

the aligned ZIKV NS2B epitope (A). Serum from rhesus macaques infected with ZIKV-

FP (animals B1, D1, and D2) and ZIKV-MR766 (C1) was assayed for reactivity against

polyproteins of DENV serotypes 1-4 (B-E, respectively). The segment of DENV NS2B

which aligns with the ZIKV $NS2B_{1427-1451}RD25$ epitope is highlighted in grey.

Figure 4: Antibody reactivity to the ZIKV NS2B epitope of animals with recent infection

with DENV. Three macaques (F1, F2, and F3) with history of challenge and re-

challenge with ZIKV-FP 12 and 9.5 months prior were infected with DENV-2 and their

serum was assessed before infection and at 28 dpi. The $NS2B_{1427-1451}RD25$ epitope is

highlighted in grey.

Figure 5: IgG reactivity against the ZIKV NS2B epitope during pregnancy. Serum

samples taken throughout the course of six animals' pregnancies were evaluated

against the ZIKV-FP polyprotein represented on peptide microarrays. Two animals (G1

and G2) were infected with ZIKV-FP at 36-38 gd. Three animals (H1, H2, H3) were

infected with ZIKV-PR at 45-47 gd. One animal (I1) had been infected with DENV-3 nine

months prior and was infected with ZIKV-PR at 35 gd. The $NS2B_{1427-1451}RD25$ epitope is

highlighted in grey. Using the mean across this highlighted region, the change of signal

intensity from 0 dpi to 21 or 29 dpi was statistically significant versus no change, by a

two-tailed log-ratio t-test (p-value = 0.017 < 0.05, df = 5).

Table 1

Cohort		Animal	Public animal identifier*	Species	Sex	Flavivirus infection history	Inoculation strain	Inoculation dose and route	Time points of sample collection (days post infection, dpi)	Peptide array design
		A1	cy0879	Macaca fascicularis	female	none	SIVmac239	7,000 TCID50, intrarectal	pre-infection, 125 dpi	Offset by 4, overlap of 12
A		A2	cy0886	Macaca fascicularis	female	none	SIVmac239	7,000 TCID50, intrarectal	pre-infection, 126 dpi	Offset by 4, overlap of 12
B		B1	411359	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1	10E4 PFU, subcutaneous	pre-infection, 28, 67 and 99 dpi	Offset by 1, overlap of 15
C		C1	295022	Macaca mulatta	female	none	Zika virus/R.macaque-tc/UGA/1947/MR766-3329	10E4 PFU, subcutaneous	pre-infection, 28, 67 and 98 dpi	Offset by 1, overlap of 15
		D1	448436	Macaca mulatta	male	none	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1	10E4 PFU, subcutaneous	pre-infection, 28 and 621 dpi	Offset by 1, overlap of 15
D		D2	861138	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1	10E4 PFU, subcutaneous	pre-infection, 28 and 582 dpi	Offset by 1, overlap of 15
		F1	393422	Macaca mulatta	male	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1, 12 and 9.5 months prior	Dengue 2 New Guinea C	10E5 PFU, subcutaneous	pre-infection, 28 dpi	Offset by 4, overlap of 12
		F2	826226	Macaca mulatta	male	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1, 12 and 9.5 months prior	Dengue 2 New Guinea C	10E5 PFU, subcutaneous	pre-infection, 28 dpi	Offset by 4, overlap of 12
F		F3	912116	Macaca mulatta	male	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1, 12 and 9.5 months prior	Dengue 2 New Guinea C	10E5 PFU, subcutaneous	pre-infection, 28 dpi	Offset by 4, overlap of 12
		G1	660875	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1	10E4 PFU, subcutaneous	pre-infection, 7, 21, 43, 78, 127 dpi	Offset by 1, overlap of 15
G		G2	827577	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/FRA/2013/FrenchPolynesia-01_v1c1	10E4 PFU, subcutaneous	0, 7, 21, 35, 71, 113 days after primary infection	Offset by 4, overlap of 12
		H1	146523	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/PUR/2015/PRVABC59	10E4 PFU, subcutaneous	pre-infection, 7, 21, 70 dpi	Offset by 1, overlap of 15
		H2	858972	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/PUR/2015/PRVABC59	10E4 PFU, subcutaneous	pre-infection, 7, 21, 70 dpi	Offset by 1, overlap of 15
H		H3	419969	Macaca mulatta	female	none	Zika virus/H.sapiens-tc/PUR/2015/PRVABC59	10E4 PFU, subcutaneous	pre-infection, 7, 21, 70 dpi	Offset by 1, overlap of 15
I		I1	776301	Macaca mulatta	female	Dengue 3 Slenman/78, 9 months prior	Zika virus/H.sapiens-tc/PUR/2015/PRVABC59 bar code virus	10E4 PFU, subcutaneous	pre-infection, 8, 29, 57, 78 dpi	Offset by 4, overlap of 12

*Public animal identifiers are used in studies on the Zika Open Research Portal

(https://zika.labkey.com/project/home/begin.view?).

Table 2

Animal	Gestational days (gd) at inoculation	Pregnancy outcome
G1	36	delivery at 158 gd (122 dpi)
G2	38	delivery at 158 gd (120 dpi)
H1	46	fetectomy at 116 gd (70 dpi)
H2	47	fetectomy at 117 gd (70 dpi)
H3	43	fetectomy at 111 gd (68 dpi)
I1	35	delivery at 155 gd (120 dpi)