Title: Orexin-1 receptor signaling in ventral pallidum regulates demand for the opioid remifentanil

Running title: Orx-1 receptor signaling in VP mediates remifentanil demand

Aida Mohammadkhani, Ph.D.[1,2], Caroline B. Pantazis, B.A.[2], Hannah E. Bowrey, Ph.D.[2,3], Morgan H. James, Ph.D.[2,4], and Gary Aston-Jones, Ph.D.[2]

[1] School of Cognitive Sciences (SCS), Institute for Research in Fundamental Sciences (IPM), PO Box 1954851167, Tehran, Iran

[2] Brain Health Institute, Rutgers University, Piscataway, NJ 08854

[3] Save Sight Institute, University of Sydney, NSW Australia 2000

[4] Florey Institute of Neuroscience and Mental Health, University of Melbourne, VIC, Australia 3052

Corresponding author:
Gary Aston-Jones, Ph.D.
683 Hoes Lane West,
Piscataway NJ 08854
P: (732) 235 6077
F: (732) 235 5814
E: gsa35@ca.rutgers.edu

Submitting author:
Aida Mohammadkhani
mohammadkhani@ipm.ir

- **Number of Pages:** 25
- **Number of Words Abstract:** 254
- **Number of Words Introduction:** 628
- **Number of Words Discussion:** 1489

Funding and Disclosure:

This work was supported by financial support from the Institute for Research in Fundamental Sciences (IPM; AM), C.J. Martin Fellowships from the National Health and Medical Research Council of Australia to MHJ (No. 1072706) and HEB (No. 1128089), a U.S. Public Health Service award from the National Institute of Drug Abuse to GAJ (R01 DA006214), a NIH F31 grant to CBP (DA042588) and by the Charlotte and Murray Strongwater Endowment for Neuroscience and Brain Health (GAJ).

Acknowledgements

We would like to thank Drs. Kirsten Porter-Stranksy and Brandon Bentzley for their assistance with the remifentanil demand protocol. We would also like to thank Dr. Jennifer Fragale for her helpful guidance in interpreting the data, as well as Nupur Jain for her assistance.

Abstract:

Signaling at the orexin-1 receptor (Ox1R) is important for motivation for various drugs of abuse.

Recently, our laboratory showed that systemic blockade of Ox1Rs decreased motivation for the

potent and short-acting opioid remifentanil, as well as remifentanil consumption at null cost

(Porter-Stransky et al., 2017). However, the central site(s) through which orexin acts to regulate

opioid-seeking behaviors are not known. We investigated ventral pallidum (VP) as a potential

site of orexin action, as VP is a known mediator of opioid reward and is densely innervated by

orexin-immunoreactive fibers. We used a within-session behavioral economics (BE) paradigm in

which remifentanil price (responses/µg iv remifentanil) was sequentially increased throughout

the session. Rats were implanted with bilateral cannulae into caudal VP (cVP), through which

microinjections of SB334867 (SB), an Ox1R antagonist, were given prior to BE and

reinstatement testing. Contrary to systemic SB treatment, intra-cVP SB preferentially reduced

motivation (increased demand elasticity and reinstatement) for reminfentanil, without affecting

remifentanil consumption at null cost. These effects were specific to cVP, as control

microinjections of SB immediately dorsal to cVP did not affect remifentanil-seeking. Baseline

demand for remifentanil predicted cued reinstatement of remifentanil seeking and the efficacy

with which intra-cVP SB administration reduced motivation for remifentanil. These findings

indicate a specific role for cVP Ox1R signaling in mediating the motivational properties of the

opioid remifentanil, distinct from its consumption at null cost. Our findings also highlight the BE

paradigm as an effective biomarker for predicting opioid addiction behaviors and treatment

efficacy.

Significance Statement

Abuse of prescription opioids has risen rapidly and continues to be a major health issue. There is significant urgency to understand the neurobiological and behavioral mechanisms underlying opioid addiction. Here we determined the role of caudal ventral pallidum (cVP) orexin signaling in remifentanil seeking using a within-session behavioral economics paradigm. We report that intra-cVP infusions of the orexin-1 receptor antagonist SB334867 (SB) blocked motivated responding for remifentanil as well as cued 'relapse' of remifentanil seeking. In contrast, SB intra-cVP did not affect consumption of remifentanil under low-effort conditions. Together, our findings indicate that decreasing orexin-1 receptor signaling in cVP is a potential strategy for the treatment of opioid abuse.

Introduction:

In recent years, abuse of prescription opioids has risen rapidly and has become a major health issue (Compton and Volkow, 2006; Lewis et al., 2017; Volkow and Collins, 2017; Dobbs and Fogger, 2018). Thus, it is urgent to understand the neurobiological and behavioral mechanisms underlying opioid addiction to facilitate development of new treatments for this disorder. The hypothalamic orexin (hypocretin) system is a key modulator of drug seeking behaviors across a variety of drugs of abuse, including opioids (Harris et al., 2005; Mahler et al., 2014a; Baimel et al., 2015; James et al., 2017). Orexin-containing neurons are located in hypothalamus and project widely throughout the brain, targeting two G-protein-coupled receptors (orexin receptors 1 and 2; Ox1R and Ox2R, respectively). In particular, signaling at Ox1R has been shown to mediate motivation and addiction (for review, James et al., 2018). We found that blocking Ox1Rs can disrupt opioid seeking behaviors: systemic blockade of Ox1R signaling reduced breakpoints for heroin in a progressive ratio task (Smith and Aston-Jones, 2012) and attenuated demand (motivation) for remifentanil in a behavioral economics (BE) paradigm (Porter-Stransky et al., 2017). Systemic Ox1R blockade also reduced low-effort (fixed ratio [FR] 1) self-administration of heroin and remifentanil, indicating a potential additional role for the orexin system in mediating the low-effort consumption of opiates. Despite these data, little is known about where in the reward circuitry orexin acts to contribute to opioid reward behavior.

One region of interest is ventral pallidum (VP), as this brain structure is innervated by orexin fibers (Peyron et al., 1998; Baldo et al., 2003) and VP neurons express both Ox1Rs and Ox2Rs (Marcus et al., 2001). Human studies indicate that drug-related cues robustly activate VP (Childress et al., 2008), and inactivation of VP in rats prevents various drug behaviors, including acquisition and expression of learned preferences for environments paired with morphine reward

(Dallimore et al., 2006). Moreover, VP inactivation decreases FR5 heroin self-administration

(Hubner and Koob, 1990). Interestingly, caudal VP (cVP) contains the greatest density of orexin

fibers in the VP (Baldo et al., 2003), and previous studies found that orexin microinjection into

this region enhances the hedonic impact of sucrose (Ho and Berridge, 2013). Taken together,

these studies indicate a potential role for orexin signaling in cVP in opioid reward.

The present study was designed to test the hypothesis that orexin signaling in cVP regulates

demand (motivation or low-cost consumption) for the ultra-short acting opioid remifentanil. We

focused on remifentanil because its short half-life lends itself well to BE analysis with the

threshold procedure (Glass et al., 1999), and it contributes to prescription opioid abuse (Zacny

and Galinkin, 1999; Baylon et al., 2000; Panlilio and Schindler, 2000; Levine and Bryson, 2010).

We determined remifentanil demand using a within-session BE paradigm that provides a

quantitative measure of both the desired amount of remifentanil at low effort, and the motivation

for this drug as the required effort increases. We first sought to confirm the results of our

previous study, which found that systemic administration of the selective Ox1R antagonist

SB334867 (SB) reduced motivation (increased demand elasticity; α), and reduced free

consumption (Q_0), of remifentanil. We then compared these effects to those following local

microinfusion of SB into cVP. In agreement with our previous study, systemic SB reduced

motivation (increased demand elasticity; α) and decreased free consumption of remifentanil

(Q_0). In contrast, intra-cVP SB increased demand elasticity for remifentanil without affecting

free consumption, indicating a preferential role for orexin signaling in cVP in motivation for

remifentanil. Both systemic and local blockade of Ox1R signaling in cVP also reduced cued

reinstatement of remifentanil seeking. Together, our findings point to cVP as a critical site of

Ox1R signaling that drives motivation for remifentanil reward, and thus highlight a novel target

for therapeutics designed to treat opioid addiction.

Materials and Methods

Subjects

Male Sprague Dawley rats (n=24; initial weight 275–300 g; Charles River, Raleigh, USA) were

single-housed and maintained under a 12h reverse light/dark cycle (lights off at 08:00h) in a

temperature and humidity-controlled animal facility at Rutgers University. Food and water were

available ad libitum. All experimental procedures were approved by the Rutgers Institutional

Animal Care and Use Committee and were conducted according to the Guide for the Care and

Use of Laboratory Animals. Rats were handled daily after a 3-day acclimation period at the

facility. All experiments were performed in the rats' active (dark) phase.

Drugs

Remifentanil (obtained from the National Institute of Drug Abuse (NIDA) Drug Supply

Program, Research Triangle Park, USA) was dissolved in 0.9% sterile saline for intravenous (iv)

self-administration. SB334867 (SB), a selective antagonist of the Ox1R was supplied by the

NIDA Drug Supply Program and dissolved in sterile artificial cerebrospinal fluid (aCSF) at a

concentration of 1mM for VP microinjections, as described in our previous study (Mahler et al.,

2013). For systemic administration, SB was suspended in 10% 2-hydroxypropyl-b-cyclodextrin

in sterile water and 2% DMSO (Sigma-Aldrich, USA), as in our prior studies (Bentzley and

Aston-Jones, 2015; Porter-Stransky et al., 2017; James et al., 2018a; James et al., 2018c). SB

(30mg/kg) or vehicle was injected at a volume of 4.0ml/kg i.p. 30min prior to testing. We used

this dose of SB because we previously reported that lower doses of SB (10mg/kg) did not affect

demand elasticity, free consumption, or reinstatement responding for remifentanil (Porter-

Stransky et al., 2017). A within-subjects design was used whereby each rat received both SB and

vehicle, and the order was counterbalanced across subjects.

Intravenous catheter and stereotaxic surgery

Rats were implanted with an indwelling catheter into the jugular vein as described previously

(McGlinchey et al., 2016). Rats allocated to VP experiments were then placed in a stereotaxic

frame (Kopf, Tujunga, USA) and implanted with bilateral stainless steel guide cannulae (22G ,

11mm, Plastics One) directed 2mm dorsal to cVP (−0.8mm posterior, ±2.6mm medial–lateral,

−7.5mm ventral, relative to bregma (Paxinos and Watson, 1998). Cannulae were secured to the

skull using jeweler's screws and dental acrylic, and stylets were placed into the guide cannula to

prevent occlusion.

Dorsal control and cVP microinfusions

Animals received a mock microinjection the day prior to the first microinfusion test - injection

cannulae (28G) were bilaterally inserted into the guide cannula and kept in place for 1 min (no

infusions). On test days, rats received microinfusions (0.3μl/side) of either SB (1 mM) or vehicle

(aCSF) 5min prior to behavioral testing. Microinjections took place over a 70-sec period, and

injectors were left in place for 1min following infusions to limit backflow of the injectate. We

have found the dose of SB used here to be effective in other brain areas in prior publications

(Harris et al., 2007; Espana et al., 2010; James et al., 2011; Mahler et al., 2013). SB and vehicle

microinfusions were counterbalanced in a within-subjects design and administered via

polyethylene tubing connected to gastight 10μl Hamilton syringes (Hamilton, USA) set in an

infusion pump (Model 975, Harvard Apparatus, USA).

Control microinjections of SB were first carried out 2 mm dorsal to VP to confirm that our

observed changes in behavior were not due to dorsal diffusion of SB. This within-subjects

control was performed using injectors that projected 0.2mm below the tip of the guide cannula.

Dorsal control microinjections were performed in a session prior to VP microinjection sessions.

Each rat received a maximum of 6 VP microinfusions in a counterbalanced within-subjects

design. These included microinjections of SB and vehicle for BE performance, cue-induced

reinstatement, and locomotor activity.

Localization of injection sites

Following the final behavioral test, rats were deeply anesthetized with ketamine/xylazine

(56.6⁄8.7mg⁄kg) and received bilateral microinfusions of pontamine sky blue via the VP infusion

cannulae (0.3μl). Brains were flash-frozen in 2-methylbutane, sectioned into 40μm-thick

sections, mounted on glass slides, Nissl-stained with neutral red, and cover slipped to localize

cannula tract damage and verify injection sites. We sought to test the effect of SB in cVP (AP: -

0.1 to -0.8mm relative to bregma), so rats with misplaced VP cannulae that were located in

rostral VP (rVP; AP>0.0 mm relative to bregma; n=5) were analyzed as a separate group.

Self-administration training

The self-administration procedure was similar to a recently published study from our laboratory

(Porter-Stransky et al., 2017). All self-administration sessions occurred in standard Med

Associates operant chambers. A response on the active lever resulted in remifentanil infusion

(1μg over 4s) paired with light and tone cues. After each infusion, a 20s time-out occurred

(house light off) when additional presses did not yield remifentanil or cues. Presses on the

inactive lever had no consequences. Each session ended after 2h or when 80 infusions were

earned, whichever occurred first. Subjects were trained on a fixed ratio-1 (FR1) procedure for a

minimum of 6 sessions. After earning a minimum of 25 infusions within 2h for at least 3

consecutive sessions, animals advanced to the behavioral economics procedure.

Behavioral economics (BE) procedure

Rats were trained to self-administer remifentanil on an FR1 schedule (as described above) before

being trained on the BE task, as described previously (Porter-Stransky et al., 2017). Briefly,

during the BE task the cost of drug increased by decreasing the amount of drug infused per

active lever press every 10min. Therefore, each 110min session tested 11 doses of remifentanil

(2, 1, 0.6, 0.3, 0.2, 0.1 0.06, 0.03, 0.02, 0.01 and 0.006 μg/infusion). Each infusion was

accompanied by presentation of a light-tone cue; responses on the inactive lever had no

consequence. Demand curves were generated for each BE session as previously described

(Bentzley et al., 2013; Porter-Stransky et al., 2017) using an exponential demand equation

(Hursh and Silberberg, 2008) and least sum of squares approach, to generate estimates of

preferred remifentanil intake at zero cost (Q_0) and demand elasticity (α, the slope of the demand

curve). Larger α values indicate greater demand elasticity and are characterized by a greater

reduction in responding as drug price increases; this is interpreted as decreased motivation

(Bentzley et al., 2013). Smaller α values indicate lower demand elasticity and are symptomatic of

continued responding for drug despite increases in cost to obtain drug (increased motivation).

Subjects were trained daily on the BE procedure until stable responding was obtained (≤20%

variation in α and Q_0 over 3 consecutive days) before testing the effects of SB. Between drug

treatments, rats were re-tested on the BE procedure for a minimum of 3d and and until behavior

returned to pre-treatment levels.

 Extinction and reinstatement tests

After BE testing, rats were subjected to extinction training and locomotor activity testing

(described below). The order of extinction training and locomotor tests was counterbalanced.

During each 2h extinction session, lever presses yielded neither remifentanil nor cues. Rats

received extinction training for at least 7d and until they met the criteria of ≤15 active lever

presses for at least two consecutive sessions. Rats were given two 2h cue-induced reinstatement

tests with either SB or vehicle (counterbalanced), in which pressing the active lever yielded the

cues previously paired with remifentanil infusions, but no remifentanil. Rats received at least 2d

of extinction (≤15 active lever presses) between tests. One rat from the systemic SB group died

between BE and reinstatement testing, and thus was not included in extinction/reinstatement

analyses.

Locomotor testing

Rats were tested for locomotor activity as described previously (McGlinchey et al., 2016; James

et al., 2018b) in locomotor boxes (clear acrylic, 40 x 40 x 30 cm) with Digiscan monitors

(AccuScan Instruments). Rats were placed in a chamber for 2h to acclimate to the environment.

The next day, rats received a microinjection of either SB or aCSF into cVP 5min before being

tested in the locomotor chamber for 2h. Total, horizontal, and vertical locomotor activity were

recorded using beam beaks. A within-subjects design was utilized, and drug order was

counterbalanced across rats. Rats underwent a one-day washout period in between testing

sessions in which they were placed in the locomotor chamber but did not receive drug.

Data analysis

Analyses were performed in Graphpad Prism V6, except for multiple linear regression analyses

which were performed using SPSS Statistics (V19). Acquisition of self-administration,

individual differences in BE parameters, and the effects of SB on locomotor activity were

analyzed using repeated measures ANOVA and paired t-tests. α and Q_0 values following SB

treatment were transformed to represent percent change relative to their control injections. For

VP microinjection experiments, separate analyses were used to compare the effect of SB in cVP

and rVP relative to vehicle and dorsal control injections. The effects of SB on active/inactive

lever responding during reinstatement tests were assessed using separate 2-way repeated

measures ANOVAs with 'treatment' (extinction, vehicle, SB) and 'lever type' (active, inactive)

as the variables. Tukey post-hoc tests were applied to all significant repeated measures ANOVA

tests. A linear regression was used to correlate individual Q_0 and α values. For multiple linear

regression analyses, Q_0 and α values were set as the independent variables, with the dependent

variables being active lever responses during cued reinstatement, or the difference in responding

between SB and vehicle treatments.

Results

Experiment 1: Systemic blockade of Ox1Rs reduced economic demand, free consumption, and

cued reinstatement of remifentanil seeking

Figure 1a outlines the timeline of behavioral testing to investigate the effect of systemic

treatment with SB on remifentanil seeking behaviors. During FR1 training, rats made more

presses on the active lever than the inactive lever (main effect, $F_{(11,131)}=3.501$, p=0.0003)

beginning on the third session (Bonferroni post hoc test, p=0.0173) and throughout the remainder

of the training sessions (Fig. 1b). This was associated with an increase in remifentanil infusions

across training (one-way repeated measures ANOVA; $F_{(5,65)}=11.64$, p=0.0003; Fig. 1c). Next,

rats were trained on the BE procedure, to determine drug demand at varying costs in a within-

session design (Fig. 1d). There was a significant inverse correlation between baseline Q_0 and α in

these BE tests (r =-0.6783, p=0.0311 Fig. 1e), such that individuals with high Q_0 values exhibited

more persistent drug taking as the cost for remifentanil increased than individuals with low Q_0.

We next sought to confirm our previous findings that systemic blockade of Ox1R signaling

reduces reminfentanil demand (Porter-Stranksy et al., 2015). This provided a direct comparison

to intra-VP experiments in subsequent studies (below). Blockade of Ox1Rs with SB significantly

increased demand elasticity (decreased motivation for remifentanil, 30 mg/kg SB versus vehicle,

$t_9=2.551$, p=0.0312; Fig. 1f). Additionally, this dose of SB reduced remifentanil consumption at

low effort (free consumption) compared with vehicle (decreased Q_0, $t_9=2.382$, p=0.0411; Fig.

1g). To test the effect of systemic SB treatment on cued reinstatement behavior, lever pressing

was extinguished over a minimum period of 7d such that the number of active lever presses per

session was <15 (Fig. 1h). A 2×3 repeated measures ANOVA revealed a significant main effect

of treatment ($F_{2,\ 32}=7.720$, p=0.0018), and lever ($F_{2,32}=34.56$, p<0.0001), and a significant

treatment × lever interaction ($F_{2,32}=8.672$, p=0.001). Post-hoc analyses showed that animals

reinstated active lever-pressing to remifentanil-associated cues after vehicle pretreatment

(vehicle vs. extinction, p<0.0001), and that this was blocked by systemic administration of SB

(SB vs. vehicle, p=0.0005; Fig. 1i). There was no effect of SB (p=0.9441) or vehicle (p=0.9658)

pretreatment on inactive lever responding (Fig. 1i).

Experiment 2: Antagonism of Ox1Rs in cVP decreased motivation but not free consumption for

remifentanil, and decreased cued reinstatement of extinguished remifentanil seeking.

A separate group of animals with bilateral cannulae in cVP were used to test cVP as a potential

site of action mediating the effects of Ox1R signaling on demand for remifentanil (Fig. 2a). Rats

acquired remifentanil self-administration (Fig. 2b,c) and exhibited stable demand (α) for

remifentanil that was correlated with Q_0 values (Fig. 2d) as in Experiment 1. Microinfusion of

SB into cVP significantly increased demand elasticity (decreased motivation) compared to

microinjections of vehicle (aCSF) or dorsal microinfusions of SB ($F_{(2, 4)}$=15.02, p=0.0012; Fig.

2e). In contrast to systemic injections of SB, intra-cVP microinfusions of SB did not alter

remifentanil free consumption (Q_0; $F_{(2,4)}$=0.8576, p=0.4358; Fig. 2f). In rats with misplaced SB

injections directed at rVP, there was no effect of SB on either demand elasticity (p=0.4878) or

free consumption (p=0.8281).

Next we tested the role for Ox1R signaling in cued reinstatement behavior. Lever pressing was

extinguished as in Experiment 1 (Fig. 2g). Immediately prior to reinstatement tests, rats were

microinjected with either SB or aCSF (counterbalanced) bilaterally into cVP. A 2×3 factor

repeated measures ANOVA revealed a significant main effect of treatment ($F_{(2, 52)}$=8.117,

p=0.0009) and lever ($F_{(2,52)}$=61.38, p<0.0001), and a significant treatment×lever interaction

($F_{(2,52)}$=8.280, p=0.0008). Tukey's post-hoc analyses showed that remifentanil-associated cues

reinstated active lever pressing after vehicle pretreatment (vehicle vs. extinction, p<0.0001; Fig.

2h). SB in cVP significantly reduced responding on the active lever during reinstatement (SB vs.

vehicle, p=0.0008). Separate analyses revealed that SB also reduced reinstatement in animals

with misplaced cannulae in rVP (n=5; p=0.0061). There was no effect of SB (p=0.9604) or aCSF

(p=0.9955) in cVP on inactive lever responding (Fig. 2h). Together, these data indicate that

Ox1R signaling in cVP mediates the motivational properties of remifentanil reward behavior

(demand elasticity, cued reinstatement) but is not important for its mediating consumption under

null cost conditions.

Experiment 3: Blocking Ox1Rs in VP did not affect locomotor activity

To test for non-specific effects of intra-cVP microinjections of SB, we assessed general motor

activity in an open field apparatus. Bilateral microinfusions of SB into cVP did not alter the total

distance traveled (t_{13}=0.9848; p=0.3427; Fig. 3a), nor was there an effect on either horizontal

(t_{13}=1.032; p=0.0754; Fig 3b) or vertical (t_{13}=1.436; p=0.1747; Fig. 3c) activity over the 2h test

session. These results indicate that effects of SB on demand elasticity and reinstatement were not

due to changes in general motoric activity.

Experiment 4: Baseline demand for reminfentanil predicted cued reinstatement and efficacy of

SB

Finally, we sought to test whether 'baseline' demand predicted addiction behaviors and SB

efficacy, as we have previously reported for cocaine (Bentzley et al., 2014; James et al., 2018a).

Multiple linear regression analyses revealed that baseline demand elasticity (α; β=-0.674,

p=0.008; Fig. 5a, left), but not free consumption (Q_0; β=0.438, p=0.118; Fig. 5a, right), predicted

cue-induced reinstatement after aCSF microinfusion into VP. Thus, animals with the lowest α

(highest motivation) had greater cued reinstatement behavior. Consistent with previous evidence

from our lab indicating that demand elasticity (α) for cocaine can be used to predict those

subjects most likely to benefit from the anti-motivational effects of SB treatment (James et al.,

2018a), we found here that baseline α (β=-0.751, p=0.002) predicted the extent of reduction in

reinstatement by intra-cVP SB (Fig 5b, left). A similar, but non-significant, trend was observed

for Q_0 (β=-0.502, p=0.068; Fig. 5b, right). Together, these data indicate that 'baseline' demand

elasticity for reminfentanil is a behavioral biomarker for relapse propensity and can be used to

predict those individuals most likely to benefit from treatment with an Ox1R antagonist to block

reinstatement behavior.

Discussion

We confirm previous findings (Porter-Stransky et al., 2017) that systemic blockade of

Ox1Rs with SB increased demand elasticity (decreased motivation) for remifentanil (α),

decreased free remifentanil consumption (Q_0), and attenuated cue-induced reinstatement of drug

seeking. We extend these findings to show that motivation and reinstatement, but not 'free

consumption', for remifentanil are mediated by Ox1R signaling in cVP. Finally, we show that

rats' baseline remifentanil demand elasticity predicted cue-induced reinstatement responding

and the extent of SB effects in cVP on reinstated drug seeking. Together, these results indicate a

specific role for orexin signaling in cVP in motivated responding for the opioid remifentanil and

highlight demand elasticity as a behavioral biomarker to predict relapse vulnerability and the

efficacy of cVP Ox1R antagonism in blocking reminfentanil seeking behavior.

A clear role has been established for orexin signaling in drug seeking behavior (Harris et

al., 2005; Mahler et al., 2012; James et al., 2017). Signaling at the Ox1R preferentially mediates

high effort drug seeking/taking, or when drug motivation is augmented by external stimuli, such

as drug-associated stimuli (Harris et al., 2005; Smith and Aston-Jones, 2012; Porter-Stransky et

al., 2017). We previously found that this role extends to opioids, as systemic SB reduced

motivated responding for both heroin (Smith and Aston-Jones, 2012) and remifentanil (Porter-

Stransky et al., 2017). In addition, Fos activation of lateral hypothalamus (LH) orexin neurons is

strongly correlated with the degree of morphine seeking in a conditioned place preference

paradigm (Harris et al., 2005). However, unlike with other drugs of abuse, these studies

highlighted a role for orexin signaling in free consumption of opioids, as systemic SB treatment

reduced FR1 responding for heroin and low-cost consumption (Q_0) for remifentanil. Here, we

confirm a role for Ox1R signaling in both motivation and free consumption of remifentanil

reward by showing that systemic SB treatment both increased α and decreased Q_0 values in the

same BE test session.

We investigated cVP as a potential central site of action for SB in mediating remifentanil

seeking behaviors, as this region has been implicated in both motivational and hedonic

processing. Intra-cVP microinfusions of SB increased demand elasticity (α) but had no effect on

remifentanil intake at null cost (Q_0). These effects were specific to SB in cVP, as we observed no

effect of SB microinfusions dorsal to VP and in animals with misplaced injections. VP is a key

component of the brain circuitry regulating effort-related choice behavior, as VP inactivation

reduces willingness to work on an instrumental task to obtain sucrose reward (Farrar et al.,

2008). cVP receives moderate orexin input and expresses a high density of Ox1Rs (Marcus et al.,

2001; Baldo et al., 2003), indicating that this region may be a site for orexin modulation of

reward behavior. Interestingly however, there is evidence that orexin signaling in VP also

mediates the hedonic properties of reward. Microinjections of orexin-A into cVP enhanced the

hedonic responses ('liking') for sucrose, as assessed by an affective taste reactivity test (Ho and

Berridge, 2013). While we have previously argued that Q_0 might reflect an animal's hedonic set

point for drug (Bentzley et al., 2014), this is difficult to test directly. Thus, we are unable to

directly draw conclusions from our data regarding the role for cVP Ox1R signaling and hedonic

processing for remifentanil reward. It is unlikely that the lack of effect of intra-VP SB on Q_0 is

due to the dose of SB used, as this dose produces robust behavioral effects in other paradigms

(Harris et al., 2007; Mahler et al., 2013) and higher doses of SB would likely have off-target

effects (Porter et al., 2001; Smart et al., 2001). It is possible, though, that the previously reported

effects of orexin-A infusions in cVP on 'liking' for sucrose may be mediated by actions at Ox2R,

as orexin-A has similar affinity for Ox1R and Ox2R (Sakurai et al., 1998), and Ox2Rs are

expressed in cVP (Marcus et al., 2001). It is also possible that the orexin system plays distinct

roles in mediating the hedonic aspects of sucrose versus opioid reward. Clearly, further studies

are required to test these possibilities.

Systemic SB is highly effective at reducing cued reinstatement of extinguished drug

seeking (Smith et al., 2009; Smith and Aston-Jones, 2012; Plaza-Zabala et al., 2013; Moorman et

al., 2017). Such findings led us to hypothesize that the orexin system provides motivational

activation by reward-associated stimuli (Mahler et al., 2014a; James et al., 2017). Consistent

with this view, we found that both systemic and intra-cVP SB reduced cued reinstatement of

extinguished remifentanil seeking. Importantly, VP is part of a broader reward network that

underlies reinstatement behavior (McFarland and Kalivas, 2001; Prasad and McNally, 2016);

thus, orexin may act at other sites to contribute to cued reinstatement of remifentanil seeking

(James and Dayas, 2013; Qi et al., 2013; Matzeu et al., 2018). It will therefore be important that

future studies seek to identify the relative contribution of orexin signaling in VP versus other key

reward regions in reinstatement of opioid seeking.

Importantly, here we focused on cVP, as this area is more densely innervated by orexin

terminals than rVP. However, rVP has also been implicated in motivated reward seeking (Root et

al., 2015), including for drugs of abuse (Mahler and Aston-Jones, 2012; Mahler et al., 2014b).

Interestingly, in a small number of animals (n=5) with misplaced injections directed at rVP, we

observed no effect of SB on BE parameters, but a significant effect of SB on cued reinstatement.

This may indicate a functional distinction between cVP vs rVP, where motivation is more

associated with orexin signaling in cVP, whereas stimulus-driven seeking might involve orexin

input to both regions. However, given the relatively small number of animals in this group, it is

difficult to draw comprehensive conclusions from these observations, and it will be necessary

that future studies examine this further.

It is possible that blocking Ox1R signaling in our experiments impacted general motor

behavior, owing to the role for orexin signaling in arousal (Tsujino and Sakurai, 2013; Li et al.,

2014; Li et al., 2017). However we previously reported systemic SB only modestly reduced

general locomotor activity in rats with a history of remifentanil self-administration, and that the

magnitude of this reduction was not related to SB's effects on α or Q_0 (Porter-Stransky et al.,

2017). VP has been shown to be critical for the modulation of locomotor activation (Swerdlow et

al., 1984; Williams and Herberg, 1987). However, in locomotor activity tests we observed no

differences following microinjections of SB in cVP. Moreover, intra-cVP SB did not attenuate

inactive lever presses on BE or cued reinstatement tests, and other studies have failed to observe

non-specific motor deficits following local injections of SB into other reward regions (Borgland

et al., 2009; James et al., 2011; Mahler et al., 2013). We also found that intra-cVP SB had no

effect on low effort responding (Q_0) for remifentanil, further supporting our conclusion that

intra-VP SB-induced decreases in demand were not due to motor deficits.

The orexin system is important for motivated responding for natural rewards (Cason and

Aston-Jones, 2013b, a) and thus it is possible that the effect of intra-cVP infusions on motivation

for remifentanil would also extend to motivated food seeking. Importantly however, evidence

indicates that doses of SB that are sufficient to block drug seeking (such as those used here) have

no effect on regular chow intake in ad-libitum fed animals (Borgland et al., 2009; Choi et al.,

2010; Alcaraz-Iborra et al., 2014), indicating that any potential orexin-based therapy designed to

reduce craving and motivation for opioid reward may not interfere with homeostatic feeding.

We previously reported that baseline demand elasticity is a strong predictor of several

addiction-relevant endophenotypes (Bentzley et al., 2014; James et al., 2018a). Moreover, we

showed that SB is most effective at reducing motivation for cocaine and alcohol in highly

motivated animals (Lopez et al., 2016; Moorman et al., 2017; James et al., 2018a; James et al.,

2018c). Here, we show that animals with high baseline motivation for remifentanil also have

strong cued reinstatement behavior. We also show that high motivation animals exhibited the

largest reduction in reinstatement after SB injection, indicating that demand elasticity might also

be useful for identifying those individuals most likely to benefit from the therapeutic effects of

Ox1R blockade. Given that the BE paradigm can be readily applied in clinical populations

(MacKillop et al., 2018), our findings point to the potential utility of this approach for identifying

those individuals most likely to relapse following abstinence, and to benefit from

pharmacotherapies that interfere with orexin signaling.

In conclusion, our results add to a growing body of literature implicating the orexin

system as a target for treating opioid abuse and provide new insights into the corresponding

neural substrates. We also show that BE provides a predictive behavioral marker of remifentanil

seeking behaviors and the anti-drug seeking efficacy of SB, and thus could be applied clinically

to determine opioid addiction liability and optimize pharmacological treatment strategies.

References

Alcaraz-Iborra M, Carvajal F, Lerma-Cabrera JM, Valor LM, Cubero I (2014) Binge-like consumption of caloric and non-caloric palatable substances in ad libitum-fed C57BL/6J mice: pharmacological and molecular evidence of orexin involvement. Behavioural brain research 272:93-99.

Baimel C, Bartlett SE, Chiou LC, Lawrence AJ, Muschamp JW, Patkar O, Tung LW, Borgland SL (2015) Orexin/hypocretin role in reward: implications for opioid and other addictions. British journal of pharmacology 172:334-348.

Baldo BA, Daniel RA, Berridge CW, Kelley AE (2003) Overlapping distributions of orexin/hypocretin- and dopamine-beta-hydroxylase immunoreactive fibers in rat brain regions mediating arousal, motivation, and stress. The Journal of comparative neurology 464:220-237.

Baylon GJ, Kaplan HL, Somer G, Busto UE, Sellers EM (2000) Comparative abuse liability of intravenously administered remifentanil and fentanyl. Journal of clinical psychopharmacology 20:597-606.

Bentzley BS, Aston-Jones G (2015) Orexin-1 receptor signaling increases motivation for cocaine-associated cues. The European journal of neuroscience 41:1149-1156.

Bentzley BS, Fender KM, Aston-Jones G (2013) The behavioral economics of drug self-administration: a review and new analytical approach for within-session procedures. Psychopharmacology 226:113-125.

Bentzley BS, Jhou TC, Aston-Jones G (2014) Economic demand predicts addiction-like behavior and therapeutic efficacy of oxytocin in the rat. Proceedings of the National Academy of Sciences of the United States of America 111:11822-11827.

Borgland SL, Chang SJ, Bowers MS, Thompson JL, Vittoz N, Floresco SB, Chou J, Chen BT, Bonci A (2009) Orexin A/hypocretin-1 selectively promotes motivation for positive reinforcers. The Journal of neuroscience : the official journal of the Society for Neuroscience 29:11215-11225.

Cason AM, Aston-Jones G (2013a) Attenuation of saccharin-seeking in rats by orexin/hypocretin receptor 1 antagonist. Psychopharmacology 228:499-507.

Cason AM, Aston-Jones G (2013b) Role of orexin/hypocretin in conditioned sucrose-seeking in rats. Psychopharmacology 226:155-165.

Childress AR, Ehrman RN, Wang Z, Li Y, Sciortino N, Hakun J, Jens W, Suh J, Listerud J, Marquez K, Franklin T, Langleben D, Detre J, O'Brien CP (2008) Prelude to passion: limbic activation by "unseen" drug and sexual cues. PloS one 3:e1506.

Choi DL, Davis JF, Fitzgerald ME, Benoit SC (2010) The role of orexin-A in food motivation, reward-based feeding behavior and food-induced neuronal activation in rats. Neuroscience 167:11-20.

Compton WM, Volkow ND (2006) Major increases in opioid analgesic abuse in the United States: concerns and strategies. Drug and alcohol dependence 81:103-107.

Dallimore JE, Mickiewicz AL, Napier TC (2006) Intra-ventral pallidal glutamate antagonists block expression of morphine-induced place preference. Behavioral neuroscience 120:1103-1114.

Dobbs GC, Fogger SA (2018) Opiate Dependence or Addiction?: A Review of the Centers for Disease Control and Prevention Guidelines for Management of Chronic Pain. J Addict Nurs 29:57-61.

Espana RA, Oleson EB, Locke JL, Brookshire BR, Roberts DC, Jones SR (2010) The
    hypocretin-orexin system regulates cocaine self-administration via actions on the
    mesolimbic dopamine system. The European journal of neuroscience 31:336-348.

Farrar AM, Font L, Pereira M, Mingote S, Bunce JG, Chrobak JJ, Salamone JD (2008) Forebrain
    circuitry involved in effort-related choice: Injections of the GABAA agonist muscimol
    into ventral pallidum alter response allocation in food-seeking behavior. Neuroscience
    152:321-330.

Glass PS, Gan TJ, Howell S (1999) A review of the pharmacokinetics and pharmacodynamics of
    remifentanil. Anesthesia and analgesia 89:S7-14.

Harris GC, Wimmer M, Aston-Jones G (2005) A role for lateral hypothalamic orexin neurons in
    reward seeking. Nature 437:556-559.

Harris GC, Wimmer M, Randall-Thompson JF, Aston-Jones G (2007) Lateral hypothalamic
    orexin neurons are critically involved in learning to associate an environment with
    morphine reward. Behavioural brain research 183:43-51.

Ho CY, Berridge KC (2013) An orexin hotspot in ventral pallidum amplifies hedonic 'liking' for
    sweetness. Neuropsychopharmacology : official publication of the American College of
    Neuropsychopharmacology 38:1655-1664.

Hubner CB, Koob GF (1990) The ventral pallidum plays a role in mediating cocaine and heroin
    self-administration in the rat. Brain research 508:20-29.

Hursh SR, Silberberg A (2008) Economic demand and essential value. Psychological review
    115:186.

James MH, Dayas CV (2013) What about me...? The PVT: a role for the paraventricular
    thalamus (PVT) in drug-seeking behavior. Front Behav Neurosci 7:18.

James MH, Mahler SV, Moorman DE, Aston-Jones G (2017) A Decade of Orexin/Hypocretin
    and Addiction: Where Are We Now? Current topics in behavioral neurosciences 33:247-
    281.

James MH, Bowrey HE, Stopper CM, Aston-Jones G (2018a) Demand elasticity predicts
    addiction endophenotypes and the therapeutic efficacy of an orexin/hypocretin-1 receptor
    antagonist in rats. The European journal of neuroscience.

James MH, McGlinchey EM, Vattikonda A, Mahler SV, Aston-Jones G (2018b) Cued
    Reinstatement of Cocaine but Not Sucrose Seeking Is Dependent on Dopamine Signaling
    in Prelimbic Cortex and Is Associated with Recruitment of Prelimbic Neurons That
    Project to Contralateral Nucleus Accumbens Core. International Journal of
    Neuropsychopharmacology 21:89-94.

James MH, Stopper CM, Zimmer BA, Koll NE, Bowrey HE, Aston-Jones G (2018c) Increased
    number and activity of a lateral subpopulation of hypothalamic orexin/hypocretin neurons
    underlies the expression of an addicted state in rats. Biological psychiatry.

James MH, Charnley JL, Levi EM, Jones E, Yeoh JW, Smith DW, Dayas CV (2011) Orexin-1
    receptor signalling within the ventral tegmental area, but not the paraventricular
    thalamus, is critical to regulating cue-induced reinstatement of cocaine-seeking. The
    international journal of neuropsychopharmacology / official scientific journal of the
    Collegium Internationale Neuropsychopharmacologicum 14:684-690.

Levine AI, Bryson EO (2010) Intranasal self-administration of remifentanil as the foray into
    opioid abuse by an anesthesia resident. Anesthesia and analgesia 110:524-525.

Lewis L, Carpenter CR, Schwarz ES, Jotte RS, Waller C, Winograd R, Williams R, Stenger S, Rehder H, Governick S, Giuffra L (2017) The Opioid Crisis in Missouri: A Call to Action for Physicians, Legislators, and Society. Missouri medicine 114:440-446.

Li J, Hu Z, de Lecea L (2014) The hypocretins/orexins: integrators of multiple physiological functions. British journal of pharmacology 171:332-350.

Li SB, Giardino WJ, de Lecea L (2017) Hypocretins and Arousal. Current topics in behavioral neurosciences 33:93-104.

Lopez MF, Moorman DE, Aston-Jones G, Becker HC (2016) The highly selective orexin/hypocretin 1 receptor antagonist GSK1059865 potently reduces ethanol drinking in ethanol dependent mice. Brain research 1636:74-80.

MacKillop J, Goldenson NI, Kirkpatrick MG, Leventhal AM (2018) Validation of a behavioral economic purchase task for assessing drug abuse liability. Addiction biology.

Mahler SV, Aston-Jones GS (2012) Fos activation of selective afferents to ventral tegmental area during cue-induced reinstatement of cocaine seeking in rats. The Journal of neuroscience : the official journal of the Society for Neuroscience 32:13309-13326.

Mahler SV, Smith RJ, Aston-Jones G (2013) Interactions between VTA orexin and glutamate in cue-induced reinstatement of cocaine seeking in rats. Psychopharmacology 226:687-698.

Mahler SV, Smith RJ, Moorman DE, Sartor GC, Aston-Jones G (2012) Multiple roles for orexin/hypocretin in addiction. Progress in brain research 198:79-121.

Mahler SV, Moorman DE, Smith RJ, James MH, Aston-Jones G (2014a) Motivational activation: a unifying hypothesis of orexin/hypocretin function. Nature neuroscience 17:1298-1303.

Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kaufling J, Wilson SP, Deisseroth K, Woodward JJ, Aston-Jones G (2014b) Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. Nature neuroscience 17:577-585.

Marcus JN, Aschkenasi CJ, Lee CE, Chemelli RM, Saper CB, Yanagisawa M, Elmquist JK (2001) Differential expression of orexin receptors 1 and 2 in the rat brain. The Journal of comparative neurology 435:6-25.

Matzeu A, Kallupi M, George O, Schweitzer P, Martin-Fardon R (2018) Dynorphin Counteracts Orexin in the Paraventricular Nucleus of the Thalamus: Cellular and Behavioral Evidence. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 43:1010-1020.

McFarland K, Kalivas PW (2001) The circuitry mediating cocaine-induced reinstatement of drug-seeking behavior. The Journal of neuroscience : the official journal of the Society for Neuroscience 21:8655-8663.

McGlinchey EM, James MH, Mahler SV, Pantazis C, Aston-Jones G (2016) Prelimbic to Accumbens Core Pathway Is Recruited in a Dopamine-Dependent Manner to Drive Cued Reinstatement of Cocaine Seeking. The Journal of neuroscience : the official journal of the Society for Neuroscience 36:8700-8711.

Moorman DE, James MH, Kilroy EA, Aston-Jones G (2017) Orexin/hypocretin-1 receptor antagonism reduces ethanol self-administration and reinstatement selectively in highly-motivated rats. Brain research 1654:34-42.

Panlilio LV, Schindler CW (2000) Self-administration of remifentanil, an ultra-short acting opioid, under continuous and progressive-ratio schedules of reinforcement in rats. Psychopharmacology 150:61-66.

Paxinos G, Watson C (1998) A stereotaxic atlas of the rat brain. New York: Academic.
Peyron C, Tighe DK, van den Pol AN, de Lecea L, Heller HC, Sutcliffe JG, Kilduff TS (1998)
Neurons containing hypocretin (orexin) project to multiple neuronal systems. The Journal
of neuroscience : the official journal of the Society for Neuroscience 18:9996-10015.
Plaza-Zabala A, Flores A, Martin-Garcia E, Saravia R, Maldonado R, Berrendero F (2013) A
role for hypocretin/orexin receptor-1 in cue-induced reinstatement of nicotine-seeking
behavior. Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology 38:1724-1736.
Porter-Stransky KA, Bentzley BS, Aston-Jones G (2017) Individual differences in orexin-I
receptor modulation of motivation for the opioid remifentanil. Addiction biology 22:303-
317.
Porter RA, Chan WN, Coulton S, Johns A, Hadley MS, Widdowson K, Jerman JC, Brough SJ,
Coldwell M, Smart D, Jewitt F, Jeffrey P, Austin N (2001) 1,3-Biarylureas as selective
non-peptide antagonists of the orexin-1 receptor. Bioorganic & medicinal chemistry
letters 11:1907-1910.
Prasad AA, McNally GP (2016) Ventral Pallidum Output Pathways in Context-Induced
Reinstatement of Alcohol Seeking. The Journal of neuroscience : the official journal of
the Society for Neuroscience 36:11716-11726.
Qi K, Wei C, Li Y, Sui N (2013) Orexin receptors within the nucleus accumbens shell mediate
the stress but not drug priming-induced reinstatement of morphine conditioned place
preference. Frontiers in behavioral neuroscience 7:144.
Root DH, Melendez RI, Zaborszky L, Napier TC (2015) The ventral pallidum: Subregion-
specific functional anatomy and roles in motivated behaviors. Progress in neurobiology
130:29-70.
Sakurai T et al. (1998) Orexins and orexin receptors: a family of hypothalamic neuropeptides
and G protein-coupled receptors that regulate feeding behavior. Cell 92.
Smart D, Sabido-David C, Brough SJ, Jewitt F, Johns A, Porter RA, Jerman JC (2001) SB-
334867-A: the first selective orexin-1 receptor antagonist. British journal of
pharmacology 132:1179-1182.
Smith RJ, Aston-Jones G (2012) Orexin / hypocretin 1 receptor antagonist reduces heroin self-
administration and cue-induced heroin seeking. The European journal of neuroscience
35:798-804.
Smith RJ, See RE, Aston- Jones G (2009) Orexin/hypocretin signaling at the orexin 1 receptor
regulates cue- elicited cocaine- seeking. European Journal of Neuroscience 30:493-503.
Swerdlow NR, Swanson LW, Koob GF (1984) Electrolytic lesions of the substantia innominata
and lateral preoptic area attenuate the 'supersensitive' locomotor response to apomorphine
resulting from denervation of the nucleus accumbens. Brain research 306:141-148.
Tsujino N, Sakurai T (2013) Role of orexin in modulating arousal, feeding, and motivation.
Front Behav Neurosci 7:28.
Volkow ND, Collins FS (2017) The Role of Science in Addressing the Opioid Crisis. The New
England journal of medicine 377:391-394.
Williams SF, Herberg LJ (1987) Motivational vs. motor effects of striatal and pallidal gabergic
projections to subthalamic and entopeduncular nuclei, ventromedial thalamus, and ventral
globus pallidus. Pharmacology, biochemistry, and behavior 26:49-55.
Zacny JP, Galinkin JL (1999) Psychotropic drugs used in anesthesia practice: abuse liability and
epidemiology of abuse. Anesthesiology 90:269-288.

Figure 1. Effects of systemic Ox1R blockade on demand and reinstatement for remifentanil. (a) Timeline for behavioral procedure. Rats (n=10) were trained to self-administer remifentanil and then underwent BE training. Lever pressing was extinguished, and animals were then tested in a 2h cue-induced reinstatement session. Arrows in timeline indicate 30 mg/kg SB or vehicle treatments (counterbalanced). **(b-c)** Mean behavioral performance during remifentanil self-administration training. Data reflect the final 6 days of self-administration training. $*p<0.05$; $**p<0.01$ active vs. inactive lever. **(d)** A representative demand curve for a single animal during a remifentanil BE session. **(e)** Consumption of remifentanil at null cost (Q_0) negatively correlated with demand elasticity (α). Each point represents data from a single animal. **(f-g)** SB (30mg/kg, ip) increased α (decreased motivation; f) and reduced Q_0 (free consumption; g) for remifentanil during BE sessions. **(h)** Mean number of active and inactive lever presses during 7 days of extinction training (n=9). $*p<0.05$. **(i)** Average active lever presses during the last 3 days of extinction training and 2 h cue-induced reinstatement testing 30 min after pretreatment with 30 mg/kg SB or vehicle. SB had no effect on inactive lever presses. $***p<0.001$; $****p<0.0001$. Bar graphs represent mean ± standard error of mean (SEM).

Figure 2. Ox1R antagonism in VP decreases motivation for remifentanil. (a) Experimental design of behavioral training and testing, indicating the order and number of injections in VP. **(b-c)** Mean lever pressing during remifentanil FR-1 self-administration (n=14). Data show average number of infusions, and active and inactive lever presses during the final 6 days of FR1 sessions. Rats made significantly more lever presses on the active (drug associated) lever than the inactive lever, beginning on the second session and throughout the remainder of the training sessions. This was associated with an increase in remifentanil (Remi) intake. $*p<0.05$, $**p<0.01$

vs. inactive lever. **(d)** There was a significant correlation between α and Q_0 such that high takers

at null effort (high Q_0) displayed significantly higher motivation (low α) for remifentanil than

low takers. **(e)** Demand elasticity (α) was significantly increased (motivation was decreased)

following intra-cVP delivery of SB compared to microinfusion of aCSF or SB just dorsal to VP

(n=14). No change in α values was observed in animals that received misplaced injections

located in rVP (n=5). **$p<0.01$ compared with dorsal SB. **(f)** Consumption of remifentanil at

null cost (Q_0) was not altered by intra-cVP SB microinfusion, indicating that Ox1R blockade in

VP does not alter remifentanil intake at low cost. Similarly, no effect of SB was observed in

animals with misplaced (rVP) SB injections (n=5). **(g)** Mean number of active and inactive lever

presses during 7 days of extinction training after BE training and testing (n=14). **(h)**

Microinfusion of SB into cVP reduced active lever responding during cue-induced reinstatement

of remifentanil seeking compared to after aCSF microinjections (n=14). There was also a

significant reduction in reinstatement scores in animals with misplaced (rVP) SB injections

(n=5). There was no effect of SB vs. aCSF microinjections on inactive lever presses.

***$p<0.001$; ****$p<0.0001$. (i) Representative schematic (upper panel) and photomicrograph

(lower panel; neutral red Nissl stain, frontal section, midline at center) of cannula injection sites

from animals that received microinjections in cVP. Injection sites are marked by blue dye

microinfused from cannulae at end of experiments. Bilateral injection sites across all rats (black

circles; n=14). Bar graphs represent mean ± standard error of mean (SEM). n.s.=not statistically

different.

Figure 3. Blockade of Ox1R signaling in cVP had no effect on general locomotor activity.

(**a**) Rats displayed no differences in total distance traveled in locomotor chambers after SB

microinjection into VP compared to vehicle (n=14). Microinjection of SB into VP had no effect

on (**b**) horizontal or (**c**) vertical activity. Bar graphs represent mean ± standard error of mean

(SEM). *p< 0.05; ***p<0.001; n.s.=not statistically different.

Figure 4. Baseline demand elasticity predicted reinstatement propensity and efficacy of SB

microinfusion into VP. (a) Baseline demand elasticity (α), but not free remifentanil

consumption (Q_0), significantly correlated with the magnitude of cue-induced reinstatement with

aCSF microinjection, (Q_0. (b) Baseline demand elasticity (α) correlated with the magnitude of

the reduction of cue-induced reinstatement after SB microinjection into cVP. Free remifentanil

consumption also correlated with the magnitude of the reduction of cue-induced reinstatement

with SB microinjection into VP, but this failed to reach statistical significance (Q_0). n=14 for all

correlations.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 1

Figure 2

Figure 3

Figure 4