1   **Nucleotide and codon background mutability shape cancer mutational spectrum**
2   **and advance driver mutation identification**
3

4   Minghui Li[1,#], Anna-Leigh Brown[2,#], Alexander Goncearenco[2*], Anna R. Panchenko[2*]
5

[1]   School of Biology and Basic Medical Sciences, Soochow University, Suzhou 215123,
China
[2]   National Center for Biotechnology Information, NLM, NIH, Bethesda, MD 20894, United
States

# - co-first author with equal contribution

* - corresponding authors:

Anna Panchenko, panch@ncbi.nlm.nih.gov

Alexander Goncearenco, alexandr.goncearenco@nih.gov

Abstract

To decipher the contribution of background DNA mutability in shaping observed mutational spectrum in cancer, we developed probabilistic models to estimate DNA context-dependent background mutability per nucleotide and codon substitution as a result of mutagenic processes. We showed that observed frequency of cancer mutations follows expected mutability, especially for tumor suppressor genes. In oncogenes, highly recurring mutations were characterized by a lower mutability, showing a U-shape trend. Mutations not occurring in tumors were shown to have lower mutability than observed mutations, indicating that DNA mutability could be a limiting step in mutation occurrence. To check whether mutability helps to discriminate driver mutations from passengers, we compiled a set of missense mutations with experimentally validated functional and transforming impacts. Accounting for background mutability significantly improved performance of classification into cancer driver and passenger mutations compared to mutation recurrence alone, performing similar to state-of-the-art machine-learning methods, even though no training was involved.

Introduction

Cancer is driven by changes at the gene, chromatin, and cellular levels. Somatic cells may rapidly acquire mutations, one or two orders of magnitude faster than germline cells [1]. The majority of these mutations are largely neutral (passenger mutations) in comparison to a few driver mutations that give cells the selective advantage leading to their proliferation [2]. Such a binary driver-passenger model can be adjusted by taking into account additive pleiotropic effect of mutations [3]. Mutations might have different functional consequences in various cancer types and patients, they can lead to activation or deactivation of proteins and dysregulation of a variety of cellular processes. This gives rise to high mutational, biochemical, and histological intra- and inter-tumor heterogeneity that explains the resistance of cancer to therapies and complicates the identification of driving events in cancer [4,5].

Point DNA mutations can arise from various forms of infidelity of DNA replication and repair machinery, endogenous, and exogenous mutagens [6-8]. There is an intricate interplay between processes leading to DNA damage and those maintaining genome integrity. The resulting mutation rate can vary throughout the genome by more than two orders of magnitude [9,10] due to many causes including timing of replication, nucleosome positioning, histone modifications levels, and other factors operating on local and large scales [11-14]. Many studies support point mutation rate dependence on the local DNA sequence context for various types of germline and somatic mutations [8,10,12,15]. Moreover, local DNA sequence context has been identified as a dominant factor explaining the largest proportion of mutation rate variation in germline and soma [9,16]. Additionally, differences in mutational burden between cancer types suggest tissue type and mutagen exposure as important confounding factors contributing to tumor heterogeneity.

Assessing background mutation rate is crucial in identifying significantly mutated genes [17,18], sub-gene regions [19,20], mutational hotspots[21,22], or prioritizing mutations[23]. This is especially important considering that the functional impact of the majority of changes observed in cancer is poorly understood, in particular for rarely mutated genes [24]. Despite this need, there is a

persistent lack of quantitative information on per-nucleotide background rates of cancer somatic

mutations in different gene sites in various cancer types and tissues. In this study we developed

context-dependent probabilistic models that estimate baseline mutability per nucleotide or

codon substitution in specific genomic sites. These models were applied to decipher the

contribution of background DNA mutability in the observed mutational spectrum in cancer for

missense, nonsense, and silent mutations. To complement this approach, we also analyzed

background mutability for mutations that were not observed in cancer patients. Finally, for

practical purposes, we compiled a manually curated set of cancer driver and neutral missense

mutations with experimentally validated impacts collected from multiple studies. This data set

along with the recently published data set of experimentally characterized cancer mutations[27]

was used to verify our conclusions about the properties of cancer driver and passenger

mutations. The method has been implemented and can be found online as part of the MutaGene

web-server: https://www.ncbi.nlm.nih.gov/research/mutagene/gene.

Results

Mutations not observed in cancer patients have low mutability

We analyzed all 3,241,017 theoretically possible codon substitutions that could have occurred in

520 cancer census genes and found that only about one percent of them were actually observed

in the surveyed 12,013 tumor samples (Table S1). As illustrated in Figure 1B and D, the average

mutability of codon substitutions not detected in cancer patients (μ = 1.30 x 10^{-6}) was found to

be three-fold lower compared to the mutability of observed codon substitutions for all types of

mutations (μ = 3.88 x 10^{-6}), Mann-Whitney-Wilcoxon test, $p < 0.01$. This finding holds true for

per-nucleotide mutability (μ = 1.03 x 10^{-6} versus μ = 3.34 x 10^{-6}, Mann-Whitney-Wilcoxon test,

$p < 0.01$) (Figures 1D and S1). Figure 2 shows cumulative and probability density distributions

of nucleotide mutability values for all observed and possible mutations in all cancer census genes

and for two genes in particular, CASP8 and TP53, as examples. While there are many more

possible mutations with low mutability values, mutations at the higher range of the mutability

spectrum tend to occur more often in cancer samples.

Silent mutations have highest mutability

Figures 1A,B show the distributions of codon mutability values for all possible missense, nonsense, and silent mutations accessible by single nucleotide base substitutions in the protein-coding sequences of 520 cancer census genes calculated with the pan-cancer background model. As can be seen on this figure, codon mutability spans two orders of magnitude and silent mutations have significantly higher average mutability values (μ = 5.68 x 10^{-6}) than nonsense (μ = 3.44 x 10^{-6}), or missense mutations (μ = 3.29 x 10^{-6}) according to Kruskal-Wallis test ($p < 0.01$) and Dunn's post hoc test ($p < 0.01$) for all comparisons. These differences in codon mutabilities could be a reflection of the degeneracy of genetic code, where multiple silent nucleotide substitutions in the same codon may increase its mutability (as illustrated in Figure S5). Consistent with this explanation, for nucleotide mutability shown in Figure 1C the differences between types of mutations became less pronounced. However, silent mutations were still characterized by the highest mutability values. Since observed mutational frequency of silent mutations depends on their mutability (Figures 1B,D), it is important to correct for mutability while ranking silent mutations with respect to their driver status, which is difficult to assess comprehensively. Some recurrent highly mutable silent mutations might not represent relevant candidates of drivers whereas some low mutable mutations are predicted to be drivers by our method (KDR gene p.Leu355=, NTRK1 gene p.Asn270= and others).

Background mutability may shape the observed mutational spectrum in cancer

Under the null model of all mutations arising as a result of background mutational processes, somatic mutations should accumulate with respect to their mutation rate and one would expect a positive correlation between mutability and observed mutational frequency of individual mutations. Indeed, as Figure 1 shows, this is clearly the case for silent and nonsense mutations. To further investigate this relationship, we calculated Spearman's rank (a non-parametric test taking into account that mutability is not normally distributed) and Pearson linear correlation coefficients between codon mutability and frequencies of mutations observed in 12,301 whole-exome (WES) and whole-genome (WGS) tumor samples. Pooling together all types of mutations in cancer census genes resulted in a very low but significant positive correlation between codon

117 mutability and mutation frequency ($\rho = 0.13$ and $r = 0.02$ for Spearman and Pearson

118 respectively $p < 0.01$), as shown in Figure S2A.

120 Breaking up all codon changes into silent, nonsense and missense reveals higher correlations,

121 particularly for silent ($\rho = 0.14$, $r = 0.1$, $p < 0.01$) and nonsense ($\rho = 0.20$, $r = 0.15$, $p <$

122 0.01) mutations (Figure S2A). We calculated correlation coefficients for each gene with at least

123 ten unique mutations of each type: silent, nonsense, and missense (Figures 3). Overall, we found

124 84 and 137 genes with significant ($p < 0.01$) positive Spearman and Pearson correlations,

125 respectively, for at least one mutation type (Table S2). Among the genes with significant

126 correlations, 41 belonged to tumor suppressor genes, 28 were oncogenes, and 15 genes were

127 classified as either fusion genes or both oncogene and tumor suppressor. For some genes,

128 including TP53 (second column, Figure 3B) and tumor suppressor CASP8 (third column, Figure

129 3B), a strong linear relationship between mutability and observed mutations frequency ($R^2 >$

130 0.5) was observed.

132 **Relationship between mutability and observed frequency is different for tumor suppressor and**

133 **oncogenes**

134 The effects of cancer mutations on protein function with respect to their cancer transforming

135 ability, can drastically differ in tumor suppressor genes (TSG) and oncogenes, therefore we

136 performed our analysis separately for these two categories (Figure 1E). We used COSMIC gene

137 classification separating genes into tumor suppressors and oncogenes. Mutations in tumor

138 suppressors can cause cancer through the inactivation of their products, whereas mutations on

139 oncogenes may result in protein activation. In addition, we used COSMIC classification into genes

140 with dominant or recessive mutations but overall results were similar to the ones produced using

141 classification into TSG and oncogenes (Figure S3). The strongest correlation between codon

142 mutability and mutation frequency was observed for TSG ($\rho = 0.17$, $r = 0.13$, $p < 0.01$) while

143 oncogenes showed a weak Spearman correlation, and no significant Pearson correlation ($\rho =$

144 0.13, $p < 0.01$, $r = 0$, $p = 0.61$) (Figure S2).

A U-shape trend was detected for missense and silent mutations in oncogenes: mutations with

low observed frequency (one or two samples) had the highest codon mutability on average, but

highly recurring mutations (observed in three and more samples) were characterized by a lower

average mutability (Figure 1E). In the latter case, selection may be a more important factor than

background mutation rate explaining reoccurrence of these mutation. Indeed, oncogenes are

characterized by a few gain-of-function missense mutations that are rather specific events

occurring on particular sites, compared to deactivating events in TSG that can affect a much

broader range of sites. Functional conserved sites overall were found to be more frequently

mutated in oncogenes [25], although our analysis did not find a straightforward association

between site mutability and its evolutionary conservation (data not shown).

Functional effects of cancer mutations are related to codon mutability

We calculated mutability for missense mutations which in turn were categorized based on their

experimental effect on protein function, transforming effects, and other characteristics (see

Methods). In TP53 transactivation dataset, mutability was found to differ with respect to

experimental effects with "functional" and "partially functional" mutations having significantly

higher mutability than "non-functional" (damaging) mutations (Dunn's post hoc test, $p < 0.01$

for both comparisons) (Figure 4A). No significant difference was found between functional and

partially functional mutations. Consistent with this finding, "neutral" mutations in the Martelotto

et al.[26] dataset had significantly higher mutability than "non-neutral" mutations (Mann-Whitney-

Wilcoxon test, $p < 0.01$)(Figure 4B). The same but much weaker trend was observed for BRCA1

missense mutations which were classified according to their ability to engage in homology-

directed repair (HDR) (Mann-Whitney-Wilcoxon test, $p = 0.036$)(Figure 4C). For our combined

dataset, the mutability values of "neutral" mutations ($\mu = 1.35 \times 10^{-6}$) were significantly higher

(Mann-Whitney-Wilcoxon test, $p < 0.01$) than for non-neutral mutations ($\mu = 2.59 \times 10^{-6}$)

(Figure 4D). This is consistent with our finding of high mutability of silent mutations, if we assume

that vast majority of them are benign (Figure 1). These results are also corroborated (with the

exception of skin cancers) if we use cancer-specific context-dependent background mutation

models for TP53 on Martelotto *et al.* set, although for BRCA1 set the difference between

damaging and neutral mutations is only observed using pan-cancer model (Figure S4, Table S4).

Similar to our combined dataset, activating mutations from the validation set (see methods) had

a much lower background codon mutability (μ = 1.35 x 10^{-6}) compared to neutral mutations (μ =

3.77 x 10^{-6}) (Mann-Whitney-Wilcoxon test, $p < 0.01$).

Accounting for context-dependent mutability in distinguishing driver from passenger

mutations

In the previous sections we explored the contribution of background mutational processes to

understand the observed mutational patterns in cancer, with the ultimate goal to facilitate the

detection of cancer driver mutations or provide a reasonable ranking in terms of their

carcinogenic effects. Therefore, to discover driver mutations, we tested how well different

methods differentiate between experimentally verified neutral and driver non-neutral

mutations. For this purpose, we compiled a dataset from several experimental studies which

included ten oncogenes, five tumor suppressor genes, and two genes which were classified as

both oncogenes and tumor suppressors (Table S7). In addition, we validated our results on a

recently published dataset of mutations in 50 genes [27].

We developed two measures which utilize background mutation rate. The first measure, log-

ratio (LR, equation 4), was calculated as a logarithm of a ratio of the number of observed and

expected mutations. The second measure (B-score, equation 5) was calculated from the binomial

distribution as a probability to find a given mutation in a certain number of tumor samples or

higher given the background context-dependent mutation rate (mutability). We compared the

performance of these two scores to Condel[28], a meta-predictor which incorporates several state-

of-the-art computational approaches, and FatHMM[29], a method explicitly developed to predict

cancer driver mutations which performed very well in the most recent assessment [30].

Furthermore, the performance was indirectly compared to 21 different methods on a validation

set (see Methods).

Figure 5 shows the ROC and precision-recall performance on the combined dataset. As can be

seen in the inset of Figure 5A and Table 1, B-score had the best performance in the region of

driver mutation prediction with low false positive rate and highest Matthews correlation

coefficient (MCC), a performance measure known to be more robust on unbalanced datasets.

Condel and FatHMM achieved highest AUC-ROC and a spike in true positive rate was observed in

FatHMM score, as it approached TP53 true positives, due to an extreme weight assigned by

FatHMM to TP53 gene. Importantly, correction of observed frequency by codon mutability

provided a much better performance than frequency or mutability alone. Mutability alone

performed better than random with an AUC=0.67 emphasizing the fundamental quality of non-

neutral mutations in cancer: mutability of driver mutations is lower than the mutability of

passengers (Figure 4D). As an additional validation, we compared B-Score performance to a

recent functional dataset [27] on 255 "activating" mutations and 488 "neutral" mutations. A ROC

analysis showed that B-Score performed with an AUC-ROC of 0.76 putting its performance among

the top three methods tested by Ng *et al.* Importantly to note, observed frequency performed

unexpectedly well on this test set with an AUC-ROC of 0.72, outperforming 18 of the 21 tested

computational methods tested.

Variability of mutation rates across genes

We checked if incorporating large-scale factors, allowing mutations of the same type to have

different mutational probabilities in different genes, affected retrieval performance. Several

methods have been developed to estimate gene weights, which use the overall number of

mutations, number of silent mutations affecting a gene, or other factors. We implemented and

tested all these approaches (see Methods) and in addition estimated the gene weights based on

the number of SNPs in the vicinity of a gene. We used gene weights to adjust codon mutability

values and explored whether any of the gene weight models were helpful in distinguishing

between experimentally neutral and driver non-neutral mutations. Unexpectedly, using gene

weight as an adjustment for varying background mutational rates across genes did not affect

classification performance. Only a SNP based weight affected the AUC-ROC, but the gain was very

minimal and no gene weight affected MCC (Table S6). This result held true on the validation

dataset with mutations sampled from 50 genes. Importantly, certain gene weights decreased

classification accuracy; in Figure 5 results with no gene weights are reported. In addition, we

examined the effects of several large-scale confounding factors such as gene expression levels,

replication timing, and chromatin accessibility provided in the gene covariates files for

MutSigCV[31] and found that only "no-outlier"-based weight and "silent mutation"-based weight

significantly correlated with the gene expression levels ($r = 0.66, p = 0.004$ and $r = 0.65, p =$

0.004, respectively).

Ranking of cancer point mutations in MutaGene

MutaGene webserver provides a collection of cancer specific context-dependent mutational

profiles and allows to calculate nucleotide and codon mutability for missense, nonsense and

silent mutations for a given protein coding sequence and background mutagenesis model using

the "Analyze gene" option. Following the analysis presented in this study, we added options to

provide a ranking of mutations observed in cancer samples based on B-score or the multiple-

testing adjusted q-values. Using our combined benchmark (Figure 5), we calibrated two

thresholds: the first corresponds to the maximum MCC coefficient, and the second corresponds

to 10% false positive rate (FPR). Mutations with B-score below the first threshold are predicted

to be "cancer drivers", whereas mutations with scores in between two thresholds are predicted

to be "potential drivers". All mutations with scores above the second threshold are predicted as

"passengers". Importantly, calculations can be performed using pan-cancer cohort or any

particular cancer type, the latter would result in a cancer-specific ranking of mutations and could

be useful for identification of driver mutations in a particular type of cancer. An example of

prediction of driver mutations status for EGFR is shown in Figure 6.

Discussion

To understand what processes drive point mutation accumulation in cancer, we used DNA

context-dependent probabilistic models to estimate the baseline mutability for nucleotide

mutation or codon substitution in specific genomic sites. Passenger mutations, constituting the

majority of all observed mutations, may have largely neutral functional impact and are unlikely

to be under selection pressure. For passenger mutations one would expect that mutations with

lower DNA mutability would have lower observed mutational frequency and vice versa. In
accordance with this expectation, we detected a significant positive correlation between baseline
mutability, which is an estimate of per site mutation rate, and observed frequencies of mutations
in cancer patients. In accordance with this expectation, we also found that mutations not so far
observed in cancer patients had much lower expected background nucleotide and codon
mutability compared to the observed mutations. For some genes, the observed frequency of
occurrence of mutations can be predicted purely from their mutability. Outliers of this trend are
important for inferring mutations under selection. For instance, if a mutation with low expected
mutation rate is observed in multiple patients, it is suggestive of it being potentially important in
carcinogenesis.

In this respect, reoccurring synonymous mutations with low mutability may represent interesting
cases for further investigation of potential silent drivers. Mutability of synonymous mutations
was found to be the highest among other types of mutations; and observed mutational frequency
of silent mutations scales well with their mutability. However, silent mutations that reoccur in
three or more samples in oncogenes have lower mutability than those occurring in only two
samples, suggesting that selection may be driving the high reoccurrence of these mutations.
Overall, our method predicted 102 silent mutations in 64 out of 520 cancer-associated genes to
be drivers. Indeed, it has been previously shown that some synonymous mutations might be
under selection, and synonymous codon mutations can affect the speed and accuracy of
transcription and translation, protein folding rate, and splicing [32].

There are many computational methods that aim to detect driver genes and fewer methods
trying to rank mutations with respect to their potential carcinogenicity. As many new approaches
to address this issue have been developed [33] [30], it still remains an extremely difficult task and
many driver mutations, especially in oncogenes, are not annotated as high impact or disease
related [34].

This can be explained by the scarcity of per site mutational data in WGS and WES studies for
specific cancers, difficulty to distinguish selection signal from background, and limited data on
experimental verification of driver mutational status. Increasing data in WGS and WES studies for
specific cancers thus should help to resolve the lack of cancer-specific benchmarks. While our
current attempts at including large-scale factors affecting mutation rates into gene weight
models did not improve prediction significantly in a pan-cancer benchmark, these factors are
likely tissue and cancer type specific, and should be tested in cancer-specific benchmarks.

We provide here a method to rank mutations with respect to their likelihood of being drivers.
This allows us to break ties for mutations which have been observed in the same number of
patients. For example in the TP53 gene, p.Arg181Cys, p.Arg282Gln, and p.Arg282Pro all have
been observed in two out of 9,450 pan-cancer patients. However, p.Arg181Cys (mutability 1.00
x 10^{-5}) is considered a passenger mutation, p.Arg282Gln (mutability 8.11 x 10^{-6}) as a potential
driver, and p.Arg282Pro (mutability 4.90 x 10^{-7}) as a driver mutation. Indeed, p.Arg282Pro is
annotated as driver mutation in the combined dataset, and despite the low mutation frequency,
it is correctly classified as a driver by its B-Score. While there are methods based on hotspot
identification, our ranking of mutations based on B-Score cannot be directly compared to the list
of hotspots proposed previously[21]. Hotspots are defined for sites, whereas B-Score assesses
specific mutations, and different mutations from the same hotspot can be drivers or passengers.
For instance, TP53 Tyr236 site is annotated as a hotspot in [21,35], however p.Tyr236Phe mutation
is experimentally characterized as neutral in the IARC database.

Some efforts have been focused so far on developing a comprehensive set of cancer driver
mutations verified at the levels of functional assays or animal models [36-38]. However, existing sets
often contain predictions and very few neutral passenger mutations. The vast majority of
computational prediction methods rely on machine learning algorithms trained on mutations
from a few genes and/or on recurrent mutations as estimates of driver events, in many cases the
performance is evaluated on synthetic benchmarks. In this study we collected experimental
evidence on cancer mutations, overall 3,787 neutral and 880 driver mutations from 17 genes and

488 neutral and 255 activating mutations from 50 genes. Intriguingly, we found that
experimentally annotated driver mutations have a lower background mutability than neutral
mutations, suggesting possible action of context-dependent codon bias towards less mutable
codons at critical sites for these genes, although more studies would have to be conducted to
further investigate this hypothesis.

We developed and tested two models, LR and B-Score, to adjust the number of reoccurrences of
a mutation by its expected background mutability in order to find sites which may be under
selection in cancer. We find that while including background mutability improves performance
over reoccurrence alone in distinguishing between driver and passenger mutations, the choice
of how to account for mutability is also important. B-Score integrates information about
observed rate and total sample size, something not captured by an odds ratio alone. B-Score
implements a binomial test as a probabilistic model to estimate the number of mutations
expected purely from the context-dependent background nucleotide or codon mutability, given
a certain sample size. The advantages of this model are that: (i) it is intuitive (ii) does not rely on
many parameters and (iii) does not involve explicit training on driver and passenger mutation
sets. One of the disadvantages of this model is that it requires the knowledge of a number of
patients tested. We found that B-Score exhibited a considerably improved performance of driver
mutation prediction compared to the conventional way of using mutational frequency of
occurrence. It performed comparably or better to state-of-the-art methods, many of which were
trained on existing datasets of mutations and included multiple features. Moreover, for highly
heterogeneous cancer types, the background mutational processes may differ between cohorts
of cancer patients, thus an appropriate cancer-specific model for background mutability should
be applied. Taken together, our model provides means to explore mutation rates and enables an
understanding of the differential roles that background mutation rate and selection play in
shaping the observed cancer spectrum.

Methods

Datasets of experimental functional assays and cancer mutations

Missense mutations for TP53 gene with experimentally determined functional transactivation

activities were obtained from IARC P53 database where they were classified as functional,

partially-functional, and non-functional[39]. The second dataset, referred hereinafter to as

"Martelotto *et al.*", contains experimental evidence collected from literature [26]. The

experimental evidence of impact of mutations included changes in enzymatic activity, response

to ligand binding, impacts on downstream pathways, an ability to transform human or murine

cells, tumor induction in vivo, or changes in the rates of progression-free or overall survival in

pre-clinical models. In "Martelotto *et al.*" dataset mutations were considered "damaging" if there

was literature evidence to support their impact on at least one of the above-mentioned

categories. Mutations with no significant impacts on the wild-type protein function were

classified as "neutral". Mutations with no reliable functional evidence were regarded as

"uncertain" and were not used in this study. The third dataset included experimentally verified

BRCA1 mutations and was originally collected by using deep mutational scanning to measure the

effects of missense mutations on the ability of BRCA1 to participate in homology-directed repair.

Missense mutations were categorized as either neutral or damaging [40,41]. Noteworthy, BRCA1 set

contained inherited germline as well as somatic mutations. The fourth dataset explored over

81,000 tumors to identify drivers of hypermutation in DNA polymerase epsilon and delta genes

(POLE/POLD1). Drivers of hypermutation were variants which occurred in a minimum of two

hypermutant tumors, which were never found in lowly mutated tumors, and did not co-occur

with an existing known driver mutation in the same tumor. Other variants in these genes were

considered passengers with respect to hypermutation [42].

Finally, we assembled a *combined dataset* that included mutations from these four datasets

described above. We removed redundant and conflicting entries when mutations annotated as

non-functional or neutral in one dataset were also annotated as damaging or benign in another.

All mutations were categorized as "non-neutral" (affecting function, binding or transforming) and

"neutral" (other mutations). We treated "functional" and "partially -functional" mutations in

IARC TP53 dataset as "neutral", and "non-functional" as "non-neutral". We used missense

mutations in order to compare with Condel and cancer FatHMM scores. Overall, the combined

dataset contains 4,667 mutations (3,787 neutral and 880 non-neutral) from 17 genes (Table S3,

S7).

As an additional validation on driver prediction performance, we used a recently obtained

dataset containing missense mutations which were annotated based on their effects on cell-

viability in Ba/FC and MCF10A models[27]. Activating mutations were annotated as those

mutations where the cell viability was higher than the wild-type gene, and neutral mutations

were those mutations where cell-viability was similar to the wild-type.　Ng *et al.* used these

consensus functional annotations to compare the performance of 21 different computational

tools in classifying between activating and neutral mutations using ROC analysis, with activating

mutations acting as the positive set and neutral as the negative set. Validation dataset contained

743 mutations (488 neutral and 255 activating) from 50 genes.

The Catalogue of Somatic Mutations in Cancer (COSMIC) database stores data on somatic cancer

mutations and integrates the experimental data from the full-genome (WGS) and whole-exome

(WES) sequencing studies[43]. Cancer census genes were defined according to COSMIC release v84.

Genes were binned into those with oncogene and tumor suppressor (TSG) functions. For analyses

comparing oncogenes and TSG, genes classified as only fusion genes or those with both

oncogenic and TSG activities were not used. This resulted in 205 oncogenes and 167 TSG (Table

S2). A subset of 12,013 tumors from COSMIC release v85 was used to extract observed frequency

data. COSMIC v85 samples which came from cell-lines, xenografts, or organoid cultures were

excluded. Only mutations with somatic status of "Confirmed somatic variant" were included and

mutations which were flagged as SNPs were excluded. For each cancer patient, a single sample

from a single tumor was used. Additionally, it is possible that the same patient may be assigned

different unique identifiers in different papers, and duplicate tumor samples may be erroneously

added to COSMIC database during manual curation. These samples may affect the recurrence

counts of mutations and, consequently, the B-score. We applied clustering in order to detect and

remove any redundant tumor samples. Each sample was represented as a binary vector with 1 if

a sample had a mutation in a particular genomic location and 0 otherwise. The binary vectors

were compared with Jaccard distance metric,

$J = \frac{|A \cup B| - |A \cap B|}{|A \cup B|}$

where identical samples have $J = 0$, followed by agglomerative clustering with complete linkage.

Non-singleton clusters with pairwise distance cutoff of $J = 0.3$ were extracted and only one

representative of each cluster was used, whereas other samples were discarded.

Calculation of context-dependent DNA background mutability

Context-dependent mutational profiles were constructed from the pools of mutations from

different cancer samples by counting mutations observed in a specific trinucleotide context.

Altogether, there are 64 different types of trinucleotides and three types of mutations $x \gg y$ (for

example C≫A, C≫T, C≫G and so on) in the central position of each trinucleotide which results

in 192 trinucleotide context-dependent mutation types. In a mutated double-stranded DNA both

complementary purine-pyrimidine nucleotides are substituted, and therefore we considered

only substitutions in pyrimidine (C or T) bases, resulting in 96 possible context-dependent

mutation types: m = {N [$x \gg y$] N}, where N = {A, C, G, T}. Thus, mutational profile can be

expressed as a vector of a number of mutations of certain type $(f_1, \dots, f_{96})$. Profiles were

constructed under the assumption that vast majority of cancer context-dependent mutations

have neutral effects, while only a negligible number of these mutations in specific sites are under

selection. To assure this, we removed recurrent mutations (observed twice or more times in the

same site) as these mutations might be under selection in cancer. In the current study we used

pan-cancer and cancer-specific mutational profiles for breast, lung adenocarcinoma, and skin

adenocarcinoma cancer derived from MutaGene [44].

We applied mutational profiles to build DNA context-dependent probabilistic models that

described baseline DNA mutagenesis per nucleotide or per codon. *Mutability* was defined as a

probability to observe a context-dependent nucleotide mutation purely from the baseline

mutagenic processes operating in a given group of samples used to derive the mutational profile.

Mutability is proportional to the expected mutation rate of a certain type of mutation (96

altogether) regardless of the genomic site it occurs. It was calculated using the total number

of mutations of type *x ≫ y* in a certain local context, which can be obtained from the context-

dependent mutational profile, f_m . Given the number of cancer samples used to construct

mutational profile, *N*, and the number of different trinucleotides of type *t* in a diploid human

exome, n_t (calculated from the reference genome), the mutability of a nucleotide mutation is

calculated as:

439
$$p_m^{nuc} = \frac{f_m}{N\,n_t} \qquad (1)$$

In protein-coding sequences it is practical to calculate mutation probability for a codon in its local

pentanucleotide context, given trinucleotide contexts of each nucleotide in the codon. For a

given transcript of a protein, at exon boundaries the local context of the nucleotides was taken

from the genomic context. The COSMIC consensus transcript was chosen for the transcript for

each protein. Changes in codons can lead to amino acid substitutions, synonymous and nonsense

mutations. Therefore, we calculate a probability to observe a specific type of codon change which

can be realized by single nucleotide mutations at each codon position *i* as:

448
$$p_M^{codon} = 1 - \prod_i^3\left(1 - \sum_j^k p_{ij}^{nuc}\right) \qquad (2)$$

Where *k* denotes a number of mutually exclusive mutations at codon position *i*. For example, for

Phe codon "TTT" in a given context 5'-A and G-3': three single nucleotide mutations can lead to

Phe->Leu substitution (to codons "TTG", "TTA" and "CTT" for Leu): A[T≫C]TTG in the first codon

position or mutually exclusive ATT[T≫G]G and ATT[T>>A]G in the third codon position (Figure

S5). In this case the probability of Phe→ Leu substitution in the $ATTTG$ context can be calculated

as $p_{Phe\rightarrow Leu}^{codon} = 1 - (1 - p_{A[T\rightarrow C]T})(1 - p_{T[T\rightarrow A]G} - p_{T[T\rightarrow G]G})$ where trinucleotide probabilities

were taken from the mutational profile. Amino acid substitutions corresponding to each

missense mutation are calculated by translating the mutated and wild type codons using a

standard codon table. Codon mutability strongly depends on the neighboring codons as

illustrated in Figure S5.

Gene-weight adjusted mutability

Gene weights estimate a relative probability of a gene compared to other genes to be mutated

in cancer through somatic mutagenesis. There are multiple ways the gene weights can be

calculated.

SNP-based weight was calculated using the number of SNPs in the vicinity of the gene of interest.

We used the "EnsDb.Hsapiens.v86" database to find genomic coordinates of a gene, including

introns, and extended the range in both 3' and 5' directions according to the window size (Table

S6). We then counted the number of common SNPs from dbSNP database[45] within the genomic

region. Gene weight was calculated as: $\omega_g^{SNP} = \frac{n_{SNP}}{L_{window}}$, where n_{SNP} is the number of SNPs and

L_{window} is the length of the genomic region in base pairs. We tested several window sizes for

defining the genomic regions around the gene of interest (Table S6).

Mutation-based weight was calculated using the number of nucleotide sites with reoccurring

mutations counted only once to avoid the bias that may be present due to selection on individual

sites: $\omega_g^{mut\_sites} = \frac{k_g}{n_k}$. Here k_g is the number of mutated sites and n_k is the number of base pairs

in the gene transcript.

Silent mutation-based weight was introduced previously and was shown to be superior in

assessment of significant non-synonymous mutations across genes[46]. This weight can be

calculated by taking into account only silent somatic mutations: $\omega_g^{silent} = \frac{s_g}{N\,L_g}$. Here s_g is the

total number of somatic silent mutations within the gene, N is the number of tumor samples

and L_g is the number of codons in the gene transcript.

No-outlier-based weight introduced previously[21] takes into account the number of all codon

mutations within a gene, C_g, excluding mutations in outlier codon sites bearing more than the

99[th] percentile of mutations of the gene: $\omega_g^{out} = \frac{C_g}{N\,L_g}$, normalized by gene length L_g in amino

acids and the total number of samples N.

Using gene weights, an adjusted probability per codon can be then expressed as:

491
$$p'_M = \omega_g p_M \qquad\qquad (3)$$

492   Similarly, per nucleotide probability can be calculated adjusted by gene weight.

494   ***Identification of significant mutations***

495   We introduced two measures which take into account the background DNA mutability. The first

496   measure (LR) is calculated as a ratio of the numbers of observed and expected mutations in a

497   given nucleotide site or in a given codon:

498
$$LR = \log\left(\frac{n_{obs}}{n_{exp}}\right) \qquad\qquad (4)$$

where the expected number of mutations, $n_{exp} = N p_m^{nuc}$ or $n_{exp} = N p_M^{codon}$ where *N* is the

number of tumor samples and p_m^{nuc} or p_M^{codon} are calculated using equations (1) or (2) and the

observed number of mutations n_{obs} in a given site is taken from COSMIC v85 with a pseudo count

correction ($n_{obs} + 1$) to account for mutations that have not been observed due to a limited

tumor sample collection.

The second measure uses the binomial model to calculate the probability of observing a certain

type of mutation in a given site more frequently than *k*:

508
$$B_{score} = \sum_{k=n+1}^{N} \binom{N}{k} p^k (1-p)^{N-k} \qquad\qquad (5)$$

where $p = p'_M$ or $p = p'_m$ and k is the number of observed mutations of a given type at a

particular nucleotide or codon. Depending on the dataset chosen or a particular cohort of

patients (for instance, corresponding to one cancer type), the total number of samples N and the

numbers of observed mutations k will change. While ranking mutations in a given gene, B_{score}

can further be adjusted for multiple-testing with Benjamini-Hochberg correction as implemented

on the MutaGene website.

Statistical and ROC analyses

Differences between various groups were tested with the Kruskal-Wallis, Dunn test, and Mann-

Whitney-Wilcoxon tests implemented in *R* software. Dunn's test is a non-parametric pairwise

multiple comparisons procedure based on rank sums; it is used to infer difference between

means in multiple groups and was used because it is relatively conservative post-hoc test for

Kruskal-Wallis. Associations between mutability and observed frequency (the number of

individuals with a mutation in whole-exome/genome studies from COSMIC), was tested using

Pearson as well as Spearman correlation tests since the variables were not normally distributed.

Where *R* reports the calculated p-value below 2.2×10^{-16}, the value has been shown as the alpha

level, $p < 0.01$.

To quantify the performance of scores, we performed Receiver Operating Characteristics (ROC)

and precision-recall analyses. Sensitivity or true positive rate was defined as TPR=TP/(TP + FN)

and specificity was defined as SPC=TN/(FP+TN). Additionally, in order to account for imbalances

in the labeled dataset, the quality of the predictions was described by Matthews correlation

coefficient (MCC):

534
$$\mathrm{MCC} = \frac{TP * TN + \mathrm{FP} * \mathrm{FN}}{\sqrt{(TP + FP) * (TP + FN) * (TN + FP) * (TN + FN)}}$$

ACKNOWLEDGEMENT

We thank Yuri Wolf, Alejandro Schaffer and Igor Rogozin for helpful discussions. The work was

supported by Intramural Research Programs of the National Library of Medicine, National

National Institutes of Health. Minghui Li was supported by the National Natural Science

Foundation of China (Grant No. 31701136) and Natural Science Foundation of Jiangsu Province,

China (Grant No. BK20170335)

References

1.   Lynch, M. Rate, molecular spectrum, and consequences of human mutation. *Proc Natl Acad Sci U*
544   *S A* **107**, 961-8 (2010).
545   2.   Greenman, C. *et al.* Patterns of somatic mutation in human cancer genomes. *Nature* **446**, 153-8
546   (2007).

3. Leedham, S. & Tomlinson, I. The continuum model of selection in human tumors: general paradigm or niche product? *Cancer Res* **72**, 3131-4 (2012).

4. Dagogo-Jack, I. & Shaw, A.T. Tumour heterogeneity and resistance to cancer therapies. *Nat Rev Clin Oncol* **15**, 81-94 (2018).

5. Hiley, C., de Bruin, E.C., McGranahan, N. & Swanton, C. Deciphering intratumor heterogeneity and temporal acquisition of driver events to refine precision medicine. *Genome Biol* **15**, 453 (2014).

6. Salk, J.J., Fox, E.J. & Loeb, L.A. Mutational heterogeneity in human cancers: origin and consequences. *Annu Rev Pathol* **5**, 51-75 (2010).

7. Alexandrov, L.B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415-21 (2013).

8. Rogozin, I.B. *et al.* Mutational signatures and mutable motifs in cancer genomes. *Brief Bioinform* (2017).

9. Michaelson, J.J. *et al.* Whole-genome sequencing in autism identifies hot spots for de novo germline mutation. *Cell* **151**, 1431-42 (2012).

10. Hodgkinson, A. & Eyre-Walker, A. Variation in the mutation rate across mammalian genomes. *Nat Rev Genet* **12**, 756-66 (2011).

11. Lynch, M. *et al.* Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet* **17**, 704-714 (2016).

12. Stratton, M.R. Exploring the genomes of cancer cells: progress and promise. *Science* **331**, 1553-8 (2011).

13. Schuster-Bockler, B. & Lehner, B. Chromatin organization is a major influence on regional mutation rates in human cancer cells. *Nature* **488**, 504-7 (2012).

14. Polak, P. *et al.* Cell-of-origin chromatin organization shapes the mutational landscape of cancer. *Nature* **518**, 360-364 (2015).

15. Rogozin, I.B. & Kolchanov, N.A. Somatic hypermutagenesis in immunoglobulin genes. II. Influence of neighbouring base sequences on mutagenesis. *Biochim Biophys Acta* **1171**, 11-8 (1992).

16. Chen, C., Qi, H., Shen, Y., Pickrell, J. & Przeworski, M. Contrasting Determinants of Mutation Rates in Germline and Soma. *Genetics* **207**, 255-267 (2017).

17. Gonzalez-Perez, A. & Lopez-Bigas, N. Functional impact bias reveals cancer drivers. *Nucleic Acids Res* **40**, e169 (2012).

18. Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* **171**, 1029-1041 e21 (2017).

19. Araya, C.L. *et al.* Identification of significantly mutated regions across cancer types highlights a rich landscape of functional molecular alterations. *Nat Genet* **48**, 117-25 (2016).

20. Peterson, T.A., Gauran, I.I.M., Park, J., Park, D. & Kann, M.G. Oncodomains: A protein domain-centric framework for analyzing rare variants in tumor samples. *PLoS Comput Biol* **13**, e1005428 (2017).

21. Chang, M.T. *et al.* Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. *Nat Biotechnol* **34**, 155-63 (2016).

22. Porta-Pardo, E. *et al.* Comparison of algorithms for the detection of cancer drivers at subgene resolution. *Nat Methods* **14**, 782-788 (2017).

23. Wong, W.C. *et al.* CHASM and SNVBox: toolkit for detecting biologically important single nucleotide mutations in cancer. *Bioinformatics* **27**, 2147-8 (2011).

24. Li, M. *et al.* Balancing Protein Stability and Activity in Cancer: A New Approach for Identifying Driver Mutations Affecting CBL Ubiquitin Ligase Activation. *Cancer Res* **76**, 561-71 (2016).

25. Stehr, H. *et al.* The structural impact of cancer-associated missense mutations in oncogenes and tumor suppressors. *Mol Cancer* **10**, 54 (2011).

26.   Martelotto, L.G. *et al.* Benchmarking mutation effect prediction algorithms using functionally
validated cancer-related missense mutations. *Genome Biol* **15**, 484 (2014).
27.   Ng, P.K. *et al.* Systematic Functional Annotation of Somatic Mutations in Cancer. *Cancer Cell* **33**,
450-462 e10 (2018).
28.   Gonzalez-Perez, A. & Lopez-Bigas, N. Improving the assessment of the outcome of
nonsynonymous SNVs with a consensus deleteriousness score, Condel. *Am J Hum Genet* **88**, 440-
9 (2011).
29.   Shihab, H.A. *et al.* Predicting the functional, molecular, and phenotypic consequences of amino
acid substitutions using hidden Markov models. *Hum Mutat* **34**, 57-65 (2013).
30.   Bailey, M.H. *et al.* Comprehensive Characterization of Cancer Driver Genes and Mutations. *Cell*
173, 371-385 e18 (2018).
31.   Lawrence, M.S. *et al.* Mutational heterogeneity in cancer and the search for new cancer-
associated genes. *Nature* **499**, 214-218 (2013).
32.   Supek, F., Minana, B., Valcarcel, J., Gabaldon, T. & Lehner, B. Synonymous mutations frequently
act as driver mutations in human cancers. *Cell* **156**, 1324-1335 (2014).
33.   Li, M., Goncearenco, A. & Panchenko, A.R. Annotating Mutational Effects on Proteins and Protein
Interactions: Designing Novel and Revisiting Existing Protocols. *Methods Mol Biol* **1550**, 235-260
(2017).
34.   Molina-Vila, M.A. *et al.* Activating mutations cluster in the "molecular brake" regions of protein
kinases and do not associate with conserved or catalytic residues. *Hum Mutat* **35**, 318-28 (2014).
35.   Chang, M.T. *et al.* Accelerating Discovery of Functional Mutant Alleles in Cancer. *Cancer Discov* **8**,
174-183 (2018).
36.   Ainscough, B.J. *et al.* DoCM: a database of curated mutations in cancer. *Nat Methods* **13**, 806-7
(2016).
37.   Landrum, M.J. *et al.* ClinVar: public archive of relationships among sequence variation and human
phenotype. *Nucleic Acids Res* **42**, D980-5 (2014).
38.   Tamborero, D. *et al.* Cancer Genome Interpreter annotates the biological and clinical relevance of
tumor alterations. *Genome Med* **10**, 25 (2018).
39.   Olivier, M. *et al.* The IARC TP53 database: new online mutation analysis and recommendations to
users. *Hum Mutat* **19**, 607-14 (2002).
40.   Starita, L.M. *et al.* Massively Parallel Functional Analysis of BRCA1 RING Domain Variants. *Genetics*
200, 413-22 (2015).
41.   Mahmood, K. *et al.* Variant effect prediction tools assessed using independent, functional assay-
based datasets: implications for discovery and diagnostics. *Hum Genomics* **11**, 10 (2017).
42.   Campbell, B.B. *et al.* Comprehensive Analysis of Hypermutation in Human Cancer. *Cell* **171**, 1042-
1056 e10 (2017).
43.   Forbes, S.A. *et al.* COSMIC: somatic cancer genetics at high-resolution. *Nucleic Acids Res* **45**, D777-
D783 (2017).
44.   Goncearenco, A. *et al.* Exploring background mutational processes to decipher cancer genetic
heterogeneity. *Nucleic Acids Res* **45**, W514-W522 (2017).
45.   Sherry, S.T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res* **29**, 308-11
(2001).
46.   Evans, P., Avey, S., Kong, Y. & Krauthammer, M. Adjusting for background mutation frequency
biases improves the identification of cancer driver genes. *IEEE Trans Nanobioscience* **12**, 150-7
(2013).

Figure 1. Mutability distributions by mutation type and gene role in cancer. **(A)** Cumulative distribution of
codon mutability of silent (green color), nonsense (red color) and missense (blue color) mutations. **(C)**
Cumulative distribution of nucleotide mutability for silent, nonsense and missense mutations. Inset shows
the probability density distributions of mutability by mutation type. Significance was determined by
Dunn's test; difference with $p < 0.01$). is marked with a double asterisk. **(B)** and **(D)** are codon and
nucleotide mutability respectively binned by frequency. '0', '1', '2' and '3+' refer to mutations that were
not observed (including all possible point mutations), observed once, twice, or in three or more cancer
samples. **(E)** Pooled mutations in cancer census genes were grouped for oncogene and tumor suppressor
(TSG) genes. Boxplots show codon mutability calculated with pan-cancer model. See Table S1 for counts.

Figure 2. Comparison between nucleotide mutability (expected) spectrum of all possible mutations (blue) and mutations which were observed in cancer patients (brown). **(A)** Mutations in 520 cancer-related genes; **(B)** CASP8 and **(C)** TP53 genes. Inset shows the cumulative distribution functions for both spectra. Annotations in **(A)** show nucleotide mutation types.

Figure 3. Relationship between codon mutability and frequency of mutations. Histograms show the Spearman rank correlation coefficients between observed mutations and mutability across cancer genes with at least 10 observed mutations of each type: **(A)** missense (blue), **(B)** nonsense (red) and **(C)** silent (green). Filled bars in the left column denote genes with significant correlation at $p < 0.01$. Scatterplots with regression lines and confidence intervals show the linear relationship between mutability and observed frequency of each type of mutation for several representative genes. Adjusted R^2 shown to convey goodness of fit. Bar graphs show Spearman correlation coefficient for genes with significant correlation at $p < 0.01$. Genes with bold font are tumor suppressors (TSG), underlined genes are oncogenes, and plain font were either categorized as both TSG and oncogene or fusion genes.

Figure 4. Codon mutability of missense mutations grouped by the effect on protein function. **(A)** Mutations in TP53 gene were characterized as functional, partially functional and non-functional. (B) Mutations from Martelotto *et al.* set were categorized as neutral and non-neutral. (C) Mutations in BRCA1 gene were annotated as benign and deleterious. (D) Mutations from the combined dataset were categorized as neutral and non-neutral. Significant differences with $p < 0.01$ are marked with a double asterisk and difference with $p < 0.05$ is marked with a single asterisk. Mutability was calculated with pan-cancer background model; see Supplementary Figure S4 for analysis with different background models.

Figure 5. Assessment of classification performance between neutral and non-neutral mutations in a
combined dataset. **(A)** ROC curves for B-Score, Condel, FatHMM, and observed mutational frequency.
Inset shows the performance of highlighted area corresponding to up to 10% FPR. **(B)** Precision-recall
curves for the same benchmark set. Inset shows the performance of highlighted area corresponding to
precision over 50%. The ROC for observed frequency cannot be calculated for all mutations because some
experimentally validated mutations were not observed in tumor samples.
716

Figure 6. Ranking of mutations and prediction of driver mutations based on mutational frequencies adjusted by mutability. Snapshots from MutaGene server show the results of analysis of EGFR gene with a Pan-cancer mutability model. **(A)** Scatterplot with expected mutability versus observed mutational frequencies. **(B)** Top list of mutations ranked by their B-Scores. **(C)** EGFR nucleotide and translated protein sequence shows per nucleotide site mutability (green line), per codon mutability (orange line), as well as mutabilities of nucleotide and codon substitutions (heatmaps). Mutations observed in tumors from ICGC repository are shown as circles colored by their prediction status: Driver, Potential driver, and Passenger.

726 **Table 1.** Comparison of different measures to distinguish neutral from non-neutral mutations using
727 combined experimental dataset. Scores developed in this study (B-Score and LR) are underlined. The best
728 performance is shown in bold font. AUC-ROC and AUC-PR values for observed frequency count were
729 extrapolated since some experimentally validated mutations were not observed in tumor samples.
730
731

Measure	AUC-ROC	AUC-PR	Matthews correlation	Sensitivity at 10% FPR
<u>LR</u>	0.80	0.61	0.47	0.50
<u>B-Score</u>	0.82	**0.68**	**0.58**	0.57
Condel	0.87	0.59	0.53	**0.62**
FatHMM	**0.88**	0.63	**0.58**	0.52
Mutability	0.67	0.33	0.21	0.24
Observed frequency	0.71	0.58	0.52	0.48

732
733

734 **Supplementary Figures and Tables**

738 **Figure S1.** Mutability values for observed mutations in COSMIC v85 subset cohort and all theoretically
739 possible mutations that were not observed: **(A)** Codon mutability; **(B)** Nucleotide mutability. Differences
740 on Mann-Whitney-Wilcoxon test significant at p < 0.01.

Figure S2. Relationship between codon mutability and observed mutation frequency by mutation type and role in cancer. **(A)** Scatterplot showing the observed mutation frequency for all genes and mutation types. **(B)** Scatterplots for oncogenes (n = 202) and **(C)** TSG (n = 166) for all mutation types. Different colors show scatterplots broken down by mutation type: missense (blue), nonsense (red) and silent (green). Spearman and Pearson correlation coefficient with respective p-values shown in all, significant at p < 0.01 in bold.

Figure S3. Relationship between codon mutability and observed mutation frequency by mutation type and molecular genetics. **(A)** Genes with only dominant mutations, **(E)** Genes with only recessive mutations. Different colors show scatterplots broken down by mutation type: missense (blue), nonsense (red) and silent (green). As in Figure 1E, pooled mutations in cancer census genes grouped by Dominant and Recessive mutations. Spearman and Pearson correlation coefficient with respective p-values shown in all, significant at p < 0.01 in bold. Counts summarized in Table S1.

Figure S4. Mutability of missense mutations calculated with different background models. **(A)** TP53 **(B)** BRCA1 **(C)** Martelotto et al. datasets shown in Figure 4. As in Figure 4, colors refer to functional annotation, yellow – Function/Neutral/Benign, orange – partially-functional, red – Non-functional/Non-neutral/Deleterious. Mutability values were calculated with background models of breast cancer, lung adenocarcinoma cancer, skin adenocarcinoma, and pan-cancer. Additionally, a background model for common non-pathogenic SNPs in general human population representing germline mutability was used. All models are available on the MutaGene website (https://www.ncbi.nlm.nih.gov/projects/mutagene/). Mutational profiles for each background model shown on the top. Significance tests summarized in Table S4.

$$p_M^{codon} = 1 - \prod_i^3 \left(1 - \sum_j^k p_{ij}^{nuc}\right)$$

$$p_{Phe \to Leu}^{codon} = 1 - (1 - p_{A[T \to C]T})(1 - p_{T[T \to A]G} - p_{T[T \to G]G})$$

765
766
767 **Figure S5.** Example of calculation of codon mutability from nucleotide mutabilities of all possible Phe →
768 Leu missense mutations in the Phe codon TTT with a specific pentanucleotide context.
769

770   **Table S1.** Counts for boxplots in Figure 1 and Figure S2.

		Count group			
		0 (not observed)	1	2	3+
		Codon mutations			
	Missense	2625306	31510	1898	844
	Nonsense	140706	3264	322	193
	Silent	425059	11009	770	136
		Nucleotide mutations			
	Missense	2923312	32510	1942	844
	Nonsense	174350	3418	327	197
	Silent	906077	11665	677	124
		Codon mutations - Oncogene			
	Missense	772789	10355	670	333
	Nonsense	40310	664	36	7
	Silent	124054	3890	327	47
		Codon mutations - TSG			
	Missense	1023631	13063	763	237
	Nonsense	55041	1907	224	140
	Silent	165909	4287	290	59
		Codon mutations - Dominant			
	Missense	1678008	20256	1201	440
Mutation	Nonsense	90646	1462	94	28
Type	Silent	270997	7414	554	83
		Codon mutations - Recessive			
	Missense	689246	9154	559	371
	Nonsense	36894	1605	212	163
	Silent	112121	2808	167	44
		Codon mutations - Dom/Rec			
	Missense	57126	727	44	10
	Nonsense	2818	95	8	1
	Silent	9190	278	9	1
		Codon mutations - Rec/X			
	Missense	5437	60	4	0
	Nonsense	283	19	2	0
	Silent	879	25	0	0
		Codon mutations - No Molecular Genetics Information Provided			
	Missense	195489	1313	90	23
	Nonsense	10065	83	6	1
	Silent	31872	484	40	8

771 **Table S2.** Correlation between mutability and recurrence of mutations in cancer-associated genes
772 https://www.ncbi.nlm.nih.gov/research/mutagene/static/data/TableS2.xlsx
773

774 **Table S3.** Combined dataset with experimentally annotated neutral and non-neutral mutations in 17
775 genes
776 https://www.ncbi.nlm.nih.gov/research/mutagene/static/data/TableS3.xlsx

777 **Table S4.** Comparison of mutability on three experimental datasets with different cancer-specific
778 background mutation models

	Model	P Value - Dunn Test	Comparison
TP53 IARC	Pan-cancer	2.17×10^{-9}	functional - non-functional
		0.36	functional - partially-functional
		1.41×10^{-8}	non-functional - partially-functional
	Breast cancer	1.84×10^{-9}	functional - non-functional
		0.47	functional - partially-functional
		6.11×10^{-8}	non-functional - partially-functional
	Lung adenocarcinoma	9.82×10^{-5}	functional - non-functional
		0.23	functional - partially-functional
		3.13×10^{-5}	non-functional - partially-functional
	Skin adenocarcinoma	0.33	functional - non-functional
		0.41	functional - partially-functional
		0.43	non-functional - partially-functional
	SNP	4.90×10^{-5}	functional - non-functional
		0.42	functional - partially-functional
		0.00013	non-functional - partially-functional
Martelotto et. al		**P-Value – Mann-Whitney-Wilcoxon**	
	Pan-cancer	6.24×10^{-8}	
	Breast cancer	9.06×10^{-7}	
	Lung adenocarcinoma	0.02	
	Skin adenocarcinoma	0.28	
	SNP	8.90×10^{-12}	
BRCA1 - DMS	Pan-cancer	0.04	
	Breast cancer	0.07	
	Lung adenocarcinoma	0.12	
	Skin adenocarcinoma	0.29	
	SNP	0.49	

780 **Table S5.** Performance of different measures on each individual dataset from the combined data set.
781

	AUC-ROC	AUC-PR	Matthews's correlation	Sensitivity at 10% FPR	Dataset
LR	0.57	0.35	0.25	0.29	BRCA1-DMS
B-score	0.58	0.32	0.23	0.29	
Condel	**0.78**	**0.56**	0.50	**0.53**	
FatHMM	0.68	0.52	**0.52**	0.35	
Mutability	0.57	0.33	0.24	0.29	
Frequency	0.51	0.23	0.03	0.11	
LR	0.77	0.95	0.32	0.45	Martelotto et. al
B-score	0.79	0.95	0.32	0.52	
Condel	0.85	**0.97**	0.45	**0.55**	
FatHMM	**0.88**	0.96	**0.69**	**0.56**	
Mutability	0.64	0.91	0.18	0.24	
Frequency	0.7	0.94	0.28	0.47	
LR	0.61	0.06	0.17	0.26	POLE/POLD1 Variants
B-score	0.65	**0.13**	**0.30**	0.32	
Condel	**0.90**	0.07	0.22	**0.65**	
FatHMM	0.36	0.01	0.04	0.47	
Mutability	0.57	0.02	0.05	0.24	
Frequency	0.57	0.09	0.21	0.23	
LR	0.82	0.77	0.54	0.58	TP53 Transactivation
B-score	**0.84**	**0.80**	**0.62**	**0.67**	
Condel	**0.84**	0.74	0.56	0.6	
FatHMM	0.80	0.65	0.51	0.46	
Mutability	0.61	0.47	0.19	0.18	
Frequency	0.79	0.76	0.58	0.67	

782   **Table S6.** Performance metrics for the binomial model on combined and validation datasets using
783   different values as a gene weight.

Gene Weight	AUC-ROC	AUC-PR	MCC	Sensitivity at 10% error	Dataset
SNP – 10,000 bp	0.84	0.68	0.58	0.57	
SNP – 20,000 bp	0.84	0.68	0.58	0.57	
SNP – 100,000 bp	0.84	0.69	0.58	0.58	
SNP – 200,000 bp	0.84	0.69	0.58	0.58	
No-outlier based weight	0.67	0.58	0.52	0.50	**Combined**
Mutation based weight	0.71	0.60	0.55	0.50	
No Weight	0.82	0.68	0.58	0.57	
Silent mutation based weight	0.82	0.67	0.58	0.55	
SNP – 10,000 bp	0.75	0.68	0.44	0.47	
SNP – 20,000 bp	0.75	0.68	0.43	0.47	
SNP – 100,000 bp	0.75	0.68	0.43	0.47	
SNP – 200,000 bp	0.75	0.68	0.43	0.47	
No-outlier based weight	0.74	0.67	0.44	0.45	**Validation**
Mutation based weight	0.74	0.67	0.44	0.45	
No Weight	0.76	0.69	0.44	0.49	
Silent mutation based weight	0.72	0.55	0.37	0.40	
SNP, gene weighted by the number of SNPs in a window of various base pairs, window size indicated; No-outlier-based weight, gene weight as described by Chang et. al, see Methods for description. N Mutated Sites, gene weighted by the number of mutated sites in the gene over the total number of sites in the gene; No Weight, genes given no weighting; Silent mutations, gene weighted by number of silent mutations in area					

Table S7. Number of annotated mutations in combined dataset by gene and the gene's role in cancer.

	Neutral Mutations	Non-Neutral Mutations	Role in Cancer
BRAF	0	23	Oncogene
BRCA1	369	95	TSG
BRCA2	50	12	TSG
DICER1	0	11	TSG
EGFR	0	33	Oncogene
ERBB2	5	33	Oncogene
ESR1	0	7	Oncogene, TSG, Fusion
IDH1	0	1	Oncogene
IDH2	0	3	Oncogene
KIT	1	24	Oncogene
KRAS	0	25	Oncogene
MYOD1	0	1	Oncogene
PIK3CA	1	31	Oncogene
POLD1	945	5	Oncogene
POLE	1806	29	Oncogene
SF3B1	0	7	Oncogene
TP53	610	540	Oncogene, TSG, Fusion