Non-synonymous to synonymous substitutions suggest that orthologs tend to keep their functions, while paralogs are a source of functional novelty

Juan M. Escorcia-Rodríguez[1]**, Mario Esposito**[2]**, Julio A. Freyre-González**[1]**, and Gabriel Moreno-Hagelsieb**[2,*]

[1]**Regulatory Systems Biology Research Group, Program of Systems Biology, Center for Genomic Sciences, Universidad Nacional Autońoma de Meχico (UNAM), Av. Universidad s/n, Col. Chamilpa, 62210 Cuernavaca, Morelos, Mexico.**
[2]**Wilfrid Laurier University, Department of Biology, 75 University Ave W, N2L 3C5 Waterloo, ON, Canada**

Corresponding author:
G. Moreno-hagelsieb[*]

Email address: gmorenohagelsieb@wlu.ca

ABSTRACT

Orthologs diverge after lineage separation events and paralogs after gene duplications. Thus, orthologs are expected to remain more functionally coherent across lineages, while paralogs have been proposed as a source of new functions. Because protein functional divergence follows from non-synonymous substitutions, we performed an analysis based on the ratio of non-synonymous to synonymous substitutions (dN/dS), as proxy for functional divergence. We used five working definitions of orthology, including reciprocal best hits (RBH), among other definitions based on network analyses and clustering. The results showed that orthologs, by all definitions tested, had values of dN/dS noticeably lower than those of paralogs, suggesting that, as a group, orthologs tend to be more functionally stable than paralogs. The differences in dN/dS ratios remained suggesting the functional stability of orthologs after eliminating gene comparisons with potential problems, such as genes with high codon usage biases, low coverage of either of the aligned sequences, or sequences with very high similarities. Separation by percent identity of the encoded proteins showed that the differences between the dN/dS ratios of orthologs and paralogs were more evident at high sequence identity, less so as identity dropped. While the last results partially explained the differences between dN/dS ratios of the two groups as being related to differences in protein identity, they also suggested that paralogs undergo functional divergence relatively early after duplication. Our analyses indicate that choosing orthologs as probably functionally coherent remains the right approach in comparative genomics.

INTRODUCTION

Since the beginning of comparative genomics, the assumption was made that orthologs could be expected to conserve their functions more often than paralogs (Mushegian and Koonin, 1996; Huynen and Bork, 1998; Bork et al., 1998; Tatusov et al., 2000). The expectation is based on the definitions of each homolog type: orthologs are characters diverging after speciation events, while paralogs are characters diverging after duplication events (Fitch, 2000). Given those definitions, orthologs could be considered the "same" genes in different species, while paralogy has been proposed as a mechanism for the evolution of new functions, under the argument, in very simplified terms, that one of the copies could maintain the original function, while the other copy would have some freedom to functionally change (Ohno, 1970). This neither means that orthologs cannot evolve new functions, nor that paralogs necessarily evolve new functions. However, a scenario whereby orthologs, as a group, would diverge in functions at a higher rate than paralogs seems far from parsimonious, thus very unlikely. Therefore, it has been customary to use

some working definition of orthology to infer the genes whose products most likely perform the same functions across different lineages (Mushegian and Koonin, 1996; Huynen and Bork, 1998; Bork et al., 1998; Tatusov et al., 2000; Gabaldón and Koonin, 2013).

Despite such a straightforward expectation, a report was published making the surprising claim that orthologs diverged in function more often than paralogs (Nehrt et al., 2011). The controversial article was mainly based on the comparison of Gene Ontology annotations among orthologs and paralogs from two species: humans and mice (Nehrt et al., 2011). If the report were correct, it would mean, for example, that mice myoglobin could be performing the function that human alpha-haemoglobin performs. However, data in the article showed that paralogs found within a genome, had more consistent gene ontology annotations than any homologs between both genomes. This was true even for identical proteins. Thus, rather than functional differences, it was possible that annotations of homologous genes were more consistent within a genome than between genomes. Accordingly, later work showed that gene ontologies suffered from "ascertainment bias", which made annotations more consistent within an organism than without (Thomas et al., 2012; Altenhoff et al., 2012). Later work showed gene expression data suggesting that orthologs had more coherent functions than paralogs (Kryuchkova-Mostacci and Robinson-Rechavi, 2016).

We thus wondered whether we could perform some analyses that did not suffer from annotation bias, and that could cover most of the homologs found between any pair of genomes, even if they had no functional annotations. Given that changes in protein function require changes in amino acids, analyses of non-synonymous to synonymous substitution rates, which compare the relative rates of positive and negative (purifying) selection (Ohta, 1995; Yang and Nielsen, 2000), might serve as proxies for functional divergence. The most functionally stable homologs would be expected to have lower dN/dS ratios compared to less functionally stable homologs. Thus, comparing the dN/dS distributions of orthologs and paralogs could show differences in the relative tendencies of each group to conserve their functions, if that were the case. Since most of the related works have focused on eukaryotes, we centered our analyzes on prokaryotes (Bacteria and Archaea). We used five working definitions of orthology, including RBH that is the foundation of most graph-based orthology prediction methods and is arguably the most usual working definition of orthology (Altenhoff and Dessimoz, 2009; Wolf and Koonin, 2012; Galperin et al., 2019).

While we have been working on this report Escorcia-Rodríguez et al. (2020), an article following the same basic idea, comparing dN/dS distributions between orthologs and paralogs, focusing on vertebrates, was published (David et al., 2020). Whereas the results are consistent, our work focused on the still neglected prokaryotes.

MATERIALS AND METHODS

Genome Data
We downloaded the analyzed genomes from NCBI's RefSeq Genome database (Haft et al., 2018). We performed our analyses by selecting genomes from three taxonomic phyla, using one genome within each phylum as a query genome (Table 1): *Escherichia coli* K12 MG1655 (phylum Proteobacteria, domain Bacteria, assembly ID: GCF_000005845), *Bacillus subtilis* 168 (Firmicutes, Bacteria, GCF_000009045), and *Natrialba magadii* ATCC43099 (Euryarchaeota, Archaea, GCF_000025625).

Orthologs
We used five working definitions of orthology:

Reciprocal best hits (RBH)
We compared the proteomes of each of these genomes against those of other members of their taxonomic phylum using diamond (Buchfink et al., 2015), with the $--very-sensitive$ option, and a maximum e-value of 1×10^{-6} (*-evalue 1e-6*) (Hernández-Salmerón and Moreno-Hagelsieb, 2020). We also required a minimum alignment coverage of 60% of the shortest sequence. Orthologs were defined as reciprocal best hits (RBH) as described previously (Moreno-Hagelsieb and Latimer, 2008; Ward and Moreno-Hagelsieb, 2014; Hernández-Salmerón and Moreno-Hagelsieb, 2020). Except where noted, paralogs were all matches left after finding RBH.

Table 1. Genomes used in this study

Genome ID	Class	Order	Species
Phylum Proteobacteria			
GCF_000005845	Gammaproteobacteria	Enterobacterales	*Escherichia coli*
GCF_002370525	Gammaproteobacteria	Pseudomonadales	*Acinetobacter guillouiae*
GCF_002847445	Alphaproteobacteria	Rhodobacterales	*Paracoccus zhejiangensis*
GCF_004194535	Betaproteobacteria	Neisseriales	*Iodobacter fluviatilis*
GCF_013085545	Deltaproteobacteria	Desulfovibrionales	*Desulfovibrio marinus*
GCF_013283835	Epsilonproteobacteria	Campylobacterales	*Poseidonibacter lekithochrous*
GCF_000317895	Oligoflexia	Bdellovibrionales	*Bdellovibrio bacteriovorus*
GCF_009662475	Acidithiobacillia	Acidithiobacillales	*Acidithiobacillus thiooxidans*
GCF_002795805	Zetaproteobacteria	Mariprofundales	*Mariprofundus aestuarium*
GCF_003574215	Hydrogenophilalia	Hydrogenophilales	*Hydrogenophilus thermoluteolus*
Phylum Firmcutes			
GCF_000009045	Bacilli	Bacillales	*Bacillus subtilis*
GCF_002197645	Bacilli	Lactobacillales	*Enterococcus wangshanyuanii*
GCF_000218855	Clostridia	Eubacteriales	*Clostridium acetobutylicum*
GCF_003991135	Clostridia	Halanaerobiales	*Anoxybacter fermentans*
GCF_000020005	Clostridia	Natranaerobiales	*Natranaerobius thermophilus*
GCF_003966895	Negativicutes	Selenomonadales	*Methylomusa anaerophila*
GCF_003367905	Negativicutes	Veillonellales	*Megasphaera stantonii*
GCF_012317185	Erysipelotrichia	Erysipelotrichales	*Erysipelatoclostridium innocuum*
GCF_000299355	Tissierellia	Tissierellales	*Gottschalkia acidurici*
GCF_001544015	Limnochordia	Limnochordales	*Limnochorda pilosa*
Phylum Euryarchaeota			
GCF_000025625	Halobacteria	Natrialbales	*Natrialba magadii*
GCF_000011085	Halobacteria	Halobacteriales	*Haloarcula marismortui*
GCF_000025685	Halobacteria	Haloferacales	*Haloferax volcanii*
GCF_000195895	Methanomicrobia	Methanosarcinales	*Methanosarcina barkeri*
GCF_000013445	Methanomicrobia	Methanomicrobiales	*Methanospirillum hungatei*
GCF_001433455	Thermococci	Thermococcales	*Thermococcus barophilus*
GCF_000024185	Methanobacteria	Methanobacteriales	*Methanobrevibacter ruminantium*
GCF_000006175	Methanococci	Methanococcales	*Methanococcus voltae*
GCF_000734035	Archaeoglobi	Archaeoglobales	*Archaeoglobus fulgidus*
GCF_000007185	Methanopyri	Methanopyrales	*Methanopyrus kandleri*

The query genomes were the first in each group.

Ortholog groups with inparalogs (InParanoid)

InParanoid is a graph-based tool to identify orthologs and in-paralogs from pairwise sequence comparisons (Sonnhammer and Östlund, 2015). InParanoid first runs all-versus-all blastp and identifies RBH. Then, it uses the RBH as seeds to identify co-orthologs for each gene (which the authors define as in-paralogs), proteins from the same organism that obtain better bits score than the RBH. Finally, through a series of rules, InParanoid cluster the co-orthologs to return non-overlapping groups. The authors define outparalogs as those blast-hits outside of the co-ortholog clusters (Sonnhammer and Östlund, 2015).

We ran InParanoid for each query genome against those of other members of their taxonomic order. InParanoid was run with the following parameters: double blast and 40 bits as score cutoff. The first pass run with compositional adjustment on and soft masking. This removes low complexity matches but truncates alignments (Sonnhammer and Östlund, 2015). The second pass run with compositional adjustment off to get full-length alignments. We used as in-paralogs the combinatorial of the genes of the same organism from the same cluster, and as out-paralogs those blast-hits outside of the co-ortholog clusters.

Orthologous MAtrix (OMA)

OMA is a pipeline and database that provides three different types of orthologs: pairwise orthologs, OMA groups (orthogroups), and hierarchical orthologous groups (Zahn-Zabal et al., 2020). OMA makes an effort to remove xenologs by using a third proteome as *witness of non-orthology* (Roth et al., 2008). To the best of our knowledge, OMA is the only orthology prediction method, still being maintained, able

to deal with xenology. The OMA pipeline for the identification of orthologs is based on best reciprocal Smith-Waterman hits and some tolerance for evolutionary distance that allows for co-orthology. For pairwise orthology identification, a verification step to detect xenologs is applied using a third proteome that retained both pseudo-orthologous genes (Train et al., 2017).

We ran the OMA standalone (version 2.5.0) with all the proteomes for each taxonomic group using the default parameters (Train et al., 2017). We used the pairwise orthology outputs considering the query organisms. For the identification of in-paralogs for the query organism, we used the co-orthologous genes mapping to one or more orthologs in the rest of the organisms. OMA also generates pairwise paralogy outputs, including former candidates for orthologs that did not reach the thresholds or were discarded by a third organism retaining both genes (Zahn-Zabal et al., 2020).

OrthoFinder

OrthoFinder defines an orthogroup as the set of genes derived from a single gene in the last common ancestor of the all species under consideration (Emms and Kelly, 2015). First, OrthoFinder performs all-*versus*-all blastp (Camacho et al., 2009) comparisons and uses an e-value of 1×10^{-3} as a threshold. Then, it normalizes the gene length and phylogenetic distance of the BLAST bit scores. It uses the lowest normalized value of the RBH for either gene in a gene pair as the threshold for their inclusion in an orthogroup. Finally, it weights the orthogroup graph with the normalized bit scores and clusters it using MCL. OrthoFinder outputs the orthogroups and orthology relationships, which can be many to many (co-orthology).

We ran OrthoFinder with all the proteomes for each of the taxonomic groups listed (Table 1). From the OrthoFinder outputs with the orthology relationships between every two species, we used those considering the query organism. From an orthogroup containing one or more orthology relationships, we identified the outparalogs as those genes belonging to the same orthogroup but not to the same orthology relationship. We identified the inparalogs for the query organisms as its genes belonging to the same orthogroup since they derived from a single ancestor gene.

ProteinOrtho

ProteinOrtho is a graph-based tool that implements an extended version of the RBH heuristic and is intended for the identification of ortholog groups between many organisms (Lechner et al., 2011). First, from all-*versus*-all blast results, ProteinOrtho creates subnetworks using the RBH at the seed. Then, if the second best hit for each protein is almost as good as the RBH, it is added to the graph. The algorithm claims to recover false negatives and to avoid the inclusion of false positives (Lechner et al., 2011).

We ran ProteinOrtho pairwise since we needed to identify orthologs and paralogs between the query organisms and the other members of their taxonomic data, not those orthologs shared between all the organisms. ProteinOrtho run blast with the following parameters: an E-value cutoff of 1×10^6, minimal alignment coverage of 50% of the shortest sequence, and 25% of identity. Orthologs were the genes from different genomes that belonged to the same orthogroup, and inparalogs genes from the same genome that belong to the same orthogroup. We reran ProteinOrtho with a similarity value of 75% instead of 90%, to identify outparalogs as those interactions not identified in the first run.

Non-synonymous to synonymous substitutions

To perform dN/dS estimates, we used the CODEML program from the PAML software suite (Yang, 2007). The DNA alignments were derived from the protein sequence alignments using an *ad hoc* program written in PERL. The same program ran pairwise comparisons using CODEML to produce Bayesian estimates of dN/dS (Anisimova et al., 2002; Angelis et al., 2014). The results were separated between ortholog and paralog pairs, and the density distributions were plotted using R (R Core Team, 2020). Statistical analyses were also performed with R.

Codon Adaptation Index

To calculate the Codon Adaptation Index (CAI) (Sharp and Li, 1987), we used ribosomal proteins as representatives of highly expressed genes. To find ribosomal proteins we matched the COG ribosomal protein families described by Yutin *et al.* (Yutin et al., 2012) to the proteins in the genomes under analysis using RPSBLAST (part of NCBI's BLAST+ suite) (Camacho et al., 2009). RPSBLAST was run with soft-masking (*-seg yes -soft_masking true*), a Smith-Waterman final alignment (*-use_sw_tback*), and a maximum e-value threshold of 1×10^{-3} (*-evalue 1e-3*). A minimum coverage of 60% of the COG domain

model was required. To produce the codon usage tables of the ribosomal protein-coding genes, we used the program *cusp* from the EMBOSS software suite (Rice et al., 2000). These codon usage tables were then used to calculate the CAI for each protein-coding gene within the appropriate genome using the *cai* program also from the EMBOSS software suite (Rice et al., 2000).

RESULTS AND DISCUSSION

Reciprocal best hits showed lower dN/dS ratios than paralogs

During the early stages of this research, we ran a few tests using a fast method for estimating dN/dS, namely the Yang and Nielsen approximation (Yang and Nielsen, 2000), which showed promising results. However, we chose to continue these analyses using Bayesian estimates of dN/dS, because they are considered the most robust and accurate (Anisimova et al., 2002; Angelis et al., 2014). To compare the distribution of dN/dS values between orthologs and paralogs, we plotted dN/dS density distributions using violin plots (Fig. 1). These plots demonstrated evident differences, with orthologs showing lower dN/dS ratios than paralogs, thus indicating that orthologs, as a group, have diverged in function less than paralogs. In line with the noticeable differences, Wilcoxon rank tests showed that the differences were statistically significant, with probabilities much lower than 1×10^{-9} (Table S1). Since most comparative genomics work is done using reciprocal best hits (RBH) as a working definition for orthology (Wolf and Koonin, 2012; Galperin et al., 2019), this result suggests that most research in comparative genomics has used the proteins/genes that most likely share their functions.

Differences in dN/dS resisted other working definitions of orthology

A concern with our analyses might arise from our initial focus on reciprocal best hits (RBH). However, RBH might arguably be the most usual working definition of orthology (Altenhoff and Dessimoz, 2009; Wolf and Koonin, 2012; Galperin et al., 2019). Thus, it is important to start these analyses with RBH, at least to test whether RBH are a good choice for the purpose of inferring genes most likely to have similar functions.

Analyses of the quality of RBH for inferring orthology, based on synteny, showed that RBH error rates were lower than 5% (Moreno-Hagelsieb and Latimer, 2008; Wolf and Koonin, 2012; Hernández-Salmerón and Moreno-Hagelsieb, 2020). Other analyses showed that the problem with RBH, was a slightly higher rate of false positives (paralogs mistaken for orthologs), than databases based on phylogenetic and network analyses (Altenhoff and Dessimoz, 2009). Therefore, we can assume that orthologs dominate the RBH dN/dS distributions.

Despite the above justification for focusing on RBH, we considered other four definitions of orthology (Fig. 2, Fig. S1-S4). Orthologs obtained with different working definitions, including one method dealing with xenologs (OMA), showed dN/dS ratio distributions that suggest that a higher proportion of orthologs have similar functions compared to paralogs (Fig. 2).

Differences in dN/dS persisted after testing for potential biases

While the tests above suggest that RBH separate homologs with higher tendencies to preserve their functions than other homologs, we tested for some potential biases. A potential problem could arise from comparing proteins of very different lengths. We thus filtered the dN/dS results to keep those where the pairwise alignments covered at least 80% of the length of both proteins. The results showed shorted tails in both density distributions, but the tendency for orthologs to have lower dN/dS values remained (Fig. 2, Fig. S5).

Another parameter that could bias the dN/dS results is high sequence similarity. In this case, the programs tend to produce high dN/dS ratios. While we should expect this issue to have a larger effect on orthologs, we still filtered both groups, orthologs and paralogs, to contain proteins less than 70% identical. This filter had very little effect (Fig. 2, Fig. S6).

Lateral gene transfer events might be a problem with orthology predictions. However, proper genome-wide identification of lateral gene transfer events is difficult, as xenologs are hard to distinguish from duplications events (Roth et al., 2008). Additionally, there is no good agreement between the output of different xenolog prediction methods benchmarked against real data (Ravenhall et al., 2015). In an attempt to deal with xenologs we used two approaches: We removed genes with atypical codon usage bias (see below), besides including an orthology working definition (OMA), that attempts to deal with

Figure 1. Non-synonymous to synonymous substitutions (dN/dS). The dN/dS ratios correspond to genes compared between query organisms against genomes from organisms in the same taxonomic phylum, namely: *E. coli* against other Proteobacteria, *B. subtilis* against other Firmicutes, and *N. magadii* against other Euryarchaeota. Genome identifiers are ordered from most similar to least similar to the query genome. The dN/dS distribution is higher for paralogs, suggesting that a higher proportion of orthologs have retained their functions.

xenologs. OMA uses a verification step to help reduce the number of xenologs by using a third proteome
as *witness of non-orthology* (Roth et al., 2008).

As mentioned above, to try and avoid the effect of sequences with unusual compositions, we filtered
out sequences with extreme codon usages as measured using the Codon Adaptation Index (CAI) (Sharp
and Li, 1987). For this test, we eliminated sequences with CAI values from the top and the bottom 15
percentile of the respective genome's CAI distribution. After filtering, orthologs still exhibited dN/dS
values below those of paralogs (Fig. 2, Fig. S7).

Different models for background codon frequencies can also alter the dN/dS results (Bielawski, 2013).
Thus, we performed the same tests using the Muse and Gaut model for estimating background codon
frequencies (Muse and Gaut, 1994), as advised in (Bielawski, 2013). Again, the results showed orthologs
to have lower dN/dS ratios than paralogs (Fig. 2, Fig. S8).

Figure 2. Control experiments. Left: Values of dN/dS ratios were higher for different definitions of orthology than for their paralogs. RBH were included as reference. Right: Examples of dN/dS values obtained testing for potential biases. The Goldman and Yang model for estimating codon frequencies (Goldman and Yang, 1994), included as reference, is the default. The 80 vs 80 test used data for orthologs and paralogs filtered to contain only alignments covering at least 80% of both proteins. The maximum identity test filtered out sequences more than 70% identical. The CAI test filtered out sequences having Codon Adaptation Indexes (CAI) from the top and bottom 15 percentile of the genome's CAI distribution. We also tested the effect of the Muse and Gaut model for estimating background codon frequencies (Muse and Gaut, 1994).

Differences in dN/dS ratios were more evident for genes encoding for less divergent proteins

Orthologs will normally contain more similar proteins than paralogs. Thus, a similarity test alone would naturally make orthologs appear less divergent and, apparently, less likely to have evolved new functions. While synonymous substitutions attest for the strength of purifying / negative selection in dN/dS analyses, seemingly making these ratios independent of the similarity between proteins, we still wondered whether the data changed with protein sequence divergence.

To test whether dN/dS increased against sequence divergence, we separated orthologs and paralogs into ranges of divergence of the encoded protein's percent identity. The more similar the protein sequences, the more evident were the differences between the dN/dS of orthologs and paralogs (Fig. 3). Since protein sequence identity plays a role in most working definitions of orthology, the latter results partially explained the evident disparity in dN/dS ratios between orthologs and paralogs. However, that the ratio differences were more evident at low protein sequence divergence supports the hypothesis that paralogs might be an immediate source of functional novelty. Given that redundant duplications would be expected to eventually erode (Ochman and Davalos, 2006), early functional divergence might provide paralogs with the selective pressure to survive erosion.

Figure 3. Non-synonymous to synonymous substitutions (dN/dS) and divergence. The difference between dN/dS ratios became less apparent as protein identity decreased.

CONCLUSION

The results shown above used a measure of divergence that relates to the tendencies of sequences to diverge in amino-acid composition, against their tendencies to remain unchanged. Namely, non-synonymous to synonymous substitution rates (dN/dS). Since changes in function require changes in amino-acids, this measure might suggest which sequence datasets have higher proportions that remain functionally coherent. Such proportions would show as a tendency towards lower dN/dS values. Orthologs showed evidently lower values of dN/dS than paralogs. Thus, we concluded that orthologs, as a group, could be though as more functionally stable than paralogs. It would also be proper to stop referring to the now confirmed expectation as a conjecture. Finally, we did not expect the differences to be so evident. Thus, our results also suggest that paralogs generally tend to acquire novel functions.

AVAILABILITY OF SUPPORTING DATA

A couple of programs for obtaining orthologs and dN/dS values as tested in this manuscript are available at github: https://github.com/Computational-conSequences/SequenceTools.

COMPETING INTERESTS

The authors declare that they do not have any competing interest.

FUNDING

Funding: Work supported with a Discovery Grant to GMH from the Natural Sciences and Engineering Research Council of Canada (NSERC). This work was supported by the Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica (PAPIIT-UNAM) [IN205918 and IN202421 to JAFG]. JMER is supported by PhD fellowship 959406 from CONACyT-México.

AUTHORS' CONTRIBUTIONS

JMER contributed the description and prediction of orthologs by methods other than RBH. ME performed initial comparisons of dN/dS ratios between orthologs and paralogs. JAFG directed the work by JMER, checked and edited the manuscript. GMH designed the study, wrote the manuscript, wrote the programs available at github, and performed analyses.

ACKNOWLEDGEMENTS

We are grateful to Joe Bielawski for helpful advice. We thank The Shared Hierarchical Academic Research Computing Network (SHARCNET) for computing facilities.

SUPPLEMENTARY FILES

Supplementary tables

These tables contain Wilcoxon test results.

Supplementary figures

Figure S1. Full dN/dS results obtained with InParanoid as a working definition of orthology.

Figure S2. Full dN/dS results obtained with the Orthologous Matrix (OMA) working definition of orthology.

Figure S3. Full dN/dS results obtained with the OrthoFinder working definition of orthology.

Figure S4. Full dN/dS results obtained with the ProteinOrtho working definition of orthology.

Figure S5. Full dN/dS results obtained with alignments covering at least 80% of both proteins.

Figure S6. Full dN/dS results obtained with proteins no more that 70% identical.

Figure S7. Full dN/dS results obtained with proteins within usual codon usage.

Figure S8. Full dN/dS results using Muse and Gaut's estimate of background codon frequencies.

REFERENCES

Altenhoff, A. M. and Dessimoz, C. (2009). Phylogenetic and Functional Assessment of Orthologs Inference Projects and Methods. *PLoS Computational Biology*, 5(1):e1000262.

Altenhoff, A. M., Studer, R. A., Robinson-Rechavi, M., and Dessimoz, C. (2012). Resolving the Ortholog Conjecture: Orthologs Tend to Be Weakly, but Significantly, More Similar in Function than Paralogs. *PLoS Computational Biology*, 8(5):e1002514.

Angelis, K., Reis, M. d., and Yang, Z. (2014). Bayesian Estimation of Nonsynonymous/Synonymous Rate Ratios for Pairwise Sequence Comparisons. *Molecular Biology and Evolution*, 31(7):1902–1913.

Anisimova, M., Bielawski, J. P., and Yang, Z. (2002). Accuracy and Power of Bayes Prediction of Amino Acid Sites Under Positive Selection. *Molecular Biology and Evolution*, 19(6):950–958.

Bielawski, J. P. (2013). Detecting the Signatures of Adaptive Evolution in Protein-Coding Genes. *Current Protocols in Molecular Biology*, 101(1):19.1.1–19.1.21.

Bork, P., Dandekar, T., Diaz-Lazcoz, Y., Eisenhaber, F., Huynen, M., and Yuan, Y. (1998). Predicting function: from genes to genomes and back11Edited by P. E. Wright. *Journal of Molecular Biology*, 283(4):707–725.

Buchfink, B., Xie, C., and Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, 12(1):59–60.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, 10(1):421.

David, K. T., Oaks, J. R., and Halanych, K. M. (2020). Patterns of gene evolution following duplications and speciations in vertebrates. *PeerJ*, 8:e8813.

Emms, D. M. and Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biology*, 16(1):157.

Escorcia-Rodríguez, J. M., Esposito, M., Freyre-González, J. A., and Moreno-Hagelsieb, G. (2020). Non-synonymous to synonymous substitutions suggest that orthologs tend to keep their functions, while paralogs are a source of functional novelty. *bioRxiv*.

Fitch, W. M. (2000). Homology a personal view on some of the problems. *Trends in genetics : TIG*, 16(5):227 – 231.

Gabaldón, T. and Koonin, E. V. (2013). Functional and evolutionary implications of gene orthology. *Nature Reviews Genetics*, 14(5):360–366.

Galperin, M. Y., Kristensen, D. M., Makarova, K. S., Wolf, Y. I., and Koonin, E. V. (2019). Microbial genome analysis: the COG approach. *Briefings in Bioinformatics*, 20(4):1063–1070.

Goldman, N. and Yang, Z. (1994). A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Molecular Biology and Evolution*, 11(5):725–36.

Haft, D. H., DiCuccio, M., Badretdin, A., Brover, V., Chetvernin, V., O'Neill, K., Li, W., Chitsaz, F., Derbyshire, M. K., Gonzales, N. R., Gwadz, M., Lu, F., Marchler, G. H., Song, J. S., Thanki, N., Yamashita, R. A., Zheng, C., Thibaud-Nissen, F., Geer, L. Y., Marchler-Bauer, A., and Pruitt, K. D. (2018). RefSeq: an update on prokaryotic genome annotation and curation. *Nucleic Acids Research*, 46(D1):gkx1068–.

Hernández-Salmerón, J. E. and Moreno-Hagelsieb, G. (2020). Progress in quickly finding orthologs as reciprocal best hits: comparing blast, last, diamond and MMseqs2. *BMC Genomics*, 21(1):741.

Huynen, M. A. and Bork, P. (1998). Measuring genome evolution. *Proceedings of the National Academy of Sciences of the United States of America*, 95(11):5849 – 5856.

Kryuchkova-Mostacci, N. and Robinson-Rechavi, M. (2016). Tissue-Specificity of Gene Expression Diverges Slowly between Orthologs, and Rapidly between Paralogs. *PLOS Computational Biology*, 12(12):e1005274.

Lechner, M., Findeiß, S., Steiner, L., Marz, M., Stadler, P. F., and Prohaska, S. J. (2011). Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics*, 12(1):124.

Moreno-Hagelsieb, G. and Latimer, K. (2008). Choosing BLAST options for better detection of orthologs as reciprocal best hits. *Bioinformatics*, 24(3):319–324.

Muse, S. V. and Gaut, B. S. (1994). A likelihood approach for comparing synonymous and nonsynonymous nucleotide substitution rates, with application to the chloroplast genome. *Molecular Biology and Evolution*, 11(5):715–724.

Mushegian, A. R. and Koonin, E. V. (1996). Gene order is not conserved in bacterial evolution. *Trends in genetics : TIG*, 12(8):289 – 290.

Nehrt, N. L., Clark, W. T., Radivojac, P., and Hahn, M. W. (2011). Testing the Ortholog Conjecture with Comparative Functional Genomic Data from Mammals. *PLoS Computational Biology*, 7(6):e1002073.

Ochman, H. and Davalos, L. M. (2006). The Nature and Dynamics of Bacterial Genomes. *Science*, 311(5768):1730–1733.

Ohno, S. (1970). *Evolution by gene duplication*. Springer-Verlag.

Ohta, T. (1995). Synonymous and nonsynonymous substitutions in mammalian genes and the nearly

neutral theory. *Journal of molecular evolution*, 40(1):56 – 63.

R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria.

Ravenhall, M., Škunca, N., Lassalle, F., and Dessimoz, C. (2015). Inferring Horizontal Gene Transfer.
PLOS Computational Biology, 11(5):e1004095.

Rice, P., Longden, I., and Bleasby, A. (2000). EMBOSS: the European Molecular Biology Open Software
Suite. *Trends in genetics : TIG*, 16(6):276 – 277.

Roth, A. C., Gonnet, G. H., and Dessimoz, C. (2008). Algorithm of OMA for large-scale orthology
inference. *BMC Bioinformatics*, 9(1):518.

Sharp, P. M. and Li, W. H. (1987). The codon Adaptation Index–a measure of directional synonymous
codon usage bias, and its potential applications. *Nucleic acids research*, 15(3):1281 – 1295.

Sonnhammer, E. L. and Östlund, G. (2015). InParanoid 8: orthology analysis between 273 proteomes,
mostly eukaryotic. *Nucleic Acids Research*, 43(D1):D234–D239.

Tatusov, R. L., Galperin, M. Y., Natale, D. A., and Koonin, E. V. (2000). The COG database: a tool for
genome-scale analysis of protein functions and evolution. *Nucleic Acids Research*, 28(1):33–36.

Thomas, P. D., Wood, V., Mungall, C. J., Lewis, S. E., Blake, J. A., and Consortium, G. O. (2012). On the
Use of Gene Ontology Annotations to Assess Functional Similarity among Orthologs and Paralogs: A
Short Report. *PLoS Computational Biology*, 8(2):e1002386.

Train, C.-M., Glover, N. M., Gonnet, G. H., Altenhoff, A. M., and Dessimoz, C. (2017). Orthologous Ma-
trix (OMA) algorithm 2.0: more robust to asymmetric evolutionary rates and more scalable hierarchical
orthologous group inference. *Bioinformatics*, 33(14):i75–i82.

Ward, N. and Moreno-Hagelsieb, G. (2014). Quickly Finding Orthologs as Reciprocal Best Hits with
BLAT, LAST, and UBLAST: How Much Do We Miss? *PLoS ONE*, 9(7):e101850.

Wolf, Y. I. and Koonin, E. V. (2012). A Tight Link between Orthologs and Bidirectional Best Hits in
Bacterial and Archaeal Genomes. *Genome Biology and Evolution*, 4(12):1286–1294.

Yang, Z. (2007). PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology and
Evolution*, 24(8):1586–1591.

Yang, Z. and Nielsen, R. (2000). Estimating synonymous and nonsynonymous substitution rates under
realistic evolutionary models. *Molecular Biology and Evolution*, 17(1):32 – 43.

Yutin, N., Puigbò, P., Koonin, E. V., and Wolf, Y. I. (2012). Phylogenomics of Prokaryotic Ribosomal
Proteins. *PLoS ONE*, 7(5):e36972.

Zahn-Zabal, M., Dessimoz, C., and Glover, N. M. (2020). Identifying orthologs with OMA: A primer.
F1000Res, 9:27.