Drosophila *small ovary* encodes a zinc-finger repressor required for ovarian differentiation

Leif Benner[1,2], Elias A. Castro[3], Cale Whitworth[1,4], Koen J.T. Venken[5], Haiwang Yang[1], Brian Oliver[1,6,7], Kevin R. Cook[4], and Dorothy A. Lerit[3,6]

Author affiliations:

[1]Section of Developmental Genomics, Laboratory of Cellular and Developmental Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda MD 20892. [2]Department of Biology, Johns Hopkins University, Baltimore MD 21218. [3]Department of Cell Biology, Emory University School of Medicine, Atlanta GA 30322. [4]Department of Biology, Indiana University, Bloomington, Indiana 47405. [5]Verna and Marrs McLean Department of Biochemistry and Molecular Biology, McNair Medical Institute at the Robert and Janice McNair Foundation, Dan L. Duncan Cancer Center, Center for Drug Discovery, Department of Pharmacology and Chemical Biology, Baylor College of Medicine, Houston, TX, 77030, USA

[6]**Corresponding authors:** dlerit@emory.edu briano@nih.gov

[7]**Lead contact**: briano@nih.gov

Keywords: chromatin, differentiation, gene expression, HP1, oogenesis, position-effect variegation

Abstract

Repression is essential for coordinated cell type-specific gene regulation and for controlling the expression of transposons. In the *Drosophila* ovary, stem cells regeneration and differentiation require well-controlled expression, with de-repression leading to tissue degeneration and ovarian tumors. Additionally, the ovary is acutely sensitive to deleterious consequences of transposon de-repression. The *small ovary* (*sov*) locus was identified in a female sterile screen shows dramatic effects ovarian morphogenesis. We mapped the locus to the uncharacterized gene *CG14438* and reveal it encodes a zinc-finger protein that colocalizes with the essential Heterochromatin Protein 1 (HP1a). The distinct functions of Sov we report, including repression of inappropriate signaling, transposon silencing, and effects on position-effect variegation in the eye, suggest a central role in heterochromatin stabilization.

Introduction

Multicellular organisms rely on stem cells. A combination of extrinsic and intrinsic signals function to maintain the balance between self-renewal of stem cells and the differentiation of progenitors needed for tissue homeostasis and development. The *Drosophila* adult ovary is a well-studied model for development (Fuller and Spradling 2007). It is organized into ~16 ovarioles, which are assembly lines of progressively maturing egg chambers. At the anterior tip of each ovariole is a structure called the germarium, harboring 2-3 germline stem cells (GSCs), which sustain egg development throughout the life of the animal. Surrounding the GSCs and their cystoblast daughters is a stem cell niche, a collection of nonproliferating somatic cells composed of terminal filament and cap cells. Like the GSCs, somatic stem cells divide to produce new stem cells and escort cells, which encase the germline. Each cystoblast is fated to differentiate and will undergo four rounds of cell division with incomplete cytokinesis to

Benner et al.; *sov* represses gene expression

produce a germline cyst interconnected by the fusome, a branched cytoskeletal structure.

Germline cysts are surrounded by two escort cells. Toward the posterior of the germarium, a

monolayer of somatic follicle cells replaces the escort cells to form an egg chamber.

Egg chamber development requires the careful coordination of distinct germline and

somatic cell populations through signaling within a complex microenvironment. For example,

extrinsic signals and transducers including Jak/Stat and the BMP homolog, Decapentapalegic

(Dpp), are required to maintain the proliferative potential of GSCs (Bausek 2013; Gilboa 2015;

S. Chen, Wang, and Xie 2011). Oriented divisions displace the primary cystoblast cell away

from the stem cell preserving signals and permit the expression of differentiation factors, such

as *bag-of-marbles* (*bam*). The importance of bam is revealed in bam loss-of-function mutants,

resulting in tumorous germaria filled with undifferentiated germline cells containing many

spectrosomes, the dot-fusome structure characteristic of GSCs and primary cystoblasts, as a

result of complete cytokinesis (Lin and Spradling 1995; D. McKearin and Ohlstein 1995).

Controlling this signaling requires repression in addition to activation.

Dynamic changes in chromatin landscapes within the germline and somatic cells are

critical for oogenesis (Barton et al. 2016; X. Li et al. 2017; Börner et al. 2016; Peng et al. 2016;

Soshnev et al. 2013; McConnell, Dixon, and Calvi 2012). Nucleosomes generally repress

transcription by competing for DNA binding with transcription factors (Lorch and Kornberg

2017; Kouzarides 2007; Jenuwein and Allis 2001). Further levels of repression depend largely

on histone modifications. For example, the activities of both the H3K9 methyltransferase

SETDB1, encoded by *eggless* (*egg*) (X. Wang et al. 2011; Clough et al. 2007; Clough,

Tedeschi, and Hazelrigg 2014), and the H3K4 demethylase encoded by *lysine-specific*

demethylase 1 (Di Stefano et al. 2007; Rudolph et al. 2007; Eliazer, Shalaby, and Buszczak

Benner et al.; *sov* represses gene expression

2011; Eliazer et al. 2014) are required in the somatic cells of the germarium for GSC

maintenance, normal patterns of differentiation, and germline development. Genome-wide

profiling hints at a progression from open chromatin in germline stem cells to a more closed

state during egg chamber differentiation (T. Chen and Dent 2014). Disrupting this progressive

repression in the ovary could result in stem cell over-proliferation and defective development.

There are other roles for regulated chromatin states in oogenesis. The germline has

been repeatedly hijacked to promote the vertical transmission of transposons resulting in a

loss of fitness in the *Drosophila* host (Charlesworth and Langley 1989; Pelisson et al. 2002).

This increased mobilization of transposons in gonads is countered by host responses, such as

induced heterochromatin formation, to repress transposon activity. The proximity of condensed

heterochromatin can dampen the activity of genes and transgenes (Elgin and Reuter 2013).

The conserved Heterochromatin-Protein 1a (HP1a), encoded by *Su(var)205*, is critical for

heterochromatin formation (James and Elgin 1986; Eissenberg et al. 1990; Clark and Elgin

1992).

Originally described over 40 years ago (J. D. Mohler 1977), the *small ovary* (*sov*) locus

mutants have range of phenotypes including disorganized ovariole structure, egg chamber

fusions, and undifferentiated tumors, culminating in complete ovarian degeneration (Wayne et

al. 1995). In the present work, we demonstrate *sov* encodes an unusually long 21 C_2H_2 Zinc

finger (ZnFs) nuclear protein that colocalizes with HP1. Previous work showed this ZnF protein

in complex with HP1a (Alekseyenko et al. 2014). RNA-seq analysis shows that *sov* activity is

required to repress the expression of a large number of genes in the ovary as well as

transposons. Additionally, *sov* mutations are strong dominant suppressors of Position Effect

Varigation (PEV) in the eye, a key characteristic of HP1 (Elgin and Reuter 2013), which is

Benner et al.*; sov* represses gene expression

consistent with Sov and HP1a colocalization in the nucleus. These data indicate that Sov is a

potent repressor of gene expression.

Results

small ovary is CG14438

To localize *sov*, we used complementation of female-sterile and lethal *sov* mutations with

preexisting and custom-generated deficiencies, duplications, and transposon insertions to

refine previous mapping (J. D. Mohler 1977; D. Mohler and Carroll 1984). Our results (Fig

1A,B) suggested that either the protein-coding *CG14438* gene or the intronic, noncoding

CR43496 gene was *sov*. The *sov* locus mapped to the non-complementing deletion *Df(1)sov*.

Rescue of female sterility and/or lethality of *sov^2*, *sov^{ML150}*, *sov^{EA42}* and *Df(1)sov* with

Dp(1;3)sn^{13a1} (molecularly undefined, not shown), *Dp(1;3)DC486*, and *PBac{GFP-sov}*

confirmed our mapping. We could not replicate separability of *sov^{EA42}* sterility and lethality with

Dp(1;3)sn^{13a1} (Wayne et al. 1995). Female sterile and lethal alleles of *sov* map identically.

We performed genome sequencing to determine if the lesions in *sov* alleles occur in

CG14439 or *CR43496* (Table S1). While *CR43496* contained three polymorphisms in *sov*

alleles relative to the reference, we identified the same polymorphisms in all mutant *sov* alleles

and in *sov$^+$* alleles in *w^{1118}* and *Oregon-R* lines. Thus, *CR43496* is highly unlikely to be *sov.* In

contrast, we found disruptive mutations in *CG14438* (Fig 1C), which encodes an unusually

long 3,313 residue protein with 21 C_2H_2 ZnFs, nuclear localization motifs, and coiled-coil

regions (Fig 1C). We identified a nonsense mutation (G to A at position 6,764,462) in the

CG14438 open reading frame of *sov^{EA42}* which is predicted to truncate the *sov* protein before

the ZnF domains. We found a frameshift insertion (T at position 6,769,742) located towards the

Benner et al.; *sov* represses gene expression

end of *CG14438* in *sov²* that encodes 30 novel residues followed by a stop codon within the C-

terminal ZnF, as well as removing the terminal predicted NLS motif (Fig 1C). We found a

missense mutation (C to G at position 6,763,888) in *sov^ML150^* that results in a glutamine to

glutamate substitution within a predicted coiled-coil domain. While this is a conservative

substitution, and while glutamate residues are common in coiled-coil domains, glutamine to

glutamic acid substitutions are especially disruptive in the coiled-coil region of the sigma

transcription factor (Hsieh, Tintut, and Gralla 1994). We conclude that *CG14438* is *sov*.

sov expression

To determine where *sov* is expressed and if it encodes multiple isoforms, we analyzed

expression in adult tissues by RNA-Seq. We noted that *sov* is broadly expressed as a single

mRNA isoform in adult tissues, with highest expression in ovaries (Fig 1D). The modENCODE

(Graveley et al. 2011; J. B. Brown et al. 2014) and FlyAtlas (Robinson et al. 2013; Leader et al.

2018) reference sets show a similar enrichment in the ovary, as well as in early embryos. The

enrichment of *sov* in the ovary is consistent with its role in oogenesis.

To determine if and where Sov was expressed in the ovary, we examined the

distribution of GFP-Sov in developing egg chambers (Fig 2A). We observed GFP-Sov signal in

all cell types. By pairing localization analysis of Sov with antibodies recognizing Vasa (Vas) to

label the germline (Fig 2B), we observed particularly striking nuclear localization of Sov

surrounded by perinuclear Vas in the germline cells within region 1 of the germarium, with

highest levels evident within the GSCs (dashed circles). We also used Traffic jam (Tj) to label

the soma (Fig 2C). We observed Sov nuclear staining in the Tj positive somatic and follicle

Benner et al.; *sov* represses gene expression

stem cells (dashed ovals) (Fig 2C). These data indicate that Sov is expressed in the germline

and soma in the ovary.

Sov function

To examine *sov* function in the germarium, we compared a strong allelic combination,

sov^2/sov^{EA42} to cell-type specific knockdown of *sov* using a UAS short hairpin *sov* RNAi

construct (P{TRiP.HMC04875}; hereafter, sov^{RNAi}). The germaria of heterozygous *sov* females

are wild type (Fig 3A), but the *sov* mutants often show an ovarian tumor phenotype, with

greater than the expected number of germ cells with dot spectrosomes (Fig 3B), indicating that

germ cells undergo complete cytokinesis. Ovarian tumors were observed in ~40% of *sov*

mutants (N=12/31 sov^{EA42}/sov^2 egg chambers versus N=0/47 controls). These findings are

consistent with GSC hyperproliferation and/or failed differentiation. We also observed tumors

and nurse cell nuclei residing within common follicles (Fig 3B) in about one-third of *sov*

mutants (N=11/31 sov^{EA42}/sov^2 versus N=0/24 in controls). This phenotype occurs when follicle

cells either fail to separate egg chambers, or where those chambers fuse (Goode, Wright, and

Mahowald 1992). In older *sov* ovaries, we observed extensive cell death (Fig S1). Additionally,

we found that rare sov^{EA42}/Y males that escaped lethality had no germ cells (Fig S2). Thus, *sov*

functions widely.

To examine cell-type specific functions of *sov*, we used *tj-GAL4* to express sov^{RNAi} in

somatic escort and follicle cells and *nanos-GAL4* (*nos-GAL4*) to express sov^{RNAi} (or

$mCherry^{RNAi}$ controls) in the germline. Relative to the controls, the $tj>sov^{RNAi}$ germaria (Fig

3C,D) were abnormal, showing ovarian tumor phenotypes similar to those seen in *sov*

mutants. Germariums were often filled with germline cells with dot spectrosomes and the

follicle cells encroached anteriorly. Similar results were also observed using other somatic

drivers (*c587-GAL4* and *da-GAL4*). Interestingly, *nos>sov^{RNAi}* did not impaired germaria

morphology (Fig 3E,F) and egg chambers representing all 14 morphological stages of

oogenesis appeared phenotypically normal. However, eggs produced from *nos>sov^{RNAi}*

mothers arrested during early embryogenesis, indicating that maternal *sov* is required for

embryonic development. While the expression patterns and germline RNAi suggest that *sov* is

deployed in both the soma and germline, Wayne et al. (1995) reported *sov* to be somatic line

dependent. We conducted a germline clonal analysis and found instead that *sov^+* is required in

the germline (Fig S3). Additionally, *sov^{RNAi}* resulted in embryonic germline defects in a high-

throughput study (Jankovics et al. 2014). Taken together, these data indicate that there are

both somatic and germline requirements for *sov* in oogenesis.

To explore the differentiation of the germline further, we examined the distribution of

Bam, which is expressed in germarium region 1 germ cells (D. M. McKearin and Spradling

1990; D. McKearin and Ohlstein 1995; Ohlstein and McKearin 1997). In the absence of Bam,

germ cells hyper-proliferate, resulting in tumors composed of 2-cell cysts. Given the

prevalence of undifferentiated tumorigenic germ cells upon somatic *sov^{RNAi}* knockdown, we

asked if *sov* has a non-autonomous role in promoting Bam expression in germ cells. Indeed,

somatic knockdown of *sov* results in significantly less Bam expressed in germ cells relative to

controls (Fig 3G,H; Fig S4). Both *tj-GAL4>sov^{RNAi}* and *c587-GAL4>sov^{RNAi}* resulted in strong

repression of Bam expression. Germline depletion of *sov* did not alter Bam localization (Fig

S4). The reduction in Bam expression in the germline is consistent with a non-autonomous role

of Sov function in the soma for differentiation signals directed to the germline.

Benner et al.; *sov* represses gene expression

To examine the defective follicle encapsulation of the germline cysts following *sov*

depletion (resulting in fused follicles), we used the oocyte-specific expression of Oo18 RNA

binding protein (Orb) (Lantz et al. 1994) to count the number of oocytes per cyst. Orb specifies

the future single oocyte at the posterior of the 16-germ cell egg chamber in control egg

chambers (Fig 3I, Fig S2D). When we examined ovaries where *sov* expression had been

knocked down in somatic cells, we found examples of both egg chambers with either too many

oocytes (Fig 3J) or no oocytes (Fig 3K, Fig S4). In a wild type 16-cell germline cyst, one of the

two cells that has four ring canals becomes the oocyte, and this feature can be used to

determine if extra germline divisions had occurred in cysts, or if multiple cysts were enveloped

by the follicle cells. We saw that the egg chambers with multiple Orb+ cells always had more

than 16 germ cells, and in the example shown, all three Orb+ cells had four ring canals,

indicating that egg chamber fusion had occurred (Fig 3J). Taken together, we conclude that

Sov functions in the soma to ensure the proper differentiation of the germline.

Sov represses gene expression in the ovary

To provide mechanistic insights into the function of Sov, we performed transcriptome profiling

using triplicated Poly-A$^+$ RNA-seq analyses of ovaries from *sov*, *sov*^{RNAi}, and control females.

The gene expression profiles of ovaries from sterile females were markedly different from

controls, primarily due to de-repression in the mutants (Fig 4A; Table S2). Among genes

showing differences in expression (FDR *p*adj < 0.05), we found 1,752 genes with >4-fold

increased expression in mutants, while there were only 172 genes with >4-fold decreased

expression (Table S3). To explore where these de-repressed genes are normally expressed,

we examined their expression in other female tissues and in testes in a set of quadruplicated

Benner et al.*; sov* represses gene expression

RNA-seq experiments (a lab resource for this and other studies; Fig 4B). Many of the genes

suppressed by Sov in ovaries were highly expressed in other female tissues, or in the testis,

indicating that wildtype Sov prevents ectopic gene expression.

To determine what types of genes are de-repressed in *sov* mutants, we performed

Gene Ontology (GO)(Gene Ontology Consortium 2015) term analysis (Fig S5; Table S4). The

genes with lower expression in *sov* mutants had only a few significant GO terms that were

oogenic in nature (as expected given the general lack of mature eggs in *sov* mutants). For

example, there was poor expression of the chorion genes that are required to build the

eggshell (Orr-Weaver 1991), as well as genes required for follicle cell development. In

contrast, the de-repressed genes in *sov* mutants showed that there was a significant

enrichment of genes with cell signaling GO terms, including neuronal communication. We

found that many genes repressed by Sov in the ovary are indeed expressed in the head. For

example, the *heartless* (*htl*) locus, which encodes a FGFR tyrosine kinase receptor important

for neuron/glia communication (Stork et al. 2014), was de-repressed in *sov* ovaries and highly

expressed in the head (Fig 4C). This intriguing result suggests that *sov* normally functions to

repress a host of signaling pathways. This de-repressed signaling in *sov* mutants is likely

catastrophic for communication between various somatic cells and the germline during egg

chamber development. This could well explain the variety of mutant phenotypes observed in

sov mutants.

The general repressive role of *sov* was also revealed by a dramatic and coherent

elevation of transposon expression in the mutants (Fig 4D; Table S3). 138 transposable

element classes were detected in our gene expression profiling of *sov* mutants and controls.

Of these, 91 had increased expression (FDR *p*adj < 0.05) >4-fold in *sov* mutants, while none

Benner et al.*; sov* represses gene expression

had >4-fold decreased expression. The DNA, LINE, and LTR families of transposable

elements were all de-repressed in *sov* mutants. Interestingly, the somatic knockdown of *sov*

resulted in the most dramatic de-repression of the *gypsy* and *copia* classes of transposons,

which are *Drosophila* retroviruses that develop in somatic cells and are exported to the

developing germline (Yoshioka et al. 1990). Thus, wild type *sov* may be important to protect

the germline from infection. Equally interesting, germline knockdown resulted in greatest de-

repression of the *HeT-A*, *TAHRE*, and *TART* transposons that compose the telomere (Mason,

Frydrychova, and Biessmann 2008). Poorly formed telomeres result in loss of material from

chromosome ends and telomere fusion, which results in poor anaphase separation and

aneuploidy (Cenci, Ciapponi, and Gatti 2005). This finding suggests that the general role of

sov in silencing transposon expression includes control of both transposon functions

necessary for normal cellular metabolism, exemplified by the telomere transposons, as well as

detrimental activities of retrovirus-like elements. In addition to widespread dis-regulation of

signaling and development pathways in *sov* mutants, transposon dysgenesis may contribute to

the severe defects in *sov* ovarian development.

Sov is a Suppressor of PEV and co-localizes with HP1

The repressive function of Sov is reminiscent of HP1a function as general repressor of gene

expression. HP1a was first characterized in *Drosophila* as a suppressor of PEV, a process that

reduces gene expression due to spreading of heterochromatin into a gene region (Clark and

Elgin 1992). To test the hypothesis that *sov* negatively regulates gene expression by

promoting heterochromatin formation, we examined the role of *sov* in PEV in the eye, where

patches of white[+] (red pigmented) and white[-] (not red pigmented) eye facets are easily

Benner et al.*; sov* represses gene expression

observed (Fig 5A). If, like HP1a, Sov represses gene expression by promoting

heterochromatin formation (Eissenberg et al. 1990), then it should suppress PEV, which would

be seen as increased eye pigmentation. We obtained five different variegating w^+ transgene

insertions associated with either the heterochromatic pericentric region of chromosome arm 2L

or spread along the length of the heterochromatin-rich chromosome 4. In control animals,

these insertions show a characteristic eye variegation pattern (Fig 5B). Consistent with the

allelic strengths seen in previous experiments, the weak sov^2 allele did not suppress PEV (Fig

5C) but the stronger sov^{ML150}, sov^{EA42}, and $Df(1)sov$ mutations dominantly suppressed PEV

(Fig 5D–F). These data demonstrate a role for Sov in heterochromatin formation.

The strong repressive function of Sov is reminiscent of HP1. If Sov is in complex with

HP1a (Alekseyenko et al. 2014), then Sov and HP1 should colocalize in cells. To see if this

was true, we followed HP1a-RFP and GFP-Sov localization in 1–2 hr live embryos where both

are abundantly expressed (Rudolph et al. 2007). During *Drosophila* cleavage, nuclei undergo

rapid synchronous nuclear divisions prior to cellularization at cleavage division/nuclear cycle

(NC) 14 (Foe and Alberts 1983). HP1a and Sov colocalized in all NC 10–12 embryos we

examined (Fig 6A; N=7 embryos). During prophase, both HP1a and Sov were enriched in

regions of condensed DNA (Fig 6A, 4:30). Whereas low levels of HP1a decorated DNA

throughout division, Sov was depleted during mitosis (Fig 6A, 10:00 and 11:10). Upon re-entry

into interphase, Sov localization to nuclei resumed and was coincident with HP1a. Formation

of heterochromatin is contemporaneous with or slightly proceeds HP1a apical subnuclear

localization in NC 14 (Rudolph et al. 2007; Yuan and O'Farrell 2016). At this stage, we

observed a strong colocalization of HP1a and Sov. Measuring the distribution of HP1a and Sov

(Fig 6B,B') confirmed high levels of colocalization within HP1a subnuclear domains (Fig 6C,

shaded region). These data support the idea that HP1a and Sov are deposited maternally

where they assemble into a complex.

Discussion

Sov is a novel heterochromatin-associated protein

Genomes contain large blocks of DNA of potentially mobile transposons and immobile mutated

derivatives (Vermaak and Malik 2009). The cell keeps these transposons from wreaking havoc

on the genome by actively suppressing their expression through condensation into

heterochromatin. However, some tightly regulated expression from heterochromatin is required

for normal cellular function. For example, the telomeres of *Drosophila* are maintained by

transposition of mobile elements from heterochromatic sites (Mason, Frydrychova, and

Biessmann 2008), and histone and rRNA genes are located within heterochromatic regions

(Yasuhara and Wakimoto 2006). Weakening of heterochromatin by suppressors of variegation

results in de-repression of gene expression at the edges of heterochromatin blocks,

suggesting that the boundaries between repressed and active chromatin can expand and

contract (Weiler and Wakimoto 1995; Reuter and Spierer 1992). HP1a, encoded by

Su(var)205, is a central component of heterochromatin (Ebert et al. 2006) that shows the same

strong suppression of variegation that we observed in *sov/+* flies. Similarly, HP1a is also

required to repress the expression of transposons (Vermaak and Malik 2009).

Repression is often stable and emerging themes suggest a robust set of activities,

rather than a single component, maintain a chromatin state. For example, long-term repression

is stabilized with complexes, such as the repressive Polycomb Group (PcG) which provides an

epigenetic memory function (Kassis, Kennison, and Tamkun 2017). In order to create a stable

Benner et al.*; sov* represses gene expression

epigenetic state, the PcG complexes have subunits that modify histones and subunits that bind

these modifications. In addition to being tethered to the chromatin via histones, the PcG

complexes also include DNA-binding proteins, such as the YY1-like ZnF protein Pleiohomeotic

(Pho) that further reinforce localization. The DNA anchor proteins in PcG complexes are at

least partially redundant with the histone-binding components (J. L. Brown et al. 2003),

suggesting that localization is robust due to multiple independent localization mechanisms.

In mammals, HP1a also has both histone and DNA anchoring. DNA anchors are

provided by a family of KRAB ZnF proteins that have undergone a massive radiation during

evolution (Yang, Wang, and Macfarlan 2017; Ecco, Imbeault, and Trono 2017). KRAB proteins

are essential for repression of transposons and more generally for a properly regulated

genome. The KRAB family members use a KRAB-Associated Protein (KAP1) adapter protein

to associate with HP1a and the SETDB1 methylase that modifies histones to enable HP1a

binding (Fig 7). Although KRAB proteins have not been identified in *Drosophila*, the *Drosophila*

bonus (*bon*) gene encodes a KAP1 homolog (Beckstead et al. 2005). *Drosophila* also has a

SETDB1 encoded by *egg*, and loss of *egg* results in ovarian phenotypes reminiscent of those

observed in *sov* loss-of-function females (Clough et al. 2007; Clough, Tedeschi, and Hazelrigg

2014; X. Wang et al. 2011). Like *sov* and HP1a, *bon* is a modifier of variegation, raising the

possibility that they collectively coordinate gene regulation. In support of this hypothesis, Bon,

Sov, and Egg were all identified in the same biochemical complex as HP1a (Alekseyenko et al.

2014). It is tempting to speculate that the single very long and ZnF-rich Sov protein plays the

same direct tethering role as the large family of KRAB ZnF proteins in mammals (Fig 7). If Sov

uses subsets of fingers to bind DNA, it could localize to many different sequences in a

combinatorial fashion. Additionally, the complex containing Sov and HP1a also contains RNA

Benner et al.*; sov* represses gene expression

(Alekseyenko et al. 2014) and could act as a tether via a series of RNA intermediates as

occurs in sex chromosome inactivation, for example (J. T. Lee 2009). As in the case of Pho in

the PcG complexes, Sov might be a robustness factor rather than an absolute requirement, as

we did not observe gross delocalization of HP1a following sov^{RNAi} (not shown). Further work

will be required to fully understand the relationship between Sov and HP1a.

<u>Repression by Sov promotes germline differentiation</u>

The fact that there have been many female sterile alleles of *sov* isolated suggests that

the ovary is particularly sensitive to reduced *sov* activity. The female sterility phenotype is

complex and somewhat variable, but partial *sov* loss of function is characterized by somatic-

dependent differentiation defects. Phenotypes include defective follicle cell encasement of egg

chambers, resulting in follicles with multiple oocytes, and accumulation of germline tumor cells

that undergo complete cytokinesis instead of forming mature and differentiated 16-cell egg

chambers. Stronger alleles also show a germline-dependent block in development and

lethality. Diverse functions sometimes result from isoform and/or localization diversity. While

sov functions seem diverse, the locus expresses a single major isoform and Sov seems to be

consistently located in the nucleus. It is more likely that one mechanism account for all *sov*

function. We suggest that Sov in regulates the chromatin landscape that helps repress stem

cell identity and promote differentiation. This same mechanism represses ectopic signaling and

tissue-inappropriate responses in the ovary, controls heterochromatin formation in the eye and

represses transposons.

We observed dramatic de-repression of transposons in *sov* mutants. In addition to our

observations, Czech et al (2013) found evidence for a role of *sov* in transposon repression in a

Benner et al.*; sov* represses gene expression

genome-wide screen. Transposon repression raises the possibility that Sov interacts with the

Piwi pathway (Brennecke et al. 2007; Yin and Lin 2007; Teixeira et al. 2017) which directly

interacts with HP1a to facilitate heterochromatin formation and promote transposon silencing

(Brower-Toland et al. 2007; S. H. Wang and Elgin 2011). However, genes involved more

strictly with transposable element regulation, such as Piwi, do not have strong dosage effects

on PEV (Gu and Elgin 2013), while *sov* and *HP1a*/*Su(var)205* do. This distinction suggests

that Sov has a more general effect on heterochromatin, rather than specificity for transposon

repression. Nevertheless, interplay between Piwi, Sov, and HP1a remains possible.

Conclusions

Our data show that *sov* encodes a repressor of gene expression and transposons in the

ovary. In the absence of *sov*, genes that are normally expressed at very low levels in ovary,

are activated. These same genes show dynamic patterns of expression in other adult tissues,

suggesting that they are de-repressed in the absence of Sov. The failure to restrict gene

expression results in a range of phenotypes including ovarian tumors, defective oogenesis,

and tissue degeneration. These data support the idea that Sov encodes a novel and essential

protein that is generally repressive, possibly via interactions with HP1a.

Materials and Methods

We have adopted the FlyBase-recommended resources table which includes all genetic,

biological, cell biology, genomics, manufactured reagents and algorithmic resources used in

this study (Table S5).

Benner et al.*; sov* represses gene expression

<u>Flies and genetics</u>

The *small ovaries* (*sov*) locus was defined by three non-complementing female sterile

mutations including *sov²*, which mapped to ~19 cM on the X chromosome in the 5D5-6;6E4-5

interval (D. Mohler and Carroll 1984; J. D. Mohler 1977), refined to 6BD (Wayne et al. 1995).

Cook *et al.* (2012) placed *sov* in the 4-gene *CG14438–shf* interval (Fig. 1; Table S5). *sov*

mutations complemented *shf²*, but not *P{SUPor-P}sov^KG00226* and *P{GawB}sov^NP6070*,

suggesting that *CG14438* or *CR43496* were *sov*, which we confirmed by generating *Df(1)sov*

(X:6756569..6756668; 6770708, 6C12;6C13) as a FLP-induced recombination event between

P{XP}sov^d07849 and *PBac{RB}e03842* (Parks et al. 2004; Cook et al. 2012) to remove only

CG14438 and *CR43496.*

We used the dominant female sterile technique for germline clones (Chou and Perrimon

1996). Test chromosomes were free of linked lethal mutations by male viability (*sov²*) or rescue

by *Dp(1;3)DC486* (*sov^EA42* and *Df(1)sov*), were recombined with *P{ry^+t7.2=neoFRT}19A*, and

verified by complementation tests and PCR. We confirmed *P{ry^+t7.2=neoFRT}19A* functionality

by crossing to *P{w^+mC=GMR-hid}SS1, y¹ w* P{ry^+t7.2=neoFRT}19A; P{w^+m*=GAL4-ey.H}SS5,*

P{w^+mC=UAS-FLP.D}JD2 and scoring for large eye size. We crossed females with FRT

chromosomes to *P{w^+mC=ovoD1-18}P4.1, P{ry^+t7.2=hsFLP}12, y¹ w^1118 sn³*

P{ry^+t7.2=neoFRT}19A/Y males for 24 hours of egg laying at 25°C. We heat-shocked for 1hr at

37°C on days 2 and 3. We dissected females (5d post-eclosion) to score for *ovo^D1* or wildtype

morphology.

We generated *PBac{GFP-sov}* from P[acman] BAC clone CH322-191E24 (X:6753282--

6773405) (Venken et al. 2009) grown in the SW102 strain (Warming et al. 2005). In step one,

we integrated the positive/negative marker CP6-RpsL/Kan (CP6 promoter with a bi-cistronic

Benner et al.; *sov* represses gene expression

cassette encoding the RpsL followed by the Kan) between the first two codons of *sov*, and

selected (15 µg/ml Kanamycin). We integrated at the galK operon in DH10B bacteria using

mini-lambda-mediated recombineering (Court et al. 2003). We amplified DH10B::CP6-

RpsL/Kan DNA using primers N-CG14438-CP6-RN-F and -R. Correct events were identified

by PCR, as well as resistance (15 µg/ml Kanamycin) and sensitivity (250 µg/ml Kan). In step

two, we replaced the selection markers with a multi-tag sequence (Venken et al. 2011),

tailored for N-terminal tagging (N-tag) and counter-selected (250 µg/ml Kanamycin). The N-tag

(3xFlag tag, TEV protease site, StrepII tag, superfolder GFP, FlAsH tetracysteine tag, and

flexible 4xGlyGlySer (GGS) linker) was *Drosophila* codon optimized in a R6Kγ plasmid. We

transformed plasmid into EPI300 for copy number amplification. We confirmed correct events

by PCR and Sanger DNA sequencing. Tagged P[acman] BAC clone DNA was injected into y^1

M{vas-int.Dm}ZH-2A w*; PBac{y+-attP-3B}^{VK00033} embryos, resulting in w^{1118}; *PBac{y[+mDint2]*

w^{+mC}=*FTSF.GFP-sov}^{VK00033}*.

<u>Microscopy</u>

We fixed ovaries in 4 or 5% EM-grade paraformaldehyde in PBS containing 0.1 or 0.3%

Triton X-100 (PBTX) for 10-15min, washed 3x 15 min in PBTX, and blocked >30min in 2%

normal goat serum, 0.5-1% bovine serum albumin (BSA) in PBS with 0.1% Tween-20 or 0.1%

Triton X-100. Antibodies and DAPI were diluted into blocking buffer. We incubated in primary

antibodies overnight at 4 °C and secondaries 2–3hr at room temperature. Embryos (1–2 hr)

were prepared for live imaging in halocarbon oil (Lerit et al. 2015). We imaged ovaries and

embryos using a Nikon Ti-E system or Zeiss LSM 780 microscope and eyes with a Nikon SMZ.

DNA-Seq

Genomic DNA was extracted from 30 whole flies per genotype (Huang, Rehm, and

Rubin 2009)(Sambrook and Russell 2006) to prepare DNA-seq libraries (Nextera DNA Library

Preparation Kit). We used 50 bp, single-end sequencing (Illumina HiSeq 2500, CASAVA base

calling). Sequence data are available at the SRA (SRP14438). We mapped DNAseq reads to

FlyBase r6.16 genome with Hisat2 (-k 1 --no-spliced-alignment)(Kim, Langmead, and Salzberg

2015). We used mpileup and bcftools commands from SAMtools within the genomic region

X:6756000–6771000 (H. Li et al. 2009; H. Li 2011) for variant calling and snpEFF to determine

the nature of variants in *sov* mutants (Cingolani et al. 2012).

RNA-Seq

Stranded PolyA+ RNA-seq libraries from *sov* and control ovaries (Table S5) were

created (H. Lee, Cho, et al. 2016) and are available at GEO (GSE113977). We extracted total

RNA (Qiagen RNeasy Mini Kit) in biological triplicate from 15 ovaries (4-5d post-eclosion) and

used 200 ng with 10 pg of ERCC spike-in control RNAs (pools 78A or 78B) for libraries (Jiang

et al. 2011; Zook et al. 2012; Pine et al. 2016; H. Lee, Pine, et al. 2016). We used 50 bp,

single-end sequencing as above. Tissue expression analysis are from GEO accession

GSE99574, a resource for comparing gene expression patterns.

We mapped RNA-seq reads to FlyBase r6.21 with Hisat2 (-k 1 --rna-strandness R --

dta)(Kim, Langmead, and Salzberg 2015). We determined read counts for each attribute of the

FlyBase r6.21 GTF file (with ERCC and transposable element sequences), with HTSeq-count

(Anders, Pyl, and Huber 2015). Transposon sequences were from the UCSC Genome

Browser RepeatMasker track (Casper et al. 2018; Smit, Hubley, and Green 2013-2015).

Benner et al.; *sov* represses gene expression

We conducted differential expression analysis with DESeq2 (pAdjustMethod = "fdr")

(Love, Huber, and Anders 2014). We removed genes with read counts $\leq$1 and read counts for

transposable elements with >1 location were summed for the DESeq2 analysis. *Df(1)sov/sov^2*

replicate 3 and *sov^2/w^{1118}* replicate 1 were failed. For *sov* mutant *vs.* control DESeq2 analysis,

all *sov* mutants were compared to all wild type controls. For tissue types, we compared each

sexed tissue to sexed whole organism. We used reads per kb per million reads (RPKM) for

gene-level expression.

For heatmaps, we calculated Euclidean distance, performed hierarchical cluster

analysis (agglomeration method = Ward), and mean-subtracted scaled across genotypes.

Mean sample RPKM correlation values (Table S1) were calculated by cor.test() function with

Pearson correlation coefficient in R (R Core Team 2017). We represented de-repressed genes

as mean-subtracted scaled values across tissues in the heatmap (Table S6).

For read density tracks, replicate raw read files were combined. Bedgraph files were

created with bedtools genomecov (Quinlan and Hall 2010) visualized on the UCSC genome

browser (Kent et al. 2002). Tracks were scaled by the number of reads divided by total reads

per million.

We used ClueGO (Bindea et al. 2009), with Cytoscape (Shannon et al. 2003) for one-

sided enrichment analysis (see Table S4).

Images were assembled using ImageJ and Photoshop. We used ROI tool in ImageJ,

plotted/analyzed image data (Microsoft Excel and GraphPad Prism), and calculated

significance by D'Agnostino and Pearson normality tests, followed Student's two-tailed t-test or

Mann-Whitney tests.

Benner et al.; *sov* represses gene expression

Acknowledgements

We thank Astrid Haase, Nasser Rusan, the Oliver and Lerit labs, and Dawson Mohler

(deceased) for stimulating discussions. We used resources from FlyBase, The Hybridoma

Bank, and the NIH biowulf (Table S5). This research was supported in part by the Intramural

Research Program of the NIH, NIDDK to BO; NIH Grant 5K22HL126922-02 to DAL; NIH

Graduate Partners Program to LB; Baylor College of Medicine, the Albert and Margaret Alkek

Foundation, the McNair Medical Institute at The Robert and Janice McNair Foundation, the

Cancer Prevention and Research Institute of Texas (R1313), and NIH grant 1R21HG006726 to

KJTV; and NIH grants P40OD018537 and R24RR014106, and the National Science

Foundation DBI-9816125 to KRC.

Author contributions

LB, KC, BO, and DAL designed the project and analyzed data. LB, KC, CW, and KJTV

performed genetics. LB and HY performed genomics. LB, EAC, CW and DAL performed

imaging. LB, BO, and DAL wrote the manuscript. All authors reviewed data and provided

feedback on the manuscript.

Benner et al.*; sov* represses gene expression

478
Figure 1. *sov* is *CG14438*. (A) Genes of the genomic interval X:6710000–6810000 (Gramates
et al. 2017). (B) Deficiency (*Df*) and Duplication (*Dp*) mapping. (C) Schematic of the *CG14438*
(black) and CR43496 (purple) genes. Transcription start (bent arrows), introns (thin lines) non-
coding regions (medium lines) and coding regions (thick lines) are shown. Transposon
insertions (triangles), point mutations (red lines), the region targeted by the shRNAi transgene
(base-paired), and Sov protein features are shown. Non-complementing (*sov⁻*, red) and
complementing (*sov⁺*, green) alleles and rearrangements are shown. (D) RNA expression
tracks by tissue type from female (red) or male (blue) adults.
487

488
489 **Figure 2. Sov germarium expression.** (A) Cartoon showing germarium regions I-III and
490 young egg chamber with cell types. (B,C) 1d post-eclosion, visualized for GFP-Sov (green),
491 anti-αSpectrin (red). Images show single optical sections. (B) Sov contrasted with anti-Vas
492 germline (magenta), or (C) -Tj somatic staining (magenta). GFP-Sov expression regions of
493 interest (see text) are shown (dashed lines). Bars: 10μm.
494

Benner et al.; *sov* represses gene expression

495
Figure 3. *sov* mutant defects. (A-K) images of indicated genotype (white). (A-F) Germaria (4-
5d post-eclosion) of stained with anti-Vas (blue), -Tj (green), -αSpectrin (red), and DAPI
(white). Dot spectrosomes (arrows) and displaced follicle cells (dashed lines) are shown.
Images are single optical sections. (G, H) Germaria (1d post-eclosion) stained with anti-Bam
(green), -actin (red), and DAPI (blue). Bam region of interest is shown (dashed lines). (I–K)
Germaria (1d post-eclosion) anti-Orb (green), -actin (red), and DAPI (blue). Orb+ cells shown
(*). Single optical sections (A-H) and maximum intensity projections (I-K). Bars: (A–F) 20µm,
(G–K) 10µm.
504

Figure 4. *sov* mutant transcriptome. (A) Relative expression in *sov* mutant vs *sov+* ovaries plotted against the mean expression in both sample types. Units are Log₂ normalized read counts (NRC). Data points are genes, those with >4-fold change (log₂ 2, red) or <–4 (log₂ -2, blue) and FDR *p*adj value <0.05 are highlighted. (B) Tissue-biased expression in wildtype for genes de-repressed in *sov* mutant ovaries. Heatmap from mean-subtracted ratios scaled across tissues (red=higher; blue=lower). (C) The *heartless* locus (see Fig1 for format) showing RNA-seq normalized read densities of *sov+* from head and *sov⁺* or *sov⁻* ovarian tissues. (D) Transposable element expression in *sov* mutants and control alleles (as indicated). Heatmap from mean-subtracted reads (in RPKM. red=higher; blue=lower) scaled for each transposable element (rows) across genotypes (columns). Some element subtypes are shown for DNA, non-LTR (Jockey), telomeric repeat (Telo), and LTR (Gypsy, Copia, and PAO) are indicated.

Figure 5. Sov is a dominant suppressor of position effect variegation. (A) Cartoon of
position effect variegation (PEV) in the eye. Expression of the *white* gene (bent arrow, thick
bar, red) near heterochromatin (squiggled) region can be silenced (white) by spreading. (B–F)
Eyes with variegated expression of *P{hsp26-pt-T}* transgene inserted into different
chromosomal positions (rows) in different *sov* backgrounds (columns).

Benner et al.; *sov* represses gene expression

Figure 6. Sov colocalizes with HP1a. Stills from live imaging of embryos expressing GFP-Sov and RFP-HP1a. (A) Localization of GFP-Sov and RFP-HP1a (rows) in stills from live imaging of an embryo expressing progressing from NC 10 (left) to NC 11 (right) with stages (columns, nuclear envelope breakdown=NEB) and time (min:s in merge row) shown. (B) expressing GFP-Sov and RFP-HP1a (columns) at NC 14. Boxed regions are magnified in insets below. An HP1a subnuclear domain is shown (arrows). (B') Single optical section containing peak HP1a fluorescence of the same region shown in (B). Dashed line indicates region used for linescan analysis. (A,B) Maximum projections through 1.5 μm volume at 1F/30s. Bars: 10 μm; insets, 5 μm. (C) Histogram of HP1a and Sov fluorescence intensity measured in (B'). Fluorescence levels (arbitrary units) normalized to the peak fluorescence intensity for each channel and the distance (μm) to peak HP1a signal. Line scans were scaled –3 to 3μm, where 0.0 is peak HP1a fluorescence. Half maximum HP1a fluorescence is shown (yellow shade).

540
541 **Figure 7. Working model of *sov* function.** Sov protein function and attributes are analogous
542 to the mammalian KRAB proteins (green). Like KRAB proteins, Sov is in complex with HP1a
543 (yellow), Bon (mammalian KAP-1, orange) and Egg (mammalian SETDB1, black) providing a
544 DNA tether that functions along with the H3K9me tether written by Egg (arrow, red lettering)
545 and bound by HP1, to stabilize a repressed state.
546

Benner et al.; *sov* represses gene expression

References

Alekseyenko, Artyom A., Andrey A. Gorchakov, Barry M. Zee, Stephen M. Fuchs, Peter V. Kharchenko,

and Mitzi I. Kuroda. 2014. "Heterochromatin-Associated Interactions of Drosophila HP1a with

dADD1, HIPP1, and Repetitive RNAs." *Genes & Development* 28 (13): 1445–60.

Anders, Simon, Paul Theodor Pyl, and Wolfgang Huber. 2015. "HTSeq--a Python Framework to Work

with High-Throughput Sequencing Data." *Bioinformatics* 31 (2): 166–69.

Barton, Lacy J., Kaylee E. Lovander, Belinda S. Pinto, and Pamela K. Geyer. 2016. "Drosophila Male

and Female Germline Stem Cell Niches Require the Nuclear Lamina Protein Otefin."

Developmental Biology 415 (1): 75–86.

Bausek, Nina. 2013. "JAK-STAT Signaling in Stem Cells and Their Niches in Drosophila." *JAK-STAT* 2

(3): e25686.

Beckstead, R. B., S. S. Ner, K. G. Hales, T. A. Grigliatti, B. S. Baker, and H. J. Bellen. 2005. "Bonus, a

Drosophila TIF1 Homolog, Is a Chromatin-Associated Protein That Acts as a Modifier of Position-

Effect Variegation." *Genetics* 169 (2): 783–94.

Bindea, Gabriela, Bernhard Mlecnik, Hubert Hackl, Pornpimol Charoentong, Marie Tosolini, Amos

Kirilovsky, Wolf-Herman Fridman, Franck Pagès, Zlatko Trajanoski, and Jérôme Galon. 2009.

"ClueGO: A Cytoscape Plug-in to Decipher Functionally Grouped Gene Ontology and Pathway

Annotation Networks." *Bioinformatics* 25 (8): 1091–93.

Börner, Kenneth, Dhawal Jain, Paula Vazquez-Pianzola, Sandra Vengadasalam, Natascha Steffen,

Dmitry V. Fyodorov, Pavel Tomancak, Alexander Konev, Beat Suter, and Peter B. Becker. 2016.

"A Role for Tuned Levels of Nucleosome Remodeler Subunit ACF1 during Drosophila Oogenesis."

Developmental Biology 411 (2): 217–30.

Branco, Joana, Ismael Al-Ramahi, Lubna Ukani, Alma M. Pérez, Pedro Fernandez-Funez, Diego

Rincón-Limas, and Juan Botas. 2008. "Comparative Analysis of Genetic Modifiers in Drosophila

Points to Common and Distinct Mechanisms of Pathogenesis among Polyglutamine Diseases."

Benner et al.; *sov* represses gene expression

Human Molecular Genetics 17 (3): 376–90.

Brennecke, Julius, Alexei A. Aravin, Alexander Stark, Monica Dus, Manolis Kellis, Ravi

Sachidanandam, and Gregory J. Hannon. 2007. "Discrete Small RNA-Generating Loci as Master

Regulators of Transposon Activity in Drosophila." *Cell* 128 (6): 1089–1103.

Brower-Toland, Brent, Seth D. Findley, Ling Jiang, Li Liu, Hang Yin, Monica Dus, Pei Zhou, Sarah C. R.

Elgin, and Haifan Lin. 2007. "Drosophila PIWI Associates with Chromatin and Interacts Directly

with HP1a." *Genes & Development* 21 (18): 2300–2311.

Brown, James B., Nathan Boley, Robert Eisman, Gemma E. May, Marcus H. Stoiber, Michael O. Duff,

Ben W. Booth, et al. 2014. "Diversity and Dynamics of the Drosophila Transcriptome." *Nature* 512

(7515): 393–99.

Brown, J. Lesley, Cornelia Fritsch, Jürg Mueller, and Judith A. Kassis. 2003. "The Drosophila Pho-like

Gene Encodes a YY1-Related DNA Binding Protein That Is Redundant with Pleiohomeotic in

Homeotic Gene Silencing." *Development* 130 (2): 285–94.

Casper, Jonathan, Ann S. Zweig, Chris Villarreal, Cath Tyner, Matthew L. Speir, Kate R. Rosenbloom,

Brian J. Raney, et al. 2018. "The UCSC Genome Browser Database: 2018 Update." *Nucleic Acids

Research* 46 (D1): D762–69.

Cenci, Giovanni, Laura Ciapponi, and Maurizio Gatti. 2005. "The Mechanism of Telomere Protection: A

Comparison between Drosophila and Humans." *Chromosoma* 114 (3): 135–45.

Charlesworth, B., and C. H. Langley. 1989. "The Population Genetics of Drosophila Transposable

Elements." *Annual Review of Genetics* 23: 251–87.

Chen, Shuyi, Su Wang, and Ting Xie. 2011. "Restricting Self-Renewal Signals within the Stem Cell

Niche: Multiple Levels of Control." *Current Opinion in Genetics & Development* 21 (6): 684–89.

Chen, Taiping, and Sharon Y. R. Dent. 2014. "Chromatin Modifiers and Remodellers: Regulators of

Cellular Differentiation." *Nature Reviews. Genetics* 15 (2): 93–106.

Chou, T. B., and N. Perrimon. 1996. "The Autosomal FLP-DFS Technique for Generating Germline

Mosaics in Drosophila Melanogaster." *Genetics* 144 (4): 1673–79.

Benner et al.*; sov* represses gene expression

Cingolani, Pablo, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan Wang, Susan J.

Land, Xiangyi Lu, and Douglas M. Ruden. 2012. "A Program for Annotating and Predicting the

Effects of Single Nucleotide Polymorphisms, SnpEff: SNPs in the Genome of Drosophila

Melanogaster Strain w1118; Iso-2; Iso-3." *Fly* 6 (2): 80–92.

Clark, R. F., and S. C. Elgin. 1992. "Heterochromatin Protein 1, a Known Suppressor of Position-Effect

Variegation, Is Highly Conserved in Drosophila." *Nucleic Acids Research* 20 (22): 6067–74.

Clough, Emily, Woongjoon Moon, Shengxian Wang, Kathleen Smith, and Tulle Hazelrigg. 2007.

"Histone Methylation Is Required for Oogenesis in Drosophila." *Development* 134 (1): 157–65.

Clough, Emily, Thomas Tedeschi, and Tulle Hazelrigg. 2014. "Epigenetic Regulation of Oogenesis and

Germ Stem Cell Maintenance by the Drosophila Histone Methyltransferase Eggless/dSetDB1."

Developmental Biology 388 (2): 181–91.

Cook, R. Kimberley, Stacey J. Christensen, Jennifer A. Deal, Rachel A. Coburn, Megan E. Deal, Jill M.

Gresens, Thomas C. Kaufman, and Kevin R. Cook. 2012. "The Generation of Chromosomal

Deletions to Provide Extensive Coverage and Subdivision of the Drosophila Melanogaster

Genome." *Genome Biology* 13 (3): R21.

Court, Donald L., Srividya Swaminathan, Daiguan Yu, Helen Wilson, Teresa Baker, Mikail Bubunenko,

James Sawitzke, and Shyam K. Sharan. 2003. "Mini-Lambda: A Tractable System for

Chromosome and BAC Engineering." *Gene* 315 (October): 63–69.

Di Stefano, Luisa, Jun-Yuan Ji, Nam-Sung Moon, Anabel Herr, and Nicholas Dyson. 2007. "Mutation of

Drosophila Lsd1 Disrupts H3-K4 Methylation, Resulting in Tissue-Specific Defects during

Development." *Current Biology: CB* 17 (9): 808–12.

Ebert, Anja, Sandro Lein, Gunnar Schotta, and Gunter Reuter. 2006. "Histone Modification and the

Control of Heterochromatic Gene Silencing in Drosophila." *Chromosome Research: An

International Journal on the Molecular, Supramolecular and Evolutionary Aspects of Chromosome

Biology* 14 (4): 377–92.

Ecco, Gabriela, Michael Imbeault, and Didier Trono. 2017. "KRAB Zinc Finger Proteins." *Development*

Benner et al.*; sov* represses gene expression

144 (15): 2719–29.

Eissenberg, J. C., T. C. James, D. M. Foster-Hartnett, T. Hartnett, V. Ngan, and S. C. Elgin. 1990.

"Mutation in a Heterochromatin-Specific Chromosomal Protein Is Associated with Suppression of

Position-Effect Variegation in Drosophila Melanogaster." *Proceedings of the National Academy of*

Sciences of the United States of America 87 (24): 9923–27.

Elgin, Sarah C. R., and Gunter Reuter. 2013. "Position-Effect Variegation, Heterochromatin Formation,

and Gene Silencing in Drosophila." *Cold Spring Harbor Perspectives in Biology* 5 (8): a017780.

Eliazer, Susan, Victor Palacios, Zhaohui Wang, Rahul K. Kollipara, Ralf Kittler, and Michael Buszczak.

2014. "Lsd1 Restricts the Number of Germline Stem Cells by Regulating Multiple Targets in Escort

Cells." *PLoS Genetics* 10 (3): e1004200.

Eliazer, Susan, Nevine A. Shalaby, and Michael Buszczak. 2011. "Loss of Lysine-Specific Demethylase

1 Nonautonomously Causes Stem Cell Tumors in the Drosophila Ovary." *Proceedings of the*

National Academy of Sciences of the United States of America 108 (17): 7064–69.

Foe, V. E., and B. M. Alberts. 1983. "Studies of Nuclear and Cytoplasmic Behaviour during the Five

Mitotic Cycles That Precede Gastrulation in Drosophila Embryogenesis." *Journal of Cell Science*

61 (May): 31–70.

Fuller, Margaret T., and Allan C. Spradling. 2007. "Male and Female Drosophila Germline Stem Cells:

Two Versions of Immortality." *Science* 316 (5823): 402–4.

Gene Ontology Consortium. 2015. "Gene Ontology Consortium: Going Forward." *Nucleic Acids*

Research 43 (Database issue): D1049–56.

Gilboa, Lilach. 2015. "Organizing Stem Cell Units in the Drosophila Ovary." *Current Opinion in Genetics*

& Development 32 (June): 31–36.

Goode, S., D. Wright, and A. P. Mahowald. 1992. "The Neurogenic Locus Brainiac Cooperates with the

Drosophila EGF Receptor to Establish the Ovarian Follicle and to Determine Its Dorsal-Ventral

Polarity." *Development* 116 (1): 177–92.

Gramates, L. Sian, Steven J. Marygold, Gilberto Dos Santos, Jose-Maria Urbano, Giulia Antonazzo,

Benner et al.; *sov* represses gene expression

Beverley B. Matthews, Alix J. Rey, et al. 2017. "FlyBase at 25: Looking to the Future." *Nucleic*

Acids Research 45 (D1): D663–71.

Graveley, Brenton R., Angela N. Brooks, Joseph W. Carlson, Michael O. Duff, Jane M. Landolin, Li

Yang, Carlo G. Artieri, et al. 2011. "The Developmental Transcriptome of Drosophila

Melanogaster." *Nature* 471 (7339): 473–79.

Gu, Tingting, and Sarah C. R. Elgin. 2013. "Maternal Depletion of Piwi, a Component of the RNAi

System, Impacts Heterochromatin Formation in Drosophila." *PLoS Genetics* 9 (9): e1003780.

Hsieh, M., Y. Tintut, and J. D. Gralla. 1994. "Functional Roles for the Glutamines within the Glutamine-

Rich Region of the Transcription Factor Sigma 54." *The Journal of Biological Chemistry* 269 (1):

373–78.

Huang, Audrey M., E. Jay Rehm, and Gerald M. Rubin. 2009. "Quick Preparation of Genomic DNA

from Drosophila." *Cold Spring Harbor Protocols* 2009 (4): db.prot5198.

James, T. C., and S. C. Elgin. 1986. "Identification of a Nonhistone Chromosomal Protein Associated

with Heterochromatin in Drosophila Melanogaster and Its Gene." *Molecular and Cellular Biology* 6

(11): 3862–72.

Jankovics, Ferenc, László Henn, Ágnes Bujna, Péter Vilmos, Kerstin Spirohn, Michael Boutros, and

Miklós Erdélyi. 2014. "Functional Analysis of the Drosophila Embryonic Germ Cell Transcriptome

by RNA Interference." *PloS One* 9 (6): e98579.

Jenuwein, T., and C. D. Allis. 2001. "Translating the Histone Code." *Science* 293 (5532): 1074–80.

Jiang, Lichun, Felix Schlesinger, Carrie A. Davis, Yu Zhang, Renhua Li, Marc Salit, Thomas R.

Gingeras, and Brian Oliver. 2011. "Synthetic Spike-in Standards for RNA-Seq Experiments."

Genome Research 21 (9): 1543–51.

Kassis, Judith A., James A. Kennison, and John W. Tamkun. 2017. "Polycomb and Trithorax Group

Genes in Drosophila." *Genetics* 206 (4): 1699–1725.

Kent, W. James, Charles W. Sugnet, Terrence S. Furey, Krishna M. Roskin, Tom H. Pringle, Alan M.

Zahler, and David Haussler. 2002. "The Human Genome Browser at UCSC." *Genome Research*

12 (6): 996–1006.

Kim, Daehwan, Ben Langmead, and Steven L. Salzberg. 2015. "HISAT: A Fast Spliced Aligner with

Low Memory Requirements." *Nature Methods* 12 (4): 357–60.

Kouzarides, Tony. 2007. "Chromatin Modifications and Their Function." *Cell* 128 (4): 693–705.

Lantz, V., J. S. Chang, J. I. Horabin, D. Bopp, and P. Schedl. 1994. "The Drosophila Orb RNA-Binding

Protein Is Required for the Formation of the Egg Chamber and Establishment of Polarity." *Genes &*

Development 8 (5): 598–613.

Leader, David P., Sue A. Krause, Aniruddha Pandit, Shireen A. Davies, and Julian A. T. Dow. 2018.

"FlyAtlas 2: A New Version of the Drosophila Melanogaster Expression Atlas with RNA-Seq,

miRNA-Seq and Sex-Specific Data." *Nucleic Acids Research* 46 (D1): D809–15.

Lee, Hangnoh, Dong-Yeon Cho, Cale Whitworth, Robert Eisman, Melissa Phelps, John Roote, Thomas

Kaufman, et al. 2016. "Effects of Gene Dose, Chromatin, and Network Topology on Expression in

Drosophila Melanogaster." *PLoS Genetics* 12 (9): e1006295.

Lee, Hangnoh, P. Scott Pine, Jennifer McDaniel, Marc Salit, and Brian Oliver. 2016. "External RNA

Controls Consortium Beta Version Update." *Journal of Genomics* 4 (July): 19–22.

Lee, Jeannie T. 2009. "Lessons from X-Chromosome Inactivation: Long ncRNA as Guides and Tethers

to the Epigenome." *Genes & Development* 23 (16): 1831–42.

Lerit, Dorothy A., Holly A. Jordan, John S. Poulton, Carey J. Fagerstrom, Brian J. Galletta, Mark Peifer,

and Nasser M. Rusan. 2015. "Interphase Centrosome Organization by the PLP-Cnn Scaffold Is

Required for Centrosome Function." *The Journal of Cell Biology* 210 (1): 79–97.

Li, Heng. 2011. "A Statistical Framework for SNP Calling, Mutation Discovery, Association Mapping and

Population Genetical Parameter Estimation from Sequencing Data." *Bioinformatics* 27 (21): 2987–

93.

Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo

Abecasis, Richard Durbin, and 1000 Genome Project Data Processing Subgroup. 2009. "The

Sequence Alignment/Map Format and SAMtools." *Bioinformatics* 25 (16): 2078–79.

Lin, H., and A. C. Spradling. 1995. "Fusome Asymmetry and Oocyte Determination in Drosophila."

Developmental Genetics 16 (1): 6–12.

Li, Xuanying, Christopher W. Seidel, Leanne T. Szerszen, Jeffrey J. Lange, Jerry L. Workman, and

Susan M. Abmayr. 2017. "Enzymatic Modules of the SAGA Chromatin-Modifying Complex Play

Distinct Roles in Drosophila Gene Expression and Development." *Genes & Development* 31 (15):

1588–1600.

Lorch, Yahli, and Roger D. Kornberg. 2017. "Chromatin-Remodeling for Transcription." *Quarterly*

Reviews of Biophysics 50 (January): e5.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change

and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.

Mason, James M., Radmila Capkova Frydrychova, and Harald Biessmann. 2008. "Drosophila

Telomeres: An Exception Providing New Insights." *BioEssays: News and Reviews in Molecular,*

Cellular and Developmental Biology 30 (1): 25–37.

McConnell, Kristopher H., Michael Dixon, and Brian R. Calvi. 2012. "The Histone Acetyltransferases

CBP and Chameau Integrate Developmental and DNA Replication Programs in Drosophila

Ovarian Follicle Cells." *Development* 139 (20): 3880–90.

McKearin, D. M., and A. C. Spradling. 1990. "Bag-of-Marbles: A Drosophila Gene Required to Initiate

Both Male and Female Gametogenesis." *Genes & Development* 4 (12B): 2242–51.

McKearin, D., and B. Ohlstein. 1995. "A Role for the Drosophila Bag-of-Marbles Protein in the

Differentiation of Cystoblasts from Germline Stem Cells." *Development* 121 (9): 2937–47.

Mohler, Dawson, and Andrea Carroll. 1984. "Report of Dawson Mohler and Andrea Carroll." *Drosophila*

Information Service 60 (June): 236–41.

Mohler, J. D. 1977. "Developmental Genetics of the Drosophila Egg. I. Identification of 59 Sex-Linked

Cistrons with Maternal Effects on Embryonic Development." *Genetics* 85 (2): 259–72.

Ohlstein, B., and D. McKearin. 1997. "Ectopic Expression of the Drosophila Bam Protein Eliminates

Oogenic Germline Stem Cells." *Development* 124 (18): 3651–62.

Benner et al.; *sov* represses gene expression

Orr-Weaver, T. L. 1991. "Drosophila Chorion Genes: Cracking the Eggshell's Secrets." *BioEssays:*

News and Reviews in Molecular, Cellular and Developmental Biology 13 (3): 97–105.

Parks, Annette L., Kevin R. Cook, Marcia Belvin, Nicholas A. Dompe, Robert Fawcett, Kari Huppert,

Lory R. Tan, et al. 2004. "Systematic Generation of High-Resolution Deletion Coverage of the

Drosophila Melanogaster Genome." *Nature Genetics* 36 (3): 288–92.

Pelisson, A., L. Mejlumian, V. Robert, C. Terzian, and A. Bucheton. 2002. "Drosophila Germline

Invasion by the Endogenous Retrovirus Gypsy: Involvement of the Viral Env Gene." *Insect*

Biochemistry and Molecular Biology 32 (10): 1249–56.

Peng, Jamy C., Anton Valouev, Na Liu, and Haifan Lin. 2016. "Piwi Maintains Germline Stem Cells and

Oogenesis in Drosophila through Negative Regulation of Polycomb Group Proteins." *Nature*

Genetics 48 (3): 283–91.

Pine, P. Scott, Sarah A. Munro, Jerod R. Parsons, Jennifer McDaniel, Anne Bergstrom Lucas, Jean

Lozach, Timothy G. Myers, Qin Su, Sarah M. Jacobs-Helber, and Marc Salit. 2016. "Evaluation of

the External RNA Controls Consortium (ERCC) Reference Material Using a Modified Latin Square

Design." *BMC Biotechnology* 16 (1): 54.

Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing

Genomic Features." *Bioinformatics* 26 (6): 841–42.

R Core Team. 2017. "A Language and Environment for Statistical Computing." Vienna, Austria: R

Foundation for Statistical Computing. https://www.R-project.org/.

Reuter, G., and P. Spierer. 1992. "Position Effect Variegation and Chromatin Proteins." *BioEssays:*

News and Reviews in Molecular, Cellular and Developmental Biology 14 (9): 605–12.

Robinson, Scott W., Pawel Herzyk, Julian A. T. Dow, and David P. Leader. 2013. "FlyAtlas: Database

of Gene Expression in the Tissues of Drosophila Melanogaster." *Nucleic Acids Research* 41

(Database issue): D744–50.

Rudolph, Thomas, Masato Yonezawa, Sandro Lein, Kathleen Heidrich, Stefan Kubicek, Christiane

Schäfer, Sameer Phalke, et al. 2007. "Heterochromatin Formation in Drosophila Is Initiated through

Benner et al.*; sov* represses gene expression

Active Removal of H3K4 Methylation by the LSD1 Homolog SU(VAR)3-3." *Molecular Cell* 26 (1):

103–15.

Sambrook, Joseph, and David W. Russell. 2006. "Purification of Nucleic Acids by Extraction with

Phenol:chloroform." *CSH Protocols* 2006 (1). https://doi.org/10.1101/pdb.prot4455.

Shannon, Paul, Andrew Markiel, Owen Ozier, Nitin S. Baliga, Jonathan T. Wang, Daniel Ramage, Nada

Amin, Benno Schwikowski, and Trey Ideker. 2003. "Cytoscape: A Software Environment for

Integrated Models of Biomolecular Interaction Networks." *Genome Research* 13 (11): 2498–2504.

Smit, Afa, R. Hubley, and P. Green. 2013-2015. "RepeatMasker Open-4.0."

http://www.repeatmasker.org.

Soshnev, Alexey A., Ryan M. Baxley, J. Robert Manak, Kai Tan, and Pamela K. Geyer. 2013. "The

Insulator Protein Suppressor of Hairy-Wing Is an Essential Transcriptional Repressor in the

Drosophila Ovary." *Development* 140 (17): 3613–23.

Stork, Tobias, Amy Sheehan, Ozge E. Tasdemir-Yilmaz, and Marc R. Freeman. 2014. "Neuron-Glia

Interactions through the Heartless FGF Receptor Signaling Pathway Mediate Morphogenesis of

Drosophila Astrocytes." *Neuron* 83 (2): 388–403.

Teixeira, Felipe Karam, Martyna Okuniewska, Colin D. Malone, Rémi-Xavier Coux, Donald C. Rio, and

Ruth Lehmann. 2017. "piRNA-Mediated Regulation of Transposon Alternative Splicing in the Soma

and Germ Line." *Nature* 552 (7684): 268–72.

Venken, Koen J. T., Joseph W. Carlson, Karen L. Schulze, Hongling Pan, Yuchun He, Rebecca

Spokony, Kenneth H. Wan, et al. 2009. "Versatile P[acman] BAC Libraries for Transgenesis

Studies in Drosophila Melanogaster." *Nature Methods* 6 (6): 431–34.

Venken, Koen J. T., Karen L. Schulze, Nele A. Haelterman, Hongling Pan, Yuchun He, Martha Evans-

Holm, Joseph W. Carlson, et al. 2011. "MiMIC: A Highly Versatile Transposon Insertion Resource

for Engineering Drosophila Melanogaster Genes." *Nature Methods* 8 (9): 737–43.

Vermaak, Danielle, and Harmit S. Malik. 2009. "Multiple Roles for Heterochromatin Protein 1 Genes in

Drosophila." *Annual Review of Genetics* 43: 467–92.

Benner et al.*; sov* represses gene expression

Wang, Sidney H., and Sarah C. R. Elgin. 2011. "Drosophila Piwi Functions Downstream of piRNA

Production Mediating a Chromatin-Based Transposon Silencing Mechanism in Female Germ Line."

Proceedings of the National Academy of Sciences of the United States of America 108 (52):

21164–69.

Wang, Xiaoxi, Lei Pan, Su Wang, Jian Zhou, William McDowell, Jungeun Park, Jeff Haug, Karen

Staehling, Hong Tang, and Ting Xie. 2011. "Histone H3K9 Trimethylase Eggless Controls

Germline Stem Cell Maintenance and Differentiation." *PLoS Genetics* 7 (12): e1002426.

Warming, Søren, Nina Costantino, Donald L. Court, Nancy A. Jenkins, and Neal G. Copeland. 2005.

"Simple and Highly Efficient BAC Recombineering Using galK Selection." *Nucleic Acids Research*

33 (4): e36.

Wayne, S., K. Liggett, J. Pettus, and R. N. Nagoshi. 1995. "Genetic Characterization of Small Ovaries,

a Gene Required in the Soma for the Development of the Drosophila Ovary and the Female

Germline." *Genetics* 139 (3): 1309–20.

Weiler, K. S., and B. T. Wakimoto. 1995. "Heterochromatin and Gene Expression in Drosophila."

Annual Review of Genetics 29: 577–605.

Yang, Peng, Yixuan Wang, and Todd S. Macfarlan. 2017. "The Role of KRAB-ZFPs in Transposable

Element Repression and Mammalian Evolution." *Trends in Genetics: TIG* 33 (11): 871–81.

Yasuhara, Jiro C., and Barbara T. Wakimoto. 2006. "Oxymoron No More: The Expanding World of

Heterochromatic Genes." *Trends in Genetics: TIG* 22 (6): 330–38.

Yin, Hang, and Haifan Lin. 2007. "An Epigenetic Activation Role of Piwi and a Piwi-Associated piRNA in

Drosophila Melanogaster." *Nature* 450 (7167): 304–8.

Yoshioka, K., H. Honma, M. Zushi, S. Kondo, S. Togashi, T. Miyake, and T. Shiba. 1990. "Virus-like

Particle Formation of Drosophila Copia through Autocatalytic Processing." *The EMBO Journal* 9

(2): 535–41.

Yuan, Kai, and Patrick H. O'Farrell. 2016. "TALE-Light Imaging Reveals Maternally Guided,

H3K9me2/3-Independent Emergence of Functional Heterochromatin in Drosophila Embryos."

Benner et al.; *sov* represses gene expression

Genes & Development 30 (5): 579–93.

Zook, Justin M., Daniel Samarov, Jennifer McDaniel, Shurjo K. Sen, and Marc Salit. 2012. "Synthetic

Spike-in Standards Improve Run-Specific Systematic Error Analysis for DNA and RNA

Sequencing." *PloS One* 7 (7): e41356.

Zhimulev, Igor F., and Elena S. Belyaeva. 2003. "Intercalary Heterochromatin and Genetic Silencing."

BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology 25 (11): 1040–

51.

Benner et al.; *sov* represses gene expression