***Drosophila small ovary* encodes a zinc-finger repressor required for ovarian**

differentiation

Leif Benner[1,2], Elias A. Castro[3], Cale Whitworth[1,4], Koen J.T. Venken[5], Haiwang Yang[1], Brian

Oliver[1,6,7], Kevin R. Cook[4], and Dorothy A. Lerit[3,6]

Author affiliations:

[1]Section of Developmental Genomics, Laboratory of Cellular and Developmental

Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National

Institutes of Health, Bethesda MD 20892. [2]Department of Biology, Johns Hopkins

University, Baltimore MD 21218. [3]Department of Cell Biology, Emory University School

of Medicine, Atlanta GA 30322. [4]Department of Biology, Indiana University,

Bloomington, Indiana 47405. [5]Verna and Marrs McLean Department of Biochemistry

and Molecular Biology, McNair Medical Institute at the Robert and Janice McNair

Foundation, Dan L. Duncan Cancer Center, Center for Drug Discovery, Department of

Pharmacology and Chemical Biology, Baylor College of Medicine, Houston, TX, 77030,

USA

[6]**Corresponding authors:** dlerit@emory.edu briano@nih.gov

[7]**Lead contact**: briano@nih.gov

Keywords: chromatin, differentiation, gene expression, HP1a, oogenesis, position-effect

variegation

 Summary statement

Small ovary is required for heterochromatin stabilization to repress the expression of

transposons and ectopic signaling pathways in the developing ovary.

Abstract

Repression is essential for coordinated cell type-specific gene regulation and controlling the

expression of transposons. In the *Drosophila* ovary, stem cell regeneration and differentiation

requires controlled gene expression, with derepression leading to tissue degeneration and

ovarian tumors. Likewise, the ovary is acutely sensitive to deleterious consequences of

transposon derepression. The *small ovary* (*sov*) locus was identified in a female sterile screen,

and mutants show dramatic ovarian morphogenesis defects. We mapped the locus to the

uncharacterized gene *CG14438*, which encodes a zinc-finger protein that colocalizes with the

essential Heterochromatin Protein 1 (HP1a). We demonstrate that Sov functions to repress

inappropriate cell signaling, silence transposons, and suppress position-effect variegation in

the eye, suggesting a central role in heterochromatin stabilization.

40
Introduction

Multicellular organisms rely on stem cells. A combination of extrinsic and intrinsic signals

function to maintain the balance between stem cell self-renewal and differentiation needed for

tissue homeostasis of stem cell development. The *Drosophila* adult ovary is a well-studied

model for development (Fuller and Spradling 2007). It is organized into ~16 ovarioles, which

are assembly lines of progressively maturing egg chambers. At the anterior tip of each ovariole

is a structure called the germarium, harboring 2–3 germline stem cells (GSCs), which sustain

egg production throughout the life of the animal. Surrounding the GSCs and their cystoblast

daughters is a stem cell niche, a collection of nonproliferating somatic cells composed of

terminal filament and cap cells. Like the GSCs, somatic stem cells divide to produce new stem

cells and escort cells, which encase the germline. Each cystoblast is fated to differentiate and

will undergo four rounds of cell division with incomplete cytokinesis to produce a germline cyst

interconnected by the fusome, a branched cytoskeletal structure. Germline cysts are

surrounded by two escort cells. Toward the posterior of the germarium, a monolayer of somatic

follicle cells replaces the escort cells to form an egg chamber.

Egg chamber development requires the careful coordination of distinct germline and

somatic cell populations through signaling within a complex microenvironment. For example,

extrinsic transducers and signals including Jak/Stat and the BMP homolog, Decapentapalegic

(Dpp), are required to maintain the proliferative potential of GSCs (Bausek 2013; Gilboa 2015;

S. Chen, Wang, and Xie 2011). Oriented divisions displace the primary cystoblast cell away

from the stem cell signals and permit the expression of differentiation factors, such as *bag of*

marbles (*bam*). The importance of Bam is revealed in *bam* loss-of-function mutants, which

have tumorous germaria filled with undifferentiated germline cells containing dot

spectrosomes, the dot-fusome structure characteristic of GSCs and primary cystoblasts as a

result of complete cytokinesis (Lin and Spradling 1995; D. McKearin and Ohlstein 1995). Thus,

germline differentiation requires the repression and activation of cell signaling pathways.

Dynamic changes in chromatin landscapes within germline and somatic cells are critical

for oogenesis and contribute to the regulated gene expression required for tissue homeostasis

(Barton et al. 2016; X. Li et al. 2017; Börner et al. 2016; Peng et al. 2016; Soshnev et al. 2013;

McConnell, Dixon, and Calvi 2012). Nucleosomes generally repress transcription by competing

for DNA binding with transcription factors (Lorch and Kornberg 2017; Kouzarides 2007;

Benner et al.*; sov* represses gene expression

Jenuwein and Allis 2001). The reinforcement of repression depends largely on histone

modifications. For example, the activities of both the H3K9 methyltransferase SETDB1,

encoded by *eggless* (*egg*) (Wang et al. 2011; Clough et al. 2007; Clough, Tedeschi, and

Hazelrigg 2014), and the H3K4 demethylase encoded by *lysine-specific demethylase 1* (Di

Stefano et al. 2007; Rudolph et al. 2007; Eliazer, Shalaby, and Buszczak 2011; Eliazer et al.

2014) are required in the somatic cells of the germarium for GSC maintenance, normal

patterns of differentiation, and germline development. Genomewide profiling hints at a

progression from open chromatin in stem cells to a more closed state during differentiation (T.

Chen and Dent 2014). Disrupting this progressive repression in the ovary is predicted to

contribute to stem cell overproliferation and defective oogenesis.

There are other roles for regulated chromatin states in oogenesis. The propensity of

transposons to mobilize in gonads is countered by host responses, such as induced

heterochromatin formation, to repress transposon activity. The proximity of condensed

heterochromatin also dampens the activity of genes and transgenes (Elgin and Reuter 2013).

Originally described over 40 years ago (J. D. Mohler 1977), the *small ovary* (*sov*) locus

is associated with a range of mutant phenotypes including disorganized ovarioles, egg

chamber fusions, undifferentiated tumors, and ovarian degeneration (Wayne et al. 1995). In

the present work, we define the molecular identify and function of Sov. We demonstrate *sov*

encodes an unusually long C_2H_2 zinc-finger (ZnF) nuclear protein. Previous work suggested

Sov may complex with the conserved Heterochromatin-Protein 1a (HP1a) (Alekseyenko et al.

2014), encoded by *Su(var)205* and critical for heterochromatin formation (James and Elgin

1986; Eissenberg et al. 1990; Clark and Elgin 1992). RNA-seq analysis showed that *sov*

activity is required in the ovary to repress the expression of a large number of genes and

Benner et al.*; sov* represses gene expression

transposons. Additionally, Sov functions as a dominant suppressor of position-effect

variegation (PEV) in the eye, similar to HP1a (Elgin and Reuter 2013). Moreover, Sov and

HP1a colocalize in the nucleus. These data indicate that Sov is a novel repressor of gene

expression involved in heterochromatization.

Results

small ovary is CG14438

To refine previous mapping of *sov*, we used complementation of female sterile and lethal *sov*

mutations with preexisting and custom-generated deficiencies, duplications, and transposon

insertions (J. D. Mohler 1977; D. Mohler and Carroll 1984; Wayne et al. 1995). Our results (Fig

1A, B) suggested that either the protein-coding *CG14438* gene or the intronic, noncoding

CR43496 gene is *sov* since the locus mapped to the noncomplementing deletion *Df(1)sov*.

Rescue of female sterility and/or lethality of *sov^2*, *sov^{ML150}*, *sov^{EA42}* and *Df(1)sov* with

Dp(1;3)sn^{13a1} (molecularly undefined, not shown), *Dp(1;3)DC486*, and *PBac{GFP-sov}*

confirmed our mapping. We could not replicate separability of *sov^{EA42}* sterility and lethality with

Dp(1;3)sn^{13a1} (Wayne et al. 1995). Female sterile and lethal alleles of *sov* map identically.

We performed genome sequencing to determine if the lesions in *sov* alleles occur in

CG14438 or *CR43496* (Table S1). While *CR43496* contained three polymorphisms relative to

the reference genome, we identified the same polymorphisms in all *sov* mutant and control

lines. Thus, *CR43496* is unlikely to be *sov.* In contrast, we found disruptive mutations in

CG14438 (Fig 1C), which encodes an unusually long, 3,313-residue protein with 21 C_2H_2 ZnFs

and multiple nuclear localization motifs and coiled-coil regions (Fig 1C). We identified a

nonsense mutation (G to A at position 6,764,462) in the *CG14438* open reading frame of

Benner et al.*; sov* represses gene expression

sov^{EA42} which is predicted to truncate the *sov* protein before the ZnF domains. We found a

frameshift insertion (T at position 6,769,742) located towards the end of *CG14438* in *sov²* that

encodes 30 novel residues followed by a stop codon within the C-terminal ZnF and removes

the terminally predicted NLS (Fig 1C). We found a missense mutation (C to G at position

6,763,888) in *sov^{ML150}* that results in a glutamine to glutamate substitution within a predicted

coiled-coil domain. While this is a conservative substitution and glutamate residues are

common in coiled-coil domains, glutamine to glutamate substitutions are especially disruptive

in the coiled-coil region of the sigma transcription factor (Hsieh, Tintut, and Gralla 1994). We

conclude that *CG14438* encodes *sov*.

sov is highly expressed in the ovary

To determine where *sov* is expressed and if it encodes multiple isoforms, we analyzed its

expression in adult tissues by RNA-seq. We noted that *sov* is broadly expressed as a single

mRNA isoform, with highest expression in ovaries (Fig 1D). The modENCODE (Graveley et al.

2011; J. B. Brown et al. 2014) and FlyAtlas (Robinson et al. 2013; Leader et al. 2018)

reference sets show a similar enrichment in the ovary and early embryos. The enrichment of

sov in the ovary is consistent with its reported oogenesis phenotypes.

To determine where Sov is expressed in the ovary, we generated a GFP-tagged

transgene (GFP-Sov) sufficient to rescue ovary degeneration in *sov* mutants. Examination of

the distribution of GFP-Sov in developing egg chambers (Fig 2A) revealed Sov localization in

several cell types. By pairing localization analysis of Sov with antibodies recognizing Vasa

(Vas) to label the germline, we observed particularly striking nuclear localization of Sov

surrounded by perinuclear Vas in the germline cells within region 1 of the germarium, with

highest levels evident within the GSCs (dashed circles; Fig 2B). Using Traffic jam (Tj) to label

the soma highlighted nuclear enrichment of Sov in the Tj-positive somatic and follicle stem

cells (dashed ovals; Fig 2C). These data indicate that Sov is enriched in the germline and

somatic cells of the ovary, including both stem cell populations.

Sov is required in the soma and germline for oogenesis

To examine *sov* function in the germarium, we compared a strong allelic combination,

sov^{EA42}/sov^2, to cell-type specific knockdown of *sov* using a UAS short hairpin *sov* RNAi

construct (P{TRiP.HMC04875}; hereafter, sov^{RNAi}). The germaria of heterozygous *sov* females

are wild type (Fig 3A), but *sov* mutants often show an ovarian tumor phenotype, with greater

than the expected number of germ cells with dot spectrosomes (Fig 3B), indicating that germ

cells undergo complete cytokinesis. Ovarian tumors were observed in ~40% of *sov* mutants

(N=12/31 sov^{EA42}/sov^2 egg chambers versus N=0/47 controls). These findings are consistent

with GSC hyperproliferation and/or failed differentiation. We also observed tumors and nurse

cell nuclei residing within common follicles (Fig 3B) in about one-third of *sov* mutants (N=11/31

sov^{EA42}/sov^2 versus N=0/24 in controls). This phenotype occurs when follicle cells either fail to

separate egg chambers, or where those chambers fuse (Goode, Wright, and Mahowald 1992).

Consistently, disorganization of the follicle cell monolayer was also observed (dashed lines,

Fig 3B). In older *sov* ovaries, we also observed extensive cell death (Fig S1). Additionally, we

found that rare sov^{EA42}/Y males that escaped lethality had no germ cells (Fig S2). Thus, *sov*

functions widely.

To examine cell-type specific functions of *sov*, we used *tj-GAL4* to express sov^{RNAi} (or

$mCherry^{RNAi}$ controls) in somatic escort and follicle cells and *nanos-GAL4* (*nos-GAL4*) to

Benner et al.; *sov* represses gene expression

express RNAi in the germline. Relative to the controls, the *tj>sov^RNAi* germaria (Fig 3C, D) were

abnormal, showing ovarian tumor phenotypes similar to those seen in *sov* mutants. Germaria

were often filled with germline cells with dot spectrosomes and the follicle cells encroached

anteriorly. Similar results were also observed using other somatic drivers (*c587-GAL4* and *da-*

GAL4). Interestingly, *nos>sov^RNAi* did not impair germarium morphology (Fig 3E, F) and egg

chambers representing all 14 morphological stages of oogenesis appeared phenotypically

normal. However, eggs produced from *nos>sov^RNAi* mothers arrested during early

embryogenesis, indicating that maternal *sov* is required for embryonic development. While *sov*

expression patterns and germline RNAi suggest that *sov* is deployed in both the soma and

germline, Wayne et al. (1995) reported *sov* to be somatic cell dependent. To further address a

requirement for *sov* in the germline, we conducted germline clonal analysis. Depletion of *sov*

from the germline results in an agametic phenotype (Fig S3). Additionally, *sov^RNAi* resulted in

embryonic germline defects in a high-throughput study (Jankovics et al. 2014). Taken together,

these data indicate that there are both somatic and germline requirements for *sov* in

oogenesis.

To explore the differentiation of the germline further, we examined the distribution of

Bam, which is expressed in germarium region 1 germ cells (D. M. McKearin and Spradling

1990; D. McKearin and Ohlstein 1995; Ohlstein and McKearin 1997). In the absence of Bam,

germ cells hyperproliferate, resulting in tumors composed of 2-cell cysts. Given the prevalence

of undifferentiated tumorigenic germ cells upon somatic *sov^RNAi* knockdown, we asked if *sov*

has a nonautonomous role in promoting Bam expression in germ cells. Indeed, somatic

knockdown of *sov* results in significantly less Bam expressed in germ cells relative to controls

(Fig 3G, H; Fig S4). Both *tj-GAL4>sov^RNAi* and *c587-GAL4>sov^RNAi* resulted in a strong

Benner et al.*; sov* represses gene expression

decrease in Bam expression. Germline depletion of *sov* did not alter Bam expression (Fig S4).

The reduction in Bam expression in the germline is consistent with a nonautonomous role of

Sov function in the soma for differentiation signals directed to the germline.

To examine the defective follicle encapsulation of the germline cysts following *sov*

depletion, we used the oocyte-specific expression of Oo18 RNA binding protein (Orb) (Lantz et

al. 1994) to count the number of oocytes per cyst. Orb specifies the future single oocyte at the

posterior of the egg chamber in control egg chambers (Fig 3I, Fig S2D). Somatic depletion of

sov resulted in examples of both egg chambers with either too many oocytes (Fig 3J) or no

oocytes (Fig 3K, Fig S4). In a wild type 16-cell germline cyst, one of the two cells that has four

ring canals becomes the oocyte, and this feature can be used to determine if extra germline

divisions had occurred in cysts, or if multiple cysts were enveloped by the follicle cells. We saw

that the egg chambers with multiple Orb+ cells always had more than 16 germ cells, and in the

representative example shown, all three Orb+ cells had four ring canals, indicating that egg

chamber fusion had occurred (Fig 3J). Taken together, we conclude that Sov functions in the

soma to ensure proper differentiation of the germline.

Sov represses gene expression in the ovary

To provide mechanistic insights into the function of Sov, we performed transcriptome profiling

using triplicated Poly-A$^+$ RNA-seq analyses of ovaries from *sov*, *sov*^{RNAi}, and control females.

The gene expression profiles of ovaries from sterile females were markedly different from

controls, primarily due to derepression in the mutants (Fig 4A; Table S2). Among genes

showing differences in expression (FDR $p$adj < 0.05), we found 1,752 genes with >4-fold

increased expression in mutants, while there were only 172 genes with >4-fold decreased

Benner et al.*; sov* represses gene expression

expression (Table S3). To explore where these derepressed genes are normally expressed,

we examined their expression in other female tissues and in testes in a set of quadruplicated

RNA-seq experiments (Fig 4B). Many of the genes suppressed by Sov in ovaries were highly

expressed in other female tissues, but not testes, indicating that wild type Sov prevents ectopic

gene expression.

   To determine what types of genes are derepressed in *sov* mutants, we performed Gene

Ontology (GO)(Gene Ontology Consortium 2015) term analysis (Fig S5; Table S4). The genes

with lower expression in *sov* mutants had only a few significant GO terms that were

predominantly oogenic in nature (as expected given the general lack of mature eggs). For

example, there was poor expression of the chorion genes that are required to build the

eggshell (Orr-Weaver 1991), as well as genes required for follicle cell development. In

contrast, the derepressed genes in *sov* mutants showed a significant enrichment of cell

signaling GO terms, including neuronal communication. We found that many genes repressed

by Sov in the ovary, such as the *heartless* (*htl*) locus, which encodes a FGFR tyrosine kinase

receptor important for neuron/glia communication (Stork et al. 2014), are indeed expressed in

the head (Fig 4C). These results suggest that *sov* normally functions to repress a host of

signaling pathways. Derepressed signaling in *sov* mutants is likely catastrophic for

communication between various somatic cells and the germline during egg chamber

development and could explain the variety of mutant ovarian phenotypes.

   The general repressive role of *sov* was also revealed by a dramatic and coherent

elevation of transposon expression in the mutants (Fig 4D; Table S3). Of the 138 transposable

element classes detected in our gene expression profiling of *sov* mutants and controls, 91 had

increased expression (FDR $p$adj < 0.05) >4-fold in *sov* mutants, while none had >4-fold

Benner et al.*; sov* represses gene expression

decreased expression. The DNA, LINE, and LTR families of transposable elements were all

derepressed in *sov* mutants. Interestingly, somatic knockdown of *sov* resulted in the most

dramatic derepression of the *gypsy* and *copia* classes of transposons, which are *Drosophila*

retroviruses that develop in somatic cells and are exported to the developing germline

(Yoshioka et al. 1990). Thus, wild type *sov* may be important to protect the germline from

infection. Equally interesting, germline knockdown resulted in the greatest derepression of the

HeT-A, *TAHRE*, and *TART* transposons that compose the telomere (Mason, Frydrychova, and

Biessmann 2008). This finding suggests that the general role of *sov* in silencing transposon

expression includes control of both transposon functions necessary for normal cellular

metabolism, exemplified by the telomere transposons, as well as detrimental activities of

retrovirus-like elements. In addition to widespread deregulation of signaling and development

pathways in *sov* mutants, transposon dysgenesis may contribute to the severe ovarian

development defects in *sov* mutants.

Sov is a suppressor of PEV and colocalizes with HP1

The repressive function of Sov is reminiscent of HP1a function as a general repressor of gene

expression. HP1a was first characterized in *Drosophila* as a suppressor of PEV, a process that

reduces gene expression due to spreading of heterochromatin into a gene region (Clark and

Elgin 1992). To test the hypothesis that *sov* negatively regulates gene expression by

promoting heterochromatin formation, we examined the role of *sov* in PEV in the eye, where

patches of white$^+$ (red pigmented) and white$^-$ (not red pigmented) eye facets are easily

observed (Fig 5A). If, like HP1a, Sov represses gene expression by promoting

heterochromatin formation (Eissenberg et al. 1990), then it should suppress PEV, which is

Benner et al.*; sov* represses gene expression

scored by increased eye pigmentation. We obtained five different variegating w^+ transgene

insertions associated with either the heterochromatic pericentric region of chromosome arm 2L

or spread along the length of the heterochromatin-rich chromosome 4. In control animals,

these insertions show a characteristic eye variegation pattern (Fig 5B). Consistent with the

allelic strengths seen in previous experiments, the weak sov^2 allele did not suppress PEV (Fig

5C), but the stronger sov^{ML150}, sov^{EA42}, and *Df(1)sov* mutations dominantly suppressed PEV

(Fig 5D–F). These data demonstrate a role for Sov in heterochromatin formation.

The strong repressive function of Sov is reminiscent of HP1a. Prior in vitro work

suggests Sov may complex with HP1a (Alekseyenko et al. 2014). To determine if Sov and

HP1a associate in vivo, we followed HP1a-RFP and GFP-Sov localization in 1–2 hr live

embryos. During *Drosophila* embryogenesis, nuclei undergo rapid synchronous nuclear

divisions prior to cellularization at cleavage division/nuclear cycle (NC) 14 (Foe and Alberts

1983). HP1a and Sov colocalized in all NC 10–12 embryos we examined (Fig 6A; N=7

embryos). During prophase, both HP1a and Sov were enriched in regions of condensed DNA

(Fig 6A, 4:30). Whereas low levels of HP1a decorated DNA throughout division, Sov was

depleted during mitosis (Fig 6A, 10:00 and 11:10). Upon reentry into interphase, Sov

localization to nuclei resumed and was coincident with HP1a. Formation of heterochromatin is

contemporaneous with or slightly proceeds HP1a apical subnuclear localization in NC 14

(Rudolph et al. 2007; Yuan and O'Farrell 2016). At this stage, we observed a strong

colocalization of HP1a and Sov. Measuring the distribution of HP1a and Sov (Fig 6B, B')

confirmed high levels of colocalization within HP1a subnuclear domains (Fig 6C, shaded

region). These data support the idea that HP1a and Sov are deposited maternally where they

assemble into a complex.

Benner et al.; *sov* represses gene expression

Discussion

<u>Sov is a novel heterochromatin-associated protein</u>

Genomes contain large blocks of DNA of potentially mobile transposons and immobile mutated

derivatives (Vermaak and Malik 2009). The cell keeps these transposons from wreaking havoc

on the genome by actively suppressing their expression through condensation into

heterochromatin. However, some tightly regulated expression from heterochromatin is required

for normal cellular function. For example, the telomeres of *Drosophila* are maintained by

transposition of mobile elements from heterochromatic sites (Mason, Frydrychova, and

Biessmann 2008), and histone and rRNA genes are located within heterochromatic regions

(Yasuhara and Wakimoto 2006). Weakening of heterochromatin by suppressors of variegation

results in derepression of gene expression at the edges of heterochromatin blocks, suggesting

that the boundaries between repressed and active chromatin can expand and contract (Weiler

and Wakimoto 1995; Reuter and Spierer 1992). HP1a, encoded by *Su(var)*205, is a central

component of heterochromatin (Ebert et al. 2006) that shows the same strong suppression of

variegation that we observed in *sov/+* flies. Similarly, HP1a is also required to repress the

expression of transposons (Vermaak and Malik 2009).

Repression is often stable and emerging themes suggest a robust set of activities,

rather than a single component, maintain a chromatin state. For example, long-term repression

is stabilized with complexes, such as the repressive Polycomb Group (PcG) which provides an

epigenetic memory function (Kassis, Kennison, and Tamkun 2017). In order to create a stable

epigenetic state, PcG complexes have subunits that modify histones and bind these

modifications. Chromatin tethering of PcG complexes also involves DNA-binding proteins,

Benner et al.*; sov* represses gene expression

such as the YY1-like ZnF protein Pleiohomeotic (Pho), which further reinforce localization. The

DNA anchor proteins in PcG complexes are at least partially redundant with the histone-

binding components (J. L. Brown et al. 2003), suggesting that localization is robust due to

multiple independent localization mechanisms.

In mammals, HP1a also has both histone binding and DNA anchoring requirements.

DNA anchoring is provided by a family of KRAB ZnF proteins that have undergone a massive

radiation during evolution (Yang, Wang, and Macfarlan 2017; Ecco, Imbeault, and Trono

2017). KRAB proteins are essential for repression of transposons and more generally for a

properly regulated genome. The KRAB family members use a KRAB-Associated Protein

(KAP1) adapter protein to associate with HP1a and the SETDB1 methylase that modifies

histones to enable HP1a binding (Fig 7). Although KRAB proteins have not been identified in

Drosophila, the *Drosophila bonus* (*bon*) gene encodes a KAP1 homolog (Beckstead et al.

2005). *Drosophila* also has a SETDB1 encoded by *egg*, and loss of *egg* results in ovarian

phenotypes reminiscent of those observed in *sov* loss-of-function females (Clough et al. 2007;

Clough, Tedeschi, and Hazelrigg 2014; Wang et al. 2011). Like *sov* and *HP1a*, *bon* is a

modifier of variegation, raising the possibility that they collectively coordinate gene regulation.

In support of this hypothesis, Bon, Sov, and Egg were all identified in the same biochemical

complex as HP1a (Alekseyenko et al. 2014). It is tempting to speculate that the single very

long and ZnF-rich Sov protein plays the same direct binding role as the large family of KRAB

ZnF proteins in mammals (Fig 7). If Sov uses subsets of fingers to bind DNA, it could localize

to many different sequences in a combinatorial fashion. Additionally, the complex containing

Sov and HP1a also contains RNA (Alekseyenko et al. 2014) and could act as a tether via a

series of RNA intermediates as occurs in sex chromosome inactivation, for example (J. T. Lee

Benner et al.; *sov* represses gene expression

2009). As in the case of Pho in the PcG complexes, Sov might be a robustness factor rather

than an absolute requirement, as we did not observe gross delocalization of HP1a following

sov^{RNAi} (not shown). Further work will be required to fully understand the relationship between

Sov and HP1a.

<u>Repression by Sov promotes germline differentiation</u>

The fact that there have been many female sterile alleles of *sov* suggests that the ovary

is particularly sensitive to reduced *sov* activity. The female sterility phenotype is complex and

somewhat variable, but partial *sov* loss of function is characterized by somatic-dependent

differentiation defects. Stronger alleles also show a germline-dependent block in development

and lethality. While *sov* functions seem diverse, the locus expresses a single major isoform,

and Sov seems to be consistently located in the nucleus. We propose that the

multifunctionality of Sov may be attributed to a single mechanism wherein Sov helps control

heterochromatin formation required to repress ectopic gene expression and tissue-

inappropriate responses in the ovary and repress transposons (Fig 7).

We observed dramatic derepression of transposons in *sov* mutants. Consistent with our

observations, Czech et al (2013) reported a role of *sov* in transposon repression in a

genomewide RNAi screen. That work raised the possibility that transposon repression by Sov

occurs via the piRNA pathway (Brennecke et al. 2007; Yin and Lin 2007; Teixeira et al. 2017).

However, genes involved more strictly with transposable element regulation, such as Piwi, do

not have strong dosage effects on PEV (Gu and Elgin 2013), while *sov* and *HP1a*/*Su(var)205*

do. This distinction suggests that Sov has a more general effect on heterochromatin, rather

than specificity for transposon repression. Germline knockdown of HP1a results in a strong

Benner et al.; *sov* represses gene expression

derepression of telomeric transposons (Teo et al. 2018), just like Sov. This is suggested to be

due to a reduction in piRNAs specifically targeting these transposons and thus interplay

between the piRNA pathway, Sov, and HP1a remains possible.

<u>Conclusions</u>

Our data show that *sov* encodes a repressor of gene expression and transposons in the

ovary. In the absence of *sov*, genes that are normally expressed at very low levels in the

ovary, are activated. These same genes show dynamic patterns of expression in other adult

tissues, suggesting that they are derepressed in the absence of Sov. The failure to restrict

gene expression results in a range of phenotypes including ovarian tumors, defective

oogenesis, and tissue degeneration. These data support the idea that Sov encodes a novel

and essential protein that is generally repressive, likely via interactions with HP1a.

Materials and Methods

We have adopted the FlyBase-recommended resources table which includes all genetic,

biological, cell biology, genomics, manufactured reagents and algorithmic resources used in

this study (Table S5).

<u>Flies and genetics</u>

sov was defined by three X-linked, female sterile mutations including sov^2, which

mapped to ~19 cM (Mohler 1977; Mohler and Carroll 1984), refined to 6BD (Wayne *et al*.

1995). We used existing and four custom-made deletions (*Df(1)BSC276*, *BSC285*, *BSC286*

and *BSC297*) to map *sov* to the 4-gene *CG14438–shf* interval (Fig. 1; Table S5). *sov*

Benner et al.*; sov* represses gene expression

mutations complemented shf^2, but not $P\{SUPor-P\}CG14438^{KG00226}$ or $P\{GawB\}NP6070$,

suggesting $CG14438$ or $CR43496$ is sov, which we confirmed by generating $Df(1)sov$

(X:6756569..6756668;6770708) from FLP crossover between $P\{XP\}CG14438^{d07849}$ and

$PBac\{RB\}e03842$ $(Parks\ et\ al.\ 2004;\ Cook\ et\ al.\ 2012)$ to remove only those two genes.

We used the dominant female sterile technique for germline clones (Chou and Perrimon

1996). Test chromosomes, that were free of linked lethal mutations by male viability (sov^2) or

rescue by $Dp(1;3)DC486$ (sov^{EA42} and $Df(1)sov$), were recombined with $P\{ry^{+t7.2}=neoFRT\}19A$,

and verified by complementation tests and PCR. We confirmed $P\{ry^{+t7.2}=neoFRT\}19A$

functionality by crossing to $P\{w^{+mC}=GMR-hid\}SS1,\ y^1\ w^*\ P\{ry^{+t7.2}=neoFRT\}19A;\ P\{w^{+m^*}=GAL4-$

$ey.H\}SS5,\ P\{w^{+mC}=UAS-FLP.D\}JD2$ and scoring for large eye size. We crossed females with

FRT chromosomes to $P\{w^{+mC}=ovoD1-18\}P4.1,\ P\{ry^{+t7.2}=hsFLP\}12,\ y^1\ w^{1118}\ sn^3$

$P\{ry^{+t7.2}=neoFRT\}19A/Y$ males for 24 hours of egg laying at 25^oC. We heat-shocked for 1hr at

37^oC on days 2 and 3. We dissected females (5d posteclosion) to score for ovo^{D1} or wild type

morphology.

We generated $PBac\{GFP-sov\}$ from P[acman] BAC clone CH322-191E24 (X:6753282–

6773405) (Venken et al. 2009) grown in the SW102 strain (Warming et al. 2005). In step one,

we integrated the positive/negative marker CP6-RpsL/Kan (CP6 promoter with a bi-cistronic

cassette encoding the RpsL followed by the Kan), PCR-amplified with primers N-CG14438-

CP6-RN-F and -R between the first two codons of sov, and selected (15 µg/ml Kanamycin)

We integrated at the galK operon in DH10B bacteria using mini-lambda-mediated

recombineering (Court et al. 2003). We amplified DH10B::CP6-RpsL/Kan DNA using primers

N-CG14438-CP6-RN-F and -R. Correct events were identified by PCR, as well as resistance

(15 µg/ml Kanamycin) and sensitivity (250 µg/ml Strep). In step two, we replaced the selection

Benner et al.; *sov* represses gene expression

markers with a multi-tag sequence (Venken et al. 2011), tailored for N-terminal tagging (N-tag)

and counter-selected (250 µg/ml Strep). The N-tag (3xFlag tag, TEV protease site, StrepII tag,

superfolder GFP, FlAsH tetracysteine tag, and flexible 4xGlyGlySer (GGS) linker) was

Drosophila codon optimized in a R6Kγ plasmid. We transformed plasmid into EPI300 for copy

number amplification. We confirmed correct events by PCR and Sanger DNA sequencing.

Tagged P[acman] BAC clone DNA was injected into *y^1 M{vas-int.Dm}ZH-2A w*; PBac{y+-attP-*

3B}VK00033 embryos, resulting in *w^{1118}; PBac{y[+mDint2] w^{+mC}=FTSF.GFP-sov}VK00033*.

<u>Microscopy</u>

We fixed ovaries in 4 or 5% EM-grade paraformaldehyde in PBS containing 0.1 or 0.3%

Triton X-100 (PBTX) for 10–15min, washed 3x 15 min in PBTX, and blocked >30min in 2%

normal goat serum, 0.5-1% bovine serum albumin (BSA) in PBS with 0.1% Tween-20 or 0.1%

Triton X-100. Antibodies and DAPI were diluted into blocking buffer. We incubated in primary

antibodies overnight at 4^{o}C and secondaries 2–3hr at room temperature. Embryos (1–2 hr)

were prepared for live imaging in halocarbon oil (Lerit et al. 2015). We imaged ovaries and

embryos using a Nikon Ti-E system or Zeiss LSM 780 microscope and eyes with a Nikon SMZ.

<u>DNA-Seq</u>

Genomic DNA was extracted from 30 whole flies per genotype (Huang, Rehm, and

Rubin 2009)(Sambrook and Russell 2006) to prepare DNA-seq libraries (Nextera DNA Library

Preparation Kit). We used 50 bp, single-end sequencing (Illumina HiSeq 2500, CASAVA base

calling). Sequence data are available at the SRA (SRP14438). We mapped DNAseq reads to

FlyBase r6.16 genome with Hisat2 (-k 1 --no-spliced-alignment)(Kim, Langmead, and Salzberg

Benner et al.; *sov* represses gene expression

2015). We used mpileup and bcftools commands from SAMtools within the genomic region

X:6756000–6771000 (H. Li et al. 2009; H. Li 2011) for variant calling and snpEFF to determine

the nature of variants in *sov* mutants (Cingolani et al. 2012).

<u>RNA-seq</u>

Stranded PolyA+ RNA-seq libraries from *sov* and control ovaries (Table S5) were

created (H. Lee, Cho, et al. 2016) and are available at GEO (GSE113977). We extracted total

RNA (Qiagen RNeasy Mini Kit) in biological triplicate from 15 ovaries (4–5d posteclosion) and

used 200 ng with 10 pg of ERCC spike-in control RNAs (pools 78A or 78B) for libraries (Jiang

et al. 2011; Zook et al. 2012; Pine et al. 2016; H. Lee, Pine, et al. 2016). We used 50 bp,

single-end sequencing as above. Tissue expression analysis are from GEO accession

GSE99574, a resource for comparing gene expression patterns.

We mapped RNA-seq reads to FlyBase r6.21 with Hisat2 (-k 1 --rna-strandness R --dta)

(Kim, Langmead, and Salzberg 2015). We determined read counts for each attribute of the

FlyBase r6.21 GTF file (with ERCC and transposable element sequences), with HTSeq-count

(Anders, Pyl, and Huber 2015). Transposon sequences were from the UCSC Genome

Browser RepeatMasker track (Casper et al. 2018; Smit, Hubley, and Green 2013-2015).

We conducted differential expression analysis with DESeq2 (pAdjustMethod = "fdr")

(Love, Huber, and Anders 2014). We removed genes with read counts <1 and read counts for

transposable elements with >1 locations were summed for the DESeq2 analysis. *Df(1)sov/sov^2*

replicate 3 and *sov^2/w^{1118}* replicate 1 were failed. For *sov* mutant *vs.* control DESeq2 analysis,

all *sov* mutants were compared to all wild type controls. For tissue types, we compared each

sexed tissue to sexed whole organism. We used reads per kb per million reads (RPKM) for

gene-level expression.

For heatmaps, we calculated Euclidean distance, performed hierarchical cluster

analysis (agglomeration method = Ward), and mean-subtracted scaled across genotypes.

Mean sample RPKM correlation values (Table S1) were calculated by cor.test() function with

Pearson correlation coefficient in R (R Core Team 2017). We represented derepressed genes

as mean-subtracted scaled values across tissues in the heatmap (Table S6).

For read density tracks, replicate raw read files were combined. Bedgraph files were

created with bedtools genomecov (Quinlan and Hall 2010) visualized on the UCSC genome

browser (Kent et al. 2002). Tracks were scaled by the number of reads divided by total reads

per million.

We used ClueGO (Bindea et al. 2009), with Cytoscape (Shannon et al. 2003) for one-

sided enrichment analysis (see Table S4).

Images were assembled using ImageJ and Photoshop. We used ROI tool in ImageJ,

plotted/analyzed image data (Microsoft Excel and GraphPad Prism), and calculated

significance by D'Agnostino and Pearson normality tests, followed Student's two-tailed t-test or

Mann-Whitney tests.

Acknowledgements

Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) and

Drosophila Genomics and Genetic Resources at the Kyoto Institute of Technology were used

in this study. Monoclonal antibodies were obtained from the Developmental Studies Hybridoma

Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department

Benner et al.; *sov* represses gene expression

of Biology, Iowa City, IA 52242. Sequencing was performed by the NIDDK Genomics Core,

under the direction of Harold Smith. Genetic and genomic information was obtained from

FlyBase. This work utilized the computational resources of the NIH HPC Biowulf cluster

(http://hpc.nih.gov). We thank BACPAC resources for plasmids, Norbert Perrimon for the

sov^{EA42} stock and Don Court, Neal Copeland, and Nancy Jenkins for the recombineering

strain. We thank Astrid Haase, Nasser Rusan, David Katz, and members of the Oliver and

Lerit labs for stimulating discussions. Particular thanks go to Stacy Christensen and Kimberley

Cook for generating and characterizing the *sov* deletions. This research was supported in part

by the Intramural Research Program of the NIH, National Institute of Diabetes and Digestive

and Kidney Diseases (NIDDK) to BO. DAL and EAC were supported by NIH grant

5K22HL126922-02 (DAL). LB was supported by the NIH Graduate Partners Program. KJTV

was supported by Baylor College of Medicine, the Albert and Margaret Alkek Foundation, the

McNair Medical Institute at The Robert and Janice McNair Foundation, the March of Dimes

Foundation (#1-FY14-315), the Foundation for Angelman Syndrome Therapeutics (FT2016-

002), the Cancer Prevention and Research Institute of Texas (R1313), and NIH grants

(1R21HG006726, 1R21GM110190, 1R21OD022981, and R01GM109938). CW and KRC were

supported by the Office of the NIH Director, the National Institute of General Medical Sciences

and the National Institute of Child Health and Human Development (P40OD018537). KRC was

supported by the National Center for Resource Resources (R24RR014106) and the National

Science Foundation (DBI-9816125).

Author contributions

Benner et al.; *sov* represses gene expression

484   LB, KC, BO, and DAL designed the project and analyzed data. LB, KC, CW, and KJTV

485   performed genetics. LB and HY performed genomics. LB, EAC, CW and DAL performed

486   imaging. LB, BO, and DAL wrote the manuscript. All authors reviewed data and provided

487   feedback on the manuscript.

489
Figure 1. *sov* is *CG14438*. (A) Genes of the genomic interval X:6710000–6810000 (Gramates
et al. 2017). (B) Deficiency (*Df*) and Duplication (*Dp*) mapping. Noncomplementing (*sov⁻*, red)
and complementing (*sov⁺*, green) alleles and rearrangements are shown. (C) Schematic of the
CG14438 (black) and *CR43496* (purple) genes. Transcription start (bent arrows), introns (thin
lines) non-coding regions (medium lines) and coding regions (thick lines) are shown.
Transposon insertions (triangles), point mutations (red lines), the region targeted by the
shRNAi transgene (base-paired), and Sov protein features are shown. (D) RNA expression
tracks by tissue type from female (red) or male (blue) adults.

Benner et al.; *sov* represses gene expression

498
Figure 2. Sov germarium expression. (A) Cartoon showing germarium regions I-III and
young egg chamber with cell types labeled. (B,C) 1d posteclosion, visualized for GFP-Sov
(green), anti-αSpectrin (red). Images show single optical sections. (B) Sov contrasted with anti-
Vas germline (magenta), or (C) -Tj somatic staining (magenta). GFP-Sov expression regions of
interest (see text) are shown (dashed lines). Bars: 10μm.

Benner et al.; *sov* represses gene expression

504
Figure 3. *sov* mutant defects. Immunofluorescence for the indicated probes in the noted
genotypes. (A–F) Single optical sections of germaria (4–5d posteclosion) stained with anti-Vas
(blue), -Tj (green), -αSpectrin (red), and DAPI (white) with dot spectrosomes (arrows) and
displaced follicle cells (dashed lines) shown. (G, H) Single optical sections of germaria (1d
posteclosion) stained with anti-Bam (green), -actin (red), and DAPI (blue). Bam region of
interest is shown (dashed circles). (I–K) Maximum intensity projections of germaria (1d
posteclosion) stained with anti-Orb (green), -actin (red), and DAPI (blue). Orb+ cells shown (*).
Bars: (A–F) 20μm, (G–K) 10μm.
513

514
Figure 4. *sov* mutant transcriptome. (A) Relative expression in *sov⁻* vs *sov⁺* ovaries plotted
against the mean expression in both sample types. Units are log₂ normalized read counts
(NRC). Data points are genes, those with >4-fold change (log₂ 2, red) or <–4 fold change (log₂
-2, blue) and FDR *p*adj value <0.05 are highlighted. (B) Tissue-biased expression in wild type
tissues for genes derepressed in *sov* mutant ovaries. Heatmap from mean-subtracted ratios
scaled across tissues (red=higher; blue=lower). (C) RNA-seq normalized read densities of the
heartless locus. (D) Transposable element expression in *sov* mutants (sterile) and controls
(fertile). Heatmap from mean-subtracted reads (in RPKM. red=higher; blue=lower) scaled for
each transposable element (rows) across genotypes (columns). Transposon classes for DNA,
non-LTR (Jockey), telomeric repeat (Telo), and LTR (Gypsy, Copia, and PAO) are indicated.

Benner et al.; *sov* represses gene expression

Benner_Fig5

Figure 5. Sov is a dominant suppressor of position effect variegation. (A) Cartoon of position-effect variegation (PEV) in the eye. Expression of the *white* gene (bent arrow, thick bar, red) can be silenced (white) by proximal heterochromatin (squiggled) spreading. (B–F) Eyes from adults of the indicated genotypes (columns) with variegated expression of *P{hsp26-pt-T}* transgenes inserted into the indicated chromosomal positions (rows).

Benner et al.; *sov* represses gene expression

Figure 6. Sov colocalizes with HP1a. Stills from live imaging of embryos expressing GFP-Sov and RFP-HP1a. (A) Localization of GFP-Sov and RFP-HP1a (rows) in an embryo progressing from NC 10 to 11 with cell cycle stages (columns, nuclear envelope breakdown=NEB) and time (min:s) shown. (B) NC 14 embryo. Boxed regions are magnified in insets below. An HP1a subnuclear domain is shown (arrows). (B') Single optical section containing peak HP1a fluorescence of inset from (B). Dashed line indicates region used for histogram analysis. (A,B) Maximum projections through 1.5 µm volume at 1F/30s. Bars: 10 µm; insets, 5 µm. (C) Histogram of HP1a and Sov fluorescence intensity measured in (B'). Fluorescence levels (arbitrary units) normalized to the peak fluorescence intensity for each channel and the distance (µm) to peak HP1a signal. Half maximum HP1a fluorescence is shaded (yellow).

DNA tether   **Histone tether**

545
546 **Figure 7. Working model of *sov* function.** Sov protein function and attributes are analogous
547 to the mammalian KRAB proteins (green). Like KRAB proteins, Sov is in complex with HP1a
548 (yellow), Bon (mammalian KAP-1, orange) and Egg (mammalian SETDB1, black) providing a
549 DNA tether to stabilize a repressed chromatin state.
550

References

Alekseyenko, Artyom A., Andrey A. Gorchakov, Barry M. Zee, Stephen M. Fuchs, Peter V. Kharchenko, and Mitzi I. Kuroda. 2014. "Heterochromatin-Associated Interactions of Drosophila HP1a with dADD1, HIPP1, and Repetitive RNAs." *Genes & Development* 28 (13): 1445–60.

Anders, Simon, Paul Theodor Pyl, and Wolfgang Huber. 2015. "HTSeq--a Python Framework to Work with High-Throughput Sequencing Data." *Bioinformatics* 31 (2): 166–69.

Barton, Lacy J., Kaylee E. Lovander, Belinda S. Pinto, and Pamela K. Geyer. 2016. "Drosophila Male and Female Germline Stem Cell Niches Require the Nuclear Lamina Protein Otefin." *Developmental Biology* 415 (1): 75–86.

Bausek, Nina. 2013. "JAK-STAT Signaling in Stem Cells and Their Niches in Drosophila." *JAK-STAT* 2 (3): e25686.

Beckstead, R. B., S. S. Ner, K. G. Hales, T. A. Grigliatti, B. S. Baker, and H. J. Bellen. 2005. "Bonus, a Drosophila TIF1 Homolog, Is a Chromatin-Associated Protein That Acts as a Modifier of Position-Effect Variegation." *Genetics* 169 (2): 783–94.

Bindea, Gabriela, Bernhard Mlecnik, Hubert Hackl, Pornpimol Charoentong, Marie Tosolini, Amos Kirilovsky, Wolf-Herman Fridman, Franck Pagès, Zlatko Trajanoski, and Jérôme Galon. 2009. "ClueGO: A Cytoscape Plug-in to Decipher Functionally Grouped Gene Ontology and Pathway Annotation Networks." *Bioinformatics* 25 (8): 1091–93.

Börner, Kenneth, Dhawal Jain, Paula Vazquez-Pianzola, Sandra Vengadasalam, Natascha Steffen, Dmitry V. Fyodorov, Pavel Tomancak, Alexander Konev, Beat Suter, and Peter B. Becker. 2016. "A Role for Tuned Levels of Nucleosome Remodeler Subunit ACF1 during Drosophila Oogenesis." *Developmental Biology* 411 (2): 217–30.

Brennecke, Julius, Alexei A. Aravin, Alexander Stark, Monica Dus, Manolis Kellis, Ravi Sachidanandam, and Gregory J. Hannon. 2007. "Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in Drosophila." *Cell* 128 (6): 1089–1103.

Brown, James B., Nathan Boley, Robert Eisman, Gemma E. May, Marcus H. Stoiber, Michael O. Duff, Ben W. Booth, et al. 2014. "Diversity and Dynamics of the Drosophila Transcriptome." *Nature* 512 (7515): 393–99.

Brown, J. Lesley, Cornelia Fritsch, Jürg Mueller, and Judith A. Kassis. 2003. "The Drosophila Pho-like Gene Encodes a YY1-Related DNA Binding Protein That Is Redundant with Pleiohomeotic in Homeotic Gene Silencing." *Development* 130 (2): 285–94.

Casper, Jonathan, Ann S. Zweig, Chris Villarreal, Cath Tyner, Matthew L. Speir, Kate R. Rosenbloom, Brian J. Raney, et al. 2018. "The UCSC Genome Browser Database: 2018 Update." *Nucleic Acids Research* 46 (D1): D762–69.

Chen, Shuyi, Su Wang, and Ting Xie. 2011. "Restricting Self-Renewal Signals within the Stem Cell Niche: Multiple Levels of Control." *Current Opinion in Genetics & Development* 21 (6): 684–89.

Chen, Taiping, and Sharon Y. R. Dent. 2014. "Chromatin Modifiers and Remodellers: Regulators of Cellular Differentiation." *Nature Reviews. Genetics* 15 (2): 93–106.

Chou, T. B., and N. Perrimon. 1996. "The Autosomal FLP-DFS Technique for Generating Germline Mosaics in Drosophila Melanogaster." *Genetics* 144 (4): 1673–79.

Cingolani, Pablo, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan Wang, Susan J. Land, Xiangyi Lu, and Douglas M. Ruden. 2012. "A Program for Annotating and Predicting the Effects of Single Nucleotide Polymorphisms, SnpEff: SNPs in the Genome of Drosophila Melanogaster Strain w1118; Iso-2; Iso-3." *Fly* 6 (2): 80–92.

Clark, R. F., and S. C. Elgin. 1992. "Heterochromatin Protein 1, a Known Suppressor of Position-Effect Variegation, Is Highly Conserved in Drosophila." *Nucleic Acids Research* 20 (22): 6067–74.

Clough, Emily, Woongjoon Moon, Shengxian Wang, Kathleen Smith, and Tulle Hazelrigg. 2007. "Histone Methylation Is Required for Oogenesis in Drosophila." *Development* 134 (1): 157–65.

Clough, Emily, Thomas Tedeschi, and Tulle Hazelrigg. 2014. "Epigenetic Regulation of Oogenesis and Germ Stem Cell Maintenance by the Drosophila Histone Methyltransferase Eggless/dSetDB1."

Benner et al.*; sov* represses gene expression

Developmental Biology 388 (2): 181–91.

Cook, R. Kimberley, Stacey J. Christensen, Jennifer A. Deal, Rachel A. Coburn, Megan E. Deal, Jill M. Gresens, Thomas C. Kaufman, and Kevin R. Cook. 2012. "The Generation of Chromosomal Deletions to Provide Extensive Coverage and Subdivision of the Drosophila Melanogaster Genome." *Genome Biology* 13 (3): R21.

Court, Donald L., Srividya Swaminathan, Daiguan Yu, Helen Wilson, Teresa Baker, Mikail Bubunenko, James Sawitzke, and Shyam K. Sharan. 2003. "Mini-Lambda: A Tractable System for Chromosome and BAC Engineering." *Gene* 315 (October): 63–69.

Di Stefano, Luisa, Jun-Yuan Ji, Nam-Sung Moon, Anabel Herr, and Nicholas Dyson. 2007. "Mutation of Drosophila Lsd1 Disrupts H3-K4 Methylation, Resulting in Tissue-Specific Defects during Development." *Current Biology: CB* 17 (9): 808–12.

Ebert, Anja, Sandro Lein, Gunnar Schotta, and Gunter Reuter. 2006. "Histone Modification and the Control of Heterochromatic Gene Silencing in Drosophila." *Chromosome Research: An International Journal on the Molecular, Supramolecular and Evolutionary Aspects of Chromosome Biology* 14 (4): 377–92.

Ecco, Gabriela, Michael Imbeault, and Didier Trono. 2017. "KRAB Zinc Finger Proteins." *Development* 144 (15): 2719–29.

Eissenberg, J. C., T. C. James, D. M. Foster-Hartnett, T. Hartnett, V. Ngan, and S. C. Elgin. 1990. "Mutation in a Heterochromatin-Specific Chromosomal Protein Is Associated with Suppression of Position-Effect Variegation in Drosophila Melanogaster." *Proceedings of the National Academy of Sciences of the United States of America* 87 (24): 9923–27.

Elgin, Sarah C. R., and Gunter Reuter. 2013. "Position-Effect Variegation, Heterochromatin Formation, and Gene Silencing in Drosophila." *Cold Spring Harbor Perspectives in Biology* 5 (8): a017780.

Eliazer, Susan, Victor Palacios, Zhaohui Wang, Rahul K. Kollipara, Ralf Kittler, and Michael Buszczak. 2014. "Lsd1 Restricts the Number of Germline Stem Cells by Regulating Multiple Targets in Escort Cells." *PLoS Genetics* 10 (3): e1004200.

Eliazer, Susan, Nevine A. Shalaby, and Michael Buszczak. 2011. "Loss of Lysine-Specific Demethylase 1 Nonautonomously Causes Stem Cell Tumors in the Drosophila Ovary." *Proceedings of the National Academy of Sciences of the United States of America* 108 (17): 7064–69.

Foe, V. E., and B. M. Alberts. 1983. "Studies of Nuclear and Cytoplasmic Behaviour during the Five Mitotic Cycles That Precede Gastrulation in Drosophila Embryogenesis." *Journal of Cell Science* 61 (May): 31–70.

Fuller, Margaret T., and Allan C. Spradling. 2007. "Male and Female Drosophila Germline Stem Cells: Two Versions of Immortality." *Science* 316 (5823): 402–4.

Gene Ontology Consortium. 2015. "Gene Ontology Consortium: Going Forward." *Nucleic Acids Research* 43 (Database issue): D1049–56.

Gilboa, Lilach. 2015. "Organizing Stem Cell Units in the Drosophila Ovary." *Current Opinion in Genetics & Development* 32 (June): 31–36.

Goode, S., D. Wright, and A. P. Mahowald. 1992. "The Neurogenic Locus Brainiac Cooperates with the Drosophila EGF Receptor to Establish the Ovarian Follicle and to Determine Its Dorsal-Ventral Polarity." *Development* 116 (1): 177–92.

Gramates, L. Sian, Steven J. Marygold, Gilberto Dos Santos, Jose-Maria Urbano, Giulia Antonazzo, Beverley B. Matthews, Alix J. Rey, et al. 2017. "FlyBase at 25: Looking to the Future." *Nucleic Acids Research* 45 (D1): D663–71.

Graveley, Brenton R., Angela N. Brooks, Joseph W. Carlson, Michael O. Duff, Jane M. Landolin, Li Yang, Carlo G. Artieri, et al. 2011. "The Developmental Transcriptome of Drosophila Melanogaster." *Nature* 471 (7339): 473–79.

Gu, Tingting, and Sarah C. R. Elgin. 2013. "Maternal Depletion of Piwi, a Component of the RNAi System, Impacts Heterochromatin Formation in Drosophila." *PLoS Genetics* 9 (9): e1003780.

Hsieh, M., Y. Tintut, and J. D. Gralla. 1994. "Functional Roles for the Glutamines within the Glutamine-Rich Region of the Transcription Factor Sigma 54." *The Journal of Biological Chemistry* 269 (1):

373–78.

Huang, Audrey M., E. Jay Rehm, and Gerald M. Rubin. 2009. "Quick Preparation of Genomic DNA from Drosophila." *Cold Spring Harbor Protocols* 2009 (4): db.prot5198.

James, T. C., and S. C. Elgin. 1986. "Identification of a Nonhistone Chromosomal Protein Associated with Heterochromatin in Drosophila Melanogaster and Its Gene." *Molecular and Cellular Biology* 6 (11): 3862–72.

Jankovics, Ferenc, László Henn, Ágnes Bujna, Péter Vilmos, Kerstin Spirohn, Michael Boutros, and Miklós Erdélyi. 2014. "Functional Analysis of the Drosophila Embryonic Germ Cell Transcriptome by RNA Interference." *PloS One* 9 (6): e98579.

Jenuwein, T., and C. D. Allis. 2001. "Translating the Histone Code." *Science* 293 (5532): 1074–80.

Jiang, Lichun, Felix Schlesinger, Carrie A. Davis, Yu Zhang, Renhua Li, Marc Salit, Thomas R. Gingeras, and Brian Oliver. 2011. "Synthetic Spike-in Standards for RNA-Seq Experiments." *Genome Research* 21 (9): 1543–51.

Kassis, Judith A., James A. Kennison, and John W. Tamkun. 2017. "Polycomb and Trithorax Group Genes in Drosophila." *Genetics* 206 (4): 1699–1725.

Kent, W. James, Charles W. Sugnet, Terrence S. Furey, Krishna M. Roskin, Tom H. Pringle, Alan M. Zahler, and David Haussler. 2002. "The Human Genome Browser at UCSC." *Genome Research* 12 (6): 996–1006.

Kim, Daehwan, Ben Langmead, and Steven L. Salzberg. 2015. "HISAT: A Fast Spliced Aligner with Low Memory Requirements." *Nature Methods* 12 (4): 357–60.

Kouzarides, Tony. 2007. "Chromatin Modifications and Their Function." *Cell* 128 (4): 693–705.

Lantz, V., J. S. Chang, J. I. Horabin, D. Bopp, and P. Schedl. 1994. "The Drosophila Orb RNA-Binding Protein Is Required for the Formation of the Egg Chamber and Establishment of Polarity." *Genes & Development* 8 (5): 598–613.

Leader, David P., Sue A. Krause, Aniruddha Pandit, Shireen A. Davies, and Julian A. T. Dow. 2018. "FlyAtlas 2: A New Version of the Drosophila Melanogaster Expression Atlas with RNA-Seq, miRNA-Seq and Sex-Specific Data." *Nucleic Acids Research* 46 (D1): D809–15.

Lee, Hangnoh, Dong-Yeon Cho, Cale Whitworth, Robert Eisman, Melissa Phelps, John Roote, Thomas Kaufman, et al. 2016. "Effects of Gene Dose, Chromatin, and Network Topology on Expression in Drosophila Melanogaster." *PLoS Genetics* 12 (9): e1006295.

Lee, Hangnoh, P. Scott Pine, Jennifer McDaniel, Marc Salit, and Brian Oliver. 2016. "External RNA Controls Consortium Beta Version Update." *Journal of Genomics* 4 (July): 19–22.

Lee, Jeannie T. 2009. "Lessons from X-Chromosome Inactivation: Long ncRNA as Guides and Tethers to the Epigenome." *Genes & Development* 23 (16): 1831–42.

Lerit, Dorothy A., Holly A. Jordan, John S. Poulton, Carey J. Fagerstrom, Brian J. Galletta, Mark Peifer, and Nasser M. Rusan. 2015. "Interphase Centrosome Organization by the PLP-Cnn Scaffold Is Required for Centrosome Function." *The Journal of Cell Biology* 210 (1): 79–97.

Li, Heng. 2011. "A Statistical Framework for SNP Calling, Mutation Discovery, Association Mapping and Population Genetical Parameter Estimation from Sequencing Data." *Bioinformatics* 27 (21): 2987–93.

Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo Abecasis, Richard Durbin, and 1000 Genome Project Data Processing Subgroup. 2009. "The Sequence Alignment/Map Format and SAMtools." *Bioinformatics* 25 (16): 2078–79.

Lin, H., and A. C. Spradling. 1995. "Fusome Asymmetry and Oocyte Determination in Drosophila." *Developmental Genetics* 16 (1): 6–12.

Li, Xuanying, Christopher W. Seidel, Leanne T. Szerszen, Jeffrey J. Lange, Jerry L. Workman, and Susan M. Abmayr. 2017. "Enzymatic Modules of the SAGA Chromatin-Modifying Complex Play Distinct Roles in Drosophila Gene Expression and Development." *Genes & Development* 31 (15): 1588–1600.

Lorch, Yahli, and Roger D. Kornberg. 2017. "Chromatin-Remodeling for Transcription." *Quarterly Reviews of Biophysics* 50 (January): e5.

Benner et al.*; sov* represses gene expression

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change
and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.
Mason, James M., Radmila Capkova Frydrychova, and Harald Biessmann. 2008. "Drosophila
Telomeres: An Exception Providing New Insights." *BioEssays: News and Reviews in Molecular,*
Cellular and Developmental Biology 30 (1): 25–37.
McConnell, Kristopher H., Michael Dixon, and Brian R. Calvi. 2012. "The Histone Acetyltransferases
CBP and Chameau Integrate Developmental and DNA Replication Programs in Drosophila
Ovarian Follicle Cells." *Development* 139 (20): 3880–90.
McKearin, D. M., and A. C. Spradling. 1990. "Bag-of-Marbles: A Drosophila Gene Required to Initiate
Both Male and Female Gametogenesis." *Genes & Development* 4 (12B): 2242–51.
McKearin, D., and B. Ohlstein. 1995. "A Role for the Drosophila Bag-of-Marbles Protein in the
Differentiation of Cystoblasts from Germline Stem Cells." *Development* 121 (9): 2937–47.
Mohler, Dawson, and Andrea Carroll. 1984. "Report of Dawson Mohler and Andrea Carroll." *Drosophila*
Information Service 60 (June): 236–41.
Mohler, J. D. 1977. "Developmental Genetics of the Drosophila Egg. I. Identification of 59 Sex-Linked
Cistrons with Maternal Effects on Embryonic Development." *Genetics* 85 (2): 259–72.
Ohlstein, B., and D. McKearin. 1997. "Ectopic Expression of the Drosophila Bam Protein Eliminates
Oogenic Germline Stem Cells." *Development* 124 (18): 3651–62.
Orr-Weaver, T. L. 1991. "Drosophila Chorion Genes: Cracking the Eggshell's Secrets." *BioEssays:*
News and Reviews in Molecular, Cellular and Developmental Biology 13 (3): 97–105.
Parks, Annette L., Kevin R. Cook, Marcia Belvin, Nicholas A. Dompe, Robert Fawcett, Kari Huppert,
Lory R. Tan, et al. 2004. "Systematic Generation of High-Resolution Deletion Coverage of the
Drosophila Melanogaster Genome." *Nature Genetics* 36 (3): 288–92.
Peng, Jamy C., Anton Valouev, Na Liu, and Haifan Lin. 2016. "Piwi Maintains Germline Stem Cells and
Oogenesis in Drosophila through Negative Regulation of Polycomb Group Proteins." *Nature*
Genetics 48 (3): 283–91.
Pine, P. Scott, Sarah A. Munro, Jerod R. Parsons, Jennifer McDaniel, Anne Bergstrom Lucas, Jean
Lozach, Timothy G. Myers, Qin Su, Sarah M. Jacobs-Helber, and Marc Salit. 2016. "Evaluation of
the External RNA Controls Consortium (ERCC) Reference Material Using a Modified Latin Square
Design." *BMC Biotechnology* 16 (1): 54.
Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing
Genomic Features." *Bioinformatics* 26 (6): 841–42.
R Core Team. 2017. "A Language and Environment for Statistical Computing." Vienna, Austria: R
Foundation for Statistical Computing. https://www.R-project.org/.
Reuter, G., and P. Spierer. 1992. "Position Effect Variegation and Chromatin Proteins." *BioEssays:*
News and Reviews in Molecular, Cellular and Developmental Biology 14 (9): 605–12.
Robinson, Scott W., Pawel Herzyk, Julian A. T. Dow, and David P. Leader. 2013. "FlyAtlas: Database
of Gene Expression in the Tissues of Drosophila Melanogaster." *Nucleic Acids Research* 41
(Database issue): D744–50.
Rudolph, Thomas, Masato Yonezawa, Sandro Lein, Kathleen Heidrich, Stefan Kubicek, Christiane
Schäfer, Sameer Phalke, et al. 2007. "Heterochromatin Formation in Drosophila Is Initiated through
Active Removal of H3K4 Methylation by the LSD1 Homolog SU(VAR)3-3." *Molecular Cell* 26 (1):
103–15.
Sambrook, Joseph, and David W. Russell. 2006. "Purification of Nucleic Acids by Extraction with
Phenol:chloroform." *CSH Protocols* 2006 (1). https://doi.org/10.1101/pdb.prot4455.
Shannon, Paul, Andrew Markiel, Owen Ozier, Nitin S. Baliga, Jonathan T. Wang, Daniel Ramage, Nada
Amin, Benno Schwikowski, and Trey Ideker. 2003. "Cytoscape: A Software Environment for
Integrated Models of Biomolecular Interaction Networks." *Genome Research* 13 (11): 2498–2504.
Smit, Afa, R. Hubley, and P. Green. 2013-2015. "RepeatMasker Open-4.0."
http://www.repeatmasker.org.
Soshnev, Alexey A., Ryan M. Baxley, J. Robert Manak, Kai Tan, and Pamela K. Geyer. 2013. "The

Benner et al.; *sov* represses gene expression

Insulator Protein Suppressor of Hairy-Wing Is an Essential Transcriptional Repressor in the
Drosophila Ovary." *Development* 140 (17): 3613–23.
Stork, Tobias, Amy Sheehan, Ozge E. Tasdemir-Yilmaz, and Marc R. Freeman. 2014. "Neuron-Glia
Interactions through the Heartless FGF Receptor Signaling Pathway Mediate Morphogenesis of
Drosophila Astrocytes." *Neuron* 83 (2): 388–403.
Teixeira, Felipe Karam, Martyna Okuniewska, Colin D. Malone, Rémi-Xavier Coux, Donald C. Rio, and
Ruth Lehmann. 2017. "piRNA-Mediated Regulation of Transposon Alternative Splicing in the Soma
and Germ Line." *Nature* 552 (7684): 268–72.
Teo, Ryan Yee Wei, Amit Anand, Vishweshwaren Sridhar, Katsutomo Okamura, and Toshie Kai. 2018.
"Heterochromatin Protein 1a Functions for piRNA Biogenesis Predominantly from Pericentric and
Telomeric Regions in Drosophila." *Nature Communications* 9 (1): 1735.
Venken, Koen J. T., Joseph W. Carlson, Karen L. Schulze, Hongling Pan, Yuchun He, Rebecca
Spokony, Kenneth H. Wan, et al. 2009. "Versatile P[acman] BAC Libraries for Transgenesis
Studies in Drosophila Melanogaster." *Nature Methods* 6 (6): 431–34.
Venken, Koen J. T., Karen L. Schulze, Nele A. Haelterman, Hongling Pan, Yuchun He, Martha Evans-
Holm, Joseph W. Carlson, et al. 2011. "MiMIC: A Highly Versatile Transposon Insertion Resource
for Engineering Drosophila Melanogaster Genes." *Nature Methods* 8 (9): 737–43.
Vermaak, Danielle, and Harmit S. Malik. 2009. "Multiple Roles for Heterochromatin Protein 1 Genes in
Drosophila." *Annual Review of Genetics* 43: 467–92.
Wang, Xiaoxi, Lei Pan, Su Wang, Jian Zhou, William McDowell, Jungeun Park, Jeff Haug, Karen
Staehling, Hong Tang, and Ting Xie. 2011. "Histone H3K9 Trimethylase Eggless Controls
Germline Stem Cell Maintenance and Differentiation." *PLoS Genetics* 7 (12): e1002426.
Warming, Søren, Nina Costantino, Donald L. Court, Nancy A. Jenkins, and Neal G. Copeland. 2005.
"Simple and Highly Efficient BAC Recombineering Using galK Selection." *Nucleic Acids Research*
33 (4): e36.
Wayne, S., K. Liggett, J. Pettus, and R. N. Nagoshi. 1995. "Genetic Characterization of Small Ovaries,
a Gene Required in the Soma for the Development of the Drosophila Ovary and the Female
Germline." *Genetics* 139 (3): 1309–20.
Weiler, K. S., and B. T. Wakimoto. 1995. "Heterochromatin and Gene Expression in Drosophila."
Annual Review of Genetics 29: 577–605.
Yang, Peng, Yixuan Wang, and Todd S. Macfarlan. 2017. "The Role of KRAB-ZFPs in Transposable
Element Repression and Mammalian Evolution." *Trends in Genetics: TIG* 33 (11): 871–81.
Yasuhara, Jiro C., and Barbara T. Wakimoto. 2006. "Oxymoron No More: The Expanding World of
Heterochromatic Genes." *Trends in Genetics: TIG* 22 (6): 330–38.
Yin, Hang, and Haifan Lin. 2007. "An Epigenetic Activation Role of Piwi and a Piwi-Associated piRNA in
Drosophila Melanogaster." *Nature* 450 (7167): 304–8.
Yoshioka, K., H. Honma, M. Zushi, S. Kondo, S. Togashi, T. Miyake, and T. Shiba. 1990. "Virus-like
Particle Formation of Drosophila Copia through Autocatalytic Processing." *The EMBO Journal* 9
(2): 535–41.
Yuan, Kai, and Patrick H. O'Farrell. 2016. "TALE-Light Imaging Reveals Maternally Guided,
H3K9me2/3-Independent Emergence of Functional Heterochromatin in Drosophila Embryos."
Genes & Development 30 (5): 579–93.
Zook, Justin M., Daniel Samarov, Jennifer McDaniel, Shurjo K. Sen, and Marc Salit. 2012. "Synthetic
Spike-in Standards Improve Run-Specific Systematic Error Analysis for DNA and RNA
Sequencing." *PloS One* 7 (7): e41356.

Benner et al.*; sov* represses gene expression