Recent mixing of *Vibrio parahaemolyticus* populations

Chao Yang[1#], Xiaoyan Pei[2#], Yarong Wu[1], Lin Yan[2], Yanfeng Yan[1], Yuqin Song[1], Nicola Coyle[3], Jaime Martinez-Urtaza[4], Christopher Quince[5], Qinghua Hu[6], Min Jiang[6], Edward Feil[3], Dajin Yang[2], Yajun Song[1], Dongsheng Zhou[1], Ruifu Yang[1], Daniel Falush[3*], Yujun Cui[1*]

1 State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing 100071, China;

2 National Center for Food Safety Risk Assessment, Beijing 100022, China

3 University of Bath, Bath, Somerset, United Kingdom

4 The Centre for Environment, Fisheries and Aquaculture Science, Dorset DT48UB, United Kingdom

5 Warwick Medical School, University of Warwick, Warwick, United Kingdom

6 Shenzhen Centre for Disease Control and Prevention, Shenzhen, 518055, China

[#]These authors contributed equally to the article

* Corresponding authors: D. F. (danielfalush@googlemail.com) or Y. C. (cuiyujun.new@gmail.com)

Abstract

Humans have profoundly affected the ocean environment but little is known about

anthropogenic effects on the distribution of microbes. *Vibrio parahaemolyticus* is found in

warm coastal waters and causes gastroenteritis in humans and economically significant

disease in shrimps. Based on data from 1,103 genomes, we show that *V. parahaemolyticus* is

divided into four diverse populations, VppUS1, VppUS2, VppX and VppAsia. The first two

are largely restricted to the US and Northern Europe, while the others are found worldwide,

with VppAsia making up the great majority of isolates in the seas around Asia. Patterns of

diversity within and between the populations are consistent with them having arisen by

progressive divergence via genetic drift during geographical isolation. However, we find that

there is substantial overlap in their current distribution. These observations can be reconciled

without requiring genetic barriers to exchange between populations if dispersal between

oceans has increased dramatically in the recent past. We found that VppAsia isolates from the

US have an average of 1.01% more shared ancestry with VppUS1 and VppUS2 isolates than

VppAsia isolates from Asia itself. Based on time calibrated trees of divergence within

epidemic lineages, we estimate that recombination affects about 0.017% of the genome per

year, implying that the genetic mixture has taken place within the last few decades. These

results suggest that human activity, such as shipping and aquatic products trade, are

responsible for the change of distribution pattern of this marine species.

Introduction

Hospitable environments for particular marine microbes can be separated by large distances

but whether dispersal barriers substantially influence their distribution and evolution is

unknown. There are many studies of distribution of marine microbes e.g.[1-4], but these

typically survey patterns of macro-scale diversity. Differences in species level or genus level

composition between locations are as likely to reflect environmental heterogeneity as

dispersal, making the patterns difficult to interpret. Recent spread of microbes between

continents has been documented for lineages that cause pathogenic infection of humans,

including notorious clonal groups within *V. parahaemolyticus* and *Vibrio cholerae*[5-8].

However, these lineages are unusual in using humans as vectors, which might facilitate long-

range dispersal as in the case of the Haitian cholera outbreak[9]. We currently have little

information on rates of spread of the great majority of environmental organisms that do not

colonize large-animal hosts.

V. parahaemolyticus prefers warm coastal waters and causes gastroenteritis in humans[10,11].

Disease outbreaks became common from 1990s and became global, due to spread of

particular clones which are responsible for the great majority of recognized human

infections[5], which has been attributed to factors such as El Niño and climate change[12-14]. It is

not clear to what extent this pattern is historically typical, or whether reflects better

surveillance and different patterns of usage of marine resources. These clones also make up a

small fraction of the *V. parahaemolyticus* diversity and only a very small fraction of strains

isolated during environmental sampling.

The *V. parahaemolyticus* genome undergoes high rates of homologous recombination with

other members of the species[15,16]. We have previously found evidence that the species is split

into several populations[16]. Members of a population are not necessarily particularly related at

the clonal level, for example they may have recombined their entire genomes since sharing a

common cellular ancestor, but they are nevertheless on average more similar to each other

than to members of other populations because they have acquired DNA from a common gene

pool. Previously we found evidence of a single population with a well-mixed gene pool in

Asian waters and for one or more differentiated populations in the US[16].

Here we use a larger and more broadly sampled collection of 1,103 genomes to examine the

global population structure of the species. We find four populations with different but

overlapping modern geographic distributions as well as a small number of hybrid strains.

Under the assumption that genetic exchange between strains is constrained by geography, the

current extent of overlap is too high to maintain the populations as distinct entities and we

conclude that most of this mixing is likely to have taken place within the last few decades,

possibly coinciding with the recent emergence of pandemic clones.

Results and Discussion

Distribution of *V. parahaemolyticus* populations

We analyzed genomes of 1,103 strains including 392 new strains sequenced as part of this

study. These strains were isolated from a mixture of sources during 1951-2016, and covered

24 countries (Supplementary Fig. 1 and Supplementary Table 1). Clonal relationships

between strains can be inferred from identifying long stretches of near-identity,

corresponding to regions of the genome that have been inherited by direct descent since the

strains shared a common ancestor, or, more simply, by the strains having a small number of

SNP differences between them genome wide, which can be revealed by the Neighbor-Joining

(NJ) tree (Fig. 1a). Based on criterion of high nucleotide identity, the dataset contains 13

clonal groups, with 10 associated with human disease and 3 associated with the environment

(Supplementary Table 1).

The presence of clonally related strains in the data complicates analysis of deeper population

structure, so we first removed closely related isolates to make a "non-redundancy" dataset of

469 strains, in which no sequence differed by less than 2,000 SNPs in the core genome

(Methods). We used fineSTRUCTURE to identify distinct populations[17]. In total, 115

populations were identified in this initial analysis, however most comprised only two or three

strains (Supplementary Fig. 2a). These are likely to be sets of strains that are clonally related,

so we removed all but one from each group and reran fineSTRUCTURE. After several

iterations of the same procedure, we identified four populations with between 10 and 217

members and two singletons (Supplementary Fig. 2b). These singletons might be hybrids or

representatives of otherwise unsampled populations.

The current distribution of the populations is shown in Fig. 2a. The great majority of isolates

from Asia (574/600) are assigned to VppAsia, with all but one of the remainder (VppUS1,

isolated from a shrimp farm in Thailand) being assigned to VppX. VppUS1 is found almost

entirely in the US and is most common in the Mexican Gulf, with 13 out of the 29 VppUS1

strains are isolated from there. VppUS2 is most common on the US Atlantic coast (20 of 42)

and has also been isolated several times in Northern Europe. VppX is most common on the

Pacific coast and the Northern part of the US coast. These patterns are not predominantly

determined by the spread of human disease clones, since similar patterns are observed if the

dataset is restricted to the 469 non-redundancy strains (Supplementary Fig. 3a). The

distribution of CG1, the pandemic clonal group that mostly belongs to sequence type (ST) 3[5],

is similar to that of other VppAsia isolates (Supplementary Fig. 3b), while CG2 (ST36), an

epidemic group that is abundant in US and Canada[8], has a similar distribution to that of

VppX isolates, except that it has not been isolated from Asia (Supplementary Fig. 3b). The

distribution of the four populations is also similar when analysis is restricted to strains from

the human disease (Supplementary Fig. 3c) or environment (Supplementary Fig. 3d).

The 1,103 genomes in this study have been collected for a variety of different purposes and

do not represent a defined environmental or epidemiological cohort. Furthermore, sampling

numbers in most locations are small and the coasts of Africa and Australia, for example are

almost entirely unsampled. Nevertheless, our results demonstrate that at a global scale,

geographic distributions of populations overlap considerably and that there is a substantial

difference in the frequencies of the populations in the waters of Asia and those of the US

Coast (Fig. 2a).

Relationships amongst populations

The populations have a modest level of differentiation at the nucleotide level (Supplementary

Table 2), with F_{st} values of around 0.1 approximately equivalent to that between humans

living on different continents[18], implying that most common polymorphisms are shared

between populations. VppUS1 is the most diverse and isolates are no more similar to each

other in terms of mean SNP distance than they are to members of the other populations (Fig.

1b). However, according chromosome painting, which is based on haplotype similarity and

therefore more sensitive in detecting sharing of DNA due to common descent, all of the

members show substantially higher coancestry with other members of the population than

any of the other isolates in the dataset (Fig. 2b), implying that the population consists of

isolates that share ancestry, rather than being a collection of unassignable genomes. The other

populations have consistently lower distances with members of their own populations and

VppX and VppAsia are more closely related to each other than they are to VppUS1 and

VppUS2.

One explanation for the high diversity of VppUS1 is that it has frequently absorbed genetic

material from other populations. In order to test this hypothesis, while avoiding the effect of

clonal relationships within the population itself on estimates of relationships with other

populations, we painted the chromosomes of each of its members, using the members of the

other three populations as donors. A high diversity of painting palettes was observed from

VppUS1, with between 43% and 74% assigned to VppAsia and between 15% and 49% to

VppUS2 (Supplementary Fig. 4a). By contrast, the other three populations showed lower

levels of variation in assignment fractions in analogous paintings (Supplementary Fig. 4b-d).

Thus, VppUS1 owes its high diversity to being a hub for admixture, with input from both

VppUS2 and VppAsia. The members of VppUS1 in our sample are all clearly distinct in

ancestry profile from members of other populations (Fig. 2b), justifying the distinct

population label, but if gene flow levels were higher, it seems likely that the population

would lose its distinct identity and ancestry patterns would be better described by a
continuum than discrete population labels.
156
Recent mixing of *V. parahaemolyticus* populations
The observation of distinct populations is informative about patterns of migration in the past.
Population genetic theory implies that differentiation between demes can only arise and
persist if levels of migration between them are low, specifically on the order of magnitude of
one migrant per generation or less[19]. Once a migrant arrives in a deme, it progressively
imports DNA from other strains and becomes more and more similar to the other strains in its
new deme. The intuition behind the theory is that if too many strains are migrants, the demes
will progressively lose their distinct genetic profiles and merge into a single gene pool. This
theory has been developed for outbreeding eukaryotes[20] and bacterial populations deviate
from several of the assumptions of the theory, in ways that are currently not well understood,
making quantitative predictions impossible. Nevertheless, the qualitative expectation is that
most isolates should have the ancestry profile of the region, with only a small fraction of the
isolates having part ancestry from other locations.
170
The data differs from the qualitative predictions of migration-drift equilibrium because while
there are few strains of clearly intermediate ancestry in the dataset, many locations have
multiple strains from two or more of the four distinct populations that we have identified,
making it not obvious what deme they belong to. Asia is clearly the most likely ancestral
home range of VppAsia based on its high prevalence there but it is difficult to define
boundaries of likely ancestral ranges for the other three populations with any confidence
because the isolates assigned to those populations are too dispersed and they do not make up
a clear majority anywhere. Thus the current distribution is qualitatively inconsistent with
migration-drift equilibrium.
180
There are a number of factors which can in principle maintain subdivision when members of
particular populations are found in the same location. For example, it is possible that the
mechanism by which recombination occurs results in import occurring preferentially from
members of the same population. For example, barriers to recombination due to homology
dependent mismatch repair has been proposed to account for the differentiation between
phylogroups of *Escherichia coli*, despite high overall level of recombination[21] because the
mechanism preferentially aborts recombination events between members of different

phylogroup. Other mechanisms that can generate barriers to gene flow are strain specific

phage, or differences in an ecological niche. However, the pattern of sharing of diversity is

very different in *V. parahaemolyticus* to that found in *E. coli*, with high nucleotide diversity

and low differentiation between them and there are few highly differentiated loci anywhere

within the core genome (Supplementary Fig. 5). It is difficult to conceive of a mechanistic

barrier encoded within the genomes or their phage that would effectively constrain

recombination between populations enough to explain the low number of hybrids within the

dataset as a whole, while also allowing the frequent recombination required to create the

freely mixed gene pools we see within populations. Therefore while it is difficult to rule out

this explanation, we do not consider it further.

We propose instead that barriers to movement of strains have reduced recently. Under this

hypothesis, it should be possible to approximately estimate the timescale on which mixing

has taken place, based on the amount of introgression found in locations where the different

populations now co-occur. Specifically, within our dataset, it is natural to compare the

VppAsia isolates within Asia and in North America. Since Asia has been least affected by

between continent migration (Fig. 2a), we predict that the VppAsia isolates in North America

should have more ancestry from other sources, that they have acquired recently in their new

locations. This prediction is borne out, a number of North America VppAsia isolates have

high levels of VppUS1 and VppUS2 ancestry and on average the North America VppAsia

isolates in the non-redundancy set of 469 strains have 1.01% more (in average 2.97% in

North America vs 1.96% in Asia) of their painting palette from VppUS sources than those

from Asia (Fig. 2b and Fig. 3a).

In order to provide a timescale for the acquisition of non-Asian ancestry, we examined the

evolution within the largest two clonal populations, CG1 and CG2. We removed

recombination regions, then ran BEAST[22] to estimate a clock rate of 5.5×10^{-7} per site per

year, with very similar values for the two clonal complexes (Supplementary Fig. 6). There are

about 313 bases exchanged per mutation (Supplementary Fig. 7), so this implies a rate of

recombination of 1.7×10^{-4} per site per year. Thus if all of the import into the VppAsia

bacteria was from US populations, then it would imply it would take about 59 years (with

extreme lower and upper boundaries of 32-151 years, see methods) to acquire an extra 1.01%

ancestry at this rate of import.

We also examined the origin of imports within CG1, the global pandemic clonal group. As

for the VppAsia isolates, a higher fraction of the imports was from the two US populations

amongst the isolates found in the North America than for the isolates found in Asia itself.

This small difference in ancestry, corresponding to about 0.19% of the genome in total (Fig.

3c), has arisen during around 20 years since the beginning of global spread of CG1 in 1996.

These observations are consistent with a hypothesis that barriers to migration have become

substantially weaker within the last few decades, but do not constitute direct evidence that

patterns of gene flow between populations have changed. This hypothesis is empirically

testable although we do not have a suitable strain collection to facilitate it. For example, if the

Asian bacteria have arrived in large numbers in the US recently, then DNA from VppAsia

bacteria should make up a higher proportion of recent genetic imports than older ones.

In order to explain why the pattern of dispersal has changed recently, it is necessary to first

postulate reasons why dispersal was previously limited. We hypothesis that spread of bacteria

between oceans is limited by large distances between environments that are hospitable,

making it rare that bacteria survive transportation between them. Large mammals, seabirds

and other aquatic organisms travel large distances but do not necessarily provide habitats that

V. parahaemolyticus can colonize for the days or weeks required to get from one ocean to

another. Thus, we propose that dispersal between oceans did occur but was rare.

Humans have changed several aspects of the ocean environment, creating new habitats

through effluent discharge, warming and acidifying the oceans through climate change,

providing new mobile habitats on the hulls of ship and in ballast water and transporting

copeopods and other marine organisms deliberately to facilitate aquaculture or more

accidentally through trade in marine products[23,24]. Several of these could have facilitated

transmission of bacteria between oceans. Furthermore, *V. parahaemolyticus* can adapt to

colonize copeopods[25] so that for example human-associated dispersal of species such as the

manila clam from Asia to the America and Europe[26] could be responsible for the high

frequency of Asian *V. parahaemolyticus* there. A single introduction via ballast water or

introduction of shellfish for aquaculture would typically have low values of propagule

pressure (a single event with few individuals), while recurring introductions through recently

increased human activity may contribute in a regular basis introducing trans-ocean migration

of *V. parahaemolyticus*.

Further work is required to narrow down the most important factors. To identify the

frequency of *V. parahaemolyticus* reads in extensive metagenomic sampling of the open

ocean would provide knowledge on natural transmission of this bacterium. One objection to a

direct human dispersal, rather than for example a role for climate change is that the absolute

number of bacteria transported by ships or trade is likely to be small. However, this objection

does not seem especially compelling. The absolute number of bacteria transported from one

ocean to another does not need to be very large; if bacteria are fit in their new environment,

they can multiply rapidly to constitute a substantial proportion of the bacteria in their new

habitat.

Conclusions

Our results support our earlier conclusion that *V. parahaemolyticus* is subdivided into distinct

geographical populations. We have identified 4 clearly differentiated populations, two of

which appear to have foci in the US (VppUS1 and VppUS2). A third is predominant in Asia,

while the ancestral home range of the forth VppX is difficult to guess based on current

sampling. However, these ranges pose a puzzle, in that they overlap substantially, both for

environmental and human disease causing isolates, which show approximately similar

patterns of distribution. Hybrids are rare, for example, amongst VppAsia isolates found in the

US, most have ancestry profiles indistinguishable from strains found in Asia, while a handful

have less than 10% introgression from either of the two US populations. The simplest and

most parsimonious explanation is that previous barriers to migration have been reduced

recently, allowing bacteria to disperse rapidly between continents but that because bacterial

recombine relatively slowly (about 0.017% of their genome a year on average), there has not

had sufficient time to generate hybrids.

These results have two major implications. Firstly, they suggest that recent human activity

has disrupted long-standing barriers to genetic exchange in the oceans and that this has

affected microbial population structure. Secondly, changing global patterns of *V.*

parahaemolyticus disease incidence may be directly connected to changes in dispersal of the

species, rather than being specific to the small number of clonal lineages that are responsible

for most of the major outbreaks.

Materials and Methods

Bacterial strains

Totally 1,103 strains were used in this research, including 392 newly sequenced and 711

publicly available strains (Supplementary Table 1). The newly sequenced strains were

isolated in China during daily food surveillance in 2014. The remaining 711 publicly

available strains were downloaded from the NCBI database. The genomes of newly

sequenced strains are available in GenBank with the accession numbers listed in

Supplementary Table 1.

New sequenced strains were cultured in the LB-2% NaCl agar at 37 °C, and classical

phenol/chloroform method was used to the extract genomic DNA.

Sequencing and assembly

The whole genome DNA was sequenced by using Illumina Hiseq 4000. The pair-end

sequencing library with average insert size of 350 bp were build according to the

manufacture's introduction (Illumina Inc., USA). The read length is 150 bp and in average

500 Mb raw data were generated for each strain, which is corresponding to the sequencing

depth of approximately 100 fold. The adaptor sequence and low quality reads were filtered

and the clean reads were assembled by using SOAPdenovo v2.04[27] as described before[16]. The

number of contigs and average size of assemblies are 263 and 5.1 Mb, respectively.

Variation Detection

The SNPs were identified by aligning the *V. parahaemolyticus* genomes against with the

reference genome (RIMD 2210633) by using MUMmer[28] as previously described[16], and only

bi-allelic SNPs were used in further analysis. As the number of detected SNPs would relate

with core-genome of different strain sets, we created multiple SNP sets by using different

strain sets when perform analysis in various purposes. Totally 462,214 SNPs were identified

from all 1,103 genomes, 650,683 SNPs were from 469 non-redundancy genomes, 355-8,921

SNPs were separately from 13 clonal groups.

Population structure

The NJ trees were built by using the TreeBest software

(http://treesoft.sourceforge.net/treebest.shtml) based on sequences of concatenated SNPs, and

were visualized by using online tool iTOL[29].

The population structure of *V. parahaemolyticus* was built based on the 469 non-redundancy

genomes set by using Chromosome painting and fineSTRUCTURE[17] as described before[16].

The fineSTRUCTURE result of all 469 non-redundancy genomes revealed that multiple

clonal signals still presented. Therefore we selected only one representative genome from

each clone, and combined them with the left genomes to perform another round of

Chromosome painting and fineSTRUCTURE analysis. After six iterations, we finally

obtained a set of 260 genomes with no clonal signals presented in the result (Supplementary

Fig. 2b). To balance the sampling size among different population, we selected 60 strains,

including 14-16 strains from each population and 2 hybrid strains, to repeat the

fineSTRUCTURE analysis (Supplementary Fig. 2c). The result further verify the population

structure of *V. parahaemolyticus* species. Population assignment based on fineSTRUCTURE

was consist with NJ tree (Fig. 1a) except for two strains, PCV08-7 and TUMSAT_H01_S4,

and one epidemic group, CG2. Strain PCV08-7 and TUMSAT_H01_S4 were assigned to

VppAsia and VppUS2 respectively by fineSTRUCTURE analysis, but in the NJ tree they are

more closely related with VppX strains. The CG2 strains were all assigned to VppX

populations by fineSTRUCTURE, but in NJ tree it was grouped with VppAsia strains.

The length of chunks were extracted from the output file of Chromosome painting based on

469 non-redundancy strains, to calculate the percentage of admixtures of different

populations for each non-redundancy genome (Fig. 2b).

New designation of *V. parahaemolyticus* populations

In previous study, we designated four *V. parahaemolyticus* populations, named Asia-pop, US-

pop 1, Hyb-pop 1, and Hyb-pop 2, separately, based on dataset of 157 genomes. Here with

more samples were used in distinguishing the population, we found a new population that

mostly isolated from US, and the previously defined Hyb-pop 1 were known as just several

hybrid strains, or representatives of otherwise unsampled populations. As evidences revealed

that *V. parahaemolyticus* populations are geographical clustered, we proposed novel

nomenclatures for them, which read as VppAsia, VppX, VppUS1 and VppUS2. The 'Vpp' is

abbreviation of '*V. parahaemolyticus* population'. The first three populations are

corresponding with previously defined Asia-pop, Hyb-pop 2 and US-pop 1, and the VppUS2

is the population newly identified in this study.

Inference of substitution rate using BEAST

Two clonal groups with large sample size, CG1 (n = 153, global pandemic group, also known

as O3:K6 and its sero-variants group) and CG2 (n = 92, an epidemic group that popular in

US, also known as serotype O4:K12), were selected to calculate molecular clock respectively

by using BEAST v1.83 [22]. The variations that caused by recombination inferred by our

pipeline were excluded in substitution rates analysis. There are 10 of total 153 CG1 strains

revealed too many strain-specific SNPs and revealed unusual long branches in the NJ tree

even after removing the recombination variations (Supplementary Fig. 8), and similar pattern

was observed in 1 of total 92 CG2 strains. These 11 strains with unusual high number of

SNPs, and 22 strains with unknown isolation time, were excluded from BEAST analysis. We

implemented analysis under GTR + I and relaxed clock model with constant size coalescent.

The MCMC chain was run for 10^8 and sampling for every 5,000 generations. The effective

sample sizes of all inferred parameters were above than 200 in our results. The estimated

molecular clock based on CG1 genomes is 5.6×10^{-7} with 95% confidence interval (CI) of

$4.3\text{-}6.7 \times 10^{-7}$ per site per year, and 5.4×10^{-7} with 95% CI of $3.6\text{-}7.2 \times 10^{-7}$ for CG2

genomes. Here the average value, 5.5×10^{-7}, was used as the most likely estimate of *V.*

parahaemolyticus molecular clock. The extremes of 95% CI based on two clonal groups, i.e.,

$3.6\text{-}7.2 \times 10^{-7}$, were used as lower and upper 95% boundaries for ensuring that they

encompassed the true values as much as possible.

Recombination detection and inference of recombination rate

Totally 13 clonal groups with more than 10 strains (Supplementary Table 1), defined by intra-

group paired-distance less than 2,000 SNPs, were selected to be used in detection of

recombination events. We firstly used previously pipeline to detect recombination[16]. Briefly,

we recalled the SNPs for each clonal group because different datasets had different core-

genomes, and these SNPs were used to construct a NJ tree. Then PAML software package[30]

was used to determine the SNPs of each branch. Assuming neutrality and no recombination,

the observed SNP density of a given region should follow the binomial distribution. We used

the sliding window method to identify regions that rejected the null hypothesis ($P < 0.05$) and

all SNPs in such windows were treated as recombined SNPs. We also used

ClonalFrameML[31], a software based on maximum likelihood method, to detect bacterial

recombination within the same dataset. Sequence alignments of genomes for each clonal

group and the corresponding maximum-likelihood tree constructed using PHYML with HKY

model[32], were used as input files and non-core regions were ignored during calculation. The

inferred recombination regions using ClonalFrameML are mostly consistent with our in-

house method (Supplementary Fig. 7).

Two sets of r/μ (ratio of size of recombination regions to the number of mutation sites) were

obtained through different methods. The value is 331 with 99% CI of 228-435 for our in-

house method and 295 with 99% CI of 186-404 for ClonalFrameML. The average value, 313,

was selected as r/μ of the *V. parahaemolyticus*. Concerning the molecular clock rate of $5.5 \times$

10^{-7} per site per year, the most likely recombination rate of *V. parahaemolyticus* is 1.7×10^{-4}

per site per year. We selected the extremes of r/ μ from two sets of 99% CI, 186-435, to

calculate the lower and upper boundaries of recombination rate through multiplying the

extremes of the molecular clock, which obtained the results of 6.7×10^{-5} - 3.1×10^{-4} per site

per year. Accordingly, the time to obtained 1.01% of genome fragments would be 59 years

with extreme boundaries of 32-151 years.

Identify the contribution of *V. parahaemolyticus* populations to pandemic genomes in

different geographical location

We assigned CG1 strains into two groups according to their isolated location, with one

isolated from Asia and another isolated from North America. By using ClonalFrameML, we

inferred the recombination fragments that occurred on each strain. Totally 81 fragments were

found in Asia CG1 strains and 65 fragments were found in North America CG1 strains, with

total size of 221 kb and the median length of 1035 bp. Then we identify the possible donor

genome of these recombination fragments by align them against with 468 non-redundancy

genomes (excluding the CG1 genome from the dataset) using BLASTn, with a threshold of

coverage >= 80% and identity >= 99.5%. The observed frequency of the donor genomes in

each population was calculated. For recombination fragments carried by Asia CG1 genomes,

the average value of their donor frequency in a population were taken as contribution

proportion of the corresponding population to the recipient sequences in Asia CG1 genomes,

and similarly, we obtained the contribution proportion of each population to the North

America CG1 genomes (Fig. 3c). We also try the relaxed identity thresholds (99.0%) in

BLASTn and acquired similar results.

Acknowledgements

We gratefully acknowledge Dr. Narjol Gonzalez-Escalona for contributing *V. parahaemolyticus* genomes. This work is supported by the National Key Research & Development Program of China (No. 2017YFC1601503, 2016YFC1200100 and 2017YFC1200800), Sanming Project of Medicine in Shenzhen (No. SZSM201811071) and the National Natural Science Foundation of China (No. 31770001). J. Martinez-Urtaza were funded by Natural Environment Research Council (NERC) project (No. NE/P004121/1). D.F. is funded by a Medical Research Council Fellowship as part of the MRC CLIMB consortium for microbial bioinformatics (grant number MR/M501608/1).

Author Contributions

Y. C., D. F. and R. Y. designed the study and coordinated the project; X. P., L. Y., J. M., Q. H., and D. Y. contributed strains for analysis; C. Y., X. P., Y. W, N. C., Y.Q. S., Y.J. S., Y. Y., M. J., C. Q., D. F. and Y. C. analyzed the data; E. F., J. M. and D. Z. provided insightful comments, D. F. and Y. C. wrote the manuscript. All authors approved the final version of the manuscript.

Competing Financial Interests statement

No

References

1. Brown, M.V., Ostrowski, M., Grzymski, J.J. & Lauro, F.M. A trait based perspective on the biogeography of common and abundant marine bacterioplankton clades. *Mar Genomics* **15**, 17-28 (2014).
2. Yilmaz, P., Yarza, P., Rapp, J.Z. & Glockner, F.O. Expanding the World of Marine Bacterial and Archaeal Clades. *Front Microbiol* **6**, 1524 (2015).
3. Kent, A.G., Dupont, C.L., Yooseph, S. & Martiny, A.C. Global biogeography of Prochlorococcus genome diversity in the surface ocean. *ISME J* **10**, 1856-65 (2016).
4. Hellweger, F.L. *et al.* The Role of Ocean Currents in the Temperature Selection of Plankton: Insights from an Individual-Based Model. *PLoS One* **11**, e0167010 (2016).
5. Nair, G.B. *et al.* Global dissemination of Vibrio parahaemolyticus serotype O3:K6 and its serovariants. *Clin Microbiol Rev* **20**, 39-48 (2007).
6. Mutreja, A. *et al.* Evidence for several waves of global transmission in the seventh cholera pandemic. *Nature* **477**, 462-5 (2011).
7. Weill, F.X. *et al.* Genomic history of the seventh pandemic of cholera in Africa. *Science* **358**, 785-789 (2017).

8.   Martinez-Urtaza, J. *et al.* Genomic Variation and Evolution of Vibrio
parahaemolyticus ST36 over the Course of a Transcontinental Epidemic Expansion.
MBio **8**(2017).
9.   Chin, C.S. *et al.* The origin of the Haitian cholera outbreak strain. *N Engl J Med* **364**,
33-42 (2011).
10.   Yeung, P.S. & Boor, K.J. Epidemiology, pathogenesis, and prevention of foodborne
Vibrio parahaemolyticus infections. *Foodborne Pathog Dis* **1**, 74-88 (2004).
11.   Su, Y.C. & Liu, C. Vibrio parahaemolyticus: a concern of seafood safety. *Food*
Microbiol **24**, 549-58 (2007).
12.   Ansede-Bermejo, J., Gavilan, R.G., Trinanes, J., Espejo, R.T. & Martinez-Urtaza, J.
Origins and colonization history of pandemic Vibrio parahaemolyticus in South
America. *Mol Ecol* **19**, 3924-37 (2010).
13.   Martinez-Urtaza, J., Trinanes, J., Gonzalez-Escalona, N. & Baker-Austin, C. Is El Nino a
long-distance corridor for waterborne disease? *Nat Microbiol* **1**, 16018 (2016).
14.   Baker-Austin, C., Trinanes, J., Gonzalez-Escalona, N. & Martinez-Urtaza, J. Non-
Cholera Vibrios: The Microbial Barometer of Climate Change. *Trends Microbiol* **25**,
76-84 (2017).
15.   Yan, Y. *et al.* Extended MLST-based population genetics and phylogeny of Vibrio
parahaemolyticus with high levels of recombination. *Int J Food Microbiol* **145**, 106-12
(2011).
16.   Cui, Y. *et al.* Epidemic Clones, Oceanic Gene Pools, and Eco-LD in the Free Living
Marine Pathogen Vibrio parahaemolyticus. *Mol Biol Evol* **32**, 1396-410 (2015).
17.   Lawson, D.J., Hellenthal, G., Myers, S. & Falush, D. Inference of population structure
using dense haplotype data. *PLoS Genet* **8**, e1002453 (2012).
18.   Rosenberg, N.A. *et al.* Genetic structure of human populations. *Science* **298**, 2381-5
(2002).
19.   Wright, S. Evolution in Mendelian Populations. *Genetics* **16**, 97-159 (1931).
20.   Whitlock, M.C. & McCauley, D.E. Indirect measures of gene flow and migration: FST
not equal to 1/(4Nm + 1). *Heredity (Edinb)* **82 (Pt 2)**, 117-25 (1999).
21.   Didelot, X., Meric, G., Falush, D. & Darling, A.E. Impact of homologous and non-
homologous recombination in the genomic evolution of Escherichia coli. *BMC*
Genomics **13**, 256 (2012).
22.   Drummond, A.J., Suchard, M.A., Xie, D. & Rambaut, A. Bayesian phylogenetics with
BEAUti and the BEAST 1.7. *Mol Biol Evol* **29**, 1969-73 (2012).
23.   Ruiz, G.M. *et al.* Global spread of microorganisms by ships. *Nature* **408**, 49-50 (2000).
24.   Martinez-Urtaza, J. *et al.* Spread of Pacific Northwest Vibrio parahaemolyticus strain.
N Engl J Med **369**, 1573-4 (2013).
25.   Martinez-Urtaza, J. *et al.* Ecological determinants of the occurrence and dynamics of
Vibrio parahaemolyticus in offshore areas. *ISME J* **6**, 994-1006 (2012).
26.   Chiesa, S. *et al.* A history of invasion: COI phylogeny of Manila clam Ruditapes
philippinarum in Europe. *Fisheries Research* **186**, 25-35 (2017).
27.   Luo, R. *et al.* SOAPdenovo2: an empirically improved memory-efficient short-read de
novo assembler. *Gigascience* **1**, 18 (2012).
28.   Delcher, A.L., Salzberg, S.L. & Phillippy, A.M. Using MUMmer to identify similar
regions in large sequence sets. *Curr Protoc Bioinformatics* **Chapter 10**, Unit 10 3
(2003).

29. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display
and annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**, W242-5
(2016).
30. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**,
1586-91 (2007).
31. Didelot, X. & Wilson, D.J. ClonalFrameML: efficient inference of recombination in
whole bacterial genomes. *PLoS Comput Biol* **11**, e1004041 (2015).
32. Guindon, S. & Gascuel, O. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. *Syst Biol* **52**, 696-704 (2003).
512

Figure 1. Population structure of *V. parahaemolyticus* and relationships within and between populations. (a) NJ tree of 1,103 *V. parahaemolyticus* stains based on 462,214 SNPs. Branch colors indicate populations defined by fineSTRUCTURE, green for VppAsia, orange for VppX, light blue for VppUS1, dark blue for VppUS2, purple for hybrid strains. The ring colors from inner to outer indicate isolation time and sample type, respectively. The blank indicates information not available. (b) SNP distance within and between populations based on 469 non-redundancy strains. Colors indicate populations and are consistent with branch colors of panel a.

Figure 2. Geographical distribution and admixture of *V. parahaemolyticus* populations. Colors in circle and bar plot indicate populations and are as in Figure 1. Each circle indicates the population composition of a city/country, with radius in proportion to the sample size. Bar plot indicates the ancestry composition inferred by chromosome painting of two geographical regions: Asia and North America. Each vertical bar represents one non-redundancy strain and the proportion of color indicates the contribution of each population. Different populations are separated by blank vertical bar. Only strains with information of isolation location are included in panel a (n = 1,008) and panel b (n = 422).

Figure 3. Recent mixing of *V. parahaemolyticus* populations. (a) Ancestry composition of three other *V. parahaemolyticus* populations in VppAsia strains in different geographical regions. The contribution from other populations to the VppAsia is inferred by chromosome painting. X axis indicates the proportion of contributed chunk length of a population in one strain and Y axis indicates the corresponding frequency. (b) ClonalFrameML recombination analysis of 141 CG1 strains. Left: ClonalFrameML reconstructed phylogeny. Right: dark blue horizontal bars indicate recombination events, grey areas indicate non-core regions. Two chromosomes are separated by dot line. (c) Source of recombination fragments of CG1 strains in different geographical regions. Y axis indicates the proportion of recombination fragments input from different population against core genome. Colors in (a) and (c) indicate four populations and are as in Figure 1.