DNA-uptake pilus of *Vibrio cholerae* capable of kin-

discriminated auto-aggregation

David. W. Adams, Sandrine Stutzmann, Candice Stoudmann and Melanie Blokesch*

Laboratory of Molecular Microbiology, Global Health Institute, School of Life Sciences,

Station 19, EPFL-SV-UPBLO, Swiss Federal Institute of Technology Lausanne (Ecole

Polytechnique Fédérale de Lausanne; EPFL), CH-1015 Lausanne, Switzerland.

*For correspondence email: melanie.blokesch@epfl.ch.

Abstract

Natural competence for transformation is a widely used and key mode of horizontal gene

transfer that can foster rapid bacterial evolution. Competent bacteria take-up DNA from

their environment using Type IV pili, a widespread and multi-purpose class of cell surface

polymers. However, how pili facilitate DNA-uptake has remained unclear. Here, using direct

labelling, we show that in the Gram-negative pathogen *Vibrio cholerae* DNA-uptake pili are

highly dynamic and that they retract prior to DNA-uptake. Unexpectedly, these pili can self-

interact to mediate auto-aggregation of cells into macroscopic structures. This phenotype is

conserved in disease causing pandemic strains. However, extensive strain-to-strain

variability in the major pilin subunit PilA, present in environmental isolates, controls the

ability of pili to interact without affecting transformation. We go on to show that

interactions between pili are highly specific, enabling cells producing pili composed of

different PilA subunits to discriminate between one another. On chitin surfaces, a natural

habitat of *V. cholerae*, pili connect cells within dense networks, suggesting a model whereby

DNA-uptake pili function to promote inter-bacterial interactions during surface colonisation.

Moreover, our results provide evidence that type IV pili could provide a simple and

potentially widespread mechanism for bacterial kin recognition.

Keywords

Auto-aggregation, DNA uptake, natural transformation, pilus dynamics, type IV pilus, *Vibrio*

cholerae

Introduction

How bacteria physically sense and interact with their environment is a fundamental problem in biology. Type IV pili (T4P) are cell surface polymers ideally suited to this task[1,2]. Composed of a single major pilin and assembled by widely distributed and conserved machinery, T4P exhibit extensive functional versatility, with roles in motility, DNA-uptake, surface sensing and adhesion[3-5]. Consequently, T4P are critical virulence factors for a number of important human pathogens including *Vibrio cholerae*, which causes the pandemic diarrhoeal disease cholera[6]. In Gram-negative bacteria pilins are processed at the inner-membrane, extracted by the assembly machinery and polymerised into a helical pilus fibre that exits the cell surface through a gated outer-membrane pore; the secretin[7-11]. A key feature of T4P is their ability to undergo dynamic cycles of extension and retraction[12,13], powered by the action of dedicated extension (*e.g.* PilB) and retraction (*e.g.* PilT) ATPases, which either add or liberate pilin subunits at the base[14,15]. These dynamics are essential for many T4P functions *e.g.* twitching-motility[11,12]. Thus, understanding how T4P function may yield insights valuable for understanding mechanisms of environmental survival and pathogenesis.

Despite their multifunctional potential, pandemic strains of *V. cholerae* typically encode three distinct T4P systems – two of which are well characterised. First, toxin co-regulated pili (TCP) serve a dual role as both a receptor for CTXϕ bacteriophage[16], which carries the cholera toxin genes, and as the primary human colonisation factor with multiple essential roles in infection involving adhesion and auto-aggregation on the intestinal cell surface[17-19]. Second, Mannose-sensitive haemagglutinin (MSHA) pili are involved in surface sensing and attachment and thus, are important in the initiation of biofilm formation[20-24]. Third, in its natural aquatic environment *V. cholerae* often associates with chitinous surfaces[25], which are nutritious, foster biofilm formation and likely play a role in environmental dissemination and transmission to humans[26-29]. Chitin utilisation triggers competence for natural transformation[30], a widely used mode of horizontal gene transfer

that allows bacteria to take up DNA from their environment, and which can thus, foster

rapid bacterial evolution[31]. This requires the production of the Chitin-Regulated (ChiRP) or

DNA-uptake pilus[30].

We previously showed DNA-uptake pili form *bona fide* pili composed of the major

subunit PilA and that transformation was dependent on the presumed retraction ATPase

PilT[32]. However, the pilus itself is not sufficient for transformation and requires the

concerted action of a periplasmic DNA-binding protein, ComEA[32,33]. Upon receipt of

transforming DNA ComEA switches from a diffuse to focal localisation[33,34]. These findings,

together with work in other organisms, led to a model in which pilus retraction facilitates

DNA entry into the periplasm[35], wherein ComEA acts as 'ratchet' to pull in the remaining

DNA[33]. Subsequently, DNA transport across the inner-membrane occurs via a spatially

coupled channel, ComEC[34,36]. Though this model is well supported by genetic experiments[32]

and the similarly combined action of T4P and ComEA in other organisms[37,38], direct evidence

is lacking.

Here, we visualised the DNA-uptake pilus directly using a cysteine labelling

approach, which was recently validated as a tool for labelling pili[39]. As predicted, we

demonstrate that the pili are highly dynamic and that these dynamics are PilT-dependent.

Unexpectedly, however, we discovered that DNA-uptake pili are also capable of self-

interacting, which in liquid culture results in a strong auto-aggregation phenotype.

Variability in PilA controls this activity but has no affect on transformation. Remarkably,

specific interactions allow pili composed of different PilA subunits to distinguish between

one another, enabling a simple mechanism for kin recognition.

Results

Direct observation of pilus dynamics by cysteine labelling

To avoid the limitations imposed by immuno-fluorescent methods, we employed a cysteine

labelling approach using a thiol-reactive dye[39,40]. PilA cysteine variants were created along

the length of the surface exposed αβ-loop[9,11] and tested for functionality using a previously

validated chitin-independent transformation system in which competence induction is

arabinose-inducible[41] (Tn*tfoX*; see methods) (Fig. 1A, B and Fig. S1A). A variant, PilA[S67C],

was identified that relative to the unmodified parent is fully transformable (Fig.1A),

produced at similar levels (Fig. S1B), and does not affect pilus assembly (Fig. S1C), as assayed

by a classical shearing method. Since other variants exhibited functional defects they were

not studied further.

When stained, competent cells producing PilA[S67C] exhibited visible pili (Fig. 1C).

On average 27 ± 3 % cells were piliated, with the majority of these displaying 1 or 2 pili per

cell (Fig. 1H and I). The length of pili was clustered around 1-2 μm, though pili up to 10 μm in

length were also observed (Fig. 1J). Numerous detached pili were also evident in the growth

media. Intriguingly, these pili frequently appeared to self-interact, forming large structures

composed of networks of pili (Fig. S2A). When examined by time-lapse microscopy cells

exhibited rapid pilus dynamics, with multiple assembly and retraction events immediately

apparent (Fig. 1K; Movie S1-4). Indeed, within the 1 minute time frame studied 66 ± 3 % cells

exhibited pili, with most cells producing 1-2 pili per minute (Fig. 1M). Notably, a small

subpopulation of cells was even more dynamic, elaborating ≥5 pili per minute. Consistent

with this dynamic behaviour, and in support of the hypothesis that pilus retraction precedes

DNA-uptake, when cells were provided with purified genomic DNA it was possible to

concurrently visualise pilus retraction followed by DNA-uptake, as monitored by the re-

localisation of ComEA-mCherry, which binds incoming DNA in the periplasm (Fig. S3; Movie

S5).

As expected, deletion of components required for pilus assembly (*e.g.* the assembly

ATPase PilB or the secretin PilQ) abolished piliation (Fig. 1D and E). However, despite the

absence of obvious pili the cell body still stained. Control experiments in which PilA[S67C],

but not PilA[WT], was produced independently of the normal assembly machinery also

stained similarly (Fig. S1D), suggesting that this effect results from the dye being taken up

non-specifically and retained by the inner-membrane pool of PilA[S67C].

As in other species *V. cholerae* encodes two potential retraction ATPases, PilT and

PilU. Deletion of *pilU* did not affect piliation (Fig. 1F, H and I), consistent with its

dispensability for transformation[32]. In contrast, cells lacking the presumed retraction ATPase

PilT were hyper-piliated, with essentially all cells displaying multiple static pili (Fig. G-I;

Movie S6-7). Taken together with the dynamics described above, these data are consistent

with the presence of multiple assembly complexes scattered across the cell, as previously

predicted based on the mobility of PilB and the presence of multiple PilQ foci[32,42]. This might

normally serve to facilitate rapid switching of pilus location or else might reflect a need to

produce multiple pili under certain conditions. Unexpectedly, cells blocked for retraction

were often grouped into small clusters within dense networks of pili as well as occasionally

large aggregates of pili-encased cells (Fig. S2B and C). Indeed, when grown in liquid culture

cells appeared to auto-aggregate.

Competent cells auto-aggregate in the absence of pilus retraction

To test directly whether the aggregation phenotype was an inherent property of hyper-

piliated cells or merely an artefact introduced by the cysteine variant, cells containing

unmodified *pilA* were examined (Fig. 2A-C). Remarkably, aggregation occurred specifically in

competence-induced cells lacking *pilT*, with the formation of large, multi-layered spherical

aggregates on the order of 25-100 μm, which went on to form macroscopic aggregates that

rapidly sediment (Fig. 2B, +Ara). Quantification of the ratio of cells remaining in solution to

those in the settled aggregates (Fig. 2D) revealed that upon competence induction, ≥90% of

the retraction deficient cells were present in aggregates (Fig. 2D). Notably, strains producing

the PilA[S67C] variant used for labelling behave similarly (Fig. 1B).

To rule out the possibility that the deletion of *pilT* has either a polar or otherwise

non-specific effect, an intact copy of *pilT* driven by its native promoter was placed at an

ectopic locus. Importantly, complementation of Δ*pilT* was sufficient to fully restore the

~1000-fold defect in transformation frequency and abolished the aggregation phenotype

(Fig. 2D and E). Complementation also fully counteracted the enhanced motility phenotype

of Δ*pilT* (Fig. S4), which occurs due to loss of function in the adhesive MSHA pilus and is in

agreement with the previously established shared role of PilT in MSHA pilus function[24,43].

Time-course experiments indicated that aggregation occurs abruptly, suggesting

that rather than gradually increasing in size, large aggregates form via the accumulation of

smaller ones (Fig. S5A and B). In support of this hypothesis, when two otherwise isogenic

Tn*tfoX* Δ*pilT* strains, one of which constitutively produces GFP, were allowed to aggregate

together they formed well-mixed aggregates (Fig. S5E). If, however, they were allowed to

aggregate separately before being mixed (Fig. S5C and D), they formed large composite

aggregates consisting of a patchwork of labelled and unlabelled cells (Fig. S5F). Finally, the

aggregates remained stable and did not disperse, even after prolonged culture overnight.

Aggregates form via pilus-pilus interactions

The data so far suggest that hyper-piliated cells auto-aggregate via their DNA-uptake

pili. Indeed, deletions affecting DNA-uptake pilus assembly (*pilQ* and *pilA*) were sufficient to

abolish aggregation (Fig. 2F). In contrast, deletions targeting the assembly of TCP (*tcpA*) and

MSHA (*mshA*) pili, the *Vibrio* polysaccharide matrix (*vpsA*) required for biofilm formation[44] or

the flagellum (*flaA*), were dispensable for aggregation, both individually and in combination

(Fig. 2F). Similarly, additional genes tested with roles in biofilm formation (*bap1*, *rbmA*,

rbmEF, *vpsT*)[45,46], adhesion (*gbpA*)[47], cell shape (*crvA*)[48] and an O-linked glycosylase (*VC0393*

or *pgIL_Vc*)[49] that could be involved in pilin modification all aggregated at levels

indistinguishable from the unmodified parent (Fig. S6A). Finally, deletion of *VC0502*, an

'orphan' type IV pilin encoded on the *Vibrio* seventh pandemic island II, had no effect on

transformation, aggregation or motility (Fig. S6B-D). Additional control experiments

indicated that aggregation occurred similarly during growth under high salt conditions (*i.e.*

LB-S 20 g/L NaCl), which are often used to better reflect the natural aquatic environment of

Vibrio sp. (Fig. S6E). Likewise, the presence of Bovine serum albumin, which was previously

reported to disrupt pilus-pilus interactions in *Neisseria gonorrhoeae*[50], also had no effect

(Fig. S6E). Curiously, however, we noticed that the strength of the aggregation phenotype is

sensitive to the concentration of divalent cations (*e.g.* $CaCl_2$ or $MgCl_2$) (Fig. S6F), though the

significance of this result remains unclear.

To investigate how the pilus mediates auto-aggregation *i.e* pilus-pilus, pilus-cell or

otherwise, we co-cultured cells producing either dynamic pili (Tn*tfoX*), static pili (Tn*tfoX*,

Δ*pilT*), or blocked for both retraction and pilus production (Tn*tfoX*, Δ*pilT*, Δ*pilA*) with a

Tn*tfoX* Δ*pilT* strain constitutively producing GFP (Fig. 2G-I) and asked whether these cells

were recruited into the GFP-labelled aggregates. Strikingly, intermixed aggregates were only

observed when both cells were themselves capable of producing non-retractile pili, and

hence aggregating (Fig. 2H), indicating that aggregates form via direct pilus-pilus interactions

between DNA-uptake pili.

MSHA pili do not mediate aggregation

MSHA pili are important for surface sensing and attachment during the switch to the sessile

lifestyle that precedes biofilm formation, and pilus retraction is thought to be required for

their function[20-24]. Since MSHA pili are constitutively produced, but retraction deficient cells

only aggregate upon competence induction, the results above suggest that MSHA pili are

unable to mediate auto-aggregation. To test this hypothesis directly we grew cells

engineered to increase the levels of the second messenger c-di-GMP, which has been

reported to enhance the assembly of MSHA pili on the cell surface[43]. Control experiments

indicated that decreasing c-di-GMP levels cells did not induce aggregation (Fig. S7A and B).

In contrast, under elevated c-di-GMP levels, cells formed large clumps that appeared similar

to the aggregates described above and had a modest sedimentation phenotype (Fig. S7A

and B). However, the assembly of these structures occurred independently of *pilT*, *pilA* and

mshA, but was abolished by the loss of *vpsA* (Fig. S7C). Thus, we conclude that these

structures are not aggregates held together by pili but rather biofilms held together by

matrix polysaccharide.

PilA from pandemic strains is sufficient for aggregation in a non-pandemic strain

A1552, the strain used throughout this work, is an O1 El Tor clinical isolate representative of

the on-going 7[th] cholera pandemic. To test whether the ability to aggregate is conserved

among other pandemic strains we tested 6 additional O1 El Tor strains. In the absence of *pilT*

all aggregated to similar levels as A1552, as might be expected given that pandemic strains

are closely related (Fig. 3A). Surprisingly, however, N16961 did not aggregate unless its well-

characterised *hapR* frame-shift mutation[51], which renders it quorum-sensing (QS) defective,

was first repaired (Fig. 3A). Since transcription of the genes required for pilus assembly are

not regulated by QS[41] this finding suggests that there may be an additional layer of post-

transcriptional control.

PilA is known to vary considerably between different environmental strains of *V.*

cholerae[52], whereas the majority of clinical isolates carry an identical *pilA*. Interesting

exceptions are the toxigenic O37-serogroup strains V52 and ATCC25872, which were

responsible for limited epidemics of a cholera-like illness in the 1960s[53,54]. These strains are

thought to derive from a pandemic O1 classical strain through the exchange of the O-antigen

cluster by horizontal gene transfer[55,56]. However, they both encode the same PilA version

that is only 50% identical to that typically carried by pandemic strains (Fig. S8). Since V52 is

QS deficient, and transformation is QS dependent, we investigated the functionality of this

PilA in ATCC25872, which has a functional QS pathway. Indeed, ATCC25872 is transformable

at similar levels to that of A1552, and as expected, transformation is PilT-dependent (Fig.

3B). However, in contrast to A1552, competence-induced cells lacking *pilT* did not aggregate

and were excluded from aggregates formed by A1552 (Fig. 3C-E, H and I).

Given that all the other known components required for pilus assembly and

transformation are highly conserved in ATCC25872, we tested whether PilA itself was

responsible for this phenotype by exchanging the endogenous *pilA* open reading frame for

that of A1552 (*pilA*ex). As expected, the ATCC25872 *pilA*ex strain was fully transformable

(Fig. 3B). Importantly, however, in the absence of *pilT*, competence-induced cells of this

strain were now able to form large aggregates, albeit at lower levels than in A1552 (Fig. 3C, F

and G), and intermix within aggregates formed by A1552 (Fig. 3J and K).

PilA variability governs auto-aggregation and enables kin-recognition

The results above suggest that the ability to aggregate is dependent on the particular PilA

variant carried. Indeed, BLAST analyses of 647 *V. cholerae* genomes deposited in NCBI

indicated that PilA exhibits extensive variation, whereas the other proteins encoded within

its operon, as well as those from neighbouring genes, are all highly conserved (Fig. S9). Of

the 636 intact *pilA* genes identified, the majority (492/636) encode a PilA identical to that of

A1552, which is likely due to an over-representation of patient-derived pandemic strains in

the database. Next, we extracted the unique PilA coding sequences (56/636) (Fig. S10A) and

combined them with those of an in-house collection of various environmental and patient

isolates. The resulting phylogenetic tree consists of around 12 distinct groups and was used

to sample PilA diversity (Fig. 4A). As expected, the N-terminal ~50 residues, which contain

sequences required for membrane trafficking and pre-pilin peptidase recognition, and that

goes on to form the long structural alpha helix (α1), are well conserved (Fig. S10A and B). In

contrast, apart from a few clusters of key conserved residues, including the C-terminal

cysteines that form the characteristic disulphide-bonded loop, the C-terminal domains are

highly variable (Fig. S10A and B).

To test how PilA variability affects pilus function but avoid the potential problems of

working in various strain backgrounds, we inserted new *pilA* alleles at the native *pilA* locus in

strain A1552, and used a short (30 bp) duplication of 3' end of the original *pilA* to maintain

any regulation of the downstream genes. We validated this *pilA* replacement (*pilA*rep)

approach using A1552 PilA (*i.e.* Tn*tfoX*, *pilA*rep[A1552]), which is fully transformable, and in

the absence of *pilT* aggregates at levels similar to the unmodified parent (Fig. 4B and C). We

then tested 16 different PilA sequences from across the tree (Fig. 4A and Fig. S10B).

Interestingly, all were equally capable of supporting transformation (Fig. 4B). In contrast, the

ability to aggregate varied depending on the particular PilA. Indeed, in the absence of

retraction, 9/16 PilAs supported the aggregation phenotype, though PilA DRC186-4 was

intermediate, whereas 7/16 did not detectably aggregate (Fig. 4A and C).

Given that aggregation occurs via direct pilus-pilus interactions we hypothesised

that the variability between the different aggregation-proficient PilA might allow pili

composed of different PilA to distinguish between one another. To test this idea we used

the same co-culture approach as before, using strains with and without constitutive GFP

production, and examined all possible combinations (Fig. 4D). As expected, cells producing

pili composed of an identical PilA always exhibited uniform mixing (Fig. 4D and E).

Remarkably, however, in 41/45 possible unique combinations the interactions between

different pili were highly specific (Fig. 4D). Indeed, cells of these strains aggregated in a pilin-

specific manner, preferentially forming aggregates with cells producing pili composed of the

same PilA. This resulted in aggregates that were either almost exclusively fluorescent or

non-fluorescent (Fig. 4F). In 4/45 cases a partial cross-interaction was observed, resulting in

an intermediate mixing phenotype, with aggregates composed of smaller but still segregated

groups of cells (Fig. 4D and G). Overall these data indicate that pili composed of different

PilA are able to discriminate between one another, probably via specific PilA-PilA

interactions. Moreover, these data demonstrate that PilA variability not only determines the

ability to aggregate, but also provides a mechanism for kin-recognition.

The TCP is also capable of aggregation under conditions mimicking virulence

induction[18,57]. Indeed, in strains artificially induced for virulence TCP-mediated aggregates

readily formed that appeared similar to those described in this work (Fig. S11A and C). As

expected, they did not intermix with aggregates formed by DNA-uptake pili (Fig. S11B-D). Of

note, however, is that TCP are largely limited to pandemic lineages with 2 variants of its

major pilin TcpA: Classical and El Tor[58-60]. In contrast to the results above and in line with

previous work[58], a classical and El Tor strain formed uniformly mixed TCP aggregates (Fig.

S11E), indicating that the different TcpA do not discriminate between one another.

278   The unusual tail of ATCC25872/V52 PilA is an inbuilt inhibitor of aggregation

To rule out the possibility that strains unable to aggregate simply fail to make sufficient

numbers of pili, we made labelable *pilA*rep cysteine variants of A1552 PilA (*pilA*rep[A1552;

S67C]) and 2 non-aggregating alleles (*i.e. pilA*rep[Sa5Y; S67C] and *pilA*rep[V52; N67C]). All

constructs behaved similarly to the equivalent non-cysteine variant, though the

*pilA*rep[Sa5Y; S67C] has a modest transformation defect (Fig. S12A and B). Both the

PilA[Sa5Y] and PilA[V52] cysteine variants were piliated at similar levels to the A1552 control

and in the absence of *pilT* were hyper-piliated, although pili composed of PilA[Sa5Y] were

generally very short (Fig. 5A and B). The hyper-piliation was especially clear for the V52

cysteine variant and provides direct evidence that the inability of some alleles to aggregate

is likely not due to a failure to produce sufficient numbers of pili but reflects a specific

property of the pilin itself. Notably, sheared pili of the *pilA*rep[A1552; S67C] variant

assembled into large structures, as above, whereas those of the others did not (Fig. S12C).

Strains V52 and ATCC25872 encode an identical PilA with an unusual repetitive C-

terminal extension (SGSGSGSGSGSGSGSGSGSGN), or 'tail'. Among the PilA sequences analysed

here, there were 9 examples of this type of tail clustered into 2 well-separated phylogenetic

groups (Fig. S8). Moreover, of the *pilA*rep PilAs unable to support aggregation 3/7 contain a

tail (Fig. S8B). Thus, to test if the tail affected V52 PilA function we created a truncated

variant *i.e. pilA*rep[V52Δtail], which was similarly transformable to the equivalent tailed

version (Fig. 5C). Remarkably, however, removal of the tail restored the ability to aggregate

in the absence of *pilT*, at levels indistinguishable from that of the PilA[A1552] controls (Fig.

5D), and demonstrated a similar ability to 'recognise' itself in co-culture experiments (Fig.

5E). These data suggest that the tail inhibits the ability of pili to aggregate, possibly by

masking the site of pilus-pilus interaction. Indeed, a strain in which this tail was transplanted

onto PilA of A1552 (*i.e. pilA*rep[A1552+tail] remained highly transformable but was unable

to aggregate (Fig. 5C and D). However, since efforts to label this PilA variant have so far been

unsuccessful we cannot exclude another, non-specific effect.

DNA-uptake pili form networks on chitin surfaces

The data so far indicates that interactions between pili can mediate intercellular contacts.

However, in liquid culture this relies on blocking pilus retraction. We reasoned that if these

interactions are relevant to the normal ecology of *V. cholerae* then we might be able to

detect them under more realistic conditions. Therefore, we visualised pili produced by

otherwise WT cells upon cultivation on chitinous surfaces (Fig. 6A). Strikingly, under these

conditions of natural induction, cells colonising the chitin surface were generally found

embedded within dense networks of pili that were often overlaid by larger pili structures

(Fig. 6A). In contrast to the chitin-independent liquid culture experiments, cells on the chitin

surface often appeared to possess multiple pili simultaneously (Fig. 6B), though the crowded

nature of the cells precluded direct quantification. Furthermore, since the washing steps

required for pilus staining tend to remove cells, but not pili, from the surface, the number of

cells engaged in these networks is likely underestimated. Control experiments confirmed

that labelling on surfaces is specific and that these structures are DNA-uptake pili (Fig. S13A-

C). Notably, ΔpilT cells are heavily hyper-piliated under these natural induction conditions

(Fig. S13D), indicating that the phenotypes observed in liquid culture are not an artefact of

artificial competence induction.

Next, using the pilArep cysteine variants described above, we directly compared the

pili assembled on chitin surfaces by cells producing the aggregation-proficient PilA[A1552]

with those producing the aggregation-deficient PilA[V52]. Consistent with its functionality in

the various assays above, cells carrying pilArep[A1552; S67C] were also found within similar

pili networks (Fig. 6C and E). In contrast, such networks were never observed for cells

carrying pilArep[V52; N67C] (Fig. 6D and F). Instead, the surfaces were coated in what

appeared to be mostly individual pili. This difference was particularly clear at later time-

points when cells presumably began to exhaust the chitin surface. Cells producing

PilA[A1552] covered surfaces with extensive and long-range networks of pili that were

maintained within detached pieces of biofilm (Fig. S14A and C). In contrast, for cells

producing PilA[V52] the surfaces were coated with individual pili and never large networks

(Fig. S14B and D). In summary, these data suggest that the aggregation phenotypes

observed in liquid culture experiments report the natural ability of pili to interact on chitin

surfaces.

Discussion

Here we demonstrate that DNA-uptake pili are highly dynamic, that these dynamics are PilT-

dependent, and that cells lacking pilT are minimally transformable, providing direct evidence

for the longstanding model, whereby pilus retraction facilitates DNA-uptake. Indeed, our

results on pilus dynamics are in close agreement with those recently reported by Ellison *et*

al., who notably, went on to demonstrate that the pilus binds directly to DNA[61]. The major

finding of this work, however, is that DNA-uptake pili are able to interact and distinguish

between one another in a specific manner depending on their sequence. In liquid culture

when retraction is blocked this manifests as an exaggerated auto-aggregation phenotype.

Since only a subpopulation of cells is piliated at any one time, and these pili are dynamic,

blocking retraction likely facilitates auto-aggregation by producing a homogenous

population of hyper-piliated cells, thereby increasing the chances of interactions between

pili. Work in *Neisseria meningitidis*, which auto-aggregates naturally at low levels but is

dramatically enhanced by the deletion of *pilT*, supports this idea[62,63]. Importantly, however,

under natural induction conditions on chitin surfaces, cells producing pili capable of

aggregation elaborate multiple pili and form dense pili networks in an otherwise unmodified

background, indicating that the chitin surface likely promotes interactions between pili. The

crowded surface environment might inherently foster these interactions. Alternatively, the

altered physiology of cells growing on chitin might also impact pilus assembly via effects on

the extension/retraction motors.

The TCP of *V. cholerae* is induced during virulence and is essential for

colonisation[18,57,64]. Under laboratory conditions TCP production results in auto-aggregation

similar to that described here[17,19,58]. Moreover, in an infection model cells on the intestinal

cell surface were found encased within networks of TCP, that were suggested to protect

cells from host defences[65]. Given the similarities between the two systems, especially our

observation of dense pili networks on colonised chitin surfaces, we propose that the DNA-

uptake pilus might play an analogous role in the aquatic environment. Indeed, as

hypothesised elsewhere[29,66], since pilus production is dependent on an intact chitin-

utilisation pathway, colonisation mechanisms using DNA-uptake pili would be inherently

selective for (i) nutritious chitinous surfaces and (ii) favour the recruitment and retention of

productive cells while excluding non-productive cells unable to make pili. Thus, intercellular

interactions between DNA-uptake pili could act at multiple stages of colonisation, but might

be particularly advantageous during early stages to bring smaller chitin particles together,

and thus provide resistance to protozoan grazers, as well as at later stages to keep cells

together during biofilm dispersal. It remains possible, however, that these interactions

represent an ancient colonisation mechanism that has since been replaced. Arguing against

this idea is the observation that all pandemic strains retain the same interaction-proficient

PilA even though other environmental strains have been free to vary their PilA without

impacting its ability to mediate transformation. Interestingly, ingestion of colonised chitin

particles is thought to facilitate transmission to humans[26] and recovering cholera patients

exhibit a strong immune response to PilA[67]. One possibility that should be investigated is

that the networks of DNA-uptake pili we observed on chitin surfaces protect cells during this

process.

In *Neisseria gonorrhoeae*, artificially varying the density or post-translation

modification state of pili leads to a form of cell sorting based on differential interaction

forces between pili[68]. This effect is likely related to aggregate dispersal during its infective

lifestyle but does not permit specific recognition *per se*[69]. In contrast, the discovery here that

the natural variability of PilA controls the ability of pili to self-interact, and creates highly

specific interactions, provides a direct mechanism for kin recognition. The best-studied

examples of kin recognition in microorganisms all involve adhesins and some form of

aggregation (*e.g.* Flo1; *Saccharomyces cerevisiae*[70], TgrB1-TgrC1; *Dictyostelium discoideum*[71],

and TraA; Myxobacteria[72]). In evolutionary terms, these recognition mechanisms are

classified as 'greenbeards' because the cue, recognition of the cue and the resulting

cooperative activity are all encoded by the same gene[73,74]. The ability of DNA-uptake pili to

recognise and interact with pili composed of the same kind of PilA fits this classification and

is therefore a specific form of greenbeard recognition[73,74]. However, this form of recognition

implies close identity only at the greenbeard locus and so is better referred to as kind

recognition[73,74].

396   An important question going forward will be to understand what drives PilA diversity

and how this is related to the type VI secretion system, which acts to kill non-kin

bacteria[75,76]. Similarly, the apparent acquisition of an inhibitor of pilus interactions by some

PilA (*e.g.* ATCC25872/V52) hints that the ability to interact may not always be beneficial.

Therefore, future work should focus on how the pili networks we observed on chitin

surfaces contribute to the ecology of *V. cholerae*, especially under environmental conditions.

Indeed, we still know relatively little about the natural lifestyle of *V. cholerae* on chitin, in

part due to the inherent technical difficulties associated with manipulating these surfaces.

Nevertheless, the demonstration that in liquid media, specific interactions between pili

composed of different major pilins is sufficient to enable segregation, provides a robust

proof-of-concept that T4P have the ability to function as a recognition mechanism. Finally,

the fact that (i) T4P are widespread, (ii) auto-aggregation via T4P has been reported in

multiple species[77] and (iii) the major pilin subunit often varies[4,52], raises the possibility that

specific interactions between T4P might be quite common and therefore represent an

important contribution to bacterial kin recognition worthy of continued investigation.

Materials and Methods

Bacterial strains and plasmids

The bacterial strains used in this study are shown in Supplementary Table S1, together with

the plasmids used and their construction. A1552, the *V. cholerae* strain used throughout this

work[78], is a toxigenic O1 El Tor Inaba strain representative of the on-going 7[th] cholera

pandemic, and was derived from a traveller entering the United States after being infected

on a commercial aeroplane that took off in Peru[79].

General methods

Bacterial cultures were grown aerobically at 30˚C or 37˚C, as required. Liquid medium used

for growing bacterial strains was Lysogeny Broth (LB-Miller; 10 g/L NaCl, Carl Roth,

Switzerland) and solid medium was LB agar. Where indicated, LB-S contained 20 g/L NaCl.

Ampicillin (Amp; 100 μg/mL), gentamicin (Gent; 50 μg/mL), kanamycin (Kan; 75 μg/mL),

streptomycin (Str; 100 μg/mL) and rifampicin (Rif; 100 μg/mL) were used for selection in *E.*

coli and *V. cholerae*, as required. To induce expression from the P_{BAD} promoter, cultures

were grown in media supplemented with 0.2% L-arabinose. Natural transformation of *V.*

cholerae on chitin flakes was done in 0.5x DASW (Defined artificial seawater), supplemented

with vitamins (MEM, Gibco) and 50 mM HEPES, as previously described[30]. Counter-selection

of phenylalanyl-tRNA synthetase (*pheS**) insertions (Trans2 method; see below) was done

on medium supplemented with 20 mM 4-chloro-phenylalanine (cPhe; Sigma-Aldrich,

Switzerland). Thiosulfate citrate bile salts sucrose (TCBS; Sigma-Aldrich, Switzerland) agar

was used to counter-select for *E. coli* following bacterial mating. SacB-based counter-

selection was done on NaCl-free medium containing 10 % sucrose.

Strain construction

DNA manipulations and *E. coli* transformations were carried out using standard methods[80],

and all constructs were verified by PCR and Sanger sequencing (Microsynth AG, Switzerland).

Genetic engineering of *V. cholerae* was done using a combination of natural transformation

and FLP-recombination; Trans-FLP[81,82], *pheS**-based counter-selection; Trans2[83,84], and allelic

exchange using bi-parental mating and the counter-selectable plasmid pGP704-Sac28[29]. The

mini-Tn7 transposon carrying *araC* and various P_{BAD}-driven genes was integrated into the

large chromosome by tri-parental mating, as previously described[85].

Chitin-independent competence induction

Chitin oligosaccharides resulting from growth on chitin trigger natural competence induction via the production of a master regulator, TfoX[29,30,86-88]. High cell-density, as sensed by quorum sensing via HapR, results in the production of an intermediate regulator QstR, which acts in concert with TfoX to regulate the transcription of a subset of the competence genes[30,41,89]. Therefore, to induce natural competence in liquid culture we used a well characterised and already validated chitin-independent approach that results in low levels of TfoX production[41]. This approach is based on the integration of a mini-Tn7 transposon into the large chromosome of *V. cholerae* containing an arabinose-inducible copy of *tfoX* (*i.e. araC*, P_{BAD}-*tfoX*), which we refer to as Tn*tfoX*. In the presence of inducer, strains carrying Tn*tfoX* turn on the expression of the competence genes according to the known regulatory pathways and upon reaching high cell-density are transformable at levels similar to those seen on chitin[41]. In the absence of inducer, competence genes are not produced and strains are non-transformable[41].

Transformation frequency assay

Diverse strains harbouring Tn*tfoX* were tested for transformation using a chitin-independent transformation frequency assay, as previously described[32,41]. Briefly, overnight cultures were back-diluted 1:100 in fresh media with and without arabinose, as indicated, and grown 3h at 30˚C with shaking (180 rpm). 0.5 mL aliquots of the cultures were mixed with 1 µg genomic DNA (GC#135; A1552-lacZ-Kan) in 1.5 mL eppendorf tubes and incubated 5h at 30˚C with shaking (180 rpm), prior to serial dilution in PBS (Phosphate buffered saline) and enumeration after overnight growth on LB media in the absence and presence of kanamycin. Transformation frequency was calculated as the number of transformants divided by the total number of bacteria.

Pilus shearing assay

Cultures were grown for 6h at 30˚C with shaking (180 rpm) in 25 mL LB + 0.2% arabinose within a 125 mL Erlenmeyer flask. To shear pili from the cell surface, 10 mL culture was removed, vortexed at max speed for 1 min, and cells removed by three sequential centrifugation steps (10min; 4000 x g; 4˚C). To the resulting supernatant saturated ammonium sulphate was added to 40% and incubated on ice for 1h. Precipitated proteins were recovered by centrifugation (30min; 20,000 x g; 4˚C) and washed once with PBS. Samples were then re-suspended in 2x Laemmli buffer, boiled (15min; 95˚C) and stored at -20˚C until needed. To compare PilA levels between samples the re-suspension volume was normalised according to the optical density of the starting culture. Total protein controls were intact cell lysates. The relative amount of PilA in each sample was determined by Western blotting.

Aggregation assay

Overnight cultures were back-diluted 1:100 in the absence and presence of arabinose, as needed, and grown in 14 mL round bottom polystyrene test tubes (Falcon, Corning) on a carousel style rotary wheel (40 rpm) at 30˚C. After 6h growth, aggregates were allowed to settle by standing the tube at RT for 30 min. The optical density at 600 nm (O.D.$_{600}$) of the culture was then measured before and after mechanical disruption (vortex max speed; ~5 sec), which served to disperse any settled aggregates and return them to solution. Aggregation is expressed as the ratio of the O.D.$_{600}$ Pre/Post-vortexing. For time-course experiments the standing time was reduced to 5 min. To visualise aggregates by microscopy overnight cultures were back-diluted either 1:100 individually or 1:200 when mixed, as needed, and were grown for 4h, as described above.

Microscopy

Cells were mounted on microscope slides coated with a thin agarose pad (1.2% w/v in PBS), covered with a #1 cover-slip, and were observed using a Zeiss Axio Imager M2 epi-fluorescence microscope attached to an AxioCam MRm camera and controlled by Zeiss AxioVision software. Image acquisition was done using a Plan-Apochromat 100×/1.4 Ph3 oil objective illuminated by an HXP120 lamp. Images were analysed and prepared for publication using ImageJ (http://rsb.info.nih.gov/ij).

Pilus staining and quantification

Alexa Fluor™ 488 C_5 Maleimide (AF-488-Mal; Thermo Fisher Scientific; Cat# A10254) was dissolved in DMSO, aliquoted and stored at -20˚C protected from light. Cultures used for staining were grown for 3.5h at 30˚C on the rotary wheel, as above, in the absence and presence of competence induction, as required. To stain cells 100 µL of culture was mixed with dye at a final concentration of 25 µg/mL[39] and incubated at RT for 5 min in the dark. Stained cells were harvested by centrifugation (5000 x *g*; 1 min), washed once with LB, re-suspended in 200 µL LB and imaged immediately. For quantification of piliation in snapshot imaging approximately 2000 cells per strain were analysed in each of three independent repeats. A subset of this data set was also analysed for the number of pili per cell and pilus length, as indicated in the text. For time-lapse analysis, images were acquired at RT at 10 sec intervals for 1 min (*i.e.* 7 frames). To analyse the number of pili produced per cell, per min, 5 fields of cells were analysed for each of the three independent repeats, yielding an analysis of 1947 cells in total.

Imaging on chitin surfaces

Visualisation of pilus production on chitin surfaces was done using chitin beads (NEB; Cat#S6651), which have previously been validated as a useful analogue for the natural chitin

surface[41,75]. Prior to use, chitin beads were washed 5x with 0.5x DASW + 50 mM HEPES +

vitamins. Colonisation of chitin beads was done in 12-well culture plates (Cellstar) by mixing

0.1mL washed O/N culture with 0.1 mL washed beads in a final volume of 1 mL 0.5x DASW +

50 mM HEPES + vitamins, and incubated at 30˚C; 48h. Pilus staining was done as described

above except that the final re-suspension volume was reduced to 50 µL. Stained beads were

mounted directly on glass microscope slides, covered with a #1 cover-slip and imaged using

either a Plan-Apochromat 100x/1.4 Ph3 or Plan-Neofluar 40x/1.3 Ph3 oil objective, as

needed. To image cells after prolonged incubation (≥72h), plates were incubated within a

homemade humidified chamber. To avoid damaging the beads manipulations were done

using wide bore tips.

Western blotting

Cell lysates were prepared by suspending harvested cells in 2x Laemmli buffer (100 µL buffer

per O.D. unit) before boiling at 95˚C for 15 min. Proteins were separated by SDS-PAGE using

a 15% resolving gel and blotted onto PVDF membranes using a wet-transfer apparatus.

Immuno-detection was performed as described previously[41]. Primary anti-PilA antibodies

were raised in rabbits against synthetic peptides of A1552 PilA (Eurogentec, Belgium;

#1510525) and used at a dilution of 1:5000. Anti-Rabbit IgG HRP (Sigma; Cat# A9169) diluted

1:5000 was used as a secondary antibody. Sample loading was verified with anti-RNA

Sigma70-HRP (BioLegend; Cat# 663205) diluted 1:10,000.

Motility assay

To quantity motility phenotypes 2 µL overnight culture was spotted soft LB agar (0.3%)

plates (two technical replicates) and incubated at RT for 24h prior to photography. The

swarming diameter (cm) was measured and is expressed as the mean of three independent

biological repeats. A flagellin-deficient (Δ*flaA*) non-motile strain was used as a negative

control.

Bioinformatics of PilA diversity

Vibrio cholerae genomes were obtained from NCBI (National Center for Biotechnology

Information), and are listed in Supplementary File S1. Geneious software (10.2.3)[90] was used

to perform custom BLAST analyses and identify *pilA*. Unique PilA sequences were extracted

and combined with the PilA sequences from strain A1552 and a collection of environmental

isolates, as deduced by Sanger sequencing (Supplementary File S2), as indicated in the text.

PilA sequences were aligned with Muscle and a consensus neighbour-joining tree

constructed using the Jukes-Cantor substitution model, resampled with 100 bootstrap

replicates. MshA from strain A1552 was used as an outgroup.

Reproducibility

All data shown are representative of the results of at least three independent biological

repeats.

Acknowledgements

We thank Ivan Mateus-Gonzalez for assistance with bioinformatics analyses. We further

thank A. Boehm, S. Pukatzki, J. Mekalanos, and members of the Institut National de

Recherche Biomédicale of the Democratic Republic of the Congo, for providing *V. cholerae*

strains and V. Pelicic for advice on *pheS*-mediated counter-selection. Work on this problem

was supported by a Marie Skłodowska-Curie Individual Fellowship (703340; CMDNAUP) to

D.W.A. and by EPFL intramural funding and a Starting Grant from the European Research

Council (ERC; 309064-VIR4ENV) to MB. M.B. is a Howard Hughes Medical Institute (HHMI)

International Research Scholar (Grant# 55008726).

Author contributions

Conception, design and analysis: D.W.A and M.B. Performed research: D.W.A, S.S, C.S and

M.B. Wrote the manuscript: D.W.A and M.B.

Declaration of Interests

The authors declare no competing interests.

580   **References**

581   1   Maier, B. & Wong, G. C. L. How Bacteria Use Type IV Pili Machinery on Surfaces.
582   *Trends in microbiology* **23**, 775-788, doi:10.1016/j.tim.2015.09.002 (2015).
583   2   Persat, A. *et al.* The mechanical world of bacteria. *Cell* **161**, 988-997,
584   doi:10.1016/j.cell.2015.05.005 (2015).
585   3   Berry, J. L. & Pelicic, V. Exceptionally widespread nanomachines composed of type IV
586   pilins: the prokaryotic Swiss Army knives. *FEMS microbiology reviews* **39**, 134-154,
587   doi:10.1093/femsre/fuu001 (2015).
588   4   Giltner, C. L., Nguyen, Y. & Burrows, L. L. Type IV pilin proteins: versatile molecular
589   modules. *Microbiology and molecular biology reviews : MMBR* **76**, 740-772,
590   doi:10.1128/MMBR.00035-12 (2012).
591   5   Melville, S. & Craig, L. Type IV pili in Gram-positive bacteria. *Microbiology and*
592   *molecular biology reviews : MMBR* **77**, 323-341, doi:10.1128/MMBR.00063-12
593   (2013).
594   6   Clemens, J. D., Nair, G. B., Ahmed, T., Qadri, F. & Holmgren, J. Cholera. *Lancet* **390**,
595   1539-1549, doi:10.1016/S0140-6736(17)30559-7 (2017).
596   7   Chang, Y. W. *et al.* Architecture of the *Vibrio cholerae* toxin-coregulated pilus
597   machine revealed by electron cryotomography. *Nature microbiology* **2**, 16269,
598   doi:10.1038/nmicrobiol.2016.269 (2017).
599   8   Chang, Y. W. *et al.* Architecture of the type IVa pilus machine. *Science* **351**, aad2001,
600   doi:10.1126/science.aad2001 (2016).
601   9   Craig, L. *et al.* Type IV pilus structure by cryo-electron microscopy and
602   crystallography: implications for pilus assembly and functions. *Molecular cell* **23**,
603   651-662, doi:10.1016/j.molcel.2006.07.004 (2006).
604   10   Gold, V. A., Salzer, R., Averhoff, B. & Kuhlbrandt, W. Structure of a type IV pilus
605   machinery in the open and closed state. *eLife* **4**, doi:10.7554/eLife.07380 (2015).
606   11   Kolappan, S. *et al.* Structure of the *Neisseria meningitidis* Type IV pilus. *Nature*
607   *communications* **7**, 13015, doi:10.1038/ncomms13015 (2016).
608   12   Merz, A. J., So, M. & Sheetz, M. P. Pilus retraction powers bacterial twitching
609   motility. *Nature* **407**, 98-102, doi:10.1038/35024105 (2000).
610   13   Skerker, J. M. & Berg, H. C. Direct observation of extension and retraction of type IV
611   pili. *Proceedings of the National Academy of Sciences of the United States of America*
612   **98**, 6901-6904, doi:10.1073/pnas.121171698 (2001).
613   14   Jakovljevic, V., Leonardy, S., Hoppert, M. & Sogaard-Andersen, L. PilB and PilT are
614   ATPases acting antagonistically in type IV pilus function in *Myxococcus xanthus*.
615   *Journal of bacteriology* **190**, 2411-2421, doi:10.1128/JB.01793-07 (2008).
616   15   McCallum, M., Tammam, S., Khan, A., Burrows, L. L. & Howell, P. L. The molecular
617   mechanism of the type IVa pilus motors. *Nature communications* **8**, 15091,
618   doi:10.1038/ncomms15091 (2017).
619   16   Waldor, M. K. & Mekalanos, J. J. Lysogenic conversion by a filamentous phage
620   encoding cholera toxin. *Science* **272**, 1910-1914 (1996).
621   17   Chiang, S. L., Taylor, R. K., Koomey, M. & Mekalanos, J. J. Single amino acid
622   substitutions in the N-terminus of *Vibrio cholerae* TcpA affect colonization,
623   autoagglutination, and serum resistance. *Molecular microbiology* **17**, 1133-1142
624   (1995).
625   18   Taylor, R. K., Miller, V. L., Furlong, D. B. & Mekalanos, J. J. Use of *phoA* gene fusions
626   to identify a pilus colonization factor coordinately regulated with cholera toxin.
627   *Proceedings of the National Academy of Sciences of the United States of America* **84**,
628   2833-2837 (1987).

19 Kirn, T. J., Lafferty, M. J., Sandoe, C. M. & Taylor, R. K. Delineation of pilin domains
required for bacterial association into microcolonies and intestinal colonization by
Vibrio cholerae. *Molecular microbiology* **35**, 896-910 (2000).

20 Chiavelli, D. A., Marsh, J. W. & Taylor, R. K. The mannose-sensitive hemagglutinin of
Vibrio cholerae promotes adherence to zooplankton. *Applied and environmental*
microbiology **67**, 3220-3225, doi:10.1128/AEM.67.7.3220-3225.2001 (2001).

21 Moorthy, S. & Watnick, P. I. Genetic evidence that the *Vibrio cholerae* monolayer is a
distinct stage in biofilm development. *Molecular microbiology* **52**, 573-587,
doi:10.1111/j.1365-2958.2004.04000.x (2004).

22 Utada, A. S. *et al. Vibrio cholerae* use pili and flagella synergistically to effect motility
switching and conditional surface attachment. *Nature communications* **5**, 4913,
doi:10.1038/ncomms5913 (2014).

23 Watnick, P. I., Fullner, K. J. & Kolter, R. A role for the mannose-sensitive
hemagglutinin in biofilm formation by *Vibrio cholerae* El Tor. *Journal of bacteriology*
181, 3606-3609 (1999).

24 Watnick, P. I. & Kolter, R. Steps in the development of a *Vibrio cholerae* El Tor
biofilm. *Molecular microbiology* **34**, 586-595 (1999).

25 Tamplin, M. L., Gauzens, A. L., Huq, A., Sack, D. A. & Colwell, R. R. Attachment of
Vibrio cholerae serogroup O1 to zooplankton and phytoplankton of Bangladesh
waters. *Applied and environmental microbiology* **56**, 1977-1980 (1990).

26 Blokesch, M. Competence-induced type VI secretion might foster intestinal
colonization by *Vibrio cholerae*: Intestinal interbacterial killing by competence-
induced *V. cholerae*. *BioEssays : news and reviews in molecular, cellular and*
developmental biology **37**, 1163-1168, doi:10.1002/bies.201500101 (2015).

27 Colwell, R. R. *et al.* Reduction of cholera in Bangladeshi villages by simple filtration.
Proceedings of the National Academy of Sciences of the United States of America
100, 1051-1055, doi:10.1073/pnas.0237386100 (2003).

28 Huq, A. *et al.* A simple filtration method to remove plankton-associated *Vibrio*
cholerae in raw water supplies in developing countries. *Applied and environmental*
microbiology **62**, 2508-2512 (1996).

29 Meibom, K. L. *et al.* The *Vibrio cholerae* chitin utilization program. *Proceedings of the*
National Academy of Sciences of the United States of America **101**, 2524-2529
(2004).

30 Meibom, K. L., Blokesch, M., Dolganov, N. A., Wu, C. Y. & Schoolnik, G. K. Chitin
induces natural competence in *Vibrio cholerae*. *Science* **310**, 1824-1827,
doi:10.1126/science.1120096 (2005).

31 Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J. P. Bacterial
transformation: distribution, shared mechanisms and divergent control. *Nature*
reviews. Microbiology **12**, 181-196, doi:10.1038/nrmicro3199 (2014).

32 Seitz, P. & Blokesch, M. DNA-uptake machinery of naturally competent *Vibrio*
cholerae. *Proceedings of the National Academy of Sciences of the United States of*
America **110**, 17987-17992, doi:10.1073/pnas.1315647110 (2013).

33 Seitz, P. *et al.* ComEA is essential for the transfer of external DNA into the periplasm
in naturally transformable *Vibrio cholerae* cells. *PLoS genetics* **10**, e1004066,
doi:10.1371/journal.pgen.1004066 (2014).

34 Seitz, P. & Blokesch, M. DNA transport across the outer and inner membranes of
naturally transformable *Vibrio cholerae* is spatially but not temporally coupled. *mBio*
5, doi:10.1128/mBio.01409-14 (2014).

35 Wolfgang, M. *et al.* PilT mutations lead to simultaneous defects in competence for
natural transformation and twitching motility in piliated *Neisseria gonorrhoeae*.
Molecular microbiology **29**, 321-330 (1998).

36   Draskovic, I. & Dubnau, D. Biogenesis of a putative channel protein, ComEC, required
for DNA uptake: membrane topology, oligomerization and formation of disulphide
bonds. *Molecular microbiology* **55**, 881-896, doi:10.1111/j.1365-2958.2004.04430.x
(2005).
37   Gangel, H. *et al.* Concerted spatio-temporal dynamics of imported DNA and ComE
DNA uptake protein during gonococcal transformation. *PLoS pathogens* **10**,
e1004043, doi:10.1371/journal.ppat.1004043 (2014).
38   Laurenceau, R. *et al.* A type IV pilus mediates DNA binding during natural
transformation in *Streptococcus pneumoniae*. *PLoS pathogens* **9**, e1003473,
doi:10.1371/journal.ppat.1003473 (2013).
39   Ellison, C. K. *et al.* Obstruction of pilus retraction stimulates bacterial surface
sensing. *Science* **358**, 535-538, doi:10.1126/science.aan5706 (2017).
40   Blair, K. M., Turner, L., Winkelman, J. T., Berg, H. C. & Kearns, D. B. A molecular
clutch disables flagella in the *Bacillus subtilis* biofilm. *Science* **320**, 1636-1638,
doi:10.1126/science.1157877 (2008).
41   Lo Scrudato, M. & Blokesch, M. The regulatory network of natural competence and
transformation of *Vibrio cholerae*. *PLoS genetics* **8**, e1002778,
doi:10.1371/journal.pgen.1002778 (2012).
42   Metzger, L. C. & Blokesch, M. Composition of the DNA-uptake complex of *Vibrio*
cholerae. *Mobile genetic elements* **4**, e28142, doi:10.4161/mge.28142 (2014).
43   Jones, C. J. *et al.* C-di-GMP Regulates Motile to Sessile Transition by Modulating
MshA Pili Biogenesis and Near-Surface Motility Behavior in *Vibrio cholerae*. *PLoS*
pathogens **11**, e1005068, doi:10.1371/journal.ppat.1005068 (2015).
44   Fong, J. C., Syed, K. A., Klose, K. E. & Yildiz, F. H. Role of Vibrio polysaccharide (vps)
genes in VPS production, biofilm formation and *Vibrio cholerae* pathogenesis.
Microbiology **156**, 2757-2769, doi:10.1099/mic.0.040196-0 (2010).
45   Absalon, C., Van Dellen, K. & Watnick, P. I. A communal bacterial adhesin anchors
biofilm and bystander cells to surfaces. *PLoS pathogens* **7**, e1002210,
doi:10.1371/journal.ppat.1002210 (2011).
46   Berk, V. *et al.* Molecular architecture and assembly principles of *Vibrio cholerae*
biofilms. *Science* **337**, 236-239, doi:10.1126/science.1222981 (2012).
47   Kirn, T. J., Jude, B. A. & Taylor, R. K. A colonization factor links *Vibrio cholerae*
environmental survival and human infection. *Nature* **438**, 863-866,
doi:10.1038/nature04249 (2005).
48   Bartlett, T. M. *et al.* A Periplasmic Polymer Curves *Vibrio cholerae* and Promotes
Pathogenesis. *Cell* **168**, 172-185 e115, doi:10.1016/j.cell.2016.12.019 (2017).
49   Gebhart, C. *et al.* Characterization of exogenous bacterial oligosaccharyltransferases
in *Escherichia coli* reveals the potential for O-linked protein glycosylation in *Vibrio*
cholerae and *Burkholderia thailandensis*. *Glycobiology* **22**, 962-974,
doi:10.1093/glycob/cws059 (2012).
50   Biais, N., Ladoux, B., Higashi, D., So, M. & Sheetz, M. Cooperative retraction of
bundled type IV pili enables nanonewton force generation. *PLoS biology* **6**, e87,
doi:10.1371/journal.pbio.0060087 (2008).
51   Joelsson, A., Liu, Z. & Zhu, J. Genetic and phenotypic diversity of quorum-sensing
systems in clinical and environmental isolates of *Vibrio cholerae*. *Infection and*
immunity **74**, 1141-1147, doi:10.1128/IAI.74.2.1141-1147.2006 (2006).
52   Aagesen, A. M. & Häse, C. C. Sequence analyses of type IV pili from *Vibrio cholerae*,
Vibrio parahaemolyticus, and *Vibrio vulnificus*. *Microbial ecology* **64**, 509-524,
doi:10.1007/s00248-012-0021-2 (2012).

53 Aldova, E., Laznickova, K., Stepankova, E. & Lietava, J. Isolation of nonagglutinable
vibrios from an enteritis outbreak in Czechoslovakia. *The Journal of infectious*
diseases **118**, 25-31 (1968).
54 Chun, J. *et al.* Comparative genomics reveals mechanism for short-term and long-
term clonal transitions in pandemic *Vibrio cholerae*. *Proceedings of the National*
Academy of Sciences of the United States of America **106**, 15442-15447,
doi:10.1073/pnas.0907787106 (2009).
55 Blokesch, M. & Schoolnik, G. K. Serogroup conversion of *Vibrio cholerae* in aquatic
reservoirs. *PLoS pathogens* **3**, e81, doi:10.1371/journal.ppat.0030081 (2007).
56 Li, M., Shimada, T., Morris, J. G., Jr., Sulakvelidze, A. & Sozhamannan, S. Evidence for
the emergence of non-O1 and non-O139 *Vibrio cholerae* strains with pathogenic
potential by exchange of O-antigen biosynthesis regions. *Infection and immunity* **70**,
2441-2453 (2002).
57 DiRita, V. J., Neely, M., Taylor, R. K. & Bruss, P. M. Differential expression of the ToxR
regulon in classical and E1 Tor biotypes of *Vibrio cholerae* is due to biotype-specific
control over toxT expression. *Proceedings of the National Academy of Sciences of*
the United States of America **93**, 7991-7995 (1996).
58 Jude, B. A. & Taylor, R. K. The physical basis of type 4 pilus-mediated microcolony
formation by *Vibrio cholerae* O1. *Journal of structural biology* **175**, 1-9,
doi:10.1016/j.jsb.2011.04.008 (2011).
59 Lim, M. S. *et al. Vibrio cholerae* El Tor TcpA crystal structure and mechanism for
pilus-mediated microcolony formation. *Molecular microbiology* **77**, 755-770,
doi:10.1111/j.1365-2958.2010.07244.x (2010).
60 Rhine, J. A. & Taylor, R. K. TcpA pilin sequences and colonization requirements for
O1 and O139 *Vibrio cholerae*. *Molecular microbiology* **13**, 1013-1020 (1994).
61 Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA
uptake during natural transformation in *Vibrio cholerae*. *Nature microbiology* **3**, 773-
780, doi:10.1038/s41564-018-0174-y (2018).
62 Hélaine, S. *et al.* PilX, a pilus-associated protein essential for bacterial aggregation, is
a key to pilus-facilitated attachment of *Neisseria meningitidis* to human cells.
Molecular microbiology **55**, 65-77, doi:10.1111/j.1365-2958.2004.04372.x (2005).
63 Imhaus, A. F. & Duménil, G. The number of *Neisseria meningitidis* type IV pili
determines host cell interaction. *The EMBO journal* **33**, 1767-1783,
doi:10.15252/embj.201488031 (2014).
64 Nielsen, A. T. *et al.* A bistable switch and anatomical site control *Vibrio cholerae*
virulence gene expression in the intestine. *PLoS pathogens* **6**, e1001102,
doi:10.1371/journal.ppat.1001102 (2010).
65 Krebs, S. J. & Taylor, R. K. Protection and attachment of *Vibrio cholerae* mediated by
the toxin-coregulated pilus in the infant mouse model. *Journal of bacteriology* **193**,
5260-5270, doi:10.1128/JB.00378-11 (2011).
66 Shime-Hattori, A. *et al.* Two type IV pili of *Vibrio parahaemolyticus* play different
roles in biofilm formation. *FEMS microbiology letters* **264**, 89-97,
doi:10.1111/j.1574-6968.2006.00438.x (2006).
67 Hang, L. *et al.* Use of in vivo-induced antigen technology (IVIAT) to identify genes
uniquely expressed during human infection with *Vibrio cholerae*. *Proceedings of the*
National Academy of Sciences of the United States of America **100**, 8508-8513,
doi:10.1073/pnas.1431769100 (2003).
68 Oldewurtel, E. R., Kouzel, N., Dewenter, L., Henseler, K. & Maier, B. Differential
interaction forces govern bacterial sorting in early biofilms. *eLife* **4**,
doi:10.7554/eLife.10811 (2015).

69　Chamot-Rooke, J. *et al.* Posttranslational modification of pili upon cell contact
triggers *N. meningitidis* dissemination. *Science* **331**, 778-782,
doi:10.1126/science.1200729 (2011).

70　Smukalla, S. *et al.* FLO1 is a variable green beard gene that drives biofilm-like
cooperation in budding yeast. *Cell* **135**, 726-737, doi:10.1016/j.cell.2008.09.037
(2008).

71　Hirose, S., Benabentos, R., Ho, H. I., Kuspa, A. & Shaulsky, G. Self-recognition in
social amoebae is mediated by allelic pairs of tiger genes. *Science* **333**, 467-470,
doi:10.1126/science.1203903 (2011).

72　Pathak, D. T., Wei, X., Dey, A. & Wall, D. Molecular recognition by a polymorphic cell
surface receptor governs cooperative behaviors in bacteria. *PLoS genetics* **9**,
e1003891, doi:10.1371/journal.pgen.1003891 (2013).

73　Strassmann, J. E., Gilbert, O. M. & Queller, D. C. Kin discrimination and cooperation
in microbes. *Annual review of microbiology* **65**, 349-367,
doi:10.1146/annurev.micro.112408.134109 (2011).

74　Wall, D. Kin Recognition in Bacteria. *Annual review of microbiology* **70**, 143-160,
doi:10.1146/annurev-micro-102215-095325 (2016).

75　Borgeaud, S., Metzger, L. C., Scrignari, T. & Blokesch, M. The type VI secretion
system of *Vibrio cholerae* fosters horizontal gene transfer. *Science* **347**, 63-67,
doi:10.1126/science.1260064 (2015).

76　Ho, B. T., Dong, T. G. & Mekalanos, J. J. A view to a kill: the bacterial type VI
secretion system. *Cell host & microbe* **15**, 9-21, doi:10.1016/j.chom.2013.11.008
(2014).

77　Trunk, T., Khalil, H. S. & Leo, J. C. Bacterial autoaggregation. *AIMS Microbiology* **4**,
140-164, doi:http://dx.doi.org/10.3934/microbiol.2018.1.140 (2018).

78　Yildiz, F. H. & Schoolnik, G. K. Role of *rpoS* in stress survival and virulence of *Vibrio*
cholerae. *Journal of bacteriology* **180**, 773-784 (1998).

79　Blokesch, M. A quorum sensing-mediated switch contributes to natural
transformation of *Vibrio cholerae*. *Mobile genetic elements* **2**, 224-227,
doi:10.4161/mge.22284 (2012).

80　Sambrook, J., Fritsch, E. F. & Maniatis, T. Molecular Cloning: A Laboratory Manual.
Cold Spring Harbor: Cold Spring Harbor Laboratory Press (1989).

81　De Souza Silva, O. & Blokesch, M. Genetic manipulation of *Vibrio cholerae* by
combining natural transformation with FLP recombination. *Plasmid* **64**, 186-195,
doi:10.1016/j.plasmid.2010.08.001 (2010).

82　Marvig, R. L. & Blokesch, M. Natural transformation of *Vibrio cholerae* as a tool -
optimizing the procedure. *BMC microbiology* **10**, 155, doi:10.1186/1471-2180-10-
155 (2010).

83　Van der Henst, C. *et al.* Molecular insights into *Vibrio cholerae*'s intra-amoebal host-
pathogen interactions. *Nature communications*, doi:10.1038/s41467-018-05976-x
(2018).

84　Gurung, I., Berry, J. L., Hall, A. M. J. & Pelicic, V. Cloning-independent markerless
gene editing in *Streptococcus sanguinis*: novel insights in type IV pilus biology.
Nucleic acids research **45**, e40, doi:10.1093/nar/gkw1177 (2017).

85　Bao, Y., Lies, D. P., Fu, H. & Roberts, G. P. An improved Tn7-based system for the
single-copy insertion of cloned genes into chromosomes of gram-negative bacteria.
Gene **109**, 167-168 (1991).

86　Yamamoto, S. *et al.* Regulation of natural competence by the orphan two-
component system sensor kinase ChiS involves a non-canonical transmembrane
regulator in *Vibrio cholerae*. *Molecular microbiology* **91**, 326-347,
doi:10.1111/mmi.12462 (2014).

87 Dalia, A. B., Lazinski, D. W. & Camilli, A. Identification of a membrane-bound
transcriptional regulator that links chitin and natural competence in Vibrio cholerae.
mBio **5**, e01028-01013, doi:10.1128/mBio.01028-13 (2014).
88 Yamamoto, S. *et al.* Identification of a chitin-induced small RNA that regulates
translation of the tfoX gene, encoding a positive regulator of natural competence in
Vibrio cholerae. *Journal of bacteriology* **193**, 1953-1965, doi:10.1128/JB.01340-10
(2011).
89 Lo Scrudato, M. & Blokesch, M. A transcriptional regulator linking quorum sensing
and chitin induction to render *Vibrio cholerae* naturally transformable. *Nucleic acids
research* **41**, 3644-3658, doi:10.1093/nar/gkt041 (2013).
90 Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**,
1647-1649, doi:10.1093/bioinformatics/bts199 (2012).

Figure 1. Direct observation of dynamic DNA-uptake pili.

(**A**) Functionality of PilA cysteine variants (PilA[Cys]) in a chitin-independent transformation assay using strains carrying an arabinose-inducible copy of *tfoX* (Tn*tfoX*). Transformation frequencies are the mean of three independent biological repeats (+S.D.). < d.l., below detection limit. †, < d.l. in one repeat.

(**B**) Effect of PilA cysteine variants on the ability of retraction deficient (Tn*tfoX*, Δ*pilT*) cells to aggregate. Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated. Values are the mean of three independent biological repeats (±S.D.).

(**C**-**G**) Snapshot imaging of pili in cells of A1552-Tn*tfoX*, PilA[S67C] and its derivatives. (**C**) Cells of A1552-Tn*tfoX*, PilA[S67C] were grown in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated, and stained with AF-488-Mal. Bar = 5 μm. (D-E) Cells of A1552-Tn*tfoX*, PilA[S67C] were grown in the presence of *tfoX* induction and stained with AF-488-Mal in a (**D**) Δ*pilB*, (**E**) Δ*pilQ*, (**F**) Δ*pilU* and (**G**) Δ*pilT* background, as indicated. Bar = 5 μm.

(**H**-**J**) Quantification of piliation in snapshot imaging. Bars represent the mean of three repeats (±S.D.). (**H**) Percentage of piliated cells in the indicated backgrounds. *n* = *c.a.* 2000 cells per strain per repeat. (**I**) Histogram of number of pili per cell in piliated cells in WT parent (A1552-Tn*tfoX*, PilA[S67C]), Δ*pilU* (A1552-Tn*tfoX*, PilA[S67C], Δ*pilU*) and Δ*pilT* (A1552-Tn*tfoX*, PilA[S67C], Δ*pilT*) backgrounds, as indicated. *n* = *c.a.* 300 cells per strain per repeat. (**J**) Histogram of pilus length in cells of A1552-Tn*tfoX*, PilA[S67C]. *n* = *c.a.* 500-600 cells per repeat.

(**K**-**L**) Time-lapse series of pilus dynamics in (**K**) WT parent (A1552-Tn*tfoX*, PilA[S67C]) and (**L**) Δ*pilT* (A1552-Tn*tfoX*, PilA[S67C], Δ*pilT*) backgrounds. Cells were stained with AF-488-Mal and imaged at 10s intervals for 1 minute. Upper panels show phase-contrast (PC), lower panels show fluorescence (dye). Time in seconds (s), as indicated. Bars = 5 μm.

(**M**) Histogram showing quantification of pili produced per cell per min in the WT parent (A1552-Tn*tfoX*, PilA[S67C]) background. Bars represent the mean of three repeats (±S.D.). *n* = *c.a.* 500-700 cells per repeat.

Figure 2. Competent cells auto-aggregate in the absence of pilus retraction.

(**A**-**C**) Phase-contrast microscopy of cells of strains (**A**) A1552-Tn*tfoX*, (**B**) A1552-Tn*tfoX*, Δ*pilT* and (**C**) A1552-Tn*tfoX*, Δ*pilT*, *pilT*+, grown in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated. Scale bar = 25 μm.

(**D**-**E**) Aggregation and transformation frequency of cells of strains A1552-Tn*tfoX*, A1552-Tn*tfoX*, Δ*pilT* and A1552-Tn*tfoX*, Δ*pilT*, *pilT*+, grown in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated. A1552 (without inducible *tfoX*) was used as a negative control. (**D**) Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing. Values are the mean of three repeats (±S.D.). (**E**) Chitin-independent transformation frequency assay. Values are the mean of three repeats (+S.D.). < d.l., below detection limit.

(**F**) Effect of various deletion backgrounds on the ability of retraction deficient cells to aggregate. Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated. Δ4 = Δ*tcpA*, Δ*mshA*, Δ*vpsA*, Δ*flaA* quadruple mutant. Values are the mean of three repeats (±S.D.).

(**G**-**I**) Co-culture of fluorescent Δ*pilT* cells (A1552-Tn*tfoX*, Δ*pilT*, GFP+) producing GFP and non-fluorescent cells of the (**G**) WT parent (A1552-Tn*tfoX*), (**H**) Δ*pilT* (A1552-Tn*tfoX*, Δ*pilT*) and (**I**) Δ*pilA*, Δ*pilT* (A1552-Tn*tfoX*, Δ*pilA*, Δ*pilT*), grown in the presence of *tfoX* induction. Merged images show GFP in green and phase-contrast in red. Bar = 25 μm.

Figure 3. A1552 PilA is sufficient for aggregation in a non-pandemic strain.

(**A**) Aggregation of representative 7[th] pandemic strains of *V. cholerae*, including the effect of *hapR*[Rep] on N16961, in a Tn*tfoX* and a Tn*tfoX*, Δ*pilT* background, as indicated. Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the presence of *tfoX* induction. Values are the mean of three repeats (±S.D.).

(**B**-**C**) Transformation frequency and aggregation of *V. cholerae* strain ATCC25872-Tn*tfoX* compared to that of A1552-Tn*tfoX*. (**B**) Chitin-independent transformation assay. Transformation frequencies are the mean of three repeats (+S.D.). (**C**) Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the absence (- Ara) and presence (+ Ara) of *tfoX* induction, as indicated. Values are the mean of three repeats (±S.D.).

(**D**-**G**) Phase-contrast microscopy of ATCC25872-Tn*tfoX*, Δ*vpsA* cells carrying (**D** and **E**) their native *pilA* (*pilA* WT) and (**F** and **G**) A1552 *pilA* (*pilA*ex), in a (**D** and **F**) *pilT*+ and (**E** and **G**) Δ*pilT* background, as indicated. Strains were cultured in the absence (- Ara) and presence (+ Ara), as indicated. Bar = 25 μm. Note that ATCC25872 derivatives were co-deleted for *vpsA* to rule out any compounding effects of biofilm formation.

(**H**-**K**) Co-culture of fluorescent cells of A1552-Tn*tfoX*, Δ*pilT*, GFP+, producing GFP, and non-fluorescent cells of ATCC25872-Tn*tfoX*, Δ*vpsA* carrying (**H** and **I**) their native *pilA* (*pilA* WT) and (**J** and **K**) A1552 PilA (*pilA*ex), in a (**H** and **J**) *pilT*+ and (**I** and **K**) Δ*pilT* background, as indicated. Cells were grown in the presence of *tfoX* induction. Merged images show GFP in green and phase-contrast (PC) in red. Bar = 25 μm.

Figure 4

Figure 4. PilA variability governs auto-aggregation and enables kin-recognition.

(**A**) Consensus neighbour-joining phylogenetic tree of PilA. The tree consists of the 56 unique PilA sequences identified from NCBI, 22 sequences from an in-house collection of various environmental and patient isolates (bold), and A1552 PilA (blue) and MshA (outgroup). Values shown are consensus support values (%). Aggregation capable (green) and incapable (red) PilAs tested in the *pilA*rep experiments are highlighted. Note that ATCC25872 and V52 PilA are identical.

(**B**-**C**) Functionality of A1552-Tn*tfoX*, *pilA*rep[A1552] and 16 different PilAs assessed by transformation frequency and aggregation. (**B**) Chitin-independent transformation frequency assay. WT parent (A1552-Tn*tfoX*) and Δ*pilA* (A1552-Tn*tfoX*, Δ*pilA*) strains served as positive and negative controls. Transformation frequencies are the mean of three repeats (+S.D.). †, <d.l. in one repeat. (**C**) Aggregation was determined for the WT parent and each *pilA*rep strain in a *pilT*+ (A1552-Tn*tfoX*) and Δ*pilT* (A1552-Tn*tfoX*, Δ*pilT*) background, as indicated. Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the presence of *tfoX* induction. Values are the mean of three repeats (±S.D.).

(**D**) Interaction matrix showing the results of a pairwise analysis of all possible interactions between aggregation proficient PilA. Interactions were tested by co-culturing non-fluorescent cells of the relevant Tn*tfoX*, Δ*pilT*, pilA*rep*[] strains with a fluorescent (GFP+) derivative of each strain. Cells were grown in the presence of *tfoX* induction. Self-self combinations served as controls.

(**E**-**G**) Representative examples of (**E**) interaction, resulting in well-mixed aggregates (**F**) no interaction, resulting in fluorescent or non-fluorescent aggregates and (**G**) intermediate interactions, resulting in patterned aggregates. Cells were grown in the presence of *tfoX* induction. Merged images show GFP in green and phase-contrast (PC) in red. Bar = 25 µm.

A

B

A1552-Tn*tfoX*, *pilA*rep[ ]

C

E

A1552-Tn*tfoX*, *pilA*rep[ ]

vs. Tn*tfoX*, Δ*pilT*, GFP+

D

Figure 5. The unusual tail of ATCC25872/V52 PilA inhibits aggregation.

(**A**) Quantification of piliation in snapshot imaging of cells of strains A1552-Tn*tfoX*, *pilA*rep[A1552; S67C], A1552-Tn*tfoX*, *pilA*rep[Sa5Y; S67C] and A1552-Tn*tfoX*, *pilA*rep[V52; N67C], in the indicated backgrounds. Cells were grown with *tfoX* induction and pili stained with AF-488-Mal. Bars represent the mean of three repeats (±S.D.). *n* = *c.a.* 2000 cells per strain per repeat. (**B**) Direct observation of pili stained with AF-488-Mal in cells carrying *pilA*rep[Sa5Y; S67C] and *pilA*rep[V52; N67C], in a WT parent (A1552-Tn*tfoX*) and Δ*pilT* (A1552-Tn*tfoX*, Δ*pilT*) background, as indicated. Bar = 5 μm.

(**C**-**D**) Functionality of *pilA*rep tail variants assessed by natural transformation and aggregation. (**C**) Chitin-independent transformation assay. WT parent (A1552-Tn*tfoX*) and Δ*pilA* (A1552-Tn*tfoX*, Δ*pilA*) strains served as positive and negative controls. Transformation frequencies are the mean of three repeats (+S.D.). (**D**) Aggregation was determined for the WT parent and each *pilA*rep strain in a *pilT*+ (A1552-Tn*tfoX*) and Δ*pilT* (A1552-Tn*tfoX*, Δ*pilT*) background, as indicated. Aggregation is shown as the ratio of the culture optical density (O.D. $_{600}$) before and after vortexing, in the presence of *tfoX* induction. Values are the mean of three repeats (±S.D.).

(**E**) Co-culture of fluorescent cells of A1552-Tn*tfoX*, Δ*pilT*, GFP+, producing GFP, and non-fluorescent cells of either *pilA*rep[V52] (A1552-Tn*tfoX*, Δ*pilT*, *pilA*rep[V52]) or *pilA*rep[V52Δtail] (A1552-Tn*tfoX*, Δ*pilT*, *pilA*rep[V52Δtail]), as indicated. Note that ATCC25872 and V52 PilA are identical. Cells were grown in the presence of *tfoX* induction. Merged images show GFP in green and phase-contrast in red. Bar = 25 μm.

A

Tn*tfoX*, PilA[S67C]

Phase

Dye

B

Tn*tfoX*, *pilA*rep[Cys]

*pilA*rep[A1552; S67C]   *pilA*rep[V52; N67C]

C   **D**

E   **F**

Phase   Dye   Phase   Dye

Figure 6. DNA-uptake pili form networks on chitin surfaces.

(**A-B**) Chitin beads were stained with AF-488-Mal after incubation for 48h in defined artificial seawater (DASW) with cells of A1552-Tn*tfoX*, PilA[S67C], as indicated. The figure depicts two separate examples of colonised surfaces, bars = 25 µm. (**B**) Insets show enlarged examples of cells producing multiple pili. Bar = 5 µm.

(**C**-**F**) Chitin beads were stained with AF-488-Mal after incubation for 48h in DASW with cells of either (**C**) A1552-Tn*tfoX*, *pilA*rep[A1552; S67C] or (**D**) A1552-Tn*tfoX*, *pilA*rep[V52; N67C], as indicated. Bar = 25 µm. Panels (**E** and **F**) show enlargements of the surfaces shown in (**C** and **D**). Note the absence of large pili networks in (**F**). Bar = 5 µm.