High-resolution genetic map and QTL analysis of growth-related traits of *Hevea brasiliensis* cultivated under suboptimal temperature and humidity conditions

André Ricardo Oliveira Conson[1#], Cristiane Hayumi Taniguti[2#], Rodrigo Rampazo Amadeu[2#],
Isabela Aparecida Araújo Andreotti[1], Livia Moura de Souza[1], Luciano Henrique Braz dos Santos[1],
João Ricardo Bachega Feijó Rosa[2,3&], Camila Campos Mantello[1,4§], Carla Cristina da Silva[1], Erivaldo
José Scaloppi Junior[5], Rafael Vasconcelos Ribeiro[6], Vincent Le Guen[7], Antonio Augusto Franco
Garcia[2], Paulo de Souza Gonçalves[5], Anete Pereira de Souza[1,6*]

[1]Molecular Biology and Genetic Engineering Center, University of Campinas, Campinas, Brazil

[2]Department of Genetics, Luiz de Queiroz College of Agriculture, University of São Paulo,
Piracicaba, Brazil

[3]FTS Sementes S.A., Research and Development Center, Ponta Grossa, Brazil

[4]National Institute of Agricultural Botany (NIAB), Huntingdon Road, Cambridge CB3 0LE, UK

[5]Center of Rubber Tree and Agroforestry Systems, Agronomic Institute (IAC), Votuporanga, Brazil

[6]Department of Plant Biology, Institute of Biology, University of Campinas, Campinas, Brazil

[7]French Agricultural Research Centre for International Development (CIRAD), UMR AGAP,
Montpellier, France

[&] Present Address: FTS Sementes S.A., Research and Development Center, Ponta Grossa, Brazil

[§] Present Address: National Institute of Agricultural Botany (NIAB), Huntingdon Road, Cambridge
CB3 0LE, UK

*Correspondence:

Anete Pereira de Souza

anete@unicamp.br

[#]These authors contributed equally

Keywords

Rubber tree, linkage map, GBS, drought stress, SSR, suboptimal temperature and humidity

Abstract

Rubber tree (*Hevea brasiliensis*) cultivation is the main source of natural rubber worldwide and has
been extended to areas with suboptimal climates and lengthy drought periods; this transition affects
growth and latex production. High-density genetic maps with reliable markers support precise
mapping of quantitative trait loci (QTL), which can help reveal the complex genome of the species,
provide tools to enhance molecular breeding, and shorten the breeding cycle. In this study, QTL
mapping of the stem diameter, tree height, and number of whorls was performed for a full-sibling
population derived from a GT1 and RRIM701 cross. A total of 225 simple sequence repeats (SSRs)
and 186 single-nucleotide polymorphism (SNP) markers were used to construct a base map with 18
linkage groups and to anchor 671 SNPs from genotyping by sequencing (GBS) to produce a very
dense linkage map with small intervals between loci. The final map was composed of 1,079 markers,
spanned 3,779.7 cM with an average marker density of 3.5 cM, and showed collinearity between
markers from previous studies. Through a composite interval mapping method, a total of 38 QTLs
were identified over a 59-month evaluation period. Linkage group 4 showed the greatest number of
QTLs (7), with phenotypic explained values varying from 7.67% to 14.07%. Additionally, we
estimated segregation patterns, dominance, and additive effects for each QTL. A total of 53
significant effects for diameter were observed, and these effects were mostly related to additivity in
the GT1 clone. Associating accurate genome assemblies and genetic maps represents a promising
strategy for identifying the genetic basis of phenotypic traits in the rubber tree. Therefore, future
investigations related to the growth of *Hevea brasiliensis* will be feasible and can benefit from the
QTLs identified herein to provide a better understanding of the key determinant genes associated
with growth and drought.

Introduction

Hevea brasiliensis (Willd. ex. Adr. de Juss.) Muell-Arg., also known as the rubber tree, is one of the
most symbolic species from the Amazon basin owing to its worldwide use in natural latex
production, the global demand for which suggests expansion of the industry (Warren-Thomas *et al*.,
2015). The species is a perennial and allogamous tree belonging to the Euphorbiaceae family
(Priyadarshan and Clément-Demange, 2004) and contains 36 chromosomes (2n = 36), although
studies indicate that *H. brasiliensis* is an amphidiploid (2n = 4x = 36) that has been stabilized during
the course of evolution (Saha and Priyadarshan, 2012). Because of its genome size and high repeat
content (Lau *et al*., 2016; Tang *et al*., 2016), genetic analysis of the species represents considerable
obstacles to breeding, and several methodologies must be used to better understand its organization
(Fierst, 2015).

Rubber tree plantations are mainly located in Southeast Asia under optimal edaphoclimatic
conditions due to the occurrence of the South America Leaf Blight (SALB) disease in Latin America.
Because the species can be cultivated under suboptimal conditions, plantations in "escape areas",
such as the southeastern region of Brazil, where epidemic SALB has not been found, might be an
alternative for natural rubber production (Gonçalves *et al*., 2011; Souza *et al*., 2013; Jaimes *et al*.,
2016). However, suboptimal conditions (temperature and humidity) affect plant development,
consequently affecting rubber production (Silpi *et al*., 2006; Rao and Kole, 2016).

Developing new rubber tree clones may extend up to 20-30 years and becomes expensive. Molecular
markers can help identify allelic variants related to phenotypes adapted to these regions and can
shorten the breeding cycle (Priyadarshan and Clément-Demange, 2004). Therefore, significant results

obtained for other crops using effective and cost-efficient microsatellite markers associated with
next-generation sequencing (NGS) technologies, such as the simultaneous discovery and genotyping
of thousands of single-nucleotide polymorphisms (SNPs) through genotyping by sequencing (GBS)
(Elshire *et al.*, 2011), can be very useful for *H. brasiliensis*.

Because no inbred lines are available for the rubber tree species, the mapping methods for these
species must be adapted to full-sibling populations, once there are more alleles involved and linkage
phases to be estimated compared with the population from inbred lines. The first and most
widespread method for building genetic maps for outcrossing species is the pseudo-testcross, which
allowed the first genetic mapping studies in such species, but it considers only markers segregating
1:1 and results in separated maps for each parent (Grattapaglia and Sederoff, 1994). Later, several
methods were proposed, including maximum likelihood estimators for all types of markers
segregation (Maliepaard *et al.*, 1997) and the possibility to build integrated genetic maps using
information from all available markers with multipoint estimation (Wu *et al.*, 2002a, 2002b). In the
rubber tree, different methods have been used to obtain higher-resolution genetic maps (Pootakham
et al., 2015; Shearman *et al.*, 2015) than those obtained in previous studies (Lespinasse *et al.*, 2000;
Souza *et al.*, 2013). Methods with higher-coverage genome reference sequences of the species are
also important to detect markers associated with quantitative trait loci (QTL).

In addition to a reliable map, in QTL studies, it is also important to use a robust experimental design.
Forestry experiments require large field areas due to the needed spacing for each tree. Therefore, the
field area size limits the number of possible genotype replications in the design. Augmented blocks
are a very efficient and widely used design proposed to address reduced numbers of replicates
(Federer and Raghavarao, 1975). Joint analysis is possible using information from checks included in
all incomplete blocks. To improve the statistical power in an augmented blocks design, a possible
approach is to use replicated augmented blocks. After modeling and evaluating the experiment, it is
possible to gather the genetic values for each genotype and to use them alongside the map in QTL
mapping methods. The QTL method must also be adapted to outcrossing species. Gazaffi *et al.*
(2014) developed a composite interval mapping (CIM) model for outcrossing species considering
integrated maps. The model uses maximum likelihood and mixture models to estimate three
orthogonal contrasts, which are used to estimate additive and dominance effects and QTL segregation
patterns and linkage phases within the molecular markers. The QTL screening includes cofactors to
avoid influences from QTLs outside the mapping interval.

Here, we report an integrated high-density genetic map for *Hevea brasiliensis* obtained from a set of
high-quality SNP and simple sequence repeat (SSR) markers. Moreover, QTL mapping for complex
traits related to plant growth (stem diameter, tree height, and the number of whorls) was performed
over a 59-month evaluation period, allowing for more accurate QTL localization.

Materials and methods

Mapping population and DNA extraction

An F1 population containing 146 individuals was derived from open pollination between the
genotypes GT1 and RRIM701. The GT1 clone, which has been widely used in *Hevea* breeding since
the 1920s, is classified as a primary clone that is male sterile (Shearman *et al.*, 2015) and tolerant to
wind and cold. From the Asian breeding program, RRIM701 exhibits vigorous growth and an
increased circumference after initial tapping (Romain and Thierry, 2011).

Effective pollination was confirmed using 10 microsatellite markers that were selected based on the
polymorphism information content (PIC) and the polymorphisms between the parents. After
removing two contaminants, the mapping population was planted at the Center of Rubber Tree and
Agroforestry Systems/Agronomic Institute (IAC) in the northwest region of São Paulo state (20º 25'
00" S and 49º 59' 00" W at a 450-meter altitude), Brazil, in 2012. The climate is the Köppen Aw
type (Alvares *et al.*, 2013). Based on air temperature and rainfall, the water balance was calculated
using the method developed by Thornthwaite and Mather (1955), adopting 100 mm as the maximum
soil water holding capacity. An augmented block design was used (Federer and Raghavarao, 1975).
To improve accuracy, the trial (one augmented block design) was repeated side-by-side four times
with different randomizations. Each trial has four blocks and two plants (clones) per plot spaced at 4
m by 4 m. Hereafter, trial is considered a repetition. Therefore, each block consisted of 37 genotypes
of GT1 x RRIM701 crosses and four checks (GT1, PB235, RRIM701 and RRIM600). The
experiment has a total of 656 (41 plots x 4 blocks x 4 repetition) plots and 1,312 trees.

Genomic DNA was extracted from lyophilized leaves using 2% cetyltrimethylammonium bromide
(CTAB) as described by Doyle and Doyle (1987). DNA integrity was assessed by electrophoresis on
an ethidium bromide-stained 1% agarose gel, and its concentration was estimated with a
QuantiFluor-ST fluorometer (Promega, Madison, WI, USA).

Microsatellite analysis

A total of 1,190 SSR markers, comprising 476 genomic markers (Souza *et al.*, 2009; Le Guen *et al.*,
2011; Mantello *et al.*, 2012), 353 expressed sequence tag (EST)-SSR markers (Feng *et al.*, 2009;
Silva *et al.*, 2014), and 361 additional unpublished EST-SSRs identified in cDNA libraries
constructed from subtractive suppression hybridization libraries (Garcia *et al.*, 2011), were tested to
verify polymorphisms between the parents. Subsequently, from 569 (48%) polymorphic markers,
those with a more informative segregation pattern (Wu *et al.*, 2002a) and with a clear profile were
selected, totaling 287 SSRs for the population genotyping.

The amplification of SSRs was performed as previously described by the authors. PCR products were
visualized using three different techniques: (1) silver nitrate staining (Creste *et al.*, 2001); (2) LI-
COR 4300 DNA analyzer (LI-COR Inc., Lincoln, NE, USA), and (3) ABI 3500XL DNA sequencer
(Life Technologies, Carlsbad, CA, USA). Markers that were considered monomorphic or
homozygous for both parents and markers without clear bands were excluded.

SNP analysis

SNP markers were developed from full-length EST libraries (Silva *et al.*, 2014) and a bark
transcriptome (Mantello *et al.*, 2014). A total of 391 SNPs was selected for genotyping using the
Sequenom MassARRAY Assay Design program (Agena Bioscience®, San Diego, CA, USA) with
the following parameters: (1) a high plex preset with a multiplex level = 24, (2) an amplicon length
varying from 80 to 200 bp, and (3) number of iterations = 10. SNPs were selected for population
genotyping using the Sequenom iPLEX MassARRAY® (Sequenom Inc., San Diego, California,
USA) and were annotated within the mevalonate and 2-C-methyl-D-erythritol 4-phosphate (MEP)
pathways, which are involved in latex biosynthesis and wood synthesis (Table S1). The first
amplification reaction was performed using 1-3 ng of genomic DNA, and the subsequent steps were
performed following the manufacturer's instructions. MassArray Typer v4.0 software (Sequenom,
Inc.) was used for allele visualization.

Ninety-six other polymorphic SNPs obtained from genomic (Souza *et al.*, 2016) and transcriptomic
(Salgado *et al.*, 2014) data were also selected for genotyping using the BioMark™ Real-Time PCR
system with the 96.96 Dynamic Array™ IFC (Fluidigm Corporation, San Francisco, CA, USA). For
kompetitive allele-specific PCR (KASP) assays, 1.4 μL of KASP assay reagent, 5 μL of 2X assay
loading reagent, 3.6 μL of ultrapure water, and 2.8 μL of KASP assay mix (12 μM allele-specific
forward primer and 30 μM reverse primer) were used. Furthermore, 4.73 μL of KASP 2X reagent
mix, 0.47 μL of 20X GT sample loading reagent, 0.3 μL of ultrapure water, and 240 ng of DNA were
used for PCR. The thermal conditions for fragment amplification were 94°C for 15 minutes; 94°C for
10 seconds; 57°C for 10 seconds; 72°C for 10 seconds; 10 cycles of 94°C for 20 seconds and 65°C to
57°C for 1 minute (-0.8°C per cycle); and 26 cycles of 94°C for 20 seconds and 57°C for 1 minute.
The genotyping data were analyzed using the BioMark™ SNP Genotyping Analysis v2.1.1 program.

GBS-based SNPs

To saturate the linkage map with more SNPs, we performed the GBS protocol as described by
Elshire *et al.* (2011). GBS libraries were generated from 100 ng of genomic DNA, and the *Eco*T22I
enzyme was used to reduce the genomic complexity. Library quality checks were performed using
the Agilent DNA 1000 Kit with the Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara,
USA). Library sequencing was performed on an Illumina GAIIx platform (Illumina Inc., San Diego,
CA, USA). Quality assessment of the generated sequence data was performed with the NGS QC
Toolkit v2.3.3 (Patel and Jain, 2012).

Sequencing data analysis was performed using TASSEL-GBS v3.0 (Bradbury *et al.*, 2007). Retained
tags with a minimum count of six reads were aligned to the *H. brasiliensis* reference genome
sequence, GenBank: LVXX00000000.1 (Tang et al., 2016), using Bowtie2 version 2.1 (Langmead
and Salzberg, 2012). The obtained variant call format (VCF) files were filtered using VCFtools
(Danecek *et al.*, 2011), and only biallelic SNPs, with a maximum of 25% missing data, were retained.
Additionally, SNPs were filtered by removing redundant markers, segregation-distorted markers, and
markers homozygous for both parents using the R package OneMap v2.1.1 (Margarido *et al.*, 2007).

Genetic mapping

An integrated linkage map was developed in two stages in which all markers were tested for
segregation distortion (p-value < 0.05) with Bonferroni's multiple test correction. All of the
following logarithm of the odds (LOD) thresholds were also obtained based on Bonferroni's multiple
test correction according to the number of two point tests. First, a base map with SSR and SNP
markers (except GBS-based SNPs) was built, and the markers were grouped using an LOD value of
5.37 and a maximum recombination frequency of 0.4. The markers were ordered through a hidden
Markov chain multipoint approach (Lander and Green, 1987) considering outcrossing species (Wu *et*
al., 2002b). The most informative markers (1:1:1:1) were ordered using exhaustive search, and the
remaining markers were positioned using the TRY algorithm (Lander and Green, 1987). The RIPPLE
algorithm (Lander and Green, 1987) was used to improve final orders. The marker order of this first
map was fixed before insertion of the GBS-based markers. Later, the GBS markers were grouped
into the base map considering an LOD of 7.22 and a maximum recombination fraction of 0.4. The
best position for each marker was identified using the TRY and RIPPLE algorithms. All of the
analyses were performed using OneMap software v2.1.1 (Margarido *et al.*, 2007; Mollinari *et al.*,
2009). The Kosambi mapping function was used (Kosambi, 1943) to estimate map distances.

Phenotypic analysis

Tree height, number of whorls, and stem diameter at 50 cm above ground level were measured to evaluate the growth of individual trees, with mean values calculated based on two plants per plot. Because the early attainment of tappable girth is one of the main selection traits for *H. brasiliensis* breeding (Rao and Kole, 2016), stem diameters were measured at seven different ages between July 2013 and June 2017 (12, 17, 22, 28, 35, 47, and 59 months). The tree heights and numbers of whorls were measured at 17 and 22 months.

The statistical linear mixed model used for the analysis was as follows:

$$y = X_A a + X_R r + X_{AR} ar + Z_B b + Z_G g + \varepsilon$$

where y is the vector with the mean phenotypic trait per plot; and X_A, X_R, and X_{AR} are the incidence matrices for the fixed effects for age (a), replicate (r), and interaction age x replicate (ar). Z_B and Z_G are the incidence matrices for the random effects for block (b) and genotype (g), with $b \sim MVN(0, I_B \otimes I_R \otimes G_{AB})$ and $g \sim MVN(0, I_G \otimes G_{AG})$, where I_n is an identity matrix with order n; $\otimes$ is the Kronecker product; MVN stands for Multivariate Normal Distribution; and G_{AB} and G_{AG} are the variance-covariance matrices for the effects of replicates within blocks within age and within genotype within age, respectively. ε is the residual, where $\varepsilon \sim MVN(0, R_C \otimes R_R \otimes R_A)$, where R_C, R_R, and R_A are the variance-covariance matrices for the column, row, and age for the residual effects, respectively. Different variance-covariance structures for V_{AB}, V_{AG}, R_C, R_R, and R_A were adjusted and compared based on the Akaike information criterion (AIC) (Akaike, 1974) and the Bayesian information criterion (BIC) (Schwarz, 1978) values (Table S2). The variance components were estimated by maximizing the maximum likelihood distribution (REML). Analyses were performed using the packages ASReml-R v3 (Gilmour *et al*., 2009) and ASRemlPlus (Brien, 2016). The heritability (H^2) for each age was estimated as presented by Cullis *et al*. (2006) and Piepho and Möhring (2007), where $H^2 = 1 - \frac{\overline{v}_{BLUP}}{2\sigma_{AG}^2}$ and $\overline{v}_{BLUP}$ indicates the mean variance between the two best linear unbiased predictions (BLUPs). After the phenotypic analysis, we used the differences between estimated BLUPs of subsequent ages for QTL mapping. As an example, for a genotype with a BLUP of 10 cm of diameter at 17 months and with a BLUP of 13 cm of diameter at 22 months, the difference of BLUPs (i.e., 5 cm) was used in the QTL mapping.

QTL mapping

QTL mapping for stem diameter, tree height, and number of whorls traits was performed by applying a CIM model to full-sib families (Gazaffi *et al*., 2014). This method uses a multipoint approach that considers outcrossing segregation patterns and, through contrasts, estimates three genetic effects, one for each parent (hereafter called the additive effect) and one for their interaction (hereafter called the dominance effect). For the CIM model, QTL genotype probabilities were obtained for each 1-cM interval considering a window size of 15 cM. Cofactors were identified for each linkage group using stepwise selection based on the AIC (Akaike, 1974) to select the models. Furthermore, a maximum number of parameters, $2\sqrt{n}$, where n is the number of individuals, was used to avoid super-parametrization (Gazaffi *et al*., 2014). Subsequently, only effects with a 5% significance level were retained in the cofactor model. QTLs were defined according to a threshold based on 5% significance across distributed LOD scores obtained by selecting the second LOD profile peak from 1,000 permutations (Chen and Storey, 2006). Additionally, for each CIM model, the additive genetic effects for each parent, the dominance effects, and the linkage phases, segregation patterns, and

240 proportions of phenotypic variations explained by the QTLs (R^2) were estimated. Analyses were
241 performed using the R package fullsibQTL (Gazaffi *et al.*, 2014).

Results

Filtering and polymorphism analysis

For the linkages analysis 239 SNP markers from genomic, transcriptomic, and cDNA datasets were
retained. The marker segregation for the SSRs and SNPs are shown in Table 1. All genotyped
markers, including 33 markers (6.3%) identified with distortion from the expected Mendelian
segregation ratio, were retained in the linkage analysis.

The GBS library generated a total of 68,777,856 raw reads (90.60% Q20 bases), with an average of
474,330 reads per individual. After alignment to the *H. brasiliensis* reference genome, 81,832 SNPs
were identified. In total, 53,836 markers with more than 25% missing data and/or a non-biallelic
status, 1,587 redundant markers, 17,865 markers with two homozygous parents, and 5,546 markers
with segregation distortion were eliminated. Thus, a total of 2,998 (3.66% of the initial set of
markers) high-quality GBS-based SNPs were retained, most of which were classified in a 1:1 fashion
(2,371 SNPs).

Genetic mapping

A total of 411 markers were mapped in the base map constructed from a set of 526 markers (287
SSRs and 239 SNPs). The markers were distributed over 18 linkage groups (LG) in 2,482.3 cM
(Figure 1), and the length of each group ranged from 78.3 cM (LG4) to 191.9 cM (LG14). Linkage
group 10 showed the highest average density and the highest number of markers mapped (53),
whereas linkage group 15 showed the lowest density (Table 2).

The final genetic map, constructed using advances from the previous base map and the numerous
high-quality SNPs produced with the GBS technique, was based on 408 previously mapped markers
and 671 SNPs identified from GBS (Bioproject PRJEB25899). One SSR and two SNP markers were
removed from the base map to build the final map because there were inconsistencies in their
primers. Therefore, 1,079 mapped markers were distributed over 18 LGs and covered a total of
3,779.7 cM (Figure 2). The length of each group ranged from 109.4 cM and 44 markers (LG7) to 272
cM and 74 markers (LG5), with average marker densities of 2.5 and 3.7 cM, respectively.

The SNPs identified from GBS were uniformly distributed across the genetic map (markers in black
in Figure 2), resulting in improvements in group assessment and shortening of gaps compared with
the base map (Figure 1). Moreover, a reduction in the number of gaps greater than 15 cM to only
eight was observed, which was previously verified on 41 occasions.

Phenotypic analysis and QTL mapping

BLUP was used to perform QTL mapping, indicating the variance-covariance structures and criteria
information for the selected models shown in Table 3. Table 4 shows predicted genotypic values for
the F1 population and each parent, the H^2 for each age, and the variance components for V_{AG} and R_A.

For QTL mapping analyses of the stem diameter, tree height, and number of whorls, several QTLs
were identified by CIM for full-sibling progenies at different ages of growth. Using this method, 24
QTLs were identified for stem diameter, seven were identified for height, and seven were identified

for the number of whorls (Table 5 and Figure 3). The QTLs were located in 12 of the 18 linkage
groups, and most were in LG9 (5) for height and diameter and in LG4 (7) and LG15 (5) for all traits.
A significant number of QTLs for diameter were also observed in LG17 (4). All QTLs were named
with the prefix of the trait evaluated, the date (in months), and the linkage group mapped, for
example, Diameter12-LG2.

From the total QTLs identified in this analysis, five were identified at 12 months, 11 at 17 months
(four for height, three for the number of whorls, and four for diameter), 10 at 22 months (three for
height, four for the number of whorls, and three for diameter), four at 28 months, three at 35 months,
two at 47 months, and three at 59 months. The percentages of phenotypic variation (R^2) explained by
the QTLs ranged from 1.08% (Whorls17-LG14) to 14.07% (Height17-LG4), and the QTLs were
segregated according to the ratios 1:1:1:1, 1:2:1, 3:1, and 1:1. The obtained threshold values for LOD
scores using 1,000 permutations ranged from 3.94 (Diameter at 59 months) to 4.25 (Diameter at 22
months) and were used to declare the presence of QTLs.

QTLs associated with height occurred in seven LGs. QTL Height17-LG4 had the highest LOD score
(6.72), whereas QTL Height17-LG15 had the lowest LOD score (4.23). Additionally, QTL Height17-
LG4 had the highest proportion of phenotypic variation for this trait (14.07%), and QTL Height17-
LG9 had the lowest (6.24%). For QTL Height17-LG9, Height17-LG15, and Height22-LG11,
dominance and additive effects were observed for RRIM701, whereas dominance and additive effects
for GT1 were observed only for QTL Height22-LG6.

For the number of whorls, QTLs were detected in five LGs. The LOD scores for these QTL ranged
from 4.13 (Whorls17-LG14) to 8.59 (Whorls22-LG4), whereas the highest proportion of phenotypic
variation was observed for QTL Whorls17-LG15 (10.22%). Based on the signals of the additive and
dominance estimate effects, different segregation patterns for each of the QTLs were inferred. Up to
three main effects (one additive effect for each parent and one dominance effect) were observed for
QTL Whorls17-LG4, Whorls22-LG8, and Whorls22-LG12.

Although 24 different QTLs were mapped for the diameter trait over seven developmental stages, the
most representative linkage groups mapped were LG4 (12, 17, 35 and 47 months), LG9 (12, 17, 22,
and 59 months), and LG17 (12, 17, 35, and 47 months). The Diameter QTLs accounted for 3.01%
(Diameter35-LG17) to 13.03% (Diameter17-LG4) of the total phenotypic variation, whereas the
LOD scores ranged from 4.15 (Diameter59-LG15) to 10.06 (Diameter35-LG4). The most
representative segregation patterns observed for the QTLs were of the 1:2:1 type (12 QTLs), but all
types were inferred at least once. Among the 53 effects estimated for diameter, only a few were
observed for RRIM701 (21 additive effects for GT1, 13 additive effects for RRIM701, and 19
dominance effects). All of the QTLs identified at 12, 22, and 35 months showed significant
dominance effects.

Discussion

The mapping population was based on a cross between two important commercial clones, GT1 and
RRIM701. The construction of the mapping population was possible because GT1 is male sterile
(Shearman *et al*., 2014), and there was a large experimental area with both parental clones. Using
open-pollinated progenies from a mother tree of interest facilitated the development of progeny and
accelerated linkage map construction and QTL detection, as in sweet cherries (Guajardo *et al*., 2015).
The limited number of progenies obtained reflects the nature of the low fruit-set success of the rubber
tree, varying from no success at all to a maximum of 5-10% in controlled pollination assays
(Priyadarshan and Clément-Demange, 2004).

Based on these findings, the approach employed was based on using an expressive number of marker
loci, with the ultimate goal of having a well-saturated map. During genetic map construction, the
utilization of reliable and informative markers, such as microsatellites, might circumvent the need for
phase estimation. As many markers have already been mapped in the rubber tree (Souza *et al*., 2013),
use of the same markers from existing datasets is preferred to avoid confounding factors, as
suggested by Hodel *et al*. (2016).

Although we observed that the proportion of polymorphic SSR markers between parents of the
mapping population (48%) was consistent with that previously reported for *Hevea brasiliensis*
(Lespinasse *et al*., 2000; Souza *et al*., 2013), we observed a higher proportion of EST-SSR
polymorphisms than Nirapathpongporn *et al*. (2016). The number of observed polymorphic SSR
markers allowed the preferable selection of highly informative markers to construct the base linkage
map, e.g., the 1:1:1:1 fashion (Table 1).

Using highly informative markers together with numerous markers provided by NGS technology is
advantageous. The strategy utilized herein consisted of first building a reliable base map with
informative markers and then increasing its resolution with biallelic markers. To increase the
reliability of markers obtained from NGS technologies, numerous GBS data filtering efforts have
recently been described in several species (Peng *et al*., 2013; Ward *et al*., 2013; Guajardo *et al*.,
2015; Covarrubias-Pazaran *et al*., 2016), with the objective of avoiding false polymorphism
identification because of a low read depth. This objective is particularly important for large genomes
containing many short repeat sequences (Fierst, 2015). To improve the genomic SNP discovery
approach, a solid reference genome sequence is imperative. Thus far, a series of research studies
focusing on *Hevea brasiliensis* reference genomes has been published (Rahman *et al*., 2013; Lau *et
al*., 2016; Tang *et al*., 2016; Pootakham *et al*., 2017), which can be beneficial in linkage mapping
studies.

Compared with other species from Malpighiales, *Hevea* species contain the largest genomes but
might also have the largest repeat content (Tang *et al*., 2016). Seventy percent of the *H. brasiliensis*
genome is composed of repeat elements, which, together with the lack of information at the
chromosomal level, leads to difficulties with assembly (Rahman *et al*., 2013) and challenges in the
identification and development of SNP assays (Souza *et al*., 2016). Moreover, differences in the
estimated rubber tree genome size have been observed by microdensitometry, flow cytometry
(Bennett and Smith, 1997), and 17-mer sequence estimation (Tang *et al*., 2016) approaches.

We verified the effects of inserting non-Mendelian markers in the map before including them in the
base map. Segregation distortion is a common phenomenon in genetic analysis, although it might
affect the estimated recombination fractions between markers (Xu, 2008). Among the GBS-based
SNPs, 5,546 exhibited segregation distortion and were excluded due to the increased complexity of
such a large number of markers. The analysis of such markers with segregation distortion could help
identify whether one or both parents carry lethal or sublethal alleles and could aid in understanding
the genetic architecture of floral development and fertility genes (Covarrubias-Pazaran *et al*., 2016).

Previous genetic map studies of *H. brasiliensis* have revealed the influences of reference genome
sequences. Shortly after the first draft sequence became available for the species (Rahman *et al*.,
2013), Shearman *et al*. (2015) and Pootakham *et al*. (2015) performed SNP genotyping with high-
throughput platforms. Pootakham *et al*. (2015) used a read depth ≥ 6 and fewer than 50% missing
data to obtain 7,345 and 6,678 SNPs in two different populations. Additionally, after using more
restrictive parameters (read depth ≥ 30 and 10% missing data) and filtering uninformative markers,

2,995 and 3,124 SNPs were identified. These quantities are similar to those obtained in the present
work. It is important to mention that the map construction relies on several genetic features (e.g.,
Mendelian segregation) present on segregating populations. Therefore, after filtering, markers were
tested for these features, avoiding the presence of non-reliable markers.

The identification of SNPs in species that underwent polyploidization events, as demonstrated in
Hevea brasiliensis (Pootakham *et al.*, 2017), leads to the utilization of appropriate strategies for the
selection of useful markers for linkage mapping (Clevenger *et al.*, 2015). A reduction of the final
number of obtained SNPs (2,998 SNPs) reflects the restrictive parameters used for marker selection.
The genotyping errors and large amounts of missing data observed in NGS data (Nielsen *et al.*,
2011), particularly in GBS, could have effects on the ordering of loci and the estimation of distances
between two loci (Shields *et al.*, 1991; Hackett and Broadfoot, 2003). Thus, these factors can
critically affect the localization of regions that control quantitative traits (Doerge *et al.*, 1997).

The 2,998 SNPs obtained after filtering enabled higher marker densities than those of microsatellite-
based maps (i.e., Lespinasse *et al.*, 2000; Le Guen *et al.*, 2008; Souza *et al.*, 2013). GBS
methodology provides 1:2:1 and 1:1 markers (biallelic), with 1:1 as the most frequent (Table 1). The
later markers are less informative because they carry information for only one parent meiosis event
(Wu *et al.*, 2002a). In our study, we have the advantages of both marker types, a larger number of
SNPs, and greater information assessment of SSR.

Linkage group organization of the final map was based on previous studies of *H. brasiliensis*
(Lespinasse *et al.*, 2000; Souza *et al.*, 2013) and allowed the order of the markers to be compared
with those in common with other mapping populations. With 228.7 cM and 20 markers mapped,
linkage group 10 was the largest linkage group obtained by Souza *et al.* (2013), and this group shared
a total of 10 markers with those in our work, with the minor positioning difference between markers
HB135 and HB174. A total of nine markers were positioned in the same order by Souza *et al.* (2013)
in linkage group 14 at a greater density. In addition, previous genetic maps have demonstrated 18
LGs (Lespinasse *et al.*, 2000; Le Guen *et al.*, 2008; Pootakham *et al.*, 2015; Shearman *et al.*, 2015;
Nirapathpongporn *et al.*, 2016), corresponding to the haploid number for the species (2n = 36). The
LGs obtained with 225 SSRs and 186 SNPs support the observations by Souza *et al.* (2013),
demonstrating the necessity of mapping using different types of markers to account for additional
LGs from incomplete coverage of the *H. brasiliensis* genome.

Although our base genetic map presented a higher mean marker density than that presented by
Triwitayakorn *et al.* (2011) and Souza *et al.* (2013) when mapping 97 and 284 SSRs, respectively,
marker intervals greater than 15 cM were observed on 41 occasions. As noted by Souza *et al.* (2013),
such results can be explained by regions with low recombination frequency, with higher degrees of
homozygosity, and with a low number of polymorphic markers, reinforcing the indispensable usage
of other methodologies that provide a greater number of markers to increase resolution.

The GBS technique has promoted the discovery of thousands of polymorphic markers that are useful
for the construction of high-density linkage maps. Because increasing marker densities in low-
resolution regions can be beneficial for traditional mapping approaches and QTL mapping, GBS has
become a valid and interesting alternative. Although GBS technology has been available for *H.*
brasiliensis for several years (Pootakham *et al.*, 2015; Shearman *et al.*, 2015), no studies on the
beneficial effects of SSR markers, such as those performed on *Brassica* species (Yang *et al.*, 2016),
have been performed for *Hevea*.

Our base genetic map including 225 SSRs markers was essential for avoiding problems during
linkage group formation, ordering, and marker distance estimation. Inserting numerous GBS-based
SNP markers reduced the average marker interval in all LGs, with a mean of 3.5 cM (Table 2).
Regions with a low marker density in the base map (i.e., LG13 and LG15), with gaps up to 24.2 cM
and 32.5 cM, were saturated. As in other linkage mapping studies using large-scale genotyping
methodologies for *H. brasiliensis* (Pootakham *et al.*, 2015; Shearman *et al.*, 2015), substantial
intervals (≥ 10 cM) were observed, supporting the results obtained in different plant species
(Guajardo *et al.*, 2015; Marubodee *et al.*, 2015; Boutet *et al.*, 2016; McCallum *et al.*, 2016). The
remaining gaps are common and even expected using GBS, mainly due to centromeric regions,
which are not reached using this approach.

Our final map has a total of 1,079 markers and spans 3,779.7 cM, which reflects the molecular source
and chosen mapping methodology. Comparatively, Souza *et al.* (2013) and Lespinasse *et al.* (2000)
presented smaller maps, with 2688.8 cM and 2144 cM of total size, respectively. This size difference
can be explained by several reasons: i) the smaller number of markers (284 and 717, respectively),
cover a smaller genome region; ii) those studies used only non-NGS markers, which are likely to
have fewer genotyping errors; and iii) as well as in Pootakham *et al.* (2015), the map from
Lespinasse *et al.* (2000) uses strategies to reduce map size, removing improbable genotypes, such as
those originating from double recombinations (Hackett and Broadfoot, 2003). However, the later
strategy can also remove real recombinant events (false positives), particularly if the marker order is
inaccurate (Wu *et al.*, 2008). In our case, the consequence is to not account for those recombinant
events in the further QTL mapping studies.

The constructed linkage map allowed the anchoring of 601 scaffolds (Table S3), accounting for
approximately 8.1% of the current number of scaffolds available in the reference genome spanning
653.3 Mb (47.5% of the genome sequence). Therefore, although SNPs were not mapped in most
parts of the scaffolds, these data constitute a representative part of the genome. Sequence contiguity
of the rubber tree genome can be achieved as scaffold reviews are developed. Thus, the linkage map
presented herein and the anchored scaffolds represent a foundation for future efforts.

The identification of genes underlying growth-related traits is critical for perennial crops grown in
marginal areas (i.e., eucalyptus and rubber tree). Tree growth is determined by cell division and
expansion in apical and cambial meristems, developmental and seasonal transitions, photosynthesis
efficiency, nutrient and water uptake and transport, and responses to biotic and abiotic stresses
(Grattapaglia *et al.*, 2009). Therefore, the usage of molecular markers is not easy because such traits
are complex and governed simultaneously by many QTLs. The various numbers of growth-related
QTLs identified in several LGs under suboptimal temperature and humidity conditions herein and in
the study conducted by Souza *et al.* (2013) confirmed the numerous biological processes involved in
H. brasiliensis tree growth. Elucidating of all the factors involved is imperative considering that
rubber breeding has stagnated (Tang *et al.*, 2016) and has expanded to marginal areas (Priyadarshan,
2017). The results obtained herein provide new information regarding QTL control of growth-related
traits during almost five consecutive years in different seasons. Most importantly, characterizing
QTLs in dry seasons can be advantageous because this strategy appears to be the most important
characteristic for the identification of escape areas (Jaimes *et al.*, 2016) and limits growth and latex
production.

The climatological water balance revealed that sampling was performed in consecutive water deficit
periods, although exceptions included very brief intervals with very high precipitation levels in
March 2014 and January 2016 (Figure 4). In a suboptimal climate area with a lengthy dry season,

Chanroj *et al.* (2017) found a single SNP associated with girth accounting for 14% of the phenotypic
variance in *H. brasiliensis*. Although this was not observed herein, the QTLs identified in our work
can also be used as an initial source for marker-assisted selection because seven QTLs showed major
effects (phenotypic variance greater than 10%) (Table 5). These QTLs are supported by previous
growth-related QTLs observed for the species under seasonal water stress (Souza *et al.*, 2013).
Because loss of water vapor through leaf transpiration occurs while capturing atmospheric CO_2 for
photosynthesis, the QTLs identified in our study provide powerful information regarding the
mechanisms of water use efficiency in *H. brasiliensis*. Indeed, the search for favorable alleles for
growth must continue as in the study conducted by Coupel-Ledru *et al.* (2016), which identified four
QTLs related to reduced nighttime transpiration in *Vitis vinifera*.

Herein, the water deficit intensity was highest in 2014, and we may argue that such a situation might
help identify consistent QTLs. It is worth noting that the QTLs in linkage group 5 in May (22
months) and November 2014 (28 months) were consistent with those previously mapped by Souza *et*
al. (2013). Flanking QTL Height22-LG5, it is possible to observe a new marker developed (SNP
Hb_seq_234) that has been annotated in the hydroxymethylglutaryl-CoA reductase (NADPH)
pathway and mapped on scaffold0419.

Because all of the QTLs identified for stem diameter at 12, 22, and 35 months showed significant
dominance effects (Table 5), their contribution appeared to play a role in phenotypic variation.
Different effects, positions, and segregation patterns of QTLs detected over time must be considered
with caution, particularly for *Hevea brasiliensis*, which has a long breeding cycle. Variations could
be possible because different genes are involved in genotype-environment interactions or QTL
regulation according to the developmental stage (Conner *et al.*, 1998). In linkage group 4, QTLs with
large effects were very important because the same genomic region appeared to control tree height,
the number of whorls, and stem diameter.

Additionally, important QTLs in linkage group 6 have already been mapped by Souza *et al.* (2013)
for both girth and tree height in a different mapping population cultivated under suboptimal growth
conditions. Annotation of the sequences (Table S4) showed similarity with the *peptidyl-prolyl cis-*
trans isomerase FKBP15-1-like gene, another good candidate gene worthy of further investigation.
The activity of this isomerase has been shown to accelerate the process of protein folding both *in*
vitro and *in vivo* (Marivet *et al.*, 1994). Regulated by light in chloroplasts (Luan *et al.*, 1994), it may
also be regulated by stress and play an important role throughout the plant life cycle (Marivet *et al.*,
1994). Therefore, further research of genes associated with abiotic stress and latex production in the
rubber tree is required, as evidenced by the large number of plantations in areas with suboptimal
climates.

The recent opportunity to associate high-density genetic maps with reference genomes in rubber tree
is a key challenge to increase rubber production in marginal areas. In fact, the results of the genetic
architecture of traits of interest should improve as new studies emerge. The genetic map constructed
herein, which included several approaches to handle thousands of markers, allowed the identification
of genomic regions related to growth under water stress so that breeders can explore these resources
to apply marker-assisted selection. Additionally, rubber tree breeding programs can benefit from the
resources generated by applying genomic selection strategies.

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

Conceived and designed the experiments: AG, AS, PG, VG. Conducted the experiments and collected data: AC, CS, CM, IA, LHS, LMS. Analyzed the data: AC, AG, CT, JR, RA, RR. Collected phenotypic data: EJ. Wrote the manuscript: AC, AS, CT, RA. All of the authors read and approved the manuscript.

Funding

The authors would like to acknowledge the Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) for the financial support provided (2007/50392-1, 2012/50491-8) and PhD and Post-doctoral scholarships to LMS (2012/05473-1), CM (2011/50188-0, 2014/18755-0), CS (2009/52975-0, 2015/24346-9) and LHS (2014/11807-5, 2017/07908-9). The authors are also thankful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for PhD fellowships awarded to AC, CT, and RR; the MS fellowship awarded to RA; research fellowships awarded to AG, AS, and PG; and a post-doctoral fellowship awarded to CS. We also acknowledge the Coordenação do Aperfeiçoamento de Pessoal de Nível Superior (CAPES Computational Biology Program) for grants and post-doctoral fellowships awarded to AC, LMS, and JR and a Master fellowship awarded to IA. The authors are thankful to CAPES – Agropolis Program for the fellowship to AC to attend a four-month training internship at CIRAD, UMR AGAP, on the Rubber Tree Breeding Program, Montpellier, France. The authors would like to acknowledge the Agropolis – UNICAMP Fellowship Program for the fellowship awarded to Aline da Costa Lima Moraes, the Molecular Biology Technician at Molecular and Genetic Analysis Laboratory, CBMEG / UNICAMP, to attend a one-month training internship at CIRAD, UMR AGAP on the Rubber Tree Breeding Program, Montpellier, France.

Acknowledgements

We acknowledge Aline da Costa Lima Moraes for support with the experiments and Kaio Olímpio das Graças Dias and Jhonathan Pedroso Rigal dos Santos for helping with the phenotypic analyses.

References

Akaike, H. (1974). A new look at the statistical model identification. *IEEE Trans. Autom. Control* 19, 716–723. doi:10.1109/TAC.1974.1100705.

Alvares, C. A., Stape, J. L., Sentelhas, P. C., De Moraes Gonçalves, J. L., and Sparovek, G. (2013). Köppen's climate classification map for Brazil. *Meteorol. Zeitschrift* 22, 711–728. doi:10.1127/0941-2948/2013/0507.

Bennett, M. D., and Smith, J. B. (1997). Nuclear DNA Amounts in Angiosperms. *Philos. Trans. R. Soc. B Biol. Sci.* 274, 227–274. doi:10.1098/rstb.1976.0044.

Boutet, G., Alves Carvalho, S., Falque, M., Peterlongo, P., Lhuillier, E., Bouchez, O., et al. (2016). SNP discovery and genetic mapping using genotyping by sequencing of whole genome genomic DNA from a pea RIL population. *BMC Genomics* 17, 121. doi:10.1186/s12864-016-2447-2.

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., and Buckler, E. S. (2007). TASSEL: Software for association mapping of complex traits in diverse samples. *Bioinformatics* 23, 2633–2635. doi:10.1093/bioinformatics/btm308.

Brien, C. (2016). ASRemlPlus: Augments the Use of "ASReml" in Fitting Mixed Models. R package
version 2.0-9.

Chanroj, V., Rattanawong, R., Phumichai, T., Tangphatsornruang, S., and Ukoskit, K. (2017).
Genome-wide association mapping of latex yield and girth in Amazonian accessions of Hevea
brasiliensis grown in a suboptimal climate zone. *Genomics*, 0–1.
doi:10.1016/j.ygeno.2017.07.005.

Chen, L., and Storey, J. D. (2006). Relaxed significance criteria for linkage analysis. *Genetics* 173,
2371–2381. doi:10.1534/genetics.105.052506.

Clevenger, J., Chavarro, C., Pearl, S. a., Ozias-Akins, P., and Jackson, S. a. (2015). Single nucleotide
polymorphism identification in polyploids: A review, example, and recommendations. *Mol.*
Plant 8, 831–846. doi:10.1016/j.molp.2015.02.002.

Conner, P. J., Brown, S. K., and Weeden, N. F. (1998). Molecular-marker analysis of quantitative
traits for growth and development in juvenile apple trees. *Theor. Appl. Genet.* 96, 1027–1035.
doi:10.1007/s001220050835.

Coupel-Ledru, A., Lebon, E., Christophe, A., Gallo, A., Gago, P., Pantin, F., et al. (2016). Reduced
nighttime transpiration is a relevant breeding target for high water-use efficiency in grapevine.
Proc. Natl. Acad. Sci. 113, 201600826. doi:10.1073/pnas.1600826113.

Covarrubias-Pazaran, G., Diaz-Garcia, L., Schlautman, B., Deutsch, J., Salazar, W., Hernandez-
Ochoa, M., et al. (2016). Exploiting genotyping by sequencing to characterize the genomic
structure of the American cranberry through high-density linkage mapping. *BMC Genomics* 17,
451. doi:10.1186/s12864-016-2802-3.

Creste, S., Neto, A. T., and Figueira, A. (2001). Detection of Single Sequence Repeat Polymorphisms
in Denaturing Polyacrylamide Sequencing Gels by Silver Staining. *Plant Mol. Biol. Report.* 19,
299–306.

Cullis, B. R., Smith, A. B., and Coombes, N. E. (2006). On the design of early generation variety
trials with correlated data. *J. Agric. Biol. Environ. Stat.* 11, 381–393.
doi:10.1198/108571106X154443.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., et al. (2011). The
variant call format and VCFtools. *Bioinformatics* 27, 2156–2158.
doi:10.1093/bioinformatics/btr330.

Doerge, R. W., Zeng, Z., and Weir, B. S. (1997). Statistical issues in the search for genes affecting
quantitative traits in experimental populations. *Stat. Sci.* 12, 195–219.
doi:10.1214/ss/1030037909.

Doyle, J. J., and Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh
leaf tissue. *Phytochem. Bull.* 19, 11–15. doi:10.2307/4119796.

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. a, Kawamoto, K., Buckler, E. S., et al. (2011). A
robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS One*
6, e19379. doi:10.1371/journal.pone.0019379.

Federer, A. W. T., and Raghavarao, D. (1975). On Augmented Designs. *Biometrics* 31, 29–35.

Feng, S. P., Li, W. G., Huang, H. S., Wang, J. Y., and Wu, Y. T. (2009). Development,
characterization and cross-species/genera transferability of EST-SSR markers for rubber tree
(Hevea brasiliensis). *Mol. Breed.* 23, 85–97. doi:10.1007/s11032-008-9216-0.

Fierst, J. L. (2015). Using linkage maps to correct and scaffold de novo genome assemblies : methods
, challenges , and computational tools. *Front. Genet.* 6, 1–8. doi:10.3389/fgene.2015.00220.

Garcia, D., Carels, N., Koop, D. M., de Sousa, L. A., Andrade Junior, S. J. de, Pujade-Renaud, V., et
al. (2011). EST profiling of resistant and susceptible Hevea infected by Microcyclus ulei.
Physiol. Mol. Plant Pathol. 76, 126–136. doi:10.1016/j.pmpp.2011.07.006.

Gazaffi, R., Margarido, G. R. a., Pastina, M. M., Mollinari, M., and Garcia, A. A. F. (2014). A model
for quantitative trait loci mapping, linkage phase, and segregation pattern estimation for a full-
sib progeny. *Tree Genet. Genomes*, 1–11. doi:10.1007/s11295-013-0664-2.

Gilmour, A. R., Gogel, B. J., Cullis, B. R., and Thompson, R. (2009). ASReml user guide release 3.0.
VSN Int. Ltd, 275. doi:10.1017/CBO9781107415324.004.

Gonçalves, P. D. S., Scaloppi Júnior, E. J., Martins, M. A., Moreno, R. M. B., Branco, R. B. F., and
Gonçalves, E. C. P. (2011). Assessment of growth and yield performance of rubber tree clones
of the IAC 500 series. *Pesqui. Agropecuária Bras.* 46, 1643–1649.

Grattapaglia, D., Plomion, C., Kirst, M., and Sederoff, R. R. (2009). Genomics of growth traits in
forest trees. *Curr. Opin. Plant Biol.* 12, 148–156. doi:10.1016/j.pbi.2008.12.008.

Grattapaglia, D., and Sederoff, R. (1994). Genetic Linkage Maps of Eucalyptus grandis and
Eucalyptus urophylla using a Pseudo-Testcross: Mapping Strategy and RAPD markers. *Genetics*
137, 1121–1137.

Guajardo, V., Solís, S., Sagredo, B., Gainza, F., Muñoz, C., Gasic, K., et al. (2015). Construction of
high density sweet cherry (Prunus avium L.) linkage maps using microsatellite markers and
SNPs detected by genotyping-by-sequencing (GBS). *PLoS One* 10, 1–17.
doi:10.1371/journal.pone.0127750.

Hackett, C. a, and Broadfoot, L. B. (2003). Effects of genotyping errors, missing values and
segregation distortion in molecular marker data on the construction of linkage maps. *Heredity*
(Edinb). 90, 33–38. doi:10.1038/sj.hdy.6800173.

Hodel, R. G. J., Segovia-Salcedo, M. C., Landis, J. B., Crowl, A. A., Sun, M., Liu, X., et al. (2016).
The report of my death was an exaggeration: a review for researches using microsatellite in the
21th century. *Appl. Plant Sci.* 4, 1–13. doi:10.3732/apps.1600025.

Jaimes, Y., Rojas, J., Cilas, C., and Furtado, E. L. (2016). Suitable climate for rubber trees affected
by the South American Leaf Blight (SALB): Example for identi fi cation of escape zones in the
Colombian middle Magdalena. *Crop Prot.* 81, 99–114. doi:10.1016/j.cropro.2015.12.016.

Kosambi, D. D. (1943). The estimation of map distances from recombination values. *Ann. Eugen.* 12,
172–175. doi:10.1111/j.1469-1809.1943.tb02321.x.

Lander, E. S., and Green, P. (1987). Construction of multilocus genetic linkage maps in humans.
Proc. Natl. Acad. Sci. USA 84, 2363–7. doi:10.1073/pnas.84.8.2363.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357–359. doi:10.1038/nmeth.1923.

Lau, N., Makita, Y., Kawashima, M., Taylor, T. D., Kondo, S., Othman, A. S., et al. (2016). The
rubber tree genome shows expansion of gene family associated with rubber biosynthesis. *Sci.*
Rep. 6, 1–14. doi:10.1038/srep28594.

Le Guen, V., Gay, C., Xiong, T. C., Souza, L. M., Rodier-Goud, M., and Seguin, M. (2011).
Development and characterization of 296 new polymorphic microsatellite markers for rubber
tree (Hevea brasiliensis). *Plant Breed.* 130, 294–296. doi:10.1111/j.1439-0523.2010.01774.x.

Le Guen, V., Guyot, J., Raimundo, C. R. R., Seguin, M., and Garcia, D. (2008). Long lasting rubber
tree resistance to Microcyclus ulei characterized by reduced conidial emission and absence of
teleomorph. *Crop Prot.* 27, 1498–1503. doi:10.1016/j.cropro.2008.07.012.

Lespinasse, D., Rodier-Goud, M., Grivet, L., Leconte, A., Legnate, H., and Seguin, M. (2000). A
saturated genetic linkage map of rubber tree (Hevea spp.) based on RFLP, AFLP, microsatellite,
and isozyme markers. *Theor. Appl. Genet.* 100, 127–138. doi:10.1007/s001220050018.

Luan, S., Albers, M. W., and Schreiber, S. L. (1994). Light-regulated, tissue-specific immunophilins
in a higher plant. *Proc. Natl. Acad. Sci. U. S. A.* 91, 984–8. doi:10.1073/pnas.91.3.984.

Maliepaard, C., Jansen, J., and Van Ooijen, J. W. (1997). Linkage analysis in a full-sib family of an
outbreeding plant species: overview and consequences for applications. *Genet. Res.* 70, 237–
250. doi:10.1017/S0016672397003005.

Mantello, C. C., Cardoso-Silva, C. B., Da Silva, C. C., De Souza, L. M., Scaloppi, E. J., de
Gonçalves, P. S., et al. (2014). De Novo assembly and transcriptome analysis of the rubber tree
(Hevea brasiliensis) and SNP markers development for rubber biosynthesis pathways. *PLoS One*
9, 1–14. doi:10.1371/journal.pone.0102665.

Mantello, C. C., Suzuki, F. I., Souza, L. M., Gonçalves, P. S., and Souza, A. P. (2012). Microsatellite
marker development for the rubber tree (Hevea brasiliensis): characterization and cross-
amplification in wild Hevea species. *BMC Res. Notes* 5, 329. doi:10.1186/1756-0500-5-329.

Margarido, G. R. A., Souza, A. P., and Garcia, A. A. F. (2007). OneMap: Software for genetic
mapping in outcrossing species. *Hereditas* 144, 78–79. doi:10.1111/j.2007.0018-0661.02000.x.

Marivet, J., Margis-Pinheiro, M., Frendo, P., and Burkard, G. (1994). Bean cyclophilin gene
expression during plant development and stress conditions. *Plant Mol. Biol.* 26, 1181–1189.

Marubodee, R., Ogiso-Tanaka, E., Isemura, T., Chankaew, S., Kaga, A., Naito, K., et al. (2015).
Construction of an SSR and RAD-marker based molecular linkage map of Vigna vexillata (L.)
A. Rich. *PLoS One* 10, 1–16. doi:10.1371/journal.pone.0138942.

McCallum, S., Graham, J., Jorgensen, L., Rowland, L. J., Bassil, N. V., Hancock, J. F., et al. (2016).
Construction of a SNP and SSR linkage map in autotetraploid blueberry using genotyping by
sequencing. *Mol. Breed.* 36, 1–24. doi:10.1007/s11032-016-0443-5.

Mollinari, M., Margarido, G. R. A., Vencovsky, R., and Garcia, A. A. F. (2009). Evaluation of
algorithms used to order markers on genetic maps. *Heredity (Edinb).* 103, 494–502.
doi:10.1038/hdy.2009.96.

Nielsen, R., Paul, J. S., Albrechtsen, A., and Song, Y. S. (2011). Genotype and SNP calling from
next-generation sequencing data. *Nat. Rev. Genet.* 12, 443–451. doi:10.1038/nrg2986.

Nirapathpongporn, K., Kongsawadworakul, P., Viboonjun, U., Teerawattanasuk, K., Chrestin, H.,
Segiun, M., et al. (2016). Development and mapping of functional expressed sequence tag-
derived simple sequence repeat markers in a rubber tree RRIM600 × PB217 population. *Mol.*
Breed. 36, 39. doi:10.1007/s11032-016-0461-3.

Patel, R. K., and Jain, M. (2012). NGS QC toolkit: A toolkit for quality control of next generation
sequencing data. *PLoS One* 7. doi:10.1371/journal.pone.0030619.

Peng, Z., Lu, Y., Li, L., Zhao, Q., Feng, Q., Gao, Z., et al. (2013). The draft genome of the fast-
growing non-timber forest species moso bamboo (Phyllostachys heterocycla). *Nat. Genet.* 45,
456–61, 461e1–2. doi:10.1038/ng.2569.

Piepho, H. P., and Möhring, J. (2007). Computing heritability and selection response from
unbalanced plant breeding trials. *Genetics* 177, 1881–1888. doi:10.1534/genetics.107.074229.

Pootakham, W., Ruang-Areerate, P., Jomchai, N., Sonthirod, C., Sangsrakru, D., Yoocha, T., et al.
(2015). Construction of a high-density integrated genetic linkage map of rubber tree (Hevea
brasiliensis) using genotyping-by-sequencing (GBS). *Front. Plant Sci.* 6.
doi:10.3389/fpls.2015.00367.

Pootakham, W., Sonthirod, C., Naktang, C., Ruang-Areerate, P., Yoocha, T., Sangsrakru, D., et al.
(2017). De novo hybrid assembly of the rubber tree genome reveals evidence of paleotetraploidy
in Hevea species. *Sci. Rep.* 7, 41457. doi:10.1038/srep41457.

Priyadarshan, P. M. (2017). Refinements to Hevea rubber breeding. *Tree Genet. Genomes* 13, 20.
doi:10.1007/s11295-017-1101-8.

Priyadarshan, P. M., and Clément-Demange, A. (2004). Breeding Hevea Rubber: Formal and
Molecular Genetics. *Adv. Genet.* 52, 51–115. doi:10.1016/S0065-2660(04)52003-5.

Rahman, A. Y. A., Usharraj, A. O., Misra, B. B., Thottathil, G. P., Jayasekaran, K., Feng, Y., et al.
(2013). Draft genome sequence of the rubber tree Hevea brasiliensis. *BMC Genomics* 14, 75.
doi:10.1186/1471-2164-14-75.

Rao, G. P., and Kole, P. C. (2016). Evaluation of Brazilian wild Hevea germplasm for cold tolerance:
genetic variability in the early mature growth. *J. For. Res.* 27, 755–765. doi:10.1007/s11676-
015-0188-8.

Romain, B., and Thierry, C. (2011). RUBBERCLONES (Hevea clonal descriptions).
http://rubberclones.cirad.fr.

Saha, T., and Priyadarshan, P. M. (2012). "Genomics of Hevea Rubber," in *Genomics of Tree Crops*,
eds. R. J. Schnell and P. M. Priyadarshan (New York, NY: Springer New York), 261–298.
doi:10.1007/978-1-4614-0920-5.

Salgado, L. R., Koop, D. M., Pinheiro, D. G., Rivallan, R., Le Guen, V., Nicolás, M. F., et al. (2014).
De novo transcriptome analysis of Hevea brasiliensis tissues by RNA-seq and screening for
molecular markers. *BMC Genomics* 15, 236. doi:10.1186/1471-2164-15-236.

Schwarz, G. (1978). Estimating the dimension of a model. *Ann. Stat.* 6, 461–464.
doi:10.1214/aos/1176344136.

Shearman, J. R., Sangsrakru, D., Jomchai, N., Ruangareerate, P., Sonthirod, C., Naktang, C., et al.
(2015). SNP identification from RNA sequencing and linkage map construction of rubber tree
for anchoring the draft genome. *PLoS One* 10, 1–12. doi:10.1371/journal.pone.0121961.

Shearman, J. R., Sangsrakru, D., Ruang-Areerate, P., Sonthirod, C., Uthaipaisanwong, P., Yoocha,
T., et al. (2014). Assembly and analysis of a male sterile rubber tree mitochondrial genome
reveals DNA rearrangement events and a novel transcript. *BMC Plant Biol.* 14, 45.
doi:10.1186/1471-2229-14-45.

Shields, D. C., Collins, a, Buetow, K. H., and Morton, N. E. (1991). Error filtration, interference, and
the human linkage map. *Proc. Natl. Acad. Sci. U. S. A.* 88, 6501–6505.
doi:10.1073/pnas.88.15.6501.

Silpi, U., Thaler, P., Kasemsap, P., Lacointe, A., Chantuma, A., Adam, B., et al. (2006). Effect of
tapping activity on the dynamics of radial growth of Hevea brasiliensis trees. *Tree Physiol.* 26,
1579–1587.

Silva, C. C., Mantello, C. C., Campos, T., Souza, L. M., Gonçalves, P. S., and Souza, A. P. (2014).
Leaf-, panel- and latex-expressed sequenced tags from the rubber tree (Hevea brasiliensis) under
cold-stressed and suboptimal growing conditions: the development of gene-targeted functional
markers for stress response. *Mol. Breed.* 34, 1035–1053. doi:10.1007/s11032-014-0095-2.

Souza, L. M., Gazaffi, R., Mantello, C. C., Silva, C. C., Garcia, D., Le Guen, V., et al. (2013). QTL
Mapping of Growth-Related Traits in a Full-Sib Family of Rubber Tree (Hevea brasiliensis)
Evaluated in a Sub-Tropical Climate. *PLoS One* 8, e61238. doi:10.1371/journal.pone.0061238.

Souza, L. M., Mantello, C. C., Santos, M. O., Gonçalves, P. de S., and Souza, A. P. (2009).
Microsatellites from rubber tree (Hevea brasiliensis) for genetic diversity analysis and cross-
amplification in six Hevea wild species. *Conserv. Genet. Resour.* 1, 75–79. doi:10.1007/s12686-
009-9018-7.

Souza, L. M., Toledo-Silva, G., Cardoso-Silva, C. B., Silva, C. C., Andreotti, I. A. A., Conson, A. R.
O., et al. (2016). Development of single nucleotide polymorphism markers in the large and
complex rubber tree genome using next-generation sequence data. *Mol. Breed.* 36, 1–10.
doi:10.1007/s11032-016-0534-3.

Tang, C., Yang, M., Fang, Y., Luo, Y., Gao, S., Xiao, X., et al. (2016). The rubber tree genome
reveals new insights into rubber production and species adaptation. *Nat. Plants* 2, 16073.
doi:10.1038/nplants.2016.73.

Thornthwaite, C. W., and Mather, J. R. (1955). The water balance. *Publ. Climatol.* 8, 1–104.

Triwitayakorn, K., Chatkulkawin, P., Kanjanawattanawong, S., Sraphet, S., Yoocha, T., Sangsrakru,
D., et al. (2011). Transcriptome sequencing of Hevea brasiliensis for development of
microsatellite markers and construction of a genetic linkage map. *DNA Res.* 18, 471–82.
doi:10.1093/dnares/dsr034.

Ward, J. a, Bhangoo, J., Fernández-Fernández, F., Moore, P., Swanson, J. D., Viola, R., et al. (2013).
Saturated linkage map construction in Rubus idaeus using genotyping by sequencing and
genome-independent imputation. *BMC Genomics* 14, 2. doi:10.1186/1471-2164-14-2.

Warren-Thomas, E., Dolman, P. M., and Edwards, D. P. (2015). Increasing Demand for Natural
Rubber Necessitates a Robust Sustainability Initiative to Mitigate Impacts on Tropical
Biodiversity. *Conserv. Lett.* 8, 230–241. doi:10.1111/conl.12170.

Wu, R., Ma, C., Painter, I., and Zeng, Z.-B. (2002a). Simultaneous maximum likelihood estimation
of linkage and linkage phases in outcrossing species. *Theor. Popul. Biol.* 61, 349–63.
doi:10.1006/tpbi.2002.1577.

Wu, R., Ma, C.-X., Wu, S. S., and Zeng, Z.-B. (2002b). Linkage mapping of sex-specific differences.
Genet. Res. 79, 85–96. Available at: http://www.ncbi.nlm.nih.gov/pubmed/11974606.

Wu, Y., Bhat, P. R., Close, T. J., and Lonardi, S. (2008). Efficient and accurate construction of
genetic linkage maps from the minimum spanning tree of a graph. *PLoS Genet* 4, e1000212.
doi:10.1371/journal.pgen.1000212.

Xu, S. (2008). Quantitative trait locus mapping can benefit from segregation distortion. *Genetics* 180,
2201–2208. doi:10.1534/genetics.108.090688.

Yang, S., Chen, S., Geng, X. X., Yan, G., Li, Z. Y., Meng, J. L., et al. (2016). The first genetic map
of a synthesized allohexaploid Brassica with A , B and C genomes based on simple sequence
repeat markers. *Theor. Appl. Genet.* 129, 689–701. doi:10.1007/s00122-015-2657-z.

Figure 1. LGs (1 to 18) for the base genetic map, including 411 markers (225 SSRs and 186 SNPs).

Figure 2. LGs (1 to 18) for the final genetic map, including 1,079 markers for progenies of the cross between GT1 and RRIM701. Black: GBS-based SNP markers; red: markers from the base map.

Figure 3. LOD profile of the QTL CIM for stem diameter, number of whorls, and tree height at multiple times for the progeny of the cross between GT1 and RRIM701. The LOD score and traits are shown on the horizontal axis. The linkage group and map position in cM are shown on the vertical axis. The colors indicate different times. Magnification allows one to visualize the genetic map and marker IDs. The markers in red also belong to the base map. LOD scores above the threshold are indicated by letters, which are described in Table 5.

Figure 4. Temperature, precipitation and climatological water balance between July 2012 and August 2017 in Votuporanga, SP, Brazil. (A) Average maximum temperature, average temperature, average minimum temperature and monthly precipitation. (B) Evaluation of water excess and deficit periods. Black arrows indicate phenotypic measurements.

768 **Table 1.** Overall segregation of SSR and SNP markers mapped in the final map for progenies of the
769 GT1 and RRIM701 cross.

Cross segregation	Genomic SSR	EST-SSR	SNPs Genomics	SNPs Transcripts	SNPs GBS-based	Total
1:1:1:1	89	39	-	-	-	128
1:2:1	21	22	4	48	72	167
1:1	23	30	7	125	599	784
Total	133	91	11	173	671	1079

771 **Table 2.** Linkage group information regarding the base and final maps from the cross between GT1
772 and RRIM701. LG – linkage group; length - mean length in cM; density - mean density in cM; gap -
773 length of larger gap in cM.

LG	Base Map SSRs	SNPs	Length (cM)	Density (cM)	Gap (cM)	Final Map SSRs	SNPs	Length (cM)	Density (cM)	Gap (cM)
1	14	17	153.1	4.9	18.4	13	53	232.4	3.5	13.0
2	19	8	140.2	5.2	14.4	19	33	199.3	3.9	9.1
3	14	6	115.0	5.8	20.7	14	43	184.9	3.2	14.0
4	6	5	78.3	7.1	17.9	6	39	163.8	3.6	12.9
5	9	13	176.9	8.0	24.9	9	65	272.0	3.7	16.6
6	17	14	162.9	5.3	20.1	17	63	228.1	2.9	11.6
7	12	5	80.3	4.7	19.4	12	32	109.4	2.5	14.1
8	14	8	141.3	6.4	18.4	14	45	229.7	3.9	18.5
9	13	6	124.6	6.6	20.8	13	33	196.3	4.3	14.3
10	26	27	185.4	3.5	16.2	26	52	228.1	2.9	11.5
11	6	15	164.8	7.8	27.5	6	58	235.4	3.7	20.6
12	7	10	104.1	6.1	23.3	7	32	167.5	4.3	12.8
13	6	10	136.7	8.5	24.2	6	58	218.7	3.4	15.0
14	22	13	191.9	5.5	15.6	22	51	258.8	3.5	17.0
15	10	6	160.0	10.0	32.5	10	39	225.3	4.6	16.2
16	10	6	100.8	6.3	17.3	10	53	166.0	2.6	12.9
17	6	7	107.5	8.3	19.6	6	54	206.7	3.6	9.8
18	14	10	158.5	6.6	26.0	14	52	257.3	3.9	17.1
Total	225	186	2482.3	6.0	32.5	224	855	3779.7	3.5	20.6

Table 3. Variance-covariance structures and criteria information for the adjusted models. $G_{(AB,AG)}$: Variance-covariance matrices for effects of replicates within block within age and genotype within age; $R_{(C,R,A)}$: variance-covariance matrices for the column, row, and age for residual effects; df: degrees of freedom; Corh: matrix of heterogeneous variance with correlation; US: non-structured variance-covariance matrix; Ar1h: first-order autoregressive correlation matrix with heterogeneous variance; Ar1: first-order autoregressive correlation matrix without variance.

Trait	G_{AB}	G_{AG}	R_C	R_R	R_A	df	AIC	BIC	logREML
Tree height	US	US	Ar1	Ar1	US	11	-767.01	-707.69	394.50
Stem diameter	Corh	Corh	Ar1	Ar1	Ar1h	26	22235.43	22407.92	-11091.72
Number of whorls	US	US	Ar1	Ar1	US	11	1772.50	1831.81	-875.25

Table 4. Summary of the predicted genotypic values and estimations of additive genetic variance and of phenotypic ($\hat{\sigma}^2_{AG}$ and $\hat{\sigma}^2_{RA}$) and heritability ($\hat{H}^2$) components for tree height (TH), stem diameter (SD), and number of whorls (NW) of rubber trees.

Trait/Age	Predicted Genotypic Values					$\hat{\sigma}^2_{AG}$	$\hat{\sigma}^2_{RA}$	$\hat{H}^2$
	F1 (Minimum)	F1 (Maximum)	F1 (Mean)	GT1	RRIM701			
TH/17 months	1.52	2.63	2.19	2.28	3.33	0.07	0.23	0.54
TH/22 months	2.72	3.85	3.33	2.06	2.98	0.11	0.41	0.51
SD/12 months	14.91	18.17	16.87	16.05	15.64	0.64	11.35	0.55
SD/17 months	18.72	23.71	21.76	20.41	19.84	1.51	18.14	0.55
SD/22 months	24.95	32.81	29.64	27.74	26.69	3.63	25.23	0.57
SD/28 months	31.31	38.66	36.02	33.96	32.85	3.27	39.00	0.54
SD/35 months	42.41	57.90	51.68	47.81	46.00	13.78	64.71	0.57
SD/47 months	60.15	82.03	73.69	68.42	64.62	29.62	123.66	0.56
SD/59 months	66.21	95.10	83.58	75.48	71.93	47.02	204.38	0.56
NW/17 months	5.53	7.69	6.80	7.00	6.59	0.22	1.23	0.45
NW/22 months	7.52	10.08	9.03	9.27	8.75	0.32	1.75	0.45

788 **Table 5.** QTL mapping for tree height, number of whorls, and stem diameter for progenies of the cross GT1 x RRIM701. The phenotypes
789 were evaluated at 12, 17, 22, 28, 35, 47, and 59 months. LG - linkage group; LOD - global LOD score; R^2 - explained phenotypic variation;
790 LOD $^{(2),(3),(4)}$ - LOD values for additive and dominance effects.

QTL	ID in Fig. 3	LG	Position (cM)	LOD	R^2	Additive effect for GT1	LOD$^{(2)}$	Additive effect for RRIM701	LOD$^{(3)}$	Dominance effect	LOD$^{(4)}$	Segregation pattern
Height17-LG4	A	4	43.41	6.72	14.07	-0.049	3.256	0.054	3.965	-0.018	0.496	1:2:1
Height17-LG9	B	9	196.26	5.28	6.24	-0.002	0.006	0.038	1.881	-0.050	3.106	1:2:1
Height17-LG15	C	15	108.00	4.23	7.44	-0.004	0.020	-0.039	2.067	-0.049	2.789	1:2:1
Height17-LG18	D	18	251.44	4.60	7.38	-0.029	1.193	-0.055	3.606	-0.007	0.054	1:2:1
Height22-LG5	E	5	15.00	5.70	7.18	-0.027	3.782	-0.019	1.734	0.013	0.828	1:2:1
Height22-LG6	F	6	155.55	4.89	8.14	-0.033	1.177	-0.003	0.034	0.031	4.362	1:2:1
Height22-LG11	G	11	189.00	4.37	8.40	-0.003	0.053	0.026	3.215	0.017	1.245	1:2:1
Whorls17-LG4	A	4	49.91	7.27	8.25	-0.089	3.419	0.093	3.810	0.058	1.075	3:1
Whorls17-LG14	B	14	26.00	4.13	1.08	-0.067	1.755	-0.134	2.805	0.012	0.065	1:2:1
Whorls17-LG15	C	15	105.19	5.56	10.22	-0.032	0.366	-0.088	2.827	-0.099	3.310	1:2:1
Whorls22-LG4	D	4	6.04	8.59	7.67	0.011	0.853	-0.009	0.531	-0.029	7.036	1:1:1:1
Whorls22-LG8	E	8	71.10	4.78	3.78	0.012	1.750	-0.014	2.120	0.012	1.822	3:1
Whorls22-LG12	F	12	152.15	6.26	5.31	0.009	1.078	0.017	3.470	0.011	1.518	3:1
Whorls22-LG15	G	15	121.11	4.49	6.01	-0.006	0.406	-0.003	0.086	-0.020	4.059	1:1
Diameter12-LG2	A	2	157.93	4.22	3.80	-0.077	0.619	-0.044	0.224	0.195	4.017	1:1
Diameter12-LG4	B	4	34.77	5.69	9.73	-0.106	1.443	0.143	2.087	0.118	1.295	3:1
Diameter12-LG6	C	6	170.00	4.72	8.75	-0.140	2.258	-0.057	0.376	0.164	2.802	1:2:1
Diameter12-LG9	D	9	196.26	4.73	6.43	-0.129	1.834	-0.025	0.071	-0.189	3.800	1:2:1
Diameter12-LG17	E	17	11.05	4.71	5.13	0.274	2.179	-0.056	0.442	0.140	2.471	1:2:1
Diameter17-LG4	F	4	49.91	8.08	13.04	-0.081	3.381	0.093	4.152	0.042	0.823	1:2:1
Diameter17-LG6	G	6	163.63	5.09	6.87	-0.042	0.899	-0.040	0.867	0.087	4.057	3:1
Diameter17-LG9	H	9	193.00	5.85	4.74	-0.049	1.134	-0.012	0.062	-0.119	5.344	1:1:1:1
Diameter17-LG17	I	17	35.00	5.99	6.58	0.028	0.406	-0.019	0.205	0.110	5.068	1:1
Diameter22-LG5	J	5	6.74	4.66	8.09	-0.129	2.771	-0.027	0.123	0.116	2.000	1:2:1

Diameter22-LG9	K	9	196.26	6.75	6.61	-0.121	2.593	-0.040	0.274	-0.189	5.480	1:2:1
Diameter22-LG15	L	15	118.00	8.76	7.79	-0.107	1.395	-0.114	2.061	-0.204	6.648	3:1
Diameter28-LG5	M	5	8.00	8.21	11.40	0.072	7.216	0.030	1.224	-0.016	0.342	1:1:1:1
Diameter28-LG8	N	8	229.67	4.52	4.89	-0.035	1.480	0.036	1.926	-0.017	0.381	1:2:1
Diameter28-LG11	O	11	91.00	6.24	8.08	-0.087	3.214	-0.005	0.041	0.051	3.438	1:2:1
Diameter28-LG12	P	12	27.00	7.63	8.37	-0.086	5.561	0.027	0.749	0.074	1.800	1:2:1
Diameter35-LG4	Q	4	37.00	10.06	12.28	-0.217	1.137	0.479	4.688	0.514	4.125	3:1
Diameter35-LG12	R	12	136.00	5.74	3.99	0.190	0.826	0.013	0.004	0.490	4.580	1:1
Diameter35-LG17	S	17	24.00	6.68	3.01	1.437	3.179	0.530	2.050	0.328	2.380	1:2:1
Diameter47-LG4	T	4	28.00	6.29	10.10	-0.261	1.584	0.510	3.062	0.222	0.673	1:2:1
Diameter47-LG17	U	17	11.05	4.37	5.08	0.262	1.533	-0.196	0.913	0.301	1.723	3:1
Diameter59-LG9	V	9	144.00	5.61	9.59	-0.316	3.071	-0.173	0.921	-0.337	2.978	3:1
Diameter59-LG12	W	12	24.24	4.97	13.03	0.258	2.047	0.202	0.949	-0.538	2.712	3:1
Diameter59-LG15	X	15	224.00	4.15	4.65	0.262	2.058	0.284	1.915	-0.134	0.438	1:2:1

Table S1. SNPs developed from full-length expressed sequence tag (EST) libraries and bark
transcriptome assembled de novo.

Table S2. Variance-covariance structures adjusted and compared based on the Akaike information
criterion (AIC) (Akaike, 1974) and the Bayesian information criterion (BIC) (Schwarz, 1978).
$G_{(AB,AG)}$: variance-covariance matrices for effects of replicates within block within age and genotype
within age; $R_{(C,R,\,A)}$: variance-covariance matrices for column, row, and age for the residual effects;
df: degrees of freedom; ID: identity matrix (homogeneous genetic variance without correlation);
Diag: diagonal matrix (heterogeneous genetic variance without correlation); FA: factor analytic;
Corh: matrix of heterogeneous variance with correlation; US: non-structured variance-covariance
matrix; Ar1h: first-order autoregressive correlation matrix with heterogeneous variance; Ar1: first-
order autoregressive correlation matrix without variance.

Table S3. Consensus map anchoring the final rubber tree map with 1,079 markers for the cross GT1
x RRIM701 into the scaffolds assembled by Tang *et al*. (2016).

Table S4. Functional annotation for genes contained within the marker interval of each identified
QTL.

807