Signals of polygenic adaptation on height have been overestimated due to uncorrected population structure in genome-wide association studies

Mashaal Sohail[1,2,3+], Robert M. Maier[3,4,5+], Andrea Ganna[3,4,5,6,7], Alexander Bloemendal[3,4,5], Alicia R. Martin[3,4,5], Michael C. Turchin[8,9], Charleston W. K. Chiang[10], Joel N. Hirschhorn[3,11,12], Mark J. Daly[3,4,5,7], Nick Patterson[3,13], Benjamin M. Neale[3,4,5]*, Iain Mathieson[14]*, David Reich[3,13,15]*, Shamil R. Sunyaev[1,2,3]*

[1] Division of Genetics, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA
[2] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, USA
[3] Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[4] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[5] Analytical and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA 02114, USA
[6] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm SE-171 77, Sweden
[7] Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki FI-00014, Finland
[8] Center for Computational Molecular Biology, Brown University, Providence, RI, USA
[9] Department of Ecology and Evolutionary Biology, Brown University, Providence, RI, USA
[10] Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California, USA.
[11] Departments of Pediatrics and Genetics, Harvard Medical School, Boston, MA 02115, USA
[12] Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA 02115, USA
[13] Department of Genetics, Harvard Medical School, Boston, MA 02115, USA
[14] Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19103, USA
[15] Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115, USA

[+] Contributed Equally
[*] Co-supervised

All correspondence should be addressed to: ssunyaev@rics.bwh.harvard.edu, reich@genetics.med.harvard.edu, mathi@pennmedicine.upenn.edu, bneale@broadinstitute.org

Abstract

Genetic predictions of height differ among human populations and these differences are too large to be explained by genetic drift. This observation has been interpreted as evidence of polygenic adaptation. Differences across populations were detected using SNPs genome-wide significantly associated with height, and many studies also found that the signals grew stronger when large numbers of sub-significant SNPs were analyzed. This has led to excitement about the prospect of analyzing large fractions of the genome to detect subtle signals of selection and claims of polygenic adaptation for multiple traits. Polygenic adaptation studies of height have been based on SNP effect size measurements in the GIANT Consortium meta-analysis. Here we repeat the height analyses in the UK Biobank, a much more homogeneously designed study. Our results show that polygenic adaptation signals based on large numbers of SNPs below genome-wide significance are extremely sensitive to biases due to uncorrected population structure.

Introduction

Most human complex traits are highly polygenic.[1,2] For example, height has been estimated to be modulated by as much as 4% of human allelic variation.[2],[3] Polygenic traits are expected to evolve differently from monogenic ones, through slight but coordinated shifts in the frequencies of a large number of alleles, each with mostly small effect. In recent years, multiple methods have sought to detect selection on polygenic traits by evaluating whether shifts in the frequency of trait-associated alleles are correlated with the signed effects of the alleles estimated by genome-wide association studies (GWAS).[4−10]

Here we focus on a series of recent studies—some involving co-authors of the present manuscript—that have reported evidence of polygenic adaptation at alleles associated with height in Europeans. One set of studies observed that height-increasing alleles are systematically elevated in frequency in northern compared to southern European populations, a result that has subsequently been extended to ancient DNA.[4−11]Another study using a very different methodology (singleton density scores, SDS) found that height-increasing alleles have systematically more recent coalescent times in the United Kingdom (UK) consistent with selection for increased height over the last few thousand years.[12]

All of these studies have been based on SNP associations, in most cases with effect sizes discovered by the GIANT Consortium, which most recently combined 79 individual GWAS through meta-analysis, encompassing a total of 253,288 individuals.[13,14] Here, we show that the selection effects described in these studies are severely attenuated and in some cases no longer significant when using summary statistics derived from the UK Biobank, an independent and larger single study that includes 336,474 genetically unrelated individuals who derive their ancestry almost entirely from British Isles (identified as "white British ancestry" by the UK Biobank) (**Supplementary Table S1**). The UK Biobank analysis is based on a single cohort drawn from a relatively homogeneous population enabling excellent control of potential population stratification. Our analysis of the UK Biobank data confirms that almost all genome-wide significant loci discovered by the GIANT

93   consortium are real associations, and that the two datasets have high concordance
94   for low P value SNPs which do not reach genome-wide significance
95   (**Supplementary Figure S1**; genetic correlation between the two height studies is
96   0.94 [se=0.0078]). However, our analysis yields qualitatively different conclusions
97   with respect to signals of polygenic adaptation.
98
99   **Results**
100
We began by estimating "polygenic height scores"—sums of allele
frequencies at independent SNPs from GIANT weighted by their effect sizes—to
study population level differences among ancient and present-day European
samples. We used a set of different significance thresholds and strategies to correct
for linkage disequilibrium as employed by previous studies, and replicated their
signals for significant differences in genetic height across populations.[4–11]
(**Figure 1a, Supplementary Figure S2**). We then repeated the analysis using
summary statistics from a GWAS for height in the UK Biobank restricting to
individuals of British Isles ancestry and correcting for population stratification
based on the first ten principal components (UKB).[15] This analysis resulted in a
dramatic attenuation of differences in polygenic height scores (**Figure 1a,**
Supplementary Figures S2-S4). The differences between ancient European
populations also greatly attenuated (**Figure 1a, Supplementary Figure S5**).
Strikingly, the ordering of the scores for populations also changed depending on
which GWAS was used to estimate genetic height both within Europe (**Figure 1a,**
Supplementary Figures S2-S5) and globally (**Supplementary Figure S6**),
consistent with reports from a recent simulation study.[16] The height scores were
qualitatively similar only when we restricted to independent genome-wide
significant SNPs in GIANT and the UK Biobank ($P < 5x10^{-8}$) (**Supplementary Figure**
S2b). This replicates the originally reported significant north-south difference in the
allele frequency of the height-increasing allele[4] or in genetic height[5] across
Europe, as well as the finding of greater genetic height in ancient European steppe
pastoralists than in ancient European farmers,[6] although the signals are
attenuated even here. This suggests that tests of polygenic adaptation based on
genome-wide significant SNPs may be relatively insensitive to confounding
(**Supplementary Figure S2b**), and that confounding due to stratification is a
particular danger for sub-significant SNPs (**Figure 1a, Supplementary Figure S2a**).
128

Figure 1. **Polygenic height scores and tSDS scores based on GIANT and UK Biobank GWAS.**

Polygenic scores in present-day and ancient European populations are shown, centered by the average score across populations and standardized by the square root of the additive variance. Independent SNPs for the polygenic score from both GIANT (*red*) and the UK Biobank (*blue*) were selected by picking the SNP with the lowest P value in each of 1700 independent LD blocks similarly to refs [8,9] (see methods). Present-day populations are shown from Northern Europe (CEU, GBR) and Southern Europe (IBS, TSI) from the 1000 genomes project; Ancient populations are shown in three meta-populations (HG = Hunter-Gatherer (n=162 individuals), EF = Early Farmer (n=485 individuals), and SP = Steppe Ancestry (n=465 individuals)) (see **Supplementary Table S2**). Error bars are drawn at 95% credible intervals. See **Supplementary figures S2-S6** for polygenic height scores computed using other linkage disequilibrium pruning procedures, significance thresholds, summary statistics and populations. (b) tSDS for height-increasing allele in GIANT (left) and UK Biobank (right). The tSDS method was applied using pre-computed Singleton Density Scores for 4,451,435 autosomal SNPs obtained from 3,195 individuals from the UK10K project[12],[17] for SNPs associated with height in GIANT and the UK biobank. SNPs were ordered by GWAS P value and grouped into bins of 1000 SNPs each. The mean tSDS score within each P value bin is shown on the y-axis. The Spearman correlation coefficient between the tSDS scores and GWAS P values, as well as the correlation standard errors and P values, were computed on the un-binned data. The gray line indicates the null-expectation, and the colored lines are the linear regression fit. The correlation is significant for GIANT (Spearman r = 0.078, P = 1.55 x 10⁻⁶⁵) but not for UK Biobank (Spearman r = -0.009, P = 0.077).

Next, we looked at polygenic adaptation within the UK using the "singleton
density score" (SDS)—an independent measure that uses the local density of alleles
that occur only once in the sample as a proxy for coalescent branch lengths.[12,17]
SDS can be combined with GWAS effect sizes estimates by aligning the SDS sign to
the trait-increasing allele, after which the score is referred to as tSDS. A tSDS score
larger than zero implies that height-increasing alleles have been increasing in
frequency over time due to natural selection. We replicate the finding that tSDS
computed in the UK10K is positively rank-correlated with GIANT[12] height P
values (Spearman's ρ = 0.078, P = 1.55 x 10^{-65}, **Figure 1b**). However, this signal of
polygenic adaptation in the UK attenuated when we used UK Biobank height effect
size estimates and P values and became formally non-significant (ρ = 0.009, P =
0.077, **Figure 1b**).
We propose that the qualitative difference between the polygenic adaptation
signals in GIANT and the UK Biobank is the cumulative effect of subtle biases in each
of the contributing SNPs in GIANT. This bias can arise due to incomplete control of
the population structure in GWAS.[18] For example, if height were differentiated
along a north-south axis because of differences in environment, any variant that is
differentiated in frequency along the same axis would have an artifactually large
effect size estimated in the GWAS. Population structure is substantially less well
controlled for in the GIANT study than in the UK Biobank study, both because the
GIANT study population is more heterogeneous than that in the UK Biobank, and
because the population structure in GIANT may not have been well controlled in
some component cohorts of GIANT due to the relatively small size of individual
studies (i.e., the ability to detect and correct population structure is dependent on
sample size[19,20]). The GIANT meta-analysis also found that such stratification
effects worsen as SNPs below genome-wide significance are used to estimate height
scores,[14] consistent with our finding that the differences in genetic height
increase when including these SNPs.
To obtain further insight into our observed discrepancy between polygenic
adaptation signals in GIANT vs the UK Biobank, we repeated our analyses using
estimates of height effect sizes computed using different methods, and then
interrogated each of these for signs of population structure. Repeating our analysis
with family-based effect size estimates from an independent study (NG2015
sibs),[7] we found evidence for significant differences in polygenic scores between
northern and southern Europeans that were qualitatively similar to those obtained
using GIANT effect size estimates (**Supplementary Figure S4-S5**). Inclusion of
individuals from the UK Biobank who were not of British Isles ancestry without
controlling for population structure (UKB all no PCs) in the measurements of effect
sizes also produced this pattern (**Supplementary Figure S3-S5**). Thus, UK Biobank
estimates that retain population structure show similar patterns to GIANT and
previously published family-based estimates (NG 2015 sibs). In contrast, no
significant signals of genetic stratification of height or a strong tSDS signal are
present across populations from: 1) a genetically homogeneous sample of UK
Biobank with entirely British Isles ancestry without controlling for population
structure (UKB WB no PCs), or 2) effect size estimates based on UK Biobank families

(UKB sibs, UKB sibs WB) (**Supplementary Figures S3-S5, S7-S8**). These analyses
provide further evidence that the lack of signal in the UK Biobank analysis is
unlikely to be simply due to over-correction for structure in the original UKB
estimates.
202
203

204
Figure 2. Evidence of stratification in height summary statistics.
Top row: Pearson Correlation coefficients of (a) PC loadings and height beta coefficients
from GIANT and UKB, and (b) PC loadings and SDS (pre-computed in the UK10K) across all
SNPs. PCs were computed in all 1000 genomes phase 1 samples. Colors indicate the
correlation of each PC loading with the allele frequency difference between GBR and TSI, a
proxy for the European North-South genetic differentiation. PC 4 and 11 are most highly
correlated with the GBR - TSI allele frequency difference. Confidence intervals and P values
are based on Jackknife standard errors (1000 blocks). Open circles indicate correlations
significant at alpha = 0.05, stars indicate correlations significant after Bonferroni correction
in 20 PCs (P < 0.0025).
Bottom row: Heat map after binning all SNPs by GBR and TSI minor allele frequency of (c)
mean beta coefficients from GIANT and UKB, and (d) SDS scores for all SNPs. Only bins with
at least 300 SNPs are shown. While the stratification effect in SDS is not unexpected, it can
lead to false conclusions when applied to summary statistics that exhibit similar
stratification effects. UKB height betas exhibit stratification effects that are weaker, and in
the opposite direction of the stratification effects in GIANT (see **Supplementary Figure S9**
for a possible explanation).
222
We obtained direct confirmation that population structure is more correlated
with effect size estimates in GIANT than to those in the UK Biobank. **Figure 2a**
shows that the effect sizes estimated in GIANT are highly correlated with the SNP
loadings of several principal components of population structure (PC loadings).

Previously published family-based effect size estimates[7] (NG 2015 sibs) are
similarly correlated with the PC loadings showing that they are also affected by
population structure despite being computed within families; in other words, these
empirical analyses show that even previously published family-based effect size
estimates are not free from concerns about population structure. The within-family
strategy for eliminating concerns about population stratification is not problematic
on its own as our UK Biobank family estimates (UKB sibs, UKB sibs WB) computed
using the same method do not show any stratification effects (**Supplementary
Figures S10-S12**). We also do not see a strong correlation with PC loadings in our
UK Biobank estimates computed using unrelated individuals (UKB)(**Figure 2a**).
However, the UK Biobank estimates including individuals not of British Isles
ancestry and not correcting for population structure (UKB all no PCs) show the
same stratification effects as GIANT and NG2015 sibs (**Supplementary Figure S10-
S12**). Similarly, we find that alleles that are more common in the Great Britain
population (GBR) than in the Tuscan population from Italy (TSI) tend to be
preferentially height-increasing according to the GIANT and NG2015 sibs estimates
but not according to the UKB estimates (**Figure 2c, Supplementary Figures S11,
S12**).
The tSDS analysis should be robust to the type of population structure
discussed above.[12] However, there is a north-south cline in singleton density in
Europe due to the lower genetic diversity in northern than in southern Europeans,
with singleton density being lower in northern than in southern regions.[21] As a
consequence, SDS tends to be higher in alleles more common in GBR than in TSI
(**Figure 2d**). This cline in singleton density coincidentally parallels the phenotypic
cline in height and the major axis of genome-wide genetic variation. Therefore,
when we perform the tSDS test using GIANT-estimated effect sizes and P values, we
find fewer singletons (corresponding to higher SDS) around the inferred height-
increasing alleles which tend, due to the uncontrolled population stratification in
GIANT, to be at high frequency in northern Europe (**Figures 2c**). This effect does not
appear when we use UK Biobank summary statistics because of the much lower
level of population stratification and more modest variation in height. We find that
SDS is not only correlated with GBR-TSI allele frequency differences, but with
several principal component loadings across all SNPs (**Figure 2b**), and that these
SDS-PC correlations often coincide with correlations between GIANT-estimated
effect sizes and PC loadings (**Figure 2a**).
We further find that the tSDS signal which is observed across the whole
range of P values in some summary statistics can be mimicked by replacing SDS
with GBR-TSI allele frequency differences (**Figures 3a, 3c, Supplementary Figures
S7-S8, S13-S14**), suggesting that the tSDS signal at non-significant SNPs may be
driven in part by residual population stratification. As with the polygenic score
analysis, a small but significant effect is observed when we restrict to genome-wide
significant SNPs ($P < 5 \times 10^{-8}$). This effect persists when using UK Biobank family-
based estimates for genome-wide significant SNPs (**Figure 3b**), and is not driven by
allele frequency differences between GBR and TSI (**Figure 3d**), suggesting a true but
attenuated signal of polygenic adaptation in the UK that is driven by a much smaller
272

273
274
Figure 3. Height tSDS results for different summary statistics.
(a) Mean tSDS of the height increasing allele in each P value bin for six different summary statistics. The first two panels are computed analogously to Figure 4A and Figure S22 in Field et al. In contrast to those Figures and to **Figure 1b**, the displayed betas and P values correspond to the slope and P value of the linear regression across all un-binned SNPs (rather than the Spearman correlation coefficient and Jackknife P values). The y-axis has been truncated at 0.75, and does not show the top bin for UKB all no PCs, which has a mean tSDS of 1.5. See **Supplementary Figure S7** for other GWAS summary statistics (b) tSDS distribution of the height increasing allele in 506 LD-independent SNPs which are genome-wide significant in a UKB height GWAS, where the beta coefficient is taken from a within sibling analysis in the UKB. The gray curve represents the standard normal null distribution, and we observe a significant shift providing confirmation of a real SDS signal of polygenic adaptation for height. (c) Allele frequency difference between GBR and TSI of the height increasing allele in each P value bin for six different summary statistics. Betas and P values correspond to the slope and P value of the linear regression across all un-binned SNPs. The lowest P value bin in UKB all no PCs with a y-axis value of 0.06 has been omitted. See **Supplementary Figure S13** for other GWAS summary statistics. (d) Allele frequency difference between GBR and TSI of the height increasing allele in 329 LD-independent SNPs which are genome-wide significant in a UKB height GWAS and were intersecting with our set of 1000 genomes SNPs. There is no significant difference in frequency in these two populations, suggesting that tSDS shift at the gw-significant SNPs is not driven by population stratification.
The patterns shown here suggest that the positive tSDS values across the whole range of P values is a consequence of residual stratification. At the same time, the increase in tSDS at genome-wide significant, LD-independent SNPs in (b) cannot be explained by GBR - TSI allele frequency differences as shown in (d). Binning SNPs by P value without LD-pruning can lead to unpredictable patterns at the low P value end, as the SNPs at the low P value end are less independent of each other than higher P value SNPs (**Supplementary Figure S15**). **Supplementary Figures S8 and S14** therefore show the same data for a set of LD-pruned SNPs.

305
306 number of SNPs than previously thought. Indeed, a tSDS signal which is driven by
307 natural selection is not expected to lead to an almost linear increase over the whole
308 P value range in a well-powered GWAS. Instead, we would expect to see a greater
309 difference between highly significant SNPs and non-significant SNPs, similar to the
310 pattern observed in the UK Biobank (**Figure 3a**).
311
312

313
Figure 4. Polygenic height scores in POPRES populations show a residual albeit
attenuated signal of polygenic adaptation for height.
Standardized polygenic height scores from six summary statistics for 19 POPRES
populations with at least 10 samples per population, ordered by latitude (see
Supplementary Table S3). The grey line is the linear regression fit to the mean polygenic
scores per population. Error bars represent 95% confidence intervals and are calculated in
the same way as in **Figure 1**.
SNPs which were overlapping between each set of the summary statistics and the POPRES
SNPs were clumped using PLINK 1.9 with parameters $r^2 < 0.1$, 1 Mb distance, $P < 1$.
(Top) A number of independent SNPs was chosen for each summary statistic to match the
number of SNPs which remained when clumping UKB at $P < 0.01$.
(Bottom) A set of independent SNPs with $P < 5 \times 10^{-8}$ in the UK Biobank was selected and
used to compute polygenic scores along with effect size estimates from each of the different
summary statistics. SNP numbers for each panel ($P < 0.01$ / $P < 5 \times 10^{-8}$): GIANT: 9463 /
869; UKB all no PCs: 9341 / 359; UKB: 9341 / 1034; UKB sibs WB: 9391 / 380. The
numbers on each plot show the Q_x P value and the latitude covariance P value respectively
for each summary statistic. See **Supplementary Figures S16-S18** for other clumping
strategies and GWAS summary statistics.
332
Lastly, we asked whether any remaining differences in polygenic height
scores among populations are driven by polygenic selection by using the Q_x

framework to test against a null model of genetic drift.[5] We re-computed
polygenic height scores in the POPRES dataset for this analysis as it has larger
sample sizes of northern and southern Europeans than the 1000 Genomes
project.[22] We computed height scores using independent SNPs that are 1)
genome-wide significant in the UK Biobank ("gw-sig", $P < 5 \times 10^{-8}$) and 2) sub-
significantly associated with height ("sub-sig", $P < 0.01$) in different GWAS datasets.
For each of these, we tested if population differences were significant due to an
overall overdispersion (P_{Qx}), and if they were significant along a north-south cline
(P_{lat}) (**Figure 4, Supplementary Figure S16-S17**). Both gw-sig and sub-sig SNP-
based scores computed using GIANT effect sizes showed significant overdispersion
of height scores overall and along a latitude cline, consistent with previous results
(**Figure 4, Supplementary Figure S16-S17**). However, the signal attenuated
dramatically between sub-sig ($Q_x = 1100$, $P_{Qx} = 1 \times 10^{-220}$) and gw-sig ($Q_x = 48$, $P_{Qx} =$
2×10^{-4}) height scores. In comparison, scores that were computed using the UK
Biobank (UKB) effect sizes showed substantially attenuated differences using both
sub-sig ($Q_x = 64$, $P_{Qx} = 5 \times 10^{-7}$) and gw-sig ($Q_x = 33$, $P_{Qx} = 0.02$) SNPs, and a smaller
difference between the two scores. This suggests that the attenuation of the signal in
GIANT is not only driven by a loss of power when using fewer gw-sig SNPs, but also
reflects a decrease in stratification effects. The overdispersion signal disappeared
entirely when the UK Biobank family based effect sizes were used (**Figure 4,**
Supplementary Figure S16-S17). Moreover, Q_x P values based on randomly
ascertained SNPs and UK Biobank summary statistics are not uniformly distributed
as would be expected if the theoretical null model is valid and if population
structure is absent (**Supplementary Figure S19**). The possibility of residual
stratification effects even in the UK Biobank is also supported by a recent study [23].
Therefore, we remain cautious about interpreting any residual signals as "real"
signals of polygenic adaptation.
362
363 **Discussion**
364
We have shown that estimates of population differences in polygenic height
scores are strikingly attenuated with the UK Biobank GWAS data relative to
previous analyses. We find some evidence for population-level differences in
genetic height, but it can only be robustly seen at highly significant SNPs, because
any signal at less significant P values is dominated by the effect of residual
population structure. Even genome-wide significant SNPs in these analyses may be
subtly affected by population structure, leading to continued overestimation of the
effect. Thus, it is difficult to arrive at any quantitative conclusion regarding the
proportion of the population differences that are due to statistical biases vs.
population stratification of genetic height. It is equally challenging to test whether
differences in genetic height are due to adaptation in response to environmental
differences, migration and admixture (e.g. fraction of Steppe pastoralist ancestry),
or relaxation of negative selection. Further, estimates of the number of independent
genetic loci contributing to height variation are sensitive to and likely confounded
by residual population stratification.

We conclude that while effect estimates are highly concordant between
GIANT and the UK Biobank when measured individually (**Supplementary Tables**
S4-S6, Supplementary Figure S1), they are also influenced by residual population
stratification that can mislead inferences about polygenic selection across
populations in aggregate. Although these biases are subtle, in the context of tests for
polygenic adaptation, which are driven by small systematic shifts in allele
frequency, they can create highly significant artificial signals especially when SNPs
that are not genome-wide significant are used to estimate genetic height. In no way
do our results question the reliability of the genome-wide significant associations
discovered in the GIANT cohort or the validity of the statistical methodology used in
previously reported polygenic tests for adaptation. However, we urge caution in the
interpretation of genome-wide signals of polygenic adaption that are based on large
number of sub-significant SNPs–particularly when using effect sizes derived from
meta-analysis of heterogeneous cohorts which may be unable to fully control for
population structure.
395
396
397 **Materials and Methods**
398
Genome-wide association studies (GWAS)
We analyzed height using publicly available summary statistics that were
obtained either by meta-analysis of multiple GWAS or by a GWAS performed on a
single large population. We used results from the GIANT Consortium
(N=253,288)[14] and a GWAS performed on individuals of the UK Biobank (UKB
Neale" or simply "UK Biobank (UKB)", N=336,474)[15] who derive their ancestry
almost entirely from the British Isles (identified as "white British ancestry" by the
UK Biobank). We also used an independent GWAS that included all UK Biobank
individuals irrespective of ancestry and relatedness ("UKB Loh", N=459,327)[24].
The Neale lab's GWAS uses a linear model with sex and 10 principal components as
covariates. Loh *et al.*'s GWAS uses a BOLT-LMM Bayesian mixed model. Association
signals from the three studies are generally correlated for SNPs that are genome-
wide significant in GIANT (see [[25]]).
412
We also used previously published family-based effect size estimates[7]
("NG2015 sibs") as well as a number of test summary statistics on the UK Biobank
that we generated to study the effects of population stratification. These are: "UKB
Neale new" (Similar to UKB Neale, with less stringent ancestry definition and 20 PCs
calculated within sample), "UKB all no PCs" (All UK Biobank samples included in the
GWAS without correction by principal components), "UKB all 10 PCs" (All UK
Biobank samples included in the GWAS with correction by 10 principal
components), "UK WB no PCs" (Only "white British ancestry" samples included in
the GWAS without correction by principal components), "UKB WB 10 PCs" (Only
"white British ancestry" samples included in the GWAS with correction by 10
principal components), "UKB sibs all" (All UK Biobank siblings included in the
GWAS), "UKB sibs WB" (Only UK Biobank "white British ancestry" siblings included

in the GWAS) (Please see **Supplementary Table S1** for sample sizes and other
details).
427
Population genetic data for ancient and modern samples

We analyzed ancient and modern populations for which genotype data are
publicly available. For ancient samples[26,27], we computed scores after dividing
populations into three previously described broad ancestry labels (HG = Hunter-
Gatherer (n=162 individuals), EF = Early Farmer (n=485 individuals), and SP =
Steppe Ancestry (n=465 individuals)). For modern samples available through the
1000 genomes phase 3 release[28], we computed scores in 2 populations each from
Northern Europe (GBR, CEU), Southern Europe (IBS, TSI), Africa (YRI, LWK), South
Asia (PJL, BEB) and East Asia (CHB, JPT) (**Figure 1a**, **Supplementary Figures S2-**
S6). In total, we analyzed 1112 ancient individuals, and 1005 modern individuals
from 10 different populations in the 1000 genomes project (**Supplementary Table**
S2). We used the allele frequency differences between the GBR and TSI populations
for a number of analyses to study population stratification (**Figures 3c**, **3d**,
Supplementary Figures S11-S14). We also analyzed 19 European populations
from the POPRES[22] dataset with at least 10 samples per population (**Figure 4**,
Supplementary Table S3, Supplementary Figures S16-S18).
444
All ancient samples had 'pseudo-haploid' genotype calls at 1240k sites
generated by selecting a single sequence randomly for each individual at each SNP[9].
Thus, there is only a single allele from each individual at each site, but adjacent
alleles might come from either of the two haplotypes of the individual. We also re-
computed scores in present-day 1000 genomes individuals using only pseudo-
haploid calls at 1240k sites to allow for a fair comparison between ancient and
modern samples (**Supplementary Figure S6**).
452
Polygenic scores

454   The polygenic scores, confidence intervals and test statistics (against the null
455 model of genetic drift) were computed based on the methodology developed in refs
456 ([5],[29]). We computed the polygenic score (Z) for a trait in a population by taking
457 the sum of allele frequencies across L sites associated with the trait, weighting each
458 allele's frequency by its effect on the trait (β_l).
459

$$Z = \sum_{l}^{L} \beta_l p_l$$

460
461 Al polygenic scores are plotted in centered standardized form ($\frac{Z-\mu}{\sqrt{V_A}}$), where
462 $\mu = \sum_l \beta_l \, \overline{p}_l$, $V_A = \sum_l \beta_l^2 \, \overline{p}_l(1-\overline{p}_l)$, and $\overline{p}_l$ is the mean allele frequency across all
463 populations analyzed.
464
Polygenic scores were computed using independent GWAS SNPs associated
with height in three main ways: (1) The genome was divided into ~1700 non-

overlapping linkage disequilibrium (LD) blocks[30], and the SNP with the lowest P
value within each block was picked to give a set of ~1700 independent SNPs for
each height GWAS used (all SNPs for which effect sizes are available were
considered) similarly to ref. [29]. In (2) and (3), Plink's[31,32] clumping procedure
was used to make independent "clumps" of SNPs for each GWAS at different P value
thresholds. This procedure selects SNPs below a given P value threshold as index
SNPs to start clumps around, and then reduces all SNPs (P < 0.01) that are in LD
with these index SNPs (above an r^2 threshold, 0.1) and within a physical distance of
them (1 Mb) into clumps with them. Clumps are preferentially formed around index
SNPs with the lowest P value in a greedy manner. The index SNP from each clump is
then picked for further polygenic score analyses. The algorithm is also greedy such
that each SNP will only appear in one clump if at all. We clumped each GWAS to
obtain (2) a set of independent sub-significant SNPs associated with height (P <
0.01) similarly to ref. [7], and (3) a set of genome-wide significant SNPs associated
with height (P < 5 x 10^{-8}). The 1000 genomes phase 3 dataset was used as the
reference panel for computing LD for the clumping procedure.
483
484   The estimated effect sizes for these three sets of SNPs from each GWAS was
485 used to compute scores. Only autosomal SNPs were used for all analyses to avoid
486 creating artificial mean differences between populations with different numbers of
487 males and females.
488
The 95% credible intervals were constructed by assuming that the posterior
of the underlying population allele frequency is independent across loci and
populations and follows a beta distribution. We updated a Uniform prior
distribution with allele counts from ancient and modern populations to obtain the
posterior distribution at each locus in each population. We estimated the variance of
the polygenic score V_Z using the variance of the posterior distribution at each locus,
and computed the width of 95% credible intervals as $1.96\sqrt{V_Z}$ for each population.
496
The Q_x test statistic measures the degree of overdispersion of the mean
population polygenic score compared to a null model of genetic drift. It assumes that
the vector of mean centered mean population polygenic score follows a multivariate
normal distribution: $Z \sim MVN(0, 2 V_A F)$, where V_A is the additive genetic variance of
the ancestral population and F is a square matrix describing the population
structure. This is equivalent to the univariate case of the test statistic used in ref .
[7]. The latitude test statistic assumes that $Y'Z \sim N(0, 2 V_A Y'FY)$, where Y is a mean
centered vector of latitudes for each population.[33]
505
tSDS analysis
The Singleton Density Score (SDS) method identifies signatures of recent
positive selection based on a maximum likelihood estimate of the log-ratio of the
mean tip-branch length of the derived vs. the ancestral allele at a given SNP. The tip-
branch lengths are inferred from the average distance of each allele to the nearest
singleton SNP across all individuals in a sequencing panel. When the sign of the SDS

scores is aligned with the trait-increasing or trait-decreasing allele in the effect estimates of a GWAS, the Spearman correlation between the resulting tSDS scores and the GWAS P values has been proposed as an estimate of recent positive selection on polygenic traits.

Here, we applied the tSDS method using pre-computed Singleton Density Scores for 4,451,435 autosomal SNPs obtained from 3,195 individuals from the UK10K project [12,17] for SNPs associated with height in GIANT and the UK biobank (**Figure 1b**) and in different summary statistics (**Figure 3, Supplementary Figures S7-S8**). After normalizing SDS scores in each 1% allele frequency bin to mean zero and unit variance, excluding SNPs from the MHC region on chromosome 6 and aligning the sign of the SDS scores to the height increasing alleles (resulting in tSDS scores), we computed the Spearman correlation coefficient between the tSDS score and the GWAS P value. The tSDS Spearman correlation standard errors and P values were computed using a block-jackknife approach, where each block of 1% of all SNPs ordered by genomic location was left out and the Spearman correlation coefficient was computed on the remaining SNPs. We also compared the tSDS score distributions for only genome-wide significant SNPs (**Figure 3b**).

Population structure analysis

To compute SNP loadings of the principal components of population structure (PC loadings) in the 1000 genomes data (**Figure 2, Supplementary Figure S10**), we first computed PC scores for each individual. We used SNPs that had matching alleles in 1000 genomes, GIANT and UK Biobank, that had minor allele frequency > 5% in 1000 genomes, and that were not located in the MHC locus, the chromosome 8 inversion region, or regions of long LD. After LD pruning to SNPs with $r^2 < 0.2$ relative to each other, PCA was performed in PLINK on the 187,160 remaining SNPs. In order to get SNP PC loadings for more SNPs than those that were used to compute PC scores, we performed linear regressions of the PC scores on the genotype allele count of each SNP (after controlling for sex) and used the resulting regression coefficients as the SNP PC loading estimates.

Acknowledgements

We thank Alkes Price, Jeremy Berg, Graham Coop, Jonathan Pritchard, Matthew Robinson, Jian Yang, Peter Visscher, Hilary Finucane, John Novembre and Raymond Walters for useful discussions and comments that significantly improved the manuscript. The study was supported by National Institute of Health grants HG009088, MH101244 (M.S., R.M., B.N. and S.S.) and GM127131 (S.S.). D.R. was supported by National Institutes of Health grant GM100233 and HG006399, an Allen Discovery Center of the Paul Allen Foundation, and the Howard Hughes Medical Institute.

558 **Competing interests**
559
560 The authors declare no competing interests.
561
562
563 **References**
564

[1] Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al.
Common SNPs explain a large proportion of the heritability for human height.
Nat Genet 2010;42:565–9. doi:10.1038/ng.608.
[2] Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. Cell 2017;169:1177–86.
doi:10.1016/j.cell.2017.05.038.
[3] Zeng J, de Vlaming R, Wu Y, Robinson MR, Lloyd-Jones LR, Yengo L, et al.
Signatures of negative selection in the genetic architecture of human complex
traits. Nat Genet 2018;50:746–53. doi:10.1038/s41588-018-0101-4.
[4] Turchin MC, Chiang CWK, Palmer CD, Sankararaman S, Reich D, Hirschhorn
JN. Evidence of widespread selection on standing variation in Europe at
height-associated SNPs. Nat Genet 2012;44:1015–9. doi:10.1038/ng.2368.
[5] Berg JJ, Coop G. A Population Genetic Signal of Polygenic Adaptation. PLoS
Genet 2014;10:e1004412. doi:10.1371/journal.pgen.1004412.
[6] Mathieson I et al. Genome-wide patterns of selection in 230 ancient Eurasians.
Nature 2016;528:499–503. doi:10.1038/nature16152.Genome-wide.
[7] Robinson MR, Hemani G, Medina-Gomez C, Mezzavilla M, Esko T, Shakhbazov
K, et al. Population genetic differentiation of height and body mass index
across Europe. Nat Genet 2015;47:1357–61. doi:10.1038/ng.3401.
[8] Berg JJ, Zhang X, Coop G. Polygenic Adaptation has Impacted Multiple
Anthropometric Traits. bioRxiv 2017. doi:10.1101/167551.
[9] Racimo F, Berg JJ, Pickrell JK. Detecting polygenic adaptation in admixture
graphs. Genetics 2018;208:1565–84. doi:10.1534/genetics.117.300489.
[10] Guo J, Wu Y, Zhu Z, Zheng Z, Trzaskowski M, Zeng J, et al. Global genetic
differentiation of complex traits shaped by natural selection in humans. Nat
Commun 2018;9:1–9. doi:10.1038/s41467-018-04191-y.
[11] Simonti C, Stein J, Thompson P, Fisher SE, Dan J. Polygenic selection underlies
evolution of human brain structure and behavioral traits. BioRxiv 2017:1–51.
[12] Field Y, Boyle EA, Telis N, Gao Z, Gaulton KJ, Golan D, et al. Detection of human
adaptation during the past 2,000 years. Science 2016;354:760–4.
doi:10.1126/science.aag0776.
[13] Allen HL, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, et al.
Hundreds of variants clustered in genomic loci and biological pathways affect
human height. Nature 2010;467:832–8. doi:10.1038/nature09410.
[14] Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, et al. Defining
the role of common variation in the genomic and biological architecture of
adult human height. Nat Genet 2014;46:1173–86. doi:10.1038/ng.3097.
[15] Churchhouse C, Neale BM, Abbott L, Anttila V, Aragam K, Baumann A, et al.
Rapid gwas of thousands of phenotypes for 337,000 samples in the uk

biobank.   2017.
https://sites.google.com/broadinstitute.org/ukbbgwasresults/home?authuse
r=0 (accessed February 11, 2018).
[16] Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al.
Human Demographic History Impacts Genetic Risk Prediction across Diverse
Populations.   Am   J   Hum   Genet   2017;100:635–49.
doi:10.1016/j.ajhg.2017.03.004.
[17] Field Y, Boyle E, Telis N, Gao Z, Gaulton K, Golan D, et al. Data from: Detection
of human adaptation during the past 2000 years. Dyrad Digital Repository.
n.d. doi:https://doi.org/10.5061/dryad.kd58f.
[18] Novembre J, Barton NH. Tread Lightly Interpreting Polygenic Tests of
Selection. Genetics 2018;208:1351–5. doi:10.1534/GENETICS.118.300786.
[19] Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS
Genet 2006;2:2074–93. doi:10.1371/journal.pgen.0020190.
[20] Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D.
Principal components analysis corrects for stratification in genome-wide
association studies. Nat Genet 2006;38:904–9. doi:10.1038/ng1847.
[21] Sohail M, Vakhrusheva OA, Sul JH, Pulit SL, Francioli LC, Van Den Berg LH, et
al. Negative selection in humans and fruit flies involves synergistic epistasis.
Science 2017;356:539–42. doi:10.1126/science.aah5238.
[22] Nelson MR, Bryc K, King KS, Indap A, Boyko AR, Novembre J, et al. The
Population Reference Sample, POPRES: A Resource for Population, Disease,
and Pharmacological Genetics Research. Am J Hum Genet 2008;83:347–58.
doi:10.1016/j.ajhg.2008.08.005.
[23] Haworth S, Mitchell R, Corbin L, Wade KH, Dudding T, Budu-Aggrey A, et al.
Common genetic variants and health outcomes appear geographically
structured in the UK Biobank sample: Old concerns returning and their
implications. bioRxiv 2018:294876. doi:10.1101/294876.
[24] Loh P-R, Kichaev G, Gazal S, Schoech A, Price AL. Mixed model association for
biobank-scale data sets. bioRxiv 2017. doi:10.1101/194944.
[25] Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN, et al. Meta-
analysis of genome-wide association studies for height and body mass index
in ~700,000 individuals of European ancestry. bioRxiv 2018.
[26] Haak W, Lazaridis I, Patterson N, Rohland N, Mallick S, Llamas B, et al. Massive
migration from the steppe was a source for Indo-European languages in
Europe. Nature 2015;522:207–11. doi:10.1038/nature14317.
[27] Mathieson I, Alpaslan-Roodenberg S, Posth C, Szécsényi-Nagy A, Rohland N,
Mallick S, et al. The genomic history of southeastern Europe. Nature 2018.
doi:10.1038/nature25778.
[28] Auton A, Abecasis GR, Altshuler DM, Durbin RM, Bentley DR, Chakravarti A, et
al. A global reference for human genetic variation. Nature 2015;526:68–74.
doi:10.1038/nature15393.
[29] Berg JJ, Zhang X, Coop G. Polygenic Adaptation has Impacted Multiple
Anthropometric Traits. bioRxiv 2017:1–38. doi:10.1101/167551.
[30] Berisa T, Pickrell JK. Approximately independent linkage disequilibrium
blocks   in   human   populations.   Bioinformatics   2015;32:283–5.

650   doi:10.1093/bioinformatics/btv546.

651 [31] Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. Second-
652   generation PLINK: Rising to the challenge of larger and richer datasets.
653   Gigascience 2015;4:1–16. doi:10.1186/s13742-015-0047-8.

654 [32] Purcell S, Chang C. PLINK 1.9. Gigascience 2015. www.cog-
655   genomics.org/plink/1.9/ (accessed May 1, 2018).

656 [33] Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y,
657   et al. Reduced signal for polygenic adaptation of height in UK Biobank. bioRxiv
658   2018. doi:10.1101/354951.

Supplementary Note

Characterization of stratification effects in GIANT and UK Biobank

To better understand how stratification influences the differences observed between GIANT and UK Biobank, we grouped SNPs by their P value in GIANT and by their P value in UK Biobank (**Supplementary Figure S1**). First, we observe that SNPs with low GIANT P values, but not SNPs with low UK Biobank P values, show greater differences in estimated effect size (**Supplementary Figure S1a**). However, the relative difference in beta values decreases for lower P values, and the correlation among betas approaches one at the most significant SNPs (**Supplementary Figure S1b,c**). In GIANT, more significant SNPs exhibit a greater correlation between effect estimates and GBR-TSI allele frequency differences, while this is not observed in the UK Biobank (**Supplementary Figure S1d**). Consequently, the difference in UK Biobank and GIANT effect size estimates is more correlated to GBR-TSI allele frequency differences at more significant SNPs (**Supplementary Figure S1e**). This suggests that while stratification effects are larger at more significant SNPs, the magnitude of stratification-independent effects is even larger, which may be why polygenic score results converge when using only the most significant SNPs.

Next, we investigated how P value inflation as measured by λ_{GC} is influenced by stratification, by grouping SNPs into deciles based on their GBR-TSI allele frequency difference (**Supplementary Figure S12**). To guard against the effect of observing lower P values at more differentiated SNPs simply because those SNPs are more common on average, we restrict this analysis to SNPs with mean MAF > 20%. We find that λ_{GC} is not much increased for SNPs that are more differentiated between populations. However, in the presence of stratification, there is a large difference between λ_{GC} of height increasing alleles and λ_{GC} of height decreasing alleles (**Supplementary Figure S12b**). Similarly, there are large effects on the frequency with which a SNP is estimated to be height increasing or height decreasing. In GIANT, SNPs in the highest decile of GBR-TSI allele frequency differences are 52% more often estimated to be height increasing than height decreasing, while these rates are close to even in the UK Biobank (**Supplementary Figure S12a**).

Supplementary figures

Figure S1. Beta concordance between GIANT and UK Biobank by P value bin.

P−value bin (LD−pruned, 1000 SNPs per bin, high to low)

Figure S1 (Continued). Beta concordance between GIANT and UK Biobank by P value bin.

SNPs intersecting between GIANT and UKB were LD-pruned (using PLINK 1.9 with parameters r^2 = 0.1, window size = 1 Mb, step size 5) and grouped into P value bins of 500 SNPs each, for P values from GIANT (left) and UKB (right). Color is based on the smallest P value in each bin.

(a) Absolute beta difference. As expected, absolute beta and thus the absolute beta difference increases across P value bins.

(b) Absolute beta difference, scaled by the sum of absolute betas. The relative difference of absolute betas decreases for lower P values.

(c) Pearson correlation among betas approaches one for the lowest P values.

(d) Correlation between beta (left GIANT, right UK Biobank) and GBR-TSI allele frequency difference.

(e) Correlation between the GIANT - UK Biobank beta difference and GBR-TSI allele frequency difference.

a. clumped, $r^2 < 0.1$, 1Mb, P < 0.01

b. clumped, $r^2 < 0.1$, 1Mb, P < 5 x 10^{-8}

GIANT   UK Biobank

Figure S2. **Polygenic scores using height-associated SNPs from GIANT- and UK Biobank-based GWA studies for clumped SNPs in present-day and ancient Europeans.** Scores are shown, centered by the average score across modern and ancient populations respectively and standardized by the square root of the additive variance. SNPs were LD-pruned with plink's clumping procedure for parameters: (a) $r^2 < 0.1$, 1Mb, P < 0.01 (81,941 SNPs in UKB, 22,561 SNPs in GIANT), and (b) $r^2 < 0.1$, 1Mb, P < 5x10^{-8} (4478 SNPs in UKB, 1442 SNPs in GIANT). Modern populations are shown from Northern Europe (CEU, GBR) and Southern Europe (IBS, TSI) from the 1000 genomes project. Ancient populations are shown in three meta-populations, hunter-gatherers (HG), early farmers (EF) and Steppe Ancestry (SP). Error bars are drawn at 95% credible intervals.

a. clumped, $r^2 < 0.1$, 1Mb, P < 0.01

b. clumped, $r^2 < 0.1$, 1Mb, P < 5×10^{-8}

729
Figure S3. Polygenic height scores in 1000 genomes European populations using
clumped SNPs and effect sizes from different summary statistics.
Polygenic scores in modern European populations are shown using SNPs LD-pruned with
PLINK's clumping procedure with parameters: (a) $r^2 < 0.1$, 1Mb, P < 0.01, and (b) $r^2 < 0.1$,
1Mb, P < 5×10^{-8}. Scores are centered by the average score across populations and
standardized by the square root of the additive variance. Modern populations are shown
from Northern Europe (CEU, GBR) and Southern Europe (IBS, TSI) from the 1000 Genomes
Project. Error bars are drawn at 95% credible intervals.
738

739
Figure S4. Polygenic height scores in 1000 Genomes Project European populations
using ~1700 independent SNPs and effect sizes from different summary statistics.
Polygenic scores in modern European populations are shown using SNPs LD-pruned by
picking the SNP with the lowest P value in each of ~1700 LD-independent blocks genome-
wide. Scores are centered by the average score across populations and standardized by the
square root of the additive variance. Modern populations are shown from Northern Europe
(CEU, GBR) and Southern Europe (IBS, TSI) from the 1000 Genomes Project. Error bars are
drawn at 95% credible intervals.
748
749

750
Figure S5. Polygenic height scores in ancient populations using ~1700 independent
SNPs and effect sizes from different summary statistics.
Polygenic scores in ancient meta-populations are shown using SNPs LD-pruned by picking
the SNP with the lowest P value in each of ~1700 LD-independent blocks genome-wide.
Scores are centered by the average score across populations and standardized by the
square root of the additive variance. Error bars are drawn at 95% credible intervals. .
Ancient populations are shown in three meta-populations (HG = Hunter-Gatherer (n=162
individuals), EF = Early Farmer (n=485 individuals), and SP = Steppe Ancestry (n=465
individuals)). The y-axis is truncated at (-1.5, 1.5) for all panels – this omits two points in
the NG2015 sibs panel: HG [3.86 (CI: 3.60, 4.12)], EF [-2.18(CI: -2.34, -2.02)].
761

762
763
Figure S6. **Polygenic height scores in ancient and global modern populations using**
three different GWAS.
All scores are centered by the average score across all populations ($\mu_{GIANT} = 0.645, \mu_{LOH} =$
$-0.219, \mu_{NEALELAB} = -0.259$) and standardized by the square root of the additive variance.
Error bars are drawn at 95% credible intervals. Modern populations are shown from
Northern Europe (CEU, GBR), Southern Europe (IBS, TSI), South Asia (PJL, BEB), East Asia
(CHB, JPT) and Africa (YRI, LWK). Ancient populations are shown in three meta-
populations, hunter-gatherers (HG), early farmers (EF) and Steppe Ancestry (SP). Pseudo-
haploid genotype calls were made for modern populations before computing polygenic
scores to allow fair comparison with ancient DNA. SNPs were LD-pruned by picking the SNP
with the lowest P value in each of ~1700 LD-independent blocks genome-wide.
775

776
Figure S7. tSDS for height-increasing alleles using effect sizes from different summary
statistics.
SNPs were ordered by GWAS P value and grouped into bins of 1000 SNPs each. The mean
tSDS score within each P value bin is shown on the y-axis. In contrast to Figure 3, where
Spearman correlation coefficients and Jackknife standard errors were computed, here we
show the regression slope and P value, which were computed on the un-binned data. The
gray line indicates the null-expectation, and the colored lines are the linear regression fit.
The lowest P value bin in panel 5 with a y-axis value of 1.5 has been omitted.

785
Figure S8. tSDS for LD-pruned height-increasing alleles using effect sizes from different summary statistics.

Binning SNPs by P value can lead to spurious results at the low P value bins when SNPs are in LD (**Figure S15**). Here, LD-pruned SNPs were ordered by GWAS P value and grouped into bins of 100 SNPs each. The mean tSDS score within each P value bin is shown on the y-axis. In contrast to **Figure 3**, where Spearman correlation coefficients and Jackknife standard errors were computed, here we show the regression slope and P value, which were computed on the un-binned data. The gray line indicates the null-expectation, and the colored lines are the linear regression fit.

797
Figure S9. Height (cm) in the UKB as a function of GBR-TSI score.
We computed the relative number of GBR to TSI related alleles in each sample by multiplying the allele frequency difference by the number of alternative alleles in each sample in the UKB (GBR-TSI score). Vertical lines indicate 5th and 95th percentile of among-white British samples, showing that there is a significant negative relationship between the GBR-TSI allele sharing score and height (in cm). Among all other broadly European samples, this relationship is significantly positive across the whole range, but again significantly negative in the white British range. This can explain why stratification effects go in opposite directions in a UKB height GWAS of white British samples and a UKB height GWAS of all samples. Here, other European samples were defined as those that lie within the mean +/- 24 standard deviations along the first six principal components.

811
Figure S10. Pearson Correlation coefficients of PC loadings and height beta coefficients for different summary statistics.

PCs were computed in all 1000 genomes phase 1 samples. Colors indicate the correlation of each PC loading with the allele frequency difference between GBR and TSI, a proxy for the European North-South genetic differentiation. PC 4 and 11 are most highly correlated with the GBR - TSI allele frequency difference. Error bars indicate 95% confidence interval of the correlation coefficient, assuming 60,000 independent genetic markers. We confirmed that the resulting errors are similar to block jackknife standard errors. Open circles indicate correlations significant at alpha = 0.05, stars indicate correlations significant after Bonferroni correction in 20 PCs (P < 0.0025).

823
Figure S11. Heat map of mean beta coefficients for different summary statistics.
All SNPs are binned by GBR and TSI minor allele frequency. Only bins with at least 300 SNPs
are shown. Panel 7 (as well as 2, 3 and 4) shows stratification effects in opposite direction
to those in GIANT. **Figure S9** illustrates how these opposite-direction stratification effects
can arise.

830
831
Figure S12. Effect of GBR-TSI allele frequency difference on beta estimates and P
values.
SNPs with MAF > 0.2 (based on mean between TSI and GBR) were grouped into GBR-TSI
allele frequency difference deciles, with the first decile representing SNPs less common in
GBR and the last decile representing SNPs more common in GBR.
 (a) Fraction of height-increasing (yellow dots) vs. height-decreasing SNPs (purple dots) in
each decile. In GIANT, 59% of SNPs in the highest decile are estimated to be height-
increasing, and 41% are estimated to be height-decreasing. In the UK Biobank, this ratio is
close to 50-50.
(b) Lambda-GC in each decile for height-increasing (yellow dots) vs. height-decreasing SNPs
(purple dots). In GIANT, the median P value of SNPs in the highest decile is 2.78 for SNPs
estimated to be height-increasing and 1.83 for SNPs estimated to be height-decreasing (a
difference of 52%). In the UK Biobank, the median P value of SNPs in the highest decile is
2.65 for SNPs estimated to be height-increasing and 2.89 for SNPs estimated to be height-
decreasing (a difference of only 9%, going in the opposite direction).
847
848

849
850
851
852
853 **Figure S13. Allele frequency difference for height-increasing alleles using different**
854 **summary statistics**.
855 SNPs were ordered by GWAS P value and grouped into bins of 1000 SNPs each. The gray
856 line indicates the null-expectation, and the colored lines are the linear regression fit. The
857 lowest P value bin in panel 5 with a y-axis value of 0.06 has been omitted.
858

859
**Figure S14. Allele frequency difference for LD-pruned height-increasing alleles using
different summary statistics.**
Binning SNPs by P value can lead to spurious results at the low P value bins when SNPs are
in LD (**Figure S15**). Here, LD-pruned SNPs were ordered by GWAS P value and grouped into
bins of 100 SNPs each. The gray line indicates the null-expectation, and the colored lines are
the linear regression fit.

Figure S15. Number of independent regions per GWAS P value bin in the UK Biobank.
SDS results in Field et al. as well as in **Figure 3** in this article are visualized by grouping
non-independent SNPs into bins according to their P value. This may lead to unpredictable
patterns at the low end of the P value distribution, because the lowest P value bins do not
represent independent signals. This is demonstrated here, by grouping all UKB SNPs into
bins of 1000 SNPs each, as in the SDS plots in **Figure 1b** and **Figure 3**. Left: The number of
independent SNPs per P value bin is much lower at lower P values. Right: Neighboring P
value bins share a large fraction of 1Mb regions at lower P values. This demonstrates that
the lowest P value bins do not represent independent signals if SNPs are not LD-pruned and
can exhibit patterns that are dominated by one or a few LD-regions.

Figure S16. Polygenic height scores in POPRES for different summary statistics.
Standardized polygenic height score from diverse summary statistics for 19 POPRES populations with at least 10 samples per population, ordered by latitude (see **Table S3**). Confidence intervals and clumping procedure are the same as in (a). The gray line is the linear regression fit to the mean polygenic height score per population. The numbers on each plot show the Q_x P value, the latitude covariance P value and the number of SNPs respectively for each summary statistic. Each column shows a different selection of SNPs. clump all: clumped SNPs with no P value threshold; clump 0.01: clumped SNPs with P < 0.01 in UKB and the same number of SNPs in other summary statistics (same as **Figure 4**); clumpwindow 1.5M: genome was split into blocks of 1.5 Mb, lowest P-value SNP was picked in each bin, similar to the 1700 blocks; ldwindow 1.5Mb: genome was split into blocks of 1.5 Mb, random SNP was picked in each bin; UKB sig: LD-pruned SNPs with P < 5x10^{-8} in UKB.

893
Figure S17. Test statistics for Q_x (left) and latitude correlation (right) in the POPRES
dataset for different summary statistics.
The numbers indicate P values and the number of SNPs, and numbers in bold highlight
nominal significance ($p < 0.05$).

898
899
900

901
Figure S18. Spearman correlations between polygenic height scores in the POPRES
dataset computed from different summary statistics.
Spearman correlation coefficients of mean population polygenic score ranking for all pairs
of summary statistics at different SNP selections. Polygenic scores from independent SNPs
which are genome-wide significant in UKB lead to more consistent rankings than PRS from
other sets of SNPs, despite having lower prediction power. Each column shows a different
selection of SNPs. clump all: clumped SNPs with no P value threshold; clump 0.01: clumped
SNPs with P < 0.01 in UKB and the same number of SNPs in other summary statistics (same
as **Figure 4**); clumpwindow 1.5M: genome was split into blocks of 1.5 Mb, lowest P-value
SNP was picked in each bin, similar to the 1700 blocks; ldwindow 1.5Mb: genome was split
into blocks of 1.5 Mb, random SNP was picked in each bin; UKB sig: LD-pruned SNPs with P
< 5x10^{-8} in UKB.
914

915
916 **Figure S19. P value calibration in the POPRES dataset for Q_x and latitude covariance**
917 **tests.** Random sets of around 1700 independent markers were drawn in 100 repetitions for
918 four summary statistics and Q_x and latitude P values were computed. In UK Biobank sibling
919 estimates this resulted in a uniform P value distribution (non-significant Kolmogorov–
920 Smirnov test), while an inflation was observed for UK Biobank GWAS summary statistics.
921

Supplementary tables

name	sample size	number of SNPs (before intersecting)	type	comments	used in figures	PMID	URL
GIANT	253288	2550858	GWAS (OLS, meta-analysis)		1, 2, 3, 4	25282103	https://portals.broadinstitute.org/collaboration/giant/images/0/01/GIANT_HEIGHT_Wood_et_al_2014_publicrelease_HapMapCeuFreq.txt.gz
UKB Neale (UKB)	336474	10894596	GWAS (OLS)		1, 2, 3, 4		https://www.dropbox.com/s/sbfgb6qd5i4cxku/50.assoc.tsv.gz
UKB Neale new	360388	13791467	GWAS (OLS)	soon be available on http://www.nealelab.is/	supp only		https://storage.googleapis.com/ukbb-robert/height_ukb_giant/robert1/50.imputed_v3.results.both_sexes.tsv.gz
UKB Loh	458303	12007535	BOLT-LMM		supp only	25642633	https://data.broadinstitute.org/alkesgroup/UKBB/body_HEIGHTz.sumstats.gz
UKB all no PCs	406825	729339	GWAS (OLS)	genotyped SNPs	3, 4		
UKB all 10 PCs	406825	729339	GWAS (OLS)	genotyped SNPs	supp only		
UKB WB no PCs	337208	729339	GWAS (OLS)	genotyped SNPs	3		
UKB WB 10 PCs	337208	729339	GWAS (OLS)	genotyped SNPs	supp only		
NG2015 sibs	17500	1006257	within-family (QFAM)		3	26366552	http://cnsgenomics.com/data/robinson_et_al_2015_ng/withinfam_summary_ht_bmi_release_March2016.tar.gz
UKB sibs all	20166	729339	within-family (QFAM)	genotyped SNPs	supp only		
UKB sibs WB	17358	729339	within-family (QFAM)	genotyped SNPs	3, 4		

924
925 **Table S1. Description of 11 GWAS summary statistics.**
926

927
928

Population label	Population description	Meta-population	N
GBR	British in England & Scotland	Northern Europe	92
CEU	Utah residents with Northern & Western European Ancestry	Northern Europe	99
TSI	Toscani in Italia	Southern Europe	108
IBS	Iberian population in Spain	Southern Europe	107
PJL	Punjabi from Lahore, Pakistan	South Asia	96
BEB	Bengali from Bangladesh	South Asia	86
YRI	Yoruba in Ibadan, Nigeria	Africa	109
LWK	Luhya in Webuye, Kenya	Africa	101
JPT	Japanese in Tokyo, Japan	East Asia	104
CHB	Han Chinese in Beijing, China	East Asia	103
EF	Ancient	Early farmer	485
HG	Ancient	Hunter-gatherer	162
STP	Ancient	Steppe ancestry	465

929
930 **Table S2. Table of ancient and 1000 genomes modern populations used with sample**
931 **sizes.**
932

933
934

Population	N	Latitude	Comments
Austria	14	47.5	
Belgium	43	50.5	
Czech Republic	11	49.8	
France	92	46.2	
Germany	75	51.1	
Hungary	19	47.1	
Ireland	62	53.1	
Italy	225	41.8	
Netherlands	17	52.1	
Poland	22	51.9	
Portugal	135	39.3	
Romania	14	45.9	
Spain	137	40.4	
Sweden	11	60.1	
Swiss-French	760	46.5	Lausanne
Swiss-German	84	47.3	Zurich
Switzerland	168	46.8	
United Kingdom	390	55.3	
Yugoslavia	44	43.9	

935 **Table S3. Table of POPRES populations used with sample sizes and latitude.**
936

Name	h2	h2 SE	lambda GC	mean chisq	intercept	intercept SE	ratio	n
GIANT	0.31	0.01	2.00	2.92	1.28	0.02	0.15	253288
UKB Neale (UKB)	0.40	0.02	2.39	4.32	1.40	0.03	0.12	336474
UKB Neale new	0.42	0.02	2.48	4.69	1.43	0.03	0.12	360388
UKB Loh	0.60	0.03	3.31	7.66	1.75	0.04	0.11	458303
UKB all no PCs	13.76	0.22	55.97	65.24	9.42	0.08	0.13	406825
UKB all 10 PCs	0.48	0.02	2.37	4.69	1.39	0.04	0.11	406825
UKB WB no PCs	0.54	0.03	2.67	4.66	1.66	0.04	0.18	337208
UKB WB 10 PCs	0.51	0.03	2.17	4.19	1.30	0.04	0.09	337208
NG2015 sibs	0.49	0.03	1.07	1.08	0.90	0.01	-1.21	17500
UKB sibs all	0.45	0.05	0.93	1.08	0.93	0.01	-0.85	20166
UKB sibs WB	0.47	0.06	0.93	1.06	0.93	0.01	-1.18	17358

Table S4. LD Score regression estimates for 11 different summary statistics.
LD score regression can be used to detect residual stratification effects in summary
statistics, and so we tested whether LDSC confirms our hypothesis of residual stratification.
We detect a greatly inflated intercept estimate of 9.42 in UKB all no PCs, but only a
moderately increased intercept value in GIANT and an intercept less than one in NG2015
sibs. The relatively small GIANT intercept can be explained by cohort-wise lambda-GC
correction, while the low intercept in NG2015 sibs is possibly caused by the adaptive
permutation procedure which does not compute precise p-values for non-significant
associations. In both cases LDSC cannot be expected to pick up stratification effects, since
the generation of summary statistics is not in line with the LDSC model.
947

	GIANT	UKB Neale (UKB)	UKB Neale new	UKB Loh	UKB all no PCs	UKB all 10 PCs	UKB WB no PCs	UKB WB 10 PCs	NG2015 sibs	UKB sibs all	UKB sibs WB
GIANT	1.000										
UKB Neale (UKB)	0.660	1.000									
UKB Neale new	0.670	0.982	1.000								
UKB Loh	0.691	0.912	0.924	1.000							
UKB all no PCs	0.332	0.276	0.274	0.251	1.000						
UKB all 10 PCs	0.673	0.973	0.971	0.916	0.290	1.000					
UKB WB no PCs	0.635	0.952	0.933	0.865	0.270	0.929	1.000				
UKB WB 10 PCs	0.660	0.998	0.981	0.911	0.276	0.975	0.953	1.000			
NG2015 sibs	0.368	0.261	0.265	0.276	0.151	0.271	0.238	0.261	1.000		
UKB sibs all	0.264	0.325	0.326	0.351	0.095	0.328	0.307	0.324	0.114	1.000	
UKB sibs WB	0.250	0.312	0.312	0.336	0.088	0.311	0.296	0.311	0.108	0.928	1.000

948 **Table S5. Correlation of beta estimates at all 86,153 shared SNPs.**
949

	GIANT	UKB Neale (UKB)	UKB Neale new	UKB Loh	UKB all no PCs	UKB all 10 PCs	UKB WB no PCs	UKB WB 10 PCs	NG2015 sibs	UKB sibs all	UKB sibs WB
GIANT	1.000										
UKB Neale (UKB)	0.975	1.000									
UKB Neale new	0.976	0.999	1.000								
UKB Loh	0.972	0.986	0.986	1.000							
UKB all no PCs	0.798	0.789	0.788	0.785	1.000						
UKB all 10 PCs	0.976	0.998	0.998	0.988	0.791	1.000					
UKB WB no PCs	0.971	0.996	0.995	0.984	0.788	0.995	1.000				
UKB WB 10 PCs	0.975	1.000	0.999	0.986	0.789	0.999	0.997	1.000			
NG2015 sibs	0.847	0.828	0.829	0.840	0.675	0.829	0.822	0.828	1.000		
UKB sibs all	0.865	0.877	0.875	0.874	0.706	0.878	0.872	0.877	0.727	1.000	
UKB sibs WB	0.860	0.872	0.870	0.866	0.700	0.872	0.867	0.872	0.712	0.980	1.000

950 **Table S6. Correlation of beta estimates at 2,251 shared SNPs which are significant in**
951 **the UK Biobank.**
952