Article

Mutators as drivers of adaptation in *Streptococcus* and a risk factor for host jumps and vaccine

escape

Running Title: Mutators as drivers of pathogen evolution

Oleksandra Silayeva, Jan Engelstaedter, and Andrew C Barnes[*]

The University of Queensland, School of Biological Sciences, St Lucia Campus, Brisbane, Queensland

4072, Australia

Key words: mutators; DNA repair; pathogen evolution; immune escape; vaccination; Streptococcus;

aquaculture; host jumps, zoonoses

*Correspondence:

a.barnes@uq.edu.au

Abstract

Heritable hypermutable strains deficient in DNA repair genes (mutators) facilitate microbial adaptation as they may rapidly generate beneficial mutations. Mutators deficient in mismatch (MMR) and oxidised guanine (OG) repair are abundant in clinical samples and show increased adaptive potential in experimental infection models but their role in pathoadaptation is poorly understood. Here we investigate the role of mutators in epidemiology and evolution of the broad host pathogen, *Streptococcus iniae*, employing 80 strains isolated globally over 40 years. We determine phylogenetic relationship among *S. iniae* using 10,267 non-recombinant core genome single nucleotide polymorphisms (SNPs), estimate their mutation rate by fluctuation analysis, and detect variation in major MMR (*mutS, mutL, dnaN, recD2, rnhC*) and OG (*mutY, mutM, mutX*) genes. *S. iniae* mutation rate phenotype and genotype are strongly associated with phylogenetic diversification and variation in major streptococcal virulence determinants (capsular polysaccharide, hemolysin, cell chain length, resistance to oxidation, and biofilm formation). Furthermore, profound changes in virulence determinants observed in mammalian isolates (atypical host) and vaccine-escape isolates found in bone (atypical tissue) of vaccinated barramundi are linked to multiple MMR and OG variants and unique mutation rates. This implies that adaptation to new host taxa, new host tissue, and to immunity of a vaccinated host is promoted by mutator strains. Our findings support the importance of mutation rate dynamics in evolution of pathogenic bacteria, in particular adaptation to a drastically different immunological setting that occurs during host jump and vaccine escape events.

Importance

Host immune response is a powerful selective pressure that drives diversification of pathogenic microorganisms and, ultimately, evolution of new strains. Major adaptive events in pathogen evolution, such as transmission to a new host species or infection of vaccinated hosts, require adaptation to a drastically different immune landscape. Such adaptation may be favoured by hypermutable strains (or mutators) that are defective in normal DNA repair and consequently capable of generating multiple potentially beneficial and compensatory mutations. This permits rapid adjustment of virulence and antigenicity in a new immunological setting. Here we show that mutators, through mutations in DNA repair genes and corresponding shifts in mutation rate, are associated with major diversification events and virulence evolution in the broad host-range pathogen *Streptococcus iniae*. We show that mutators underpin infection of vaccinated hosts, transmission to new host species and the evolution of new strains.

Introduction

Mutation rates in microbial populations are a trade-off between maintenance of genetic integrity and evolvability (1, 2). High fidelity of DNA repair is advantageous in adapted populations since most mutations lead to deviation from the evolved phenotype (2, 3). Conversely, lower fidelity of DNA repair can be advantageous under stress since increased mutation supply promotes diversity and may accelerate adaptation (2, 4). Bacteria can efficiently fluctuate between high and low mutation regimes: While extremely low mutation rates are maintained in favourable conditions (5), elevation of mutation rates known as stress-induced mutagenesis (SIM) occurs in adapting populations (6, 7). One aspect of SIM is temporary hypermutability associated with changes in expression of multiple genes including error-prone DNA polymerases, recombination-preventing enzymes and enzymes involved in movement of mobile elements (8). For example transient hypermutability during RecA-mediated general stress response, SOS, is associated with up-regulation of error-prone polymerases that can by-pass DNA lesions during replication (8). However, increased fitness during SOS response stems from immediate growth advantage rather than from long-term evolution of stress resistance (9-11). Another aspect of SIM is the evolution of heritable mutator phenotypes, also referred to as constitutive, permanent, or true mutators, that contain mutator alleles and replicate over multiple generations with a high amount of error (6, 7). Heritably hypermutable strains, which periodically arise within a population via mutations in DNA repair genes (12, 13), are selected under stress indirectly via 'hitchhiking' of mutator alleles with stress-resistance mutations (14, 15), or by undefined pleiotropic effects conferred by inactivation of DNA repair (16, 17). There is much stronger theoretical and experimental evidence demonstrating that heritable mutators facilitate adaptive evolution (17). However, even heritable mutators are transient phenomena in natural populations as non-mutator variants are selected within already adapted mutator clones (18, 19). Restoration of mutation rate can occur via copy-number variation leading to full reversion of the mutator allele (20), disruption of the linkage between mutator alleles and beneficial mutations via recombination (21), anti-mutator compensatory mutations (4, 19), or prophage excision when its integration confers increase in mutation rate (22).

Bacterial mutators are abundant among clinical isolates, especially in chronic cases (23, 24). Multiple studies have attributed this to antibiotic selection (25-27), since mutators are a well-established risk factor for antimicrobial resistance (28, 29). However, the association between frequency of mutators in diagnostic samples and antibiotic treatments is weak (30, 31), and isogenic knockout mutator strains have increased ability to colonize mice in challenge models devoid of antibiotic exposure (32-34). Thus, the prevalence of mutators in clinical isolates most likely results from the selective process during

adaptation to the host (30, 35-37), where immunity acts as the ultimate selective pressure shaping the
diversity of pathogenic strains (38).

The 'immunological niche' concept states that, due to the diversity and complexity of immune
responses, every individual host represents a unique habitat that requires adaptation (39). As a
pathogen needs to adapt to every new host, the host population represents an extremely
heterogeneous environment where adaptive processes are ongoing and persistence of mutator alleles
may be favoured (40). Moreover, as a small number of infection units may enter a host, pathogen
adaptation is intrinsically associated with population bottlenecks during which the rate at which
beneficial variants arise might be crucial (41, 42). This supports the contention that bacterial evolution
is more likely to occur via multiple mutations rather than a single beneficial variant (43).

The fitness of a pathogenic strain in the host hinges on the complex, dynamic trait of virulence,
determined by a multitude of virulence factors and host responses (44). Optimal virulence is a trade-off
between acuteness and persistence with higher virulence facilitating extraction of resources and
transmission, but too much damage can kill the host and stop transmission altogether (45).
Additionally, virulence factors are often antigens so reduced virulence may facilitate immune evasion
(46). Consequently, both loss and gain of virulence can be advantageous and indicate adaptive shifts
towards acute or chronic pathogenesis respectively (47, 48). Exacerbation and attenuation of virulence
are repeatedly observed in both knockout and naturally occurring mutators (36, 49-53), thus mutator
strains/alleles may help pathogenic bacteria to evolve towards optimal virulence.

Most bacterial mutators are deficient in genes from mismatch (MMR) and oxidised guanine (OG) DNA
repair systems (23). MMR and OG genes are conserved across all domains of life (54), and single
nucleotide polymorphisms in their sequence may drastically alter mutation rate (55, 56). Nonetheless,
these genes are variable in major bacterial pathogens, implicating mutation rate dynamics in the
evolution of virulence (57, 58). MMR is coupled to replication and removes mispaired nucleotides as
well as short insertion/deletion loops, and prevents recombination between divergent sequences (59,
60). The MMR pathway in *Escherichia coli* is well-defined, in which MutS dimer protein interacts with
the replication processivity clamp DnaN (b-clamp), binds to the mismatch, recruits MutL, and the
MutS:MutL complex then activates the MutH endonuclease (54). MutH cuts the unmethylated nascent
strand at hemimethylated d(GATC) sites. MutL loads the UvrD helicase that unwinds the nicked DNA
containing the error, which is followed by excision by one of the single-strand nucleases (54). This MMR
organization, however, appears to be limited to *E. coli* and closely related Gammaproteobacteria (54,
61). In most bacteria and eukaryotes, MutH is absent and MutL has a conserved endonuclease site. dam
methylase (acting at GATC sites) is absent and the strand discrimination signal is unknown (54, 61). In

Bacillus subtilis, a model species for gram-positive bacteria, nicks created by ribonuclease *rnhC* during

removal of misincorporated ribonucleosides direct MutL endonuclease towards the newly synthethised

strand, and recD2 acts as a MMR helicase instead of uvrD (61). The OG system repairs oxidative DNA

damage throughout the cell cycle (62, 63). Among all DNA bases, guanine is most sensitive to reactive

oxygen species and its oxidised form, 8-oxo-dG or OG, is highly mutagenic as it preferentially pairs with

adenine which, if not corrected, leads to G:C to A:T transversion on a second round of replication (62,

63). OG system organization is conserved in most bacteria and eukaryotes and generally MutT

homologs (eg. MutX in *S. pneumoniae*) remove 8-oxo-dG from the nucleotide pool, MutM homologs

excise 8-oxo-dG incorporated into DNA, and MutY homologs remove adenine from OG:A mispairings

(62, 63).

We investigate the role of mutators and mutation rate dynamics in the epidemiology and evolution of

the broad host-range pathogen *Streptococcus iniae. S. iniae* is a highly adaptable species, evidenced by

its global distribution, ability to cross divergent host taxa boundaries and repeated re-infection of

previously immunised hosts during disease outbreaks (64, 65). Moreover, Streptococcal pathogens in

general are difficult candidates for immunisation in humans and animals due to high strain diversity

evidently driven by immunisation, yet there is little information on the role of mutators in this

evolutionary process (66, 67). We determine variation in mutation rate and identify mutations in MMR

(*mutS, mutL, dnaN, recD2, rnhC*) and OG (*mutY , mutM, mutX*) genes among 80 diverse strains of *S.*

iniae. We discover that both mutation rates and mutations in MMR and OG genes correlate with

phylogenetic diversification but are highly conserved within phylogenetic clades. To determine whether

variation in mutation rate phenotype and MMR/OG genotype is linked to adaptation to the host

immunity, we identify variation in major phenotypic traits that determine virulence in streptococci

(capsular polysaccharide, hemolysin, length of cell chains) and bacteria in general (resistance to reactive

oxygen species (ROS), and biofilm formation). The expression of these traits among *S. iniae* isolates is

highly consistent with phylogenetic affiliation and strongly associated with variation in MMR and OG

genes and mutation rate phenotype. This supports the contention that strain diversification driven by

adaptation to the host achieved via adjustment of virulence is facilitated by the mutator strains.

Furthermore, we find that most changes in virulence determinants and MMR and OG genes occur in

strains producing infection in unusual immunological landscapes such as mammals and bone tissue of

immunised fish, indicating that mutators may facilitate the reinfection of immunised animals and

transmission between divergent host species.

Results and Discussion

We investigated the link between mutation rate genotype and phenotype, and phylogenetic diversification among *Streptococcus iniae* strains isolated globally between 1976 and 2016 from different host taxa. Maximum likelihood phylogenetic analysis of 80 *S. iniae* isolates (Table 1) based on non-recombinant core genome SNPs derived from whole genome data resolved six major clades (A-F), one lineage with two strains (clade G), and three lineages with a single strain (Figs. 1,2). Clades C-E show a degree of geographic endemism while clades A, B, and F contain strains from diverse geographic regions. Principally, variation in MMR and GO genes occurs only between the lineages, and they are highly conserved within clades (Fig. 2, Table 2). In fact, sequences of the major MMR and OG genes are conserved even within phylogenetic groups not readily definable by geographic origin, time, or host species, and may be useful for Multilocus Sequence Typing (MLST) (68). Although even synonymous SNPs in any gene locus can affect mRNA and have pronounced phenotypic consequences (69), most SNPs in MMR and OG genes were identified in protein functional domains, and most amino acid substitutions were predicted to have a deleterious effect on protein function (Table 2). Furthermore, the number of MMR and OG variants correlates significantly with mutation rate (p = 0.0056) and with the number of atypical phenotypic traits contributing to virulence (p = 0.0331), as predicted by a Phylogenetic Generalised Least Squares (PGLS) regression model that accounts for autocorrelation occurring between closely related strains (70). In turn, mutation rate variation is highly consistent with phylogenetic affiliation: According to multiple pair-wise strain comparisons by Maximum Likelihood Ratio test (71), with probabilities corrected for multiple comparison using False Discovery Rate (72), mutation rate difference is insignificant within but significant between major phylogenetic lineages (Suppl. Table 1).

Although mutation rate variation between divergent strains is significant, the magnitude of the differences is modest (around 5-fold) (Figure 1, 2, Suppl. Table 2). This may be inherent to streptococci, since only one order of magnitude or lower differences in mutation rate distinguish mutator from non-mutator phenotypes in other streptococcal species (around 10^{-7} and 10^{-8} mutation rates respectively) (23, 73). This is in contrast to other common pathogens where 2-3 orders of magnitude differences are observed (23). On the other hand, small differences in mutation rate are consistent with transience of mutators in natural populations and restoration of low mutation rate after adaptation is gained. Evolutionary theory predicts that heritably hypermutable phenotypes are costly due to inevitable accumulation of deleterious changes thus an adapted mutator clone would evolve back towards a lower mutation rate (17, 18, 74). This may occur via compensatory, 'anti-mutator' mutations in DNA repair genes (4, 19), or via recombination (21). Our data favour the former mechanism for several reasons. Firstly, we used a maximum likelihood inference method to exclude areas of recombination

from our phylogenetic analyses (75), and the MMR and OG gene regions were not identified as

recombinant regions in this analysis. Secondly, we created a pangenome for the 80 strains in our

analysis and mapped on putative phage and prophage signatures (76), and insertion sequence elements

(77). None of the MMR or OG genes, which are all present within the core genome of *S. iniae*, occurred

in proximity to phage or ISE (Suppl. Fig. 1). Nevertheless, in laboratory *E. coli* populations,

recombination seems to be the dominant process by which alterations in MMR and OG genotype occur,

as determined by incongruence between phylogenetic trees constructed from the variant MMR/OG

gene sequences and those inferred from genome-wide data (21). We were unable to apply similar

analysis to our data set as the low number of substitutions in *S. iniae* MMR and OG genes (one or two

non-synonymous mutations per gene compared with 1-53% non-synonymous base substitutions in the

E. coli population (21)) provided insufficient support for robust phylogeny when used in isolation. The

present strains originate from complex natural ecosystems where recombination (and mutation) may

be more constrained by multifactorial selection pressures in contrast to a monoculture experiment.

Moreover, recombination rates in richly capsulated streptococci are lower than in those with reduced

capsule (78). Indeed, recombination rates within the core genome across the *S. iniae* strains in our

analysis were extremely low with mean $\mu_{r/m}$ 0.032 and only 5 from the 83 strains containing

recombination blocks within the core genome where all of the MMR and OG genes analysed are

located (Suppl. Table 5). Whilst we cannot rule out switch in MMR and OG type by recombination,

accrual of SNPs via random mutation is a more parsimonious explanation here.

To investigate whether variation in mutation rate phenotype and MMR/OG genotype is associated with

adaptation to the host, we identified variation in key virulence traits in streptococci (capsular

polysaccharide, hemolysin, length of cell chains) and bacteria in general (resistance to ROS, and biofilm

formation). Employing Phylogenetic Generalised Least Squares regression model accounting for

autocorrelation due to phylogeny, we observe significant correlation (*p* = 0.0331, PGLS) of the number

of MMR and OG variants with number of atypical phenotypes for traits contributing to virulence. This

suggests that fluctuations in mutation rate are linked to shifts in virulence, potentially facilitating major

adaptive events that lead to strain diversification (Fig. 2, Table 3). Moreover, infections in atypical hosts

(mammals) and tissue (bone) within apparently immune vaccinated hosts correlates strongly with

number of atypical virulence-associated phenotypes (p < 0.0001) and the number of variants MMR and

OG variants (p = 0.002). This implies that mutator strains facilitate multiple genetic changes leading to

profound shifts in virulence and antigenicity during adaptation to a drastically different immunological

setting.

Of the clades identified among *S. iniae*, clade A is apparently a dominant circulating clade primarily

infecting Perciform fish that has persisted globally for almost two decades (clade A contains strains

isolated from USA, Honduras, and Australia between 1999 to 2016). Mutation rates in all isolates (n=35)

among this ancestral lineage fall within a 1.5 - 2 x 10^{-8} range (Figs. 1, 2; Clade A), which is similar to the

base mutation rate estimated for other non-mutator gram positive bacteria (23, 73). All differences in

mutation rates within this clade are insignificant except three pairwise comparisons that verge on p

<0.05 (Suppl. Table 1). MMR and OG genes in these strains are identical, and only minor deviations

from wildtype traits contributing to virulence were observed (absence of capsule production and longer

cell chains in QMA0158, 216) (Figs. 1-3, Table 1). All other lineages (with the exceptions of clade C1 and

strains QMA0445-46 discussed below) have significantly different mutation rates to clade A, contain

unique and often multiple SNPs in *mut* genes, and show peculiar phenotypic variants related to

virulence (Figs. 1-3, Tables 1,2, Suppl. Table 1). Although phylogenetic relatedness between lineages is

fully resolved (Fig. 2), they originate almost simultaneously and nested lineages evolve independently

sharing little phylogenetic history (Figs. 1,2). Lack of correlation between branch length and sampling

date derived by root-to-tip regression analysis with time is evidence of non-neutral evolution amongst

this collection of *S. iniae* isolates (R^2=0.0936, Correlation coefficient 0.306, best-fit root; Suppl. Fig. 2,

(79)). Deviation from a neutral model is supported by highly skewed branching, with some branches

comprising single isolates whilst others comprise many epidemiologically unrelated isolates (Figure 1,

2). This is suggestive of frequent strong selection, presumably imposed by heterogeneity of the immune

landscape encountered during transfer between host individuals, and resembles genealogical tree

topology derived from viral evolution over similar timespans (80, 81) although we acknowledge tree

topology may also be affected by sampling bias.

Clade B contains QMA0084, a strain isolated from a black flying fox, *Pteropus alecto*, in Western

Australia (WA) and two strains isolated from ornamental fish (clown loach, *Chromobotia macracanthus*)

in USA. These isolates share a valine to isoleucine substitution in MutL predicted as neutral (PROVEAN

score -0.271), a synonymous SNP in *recD2*, and a 56 bp deletion affecting the predicted promoter of

mutY (Table 2). The latter mutation encompasses the -35 box, two binding sites of the RpoD17 general

transcription factor, and the 1^{st} bp of the -10 box, and was predicted to abolish the original promoter

(Suppl. Fig. 3). However, *mutY* might still be transcribed at some level since two other potential

promoters were identified in a nearby sequence (Suppl. Fig. 3). It appears that altered *mutY* expression

is also affected by the rest of the genetic background as isolates exhibit significant differences (Suppl.

Table 1) in mutation rate phenotype, 7 x 10^{-8} in QMA0084 and 4 x 10^{-8} in clown loach strains (Figs. 1-2).

Phenotypically, clade B has shifted towards decreased haemolytic activity (Fig. 3, B2).

Clade C shares a common ancestor with clade B and is comprised of strains obtained from barramundi

farmed in WA, Northern Territory (NT) and north QLD fish farms (nested subclades C1 and C2

respectively). While a core mutation rate of 1.5-2 x 10^{-8} is observed in subclade C1, a significantly higher

mutation rate of 7 x 10^{-8} has been maintained in subclade C2 from north QLD for almost two decades

(1995 to 2012) (Figs. 1,2 Suppl. Table 1). We speculate that the latter is attributable to rapid fluctuation

in water salinity, oxygen and temperature accompanying periodic heavy rainfall in tropical north QLD,

creating an unstable environment where adaptive processes are ongoing and high mutation rate is

advantageous. Minor variation in virulence-related phenotypes (absence of capsule production and

longer cell chains in QMA0074, 77) (Fig. 2, Table 1), are supportive that mutation rate evolution in this

lineage might be driven by outside-the-host factors. The significant difference in mutation rate between

nested clades C1 and C2 may result from variation in *dnaN*. In *Bacillus subtilis,* 90% of mismatch repair

is dependent on targeting MutS to nascent DNA via the ß sliding clamp (DnaN) zone (82, 83).

Inactivation of *dnaN1* in *B. anthracis* results in a mutator phenotype with a mutation rate equivalent to

a *mutS* mismatch repair-defective strain (84). The tyrosine to isoleucine substitution in one of the

critical residues of the loader binding interface of the ß-clamp, predicted as deleterious (PROVEAN

score -2,216), is present in all isolates of from clade C (Table 2), which may explain the high mutation

rate in clade C2 strains (Fig. 2, Table 2). In contrast, only subclade C1 harbours T1135C nucleotide

substitution in the *dnaN* that changes stop codon TAA into CAA coding for glutamine predicted as

neutral (PROVEAN score -0.005), with the next stop codon TAG found immediately downstream.

Putatively, T1135C and/or change of the *dnaN* stop codon (85) may act as compensatory to the T326I

substitution, reducing the mutation rate to a value not significantly different from the core rate found

in clade A (Fig. 1, 2) via protein elongation or by affecting translation via stop codon usage bias (85). In

addition, a synonymous variant in *recD2* that is only present in subclade C2 may affect protein

expression thereby contributing to the difference in mutation rate between C1 and C2 subclades.

Clade D consists of isolates from trout (*Oncorhynchus mykiss*) (strains obtained from Réunion and

Israel). These strains have an estimated mutation rate of 5 x 10^{-8}, contain a glutamate to aspartate

substitution in *mutM* predicted to be deleterious to protein function, a synonymous SNP in *RecD2*, and

exhibit impeded haemolytic activity (Fig 1,2, Table 2). Also, isolates from Israel form thicker and more

dense biofilm structures (Fig. 3, D2).

Isolates from humans and fish are found in clade E where multiple virulence trait phenotypes are

observed (Figs. 2-3, Table 1). This clade contains two nested subclades that share a SNP in *mutS,* but

have other unique SNPs (Fig. 2, Table 2) and exhibit mutation rates that are significantly different in

most pairwise comparisons (Suppl. Table 1). Subclade E1 contains USA isolates from humans

(QMA0133-35, 37-38) and hybrid striped bass (QMA0447-48), which have a substitution in the Shine-

Dalgarno sequence of *recD2* and a mutation rate of 6.5 x 10^{-8} (Fig 1, Table 2). Fish strains and three

human strains (QMA0135, 37-38) form shorter cell chains and are less sensitive to H_2O_2. Two human

strains QMA0133-34 produce thicker and more dense biofilm structures (Fig. 3, D2). QMA0133 is non-

encapsulated (Fig 3, A2), and QMA0134 appears to differentially express the capsular polysaccharide in

culture (Fig 3, A3). The close phylogenetic relationship presented here implicates a likely transfer from

farmed hybrid bass to humans (86), a host jump that may have been facilitated by the mutator

phenotype. Subclade E2 contains two human strains from Canada (QMA0130-31) and a tilapia strain

from USA (QMA0466). These strains have unique methionine to isoleucine substitution at the N-

terminus of MutX, and a mutation rate of 4.5 x 10^{-8} (Fig 1, Table 2). Both human strains are non-

encapsulated (Fig. 2). The tilapia strain expresses the capsule but forms short cell chains in common

with most strains from the subclade (Fig 2, Fig 3, C3). The close phylogenetic relationship again strongly

implicates transfer from tilapia farmed in USA to human patients in Canada via import into the Toronto

fish market (87) and again, the mutator state may underlie this host jump.

Clade F is not readily definable by location, time of isolation, or host species and comprises QMA0139

from unidentified fish in Canada, QMA0190 isolated from snakehead murrel in Thailand (both of which

appear on long branches), and a nested terminal clade comprised of barramundi isolates farmed in

Recirculating Aquaculture Systems (RAS) in New South Wales and South Australia (Fig. 2). The latter are

mutators and were sampled from barramundi bone lesions during a disease outbreak in vaccinated fish

where infection manifested itself as a slowly progressing osteomyelitis instead of typical acute

septicaemia and meningitis, implicating mutators in adaptation to new tissue in response to host

immunity (64, 65). Since they are distant to strains used in autogenous vaccines in clade A (QMA0155-

57, 160, QMA0250-52), the atypical outbreak might be classified as a case of vaccine-induced serotype

replacement (VISR) – spread of co-existing pathogenic strain/s after elimination of the dominant

strain/s in the vaccine (88). In contrast to natural populations where the investigation of VISR is

confounded by ecological and sampling biases, and randomised vaccine trials that lack power to detect

the population-wide effect of mass vaccination (88), barramundi RAS represent a relatively controlled

environment where vaccination against *S. iniae* fails occasionally but recurrently (65). RAS are self-

contained with minimal water exchange and high standards of water treatment (89). This ensures that

gene flow and, consequently, the probability of new serotype introduction after vaccination is reduced

to a minimum. A comparatively uniform immunized host population is exposed to dominant strains

against which it was vaccinated along with potentially co-existing lineages in the RAS. In the atypical

outbreaks in RAS in Australia, vaccine-induced immunity was partially effective in the infected fish,

evident by absence of proliferation of the new strain in typical niduses (blood, brain, and pronephros)

and cross-reactivity of vaccine-induced antibodies against bone isolates (65). This is in contrast with the

classical model of serotype replacement where a vaccine is ineffective against the replacing co-existing

strain. However, VISR can occur even if the vaccine is equally efficient against all circulating strains via

other trade-off mechanisms (88). Apparently, evasion of immune clearance and adaptation to osseous

tissue was permitted by shift towards attenuated virulence indicated by multiple phenotypic changes in

bone isolates, including absence of capsule, impeded hemolytic activity, increased cell-chain length, and

denser biofilms (Fig. 3). All of these changes have previously been associated with chronic infection and

increased potential for colonization (90-96). Multiple SNPs in repair genes (Fig. 2, Table 2) and the

unique mutation rate (Figs. 1-2, Table 1) in bone isolates suggest that shift in virulence allowing

persistence in the bone tissue might have occurred via mutator phenotype. Serine to arginine

substitution in MutL and tyrosine to isoleucine substitution in MutY occur in the ancestor of QMA0139,

QMA0190, and bone strains. Both substitutions are predicted to have a deleterious effect on protein

function (PROVEAN scores -2.050 and -2.544 respectively), but their effect on mutation rate is

combined with unique SNPs found in each branch: QMA0139 has a SNP in the binding site for the DnaA

transcription factor within the *dnaN* promoter, and a mutation rate of 4.5×10^{-8}, QMA0190 has

synonymous SNP in *mutL* and a mutation rate of 6.5×10^{-8}, and bone strains have glutamine to glycine

substitution in MutS predicted as neutral (PROVEAN score -0.635), a synonymous SNP in *mutL*, and a

mutation rate of $3-4 \times 10^{-8}$. Considering that the mutation rate of QMA0190 is significantly higher

compared to rates expressed by QMA0139 and bone isolates (Suppl. Table 1), it is likely that variants

unique to the latter strains compensate for deleterious variants shared by the isolates (4).

Clade G contains two strains from tilapia isolated in USA (QMA0445-46). Notably, despite being

phylogenetically distant from clade A, these strains have the same MMR and OG genotype and have

retained the core mutation rate and the wildtype virulence phenotype (Fig. 2, Table 1), with the

exception of capsule absence.

Three isolates of *S. iniae* were identified as independent phylogenetic lineages, with unique mutation

rate genotypes and phenotypes, and virulence traits (Figs. 1-2, Tables 1-2). The first long branch with a

single isolate contains a strain from snakehead murrel strain (*Channa striata*) from Thailand. This isolate

is non-encapsulated and weakly haemolytic and has a mutation rate of 4.5×10^{-8}, potentially

attributable to a deleterious aspartate to asparagine substitution in rnhC and SNP in the binding site of

for the *fnr* transcription factor within the *dnaN* promoter. A second long branch with a single strain

contains the oldest among the analysed strains, QMA0140 isolated in 1976 from dolphin (*Inia*

geoffrensis) (97). This strain exhibits a mutation rate of 1×10^{-8}, which is unique among the strains and

significantly lower than the core mutation rate, perhaps linked to increased translation of recD2

helicase resulting from a SNP in the ribosome binding site (Fig 2, Table 2). The longest branch on the

tree contains a second dolphin strain, QMA0141, isolated two years later in 1978 (98). This isolate is

highly divergent from the rest of strains with around 20 kB of non-recombinant SNPs in pair-wise

comparisons with other strains, accounting for around 1% genomic difference. In contrast to QMA140,

it mutates at a rate of 1×10^{-7}, the highest mutation rate phenotype determined among the isolates and

significantly different to the rest of the values. Multiple SNPs are observed in all MMR and OG genes: 3

in *dnaN* and *mutX*, 4 in *mutM*, 9 in *mutL* and *rnhC*, 15 in *recD2*, 9 in *mutL* and *rnhC*, and 68 in *mutS*. Both

dolphin isolates are non-encapsulated, show increased ability to withstand oxidative stress, and form

denser and thicker biofilms (Fig 2, 3).

The conservation of *S. iniae* mutation rate and MMR/OG genotype within phylogenetic clades and

variation between the lineages support the importance of mutation rate dynamics in pathogen

evolution. Mechanisms that underlie the flux of mutator alleles and phenotypes include random

mutations (4), tandem repeat copy-number variation (20, 58), non-homologous recombination (21),

and prophage integration/excision (22). In *S. iniae,* mutation rate dynamics appear to be primarily

driven by random mutations and compensatory evolution. Tandem-repeat copy-number variants were

not identified in *S. iniae* MMR or OG genes, supporting the contention that occurrence of mutator

phenotypes and their full reversion via this mutational type is limited to *E. coli* and closely related

species (20, 58). Indeed, the presence of tandem repeats has only been detected in *mutL* in

Proteobacteria (20, 58), and it is well recognized that *mutL* is structurally and functionally different in

gram-positive bacteria and eukaryotes (54, 61). With the exception of the large deletion affecting the

mutY promoter in clade B, all identified mutations are single nucleotide substitutions. These MMR and

OG SNPs most likely result from random mutations which can be sufficient to induce profound changes

in mutation rate (55, 56). Since, in most cases, SNPs occur in multiple genes (Fig. 2) it is likely that

compensatory evolution of mutation rate prevails in *S. iniae* where some variants increase and some

decrease the background mutation rate (4). The association of MMR/OG variation with changes in

virulence traits (Fig. 2) supports the contention that emergence and decline of mutator strains may

facilitate adaptive processes in *S. iniae* populations driven by host immune response. Furthermore, the

most profound changes in virulence-associated traits and MMR and OG variants are observed in

atypical hosts (mammals) and atypical tissue (bone) within apparently immune vaccinated fish hosts

(Fig. 2), which has major epidemiological implications. First, mutators may promote multiple adaptive

changes required for pathogen transition among divergent host taxa known as 'host jumps', such as

Osteichthyes and *Mammalia* discussed in the present study (99). Second, mutators might present a risk

factor to immune escape after vaccination and serotype replacement. Moreover, as evidenced by *S.*

iniae colonization of bone tissue of vaccinated barramundi described previously (65), vaccine escape

leading to serotype replacement might occur without major disruption of immune recognition when

multiple changes attenuating virulence and antigenicity allow persistence in a tissue with lower

immune surveillance.

We cannot eliminate the possibility that variation in mutation rate and MMR/OG genotype in *S. iniae* is

neutral and simply coincides with phylogenetic diversification and variation in traits determining

virulence. Presence of dysfunctional MMR and OG genes within the evolutionary history of bacterial

strains may leave signatures within the genome in the form of mutational bias. For example, in all but

the most AT-rich bacteria there is a predisposition for GC-AT transitions over time (100), but in a long

term evolutionary experiment in *E.coli,* defective OG repair resulted in strong bias towards AT to CG

transversions while AT to GC transitions were dominant in MMR defective mutators (101). We

quantified the mutational spectra by clade across the natural *S. iniae* isolates in this study (Fig. 4). Two

clades of non-mutator strains (A and C1) had highly similar SNP type distribution (Fig. 4). Of the mutator

clades, only clade D had mutations constrained to a single MMR/OG, a SNP in *mutM* in the OG repair

system (Fig. 2), resulting in an amino acid substitution in the DNA glycolyase domain with PROVEAN

score of -2.990 (deleterious) (Table 2). Here we found the highest frequency of AT to CG transversions,

consistent with findings in *E. coli* experimental mutators defective in OG repair, albeit at much lower

frequency in our natural *S. iniae* mutator isolates (101). Two mutator clades (B and E2) had higher

frequency of AT to GC transitions than the non-mutator clades, and indeed other mutator clades (Fig 4).

Clade B contains mutations in both *mutL* (V-I substitution, PROVEAN -0.27, Table 2) and *recD2*

(synonymous) in the MMR system. Clade B strains also contain a conserved mutation in the promoter of

mutY in the OG repair system (Table 2). We acknowledge the weakness of these signals compared to

those reported in *E.coli* (101). However, we include only natural mutator isolates in which changes occur

in multiple genes in both MMR and OG systems. Moreover, *S. iniae* genomes are already AT-rich (mean

36.55% GC compared to *E. coli* ~50%) which may constrain transition and transversion mutations that

further bias the nucleotide content. Consequently, complete conservation of MMR and OG genes linked

with consistent mutation rates within *S. iniae* lineages (Fig. 2) and the fact that most SNPs occur in

functionally relevant coding and regulatory regions and/or change amino acid sequence (Table 2),

supports that the occurrence of mutator strains is likely to precede and facilitate shifts in virulence. This

may in turn lead to the major diversification events and epidemiologically relevant incidents such as

vaccine escape outbreaks and host jumps in these bacterial lineages.

Our results support the contention that high mutation rates promote transition to a new immunological

setting requiring multiple mutations and phenotypic changes, such as adaptation to a novel host, new

host tissue, or a host with vaccine-induced adaptive immune response. Extension to other species such

as *Streptococcus pneumoniae* may be particularly interesting as serotype evolution following vaccination

programmes is well documented, but the role of mutators in this process is unknown (66, 67, 102).

Future studies should determine how generalisable the role of mutators is in the epidemiology of

bacterial pathogens, evolution of virulence and antigenicity, and emergence of new pathogenic strains.

Materials and methods

Strains and growth conditions

Eighty isolates of *S. iniae* collected in Australia, USA, Canada, Israel, Honduras, and Thailand between
1976 and 2016 from eight fish species (*Lates calcarifer, Scortum barcoo, Epalzeorhynchos frenatum,*
Epalzeorhynchos bicolor, Oreochromis sp., Channa striata, Chromobotia macracanthus, Oncorhynchus
mykiss) and three mammalian species (*Homo sapiens, Inia geoffrensis, Pteropus alecto*) were analysed
(Table 1). Strains were received from culture collections, veterinarians, or directly from fish farms
(Suppl. Table 3) and stored as master seed stocks without further subculture at -80°C in Todd-Hewitt
Broth (THB, Oxoid) + 20% glycerol. Frozen stocks were recovered on Columbia agar supplemented with
5% defibrinated sheep blood (Oxoid), and cultured at 28°C on Todd-Hewitt agar (THA) or in THB with
agitation 200 rpm unless otherwise specified.

Estimation of mutation rate phenotype by fluctuation analysis

A fluctuation analysis assay for spontaneous occurrence of rifampicin resistance was optimized
according to Rosche and Foster (103). A single broth culture was initiated from five separate colonies,
recovered on Columbia blood agar from stock, and grown overnight to late-exponential phase in THB.
Cultures were adjusted to $OD_{600} = 1$ (10^8 CFU/mL), diluted 1:100, and distributed in 200 µl aliquots into
8 wells of a sterile U-bottom 96-well plate (Greiner). Prior to dilution, 100 µl of each OD_{600}-adjusted
culture was spread onto THA containing 0.5 µg/mL rifampicin to confirm absence of resistant mutants.
Although comparatively large initial inocula (1-2 x 10^5 CFU per culture, confirmed by Miles and Misra
CFU count (104) of OD-adjusted cultures) were used, this minimised variance in final CFU number (N_t)
determined in preliminary experiments. Invariability in N_t allowed statistical comparison of mutation
rates by the Maximum Likelihood Ratio test (71). To infer the N_t in final cultures and monitor their
growth by optical density, replicate plates containing two cultures per strain were prepared and
incubated in a BMG FLUOstar OPTIMA microplate reader. When these representative cultures entered
early stationary phase, CFU counts were performed by Miles and Misra method (104) and invariably
estimated as 1-2 x 10^8 CFU in each well. Immediately after, entire 200 µL cultures (above) were plated
on THA containing 0.5 µg/mL rifampicin for selection of mutants, dried under laminar flow, and
incubated until rifampicin resistant colonies appeared. The assay was repeated four times for each
strain in blocks of eight cultures using 20 isolates haphazardly chosen from different phylogenetic
lineages in each measurement. In some cases, e.g. a single strain representing an independent
phylogenetic lineage or with significantly differences in mutation rates detected in closely related
strains, assays were repeated 1-2 more times to increase statistical power. The rifampicin-resistant
mutant counts were pooled into single data sets representing 32 to 48 cultures per strain (Suppl. Table
2).

DNA extraction, preparation and sequencing

451 Genomic DNA was extracted from cells collected from 10 mL late-exponential phase culture in Todd-
452 Hewitt broth with the DNeasy Blood & Tissue kit (Qiagen) using a modified protocol with an additional
453 lysis step as described previously (105). Sequencing was performed on the Illumina HiSeq2000 platform
454 from Nextera XT pair-end libraries at Australian Genome Research Facility, Melbourne. A reference
455 genome from strain QMA0248 was constructed using both long reads derived from a single Smrt Cell
456 using the PacBio RS II system with P4C2 chemistry and short reads from Illumina HiSeq2000 derived
457 from Nextera XT paired-end libraries as reported elsewhere (NCBI accession no: GCA_002220115.1). All
458 sequence data are deposited at NCBI under Bioproject number PRJNA417543, SRA accession
459 SRP145425. Sample numbers, accession numbers and extended metadata are provided in Suppl. Table
460 3. Assembly statistics are provided in Suppl. Table 4.

461 **Alignment, recombination detection and phylogenetic analysis**

462 Phylogeny was constructed based on core genome SNPs from *de novo* genome assemblies filtered to
463 remove recombination breakpoints. Paired-end reads from Illumina were trimmed with Nesoni clip tool
464 version 0.132 (http://www.vicbioinformatics.com/software.nesoni.shtml), with minimum read length
465 50, and the first 15 bp of each read removed as quality deterioration in this region was observed when
466 assessed with FASTQC version 0.11.5. Assembly was performed using the SPAdes assembler version
467 3.7.1 (106), with minimum read coverage cutoff set to 10. Quality of assemblies was assessed with
468 QUAST 3.2 (107). Contigs were ordered by alignment to QMA248 reference genome (CP022392.1) with
469 Mauve Contig Mover 2.4.0 (108). Genome annotation was performed using Prokka 1.11 (109). Rapid
470 alignment of core genomes was carried out using parsnp in the Harvest Tools suite version 1.2 (110),
471 and the resulting alignment provided as an input to Gubbins 1.4.7 (75) for detection and exclusion of
472 variants produced by recombination. Phylogenies were then inferred from post-filtered core genome
473 polymorphic sites by maximum likelihood using RAxML 8.2.8 (111) with the general time reversible
474 nucleotide substitution model GTRGAMMA and bootstrap support from 1000 iterations. Effect of
475 ascertainment bias associated with using only polymorphic sites on branch length during ML inference
476 was corrected using Felsenstein's correction implemented in RAxML 8 (112). The resulting phylogenetic
477 tree was visualized using Dendroscope v 3.5.7 (113) with bootstrap node support value cut-off 75. For
478 the phylogram figure, tip labels were hidden for clarity and the edge containing QMA0141 was re-
479 scaled as dotted line representing 100-fold decrease in length (Fig. 1). A cladogram based on the
480 inferred phylogeny, showing all tip labels and bootstrap support for each node, was annotated with
481 metadata using Evolview V2 (Fig. 2)(114). To determine whether there was a strong temporal signal in
482 the phylogenetic data, a root-to-tip regression of branch length against time since isolation was
483 performed in TempEst (79), using genetic distances corrected for ascertainment bias in an unrooted

484 tree estimated by maximum likelihood from the alignment of non-recombinant core-genome SNPs in
485 RAxML 8.2.8.

SNP analysis, variation in DNA repair genes and their regulatory regions

SNP analysis was performed by read-mapping with Geneious version 9.1 (115), using default settings
unless otherwise specified. Paired-end reads from each genome were trimmed, merged into a single
file with expected distances between the reads set to 250 bp, and mapped to the curated reference
genome of strain QMA0248. Mapped reads were used to detect SNPs with minimum coverage set to 10
and frequency to 0.9. A consensus "pseudogenome" was generated for every strain based on the
reference sequence incorporating any detected variants (116). Multiple alignment of these
pseudogenomes was carried out with *Geneious* aligning tool. SNP types and position per strain were
exported as a table and SNPs of each type occurring only in each clade but not in any other clade were
counted and plotted using a custom script in R. Sequences of MMR genes (*dnaN,* mut*S, muTL, recD2,*
rnhC) and OG genes (*mutY* , *mutM*, *mutX*) annotated in the QMA0248 reference genome were
identified in each strain genome sequence. Promoter regions of repair genes were determined with
BPROM (117), protein functional domains by SMART genomic (118), and effect of amino acid
substitutions on protein function by PROVEAN Protein with a sensitivity cut-off of 1.3 (119). To detect
large variants, alignment of these regions extracted from *de novo* genome assemblies was carried out,
and a 56 bp deletion detected in *mutY* promoter of clade B was confirmed by PCR with
CAGAAGGAAGAAACAGAC_F/ACCTCTATTGTAGCAAAG_R primers.

Detecting possible association between variation in DNA repair genes and mobilome

A pangenome for the 80 *S. iniae* isolates was constructed with GView server (120) using the
QMA0248 reference genbank (.gbk) file as seed by sequentially adding 79 genome assemblies by
clade in the order derived from the cladogram (Fig. 2). Phage positions in the resulting pangenome
were then determined by BLAST using Phaster (76) while IS positions were identified using ISFinder
(77) with a BLAST e-value cutoff of 1×10^{-30}. Positions of MMR and OG genes in the pangenome were
determined via manual search of the .gbk file. An image was created in BRIG (121) using the
pangenome as reference then aligning the 80 genome assemblies by BLAST with an e-value cutoff of
1×10^{-30}. The resulting image was annotated in BRIG from a spreadsheet derived from the positions
determined above, and coloured according to clade (Suppl. Fig. 1).

Phenotypic variation related to virulence and antigenicity

514 A multitude of intracellular and secreted enzymes contribute to virulence in the *Streptococcus* genus
515 (90, 122). Considering the large number of strains, we limited analysis to major, easily observable, and

516 distinct phenotypes strongly associated with virulence in streptococci (capsule, hemolysis, cell chains)

517 and bacteria in general (oxidation resistance, biofilms). All assays were performed at least in triplicate

518 per strain.

Buoyant density assay for presence of polysaccharide capsule

Presence/absence of polysaccharide capsule was estimated by Percoll buoyant density assay. Isotonic

stock Percoll (ISP) was prepared by mixing nine parts of Percoll with one part of 1.5 M NaCl. Then, 6

parts of ISP was diluted with 4 parts of 0.15 M NaCl to make final 50 % Percoll solution, which was

distributed by 3 mL into flow cytometry tubes. THB cultures (10 mL) grown to late-exponential phase

were adjusted at OD_{600} 1 (10^8 CFU/mL), centrifuged at 3220 x g for 5 min, resuspended in 0.5 mL of 0.15

M NaCl, and layered onto the Percoll. Tubes were centrifuged at 4°C in a swinging bucket rotor at 4000

x g for 3 h with low acceleration and no brake. In this assay, encapsulated cells form a clear compact

band in the Percoll gradient (Fig. 3, A1), non-encapsulated cells form a pellet in the bottom of the tube

(Figure 3, A2) and, occasionally, strains show differential expression of capsular polysaccharide

evidenced by band and a pellet (Fig. 3, A3).

Haemolytic activity assay on sheep blood agar

Rapid high-throughput detection of impaired haemolytic activity was achieved by blood-agar clearance

zone assay. Briefly, 5 mm wells were made in Columbia agar supplemented with 5% defibrinated sheep

blood (Oxoid, Australia). Bacterial cultures grown to late-exponential phase (~10^8 CFU) were diluted

1:1000 and 50 μL of diluted cultures were pipetted into punctures in the agar (initial inoculum ~$5x10^4$

CFU in each puncture). A 3 mm wide clearance zone was generally produced by bacterial lysis of sheep

erythrocytes during 24 h incubation (Fig. 3, B1). Where haemolysis was absent or fragmentary, impeded

haemolytic activity was recorded (Fig. 3, B2).

Chain formation microscopy

To assess chain formation, *S. iniae* cultures in THB were grown stationarily in 96-well plates at 28 °C for

24 h, mixed and 5 μL wet mounts prepared on a glass microscope slide. Slides were observed by bright

field microscopy under 40x objective with an Olympus BX40 microscope and captured using an

Olympus DP28 digital camera using CellSens software (Olympus Optical Co, Japan). Specimens were

observed for at least 3 min and 2-20 cell chains were generally observed (Fig. 3, C1). Where over 20

cells in a chain were repeatedly detected (Fig. 3, C2) increased chain formation was recorded, and when

more than 10 cells in a chain were not detected by similar observation and the culture was mainly

composed of detached cells (Fig. 3, C3) impeded chain formation was recorded. For the figure, 50 μL of

cultures were dried onto slides at RT, fixed with methanol, Gram stained, and images captured under

100x objective.

Oxidation resistance assay

Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) (123) of

hydrogen peroxide were measured to assess oxidative stress resistance among *S. iniae* strains. THB

cultures grown to late exponential phase were adjusted to OD_{600} = 1 (10^8 CFU/mL), diluted 1000-fold,

and distributed by 100 μL (~10^4 CFU) into wells of a U-bottom 96-well plate (Greiner). THB (100 μL) with

2-fold excess concentration of hydrogen peroxide was added to the wells and serially diluted twofold,

resulting in a range from 0 to 10 mM final peroxide concentrations. Plates were incubated stationarily

for 24h and examined for presence of observable growth to determine the MIC. Viable cell counts of

cultures without visible growth were performed to determine MBC. MIC of hydrogen peroxide was

estimated as 3 mM for all strains, and MBC as 4 mM for the majority of strains. MBC elevated to 5 mM

in dolphin and human isolates from USA was classified deviation from a wildtype phenotype.

Biofilm formation and visualisation assay

S. iniae biofilms were grown for 4 days in 8-well Lab-Tek® II Chamber Slide™ Systems. To prepare an

initial inoculum of 5 x 10^5 CFU, THB cultures grown to late exponential phase were adjusted to OD_{600} = 1

(10^8 CFU/mL), diluted 100-fold, and 0.5 mL of diluted cultures were placed into Chamber slide wells.

After 24 h incubation, THB was removed and replaced with fresh THB every 10-14 h for 3 days. For

visualisation, biofilms were washed in PBS, stained for 15 min with 1 μM fluorescent *Bac*Light Red

bacterial stain, washed in PBS, fixed for 30 min with 10% formalin, and washed twice PBS. Z-stacks were

collected by ZEISS LSM 710 Inverted Laser Scanning Confocal Microscope at 20x objective and visualised

using ZEN2012. Quantification of biofilms was performed using COMSTAT (124). Structures of biomass

larger than 3.5 and average thickness over 5 were classified as increased biofilm forming activity (Fig. 3

D).

Identifying wildtype virulence-related phenotypes

572 We consider prevalent phenotypes as a wildtype, namely: presence of polysaccharide capsule (71.25 %

573 of strains; Figure 3, A1) and haemolytic activity (85% of strains; Figure 3, B1), up to 20 cells in a chain

574 (77.5 % of strains; Figure 3, A1), 3 mM MIC and 4 mM MBC of hydrogen peroxide (91.25 % of strains),

575 and 8 -12 μm thick biofilms covering up to 30% of representative image (86.25 % of strains, Fig. 4A).

576 Other phenotypes are regarded as deviant and discussed.

Statistical analysis

Analysis of co-variance between number of MMR and OG variants, mutation rates, number of

atypical phenotype associated with virulence, and atypical places of isolation were tested with a

Phylogenetic Generalised Least Squares model that accounts for relatedness among strains under

Pagel's λ of 0.6 implemented in R (70). Estimation of mutation rates was carried out using the Ma-

Sandri-Sarkar Maximum Likelihood Estimation (MSS-MLE) method as implemented in FALCOR

fluctuation analysis calculator (125). Differences in mutation rates were compared by Likelihood

Ratio Test (LRT) using the rSalvador R package (71, 126). LRT estimates the overlap between

confidence intervals calculated for two fluctuation experiment data sets and allows to compare

fluctuation assay data with different number of cultures and resolve very small differences in

mutation rate (71, 126). The *Compare.LD* function was applied pair-wise to all strains, and obtained

p values were corrected for multiple comparisons by controlling the False Discovery Rate (72) using

p.adjust in R. Adjusted *p* values less than 0.05 were considered to indicate significant difference in

mutation rates.

Acknowledgements

This work was supported by The Australian Research Council Discovery Project Grant DP120102755

awarded to ACB. We thank the University of Queensland School of Biological Sciences for funding a

tuition fee waiver for OS. For provision of strains into our collection over 15 years we thank: Judy

Forbes-Faulkner, Oonoonba Veterinary Laboratory, QLD, Australia; Lynn Shewmaker, CDC, Atlanta, USA;

Christian Michel, INRA, France; Suresh Benedict, Berrimah Veterinary Laboratories, NT, Australia; Nicky

Buller, WA Fisheries, WA, Australia; Mark White, Tréidlia Biovet Pty (formerly Allied Biotechnology),

NSW, Australia; Matt Landos, Future Fisheries Veterinary Services, NSW, Australia; Spencer Russell,

Novartis Animal Health, Canada; Ruben Arturo Lopez Crespo, Regal Springs Tilapia, Honduras. We thank

Dr Simone Blomberg for advice on statistical analyses and Mr Nazar Rudenko for assistance with writing

software pipelines and for use of his Unix server.

References

1.   Ferenci T. 2015. Trade-off Mechanisms Shaping the Diversity of Bacteria. Trends Microbiol
605   doi:10.1016/j.tim.2015.11.009.
606   2.   Stich M, Manrubia SC, Lazaro E. 2010. Variable Mutation Rates as an Adaptive Strategy in
607   Replicator Populations. Plos One 5.
608   3.   Sniegowski P, Raynes Y. 2013. Mutation Rates: How Low Can You Go? Current Biology
609   23:R147-R149.
610   4.   Wielgoss S, Barrick JE, Tenaillon O, Wiser MJ, Dittmar WJ, Cruveiller S, Chane-Woon-Ming B,
611   Medigue C, Lenski RE, Schneider D. 2013. Mutation rate dynamics in a bacterial population
612   reflect tension between adaptation and genetic load. Proc Natl Acad Sci U S A 110:222-7.
613   5.   Wielgoss S, Barrick JE, Tenaillon O, Cruveiller S, Chane-Woon-Ming B, Medigue C, Lenski RE,
614   Schneider D. 2011. Mutation Rate Inferred From Synonymous Substitutions in a Long-Term
615   Evolution Experiment With Escherichia coli. G3 (Bethesda) 1:183-186.
616   6.   Galhardo RS, Hastings PJ, Rosenberg SM. 2007. Mutation as a stress response and the
617   regulation of evolvability. Crit Rev Biochem Mol Biol 42:399-435.
618   7.   Sundin GW, Weigand MR. 2007. The microbiology of mutability. Fems Microbiology Letters
619   277:11-20.
620   8.   Foster PL. 2007. Stress-induced mutagenesis in bacteria. Crit Rev Biochem Mol Biol 42:373-
621   97.
622   9.   Wrande M, Roth JR, Hughes D. 2008. Accumulation of mutants in "aging" bacterial colonies
623   is due to growth under selection, not stress-induced mutagenesis. Proc Natl Acad Sci U S A
624   105:11863-8.
625   10.   Torres-Barcelo C, Kojadinovic M, Moxon R, MacLean RC. 2015. The SOS response increases
626   bacterial fitness, but not evolvability, under a sublethal dose of antibiotic. Proc Biol Sci
627   282:20150885.
628   11.   Andersson DI, Hughes D. 2010. Antibiotic resistance and its cost: is it possible to reverse
629   resistance? Nat Rev Microbiol 8:260-71.
630   12.   Boe L, Danielsen M, Knudsen S, Petersen JB, Maymann J, Jensen PR. 2000. The frequency of
631   mutators in populations of Escherichia coli. Mutat Res 448:47-55.
632   13.   Miller JH. 1996. Spontaneous mutators in bacteria: insights into pathways of mutagenesis
633   and repair. Annu Rev Microbiol 50:625-43.
634   14.   Mao EF, Lane L, Lee J, Miller JH. 1997. Proliferation of mutators in A cell population. J
635   Bacteriol 179:417-22.
636   15.   Taddei F, Radman M, MaynardSmith J, Toupance B, Gouyon PH, Godelle B. 1997. Role of
637   mutator alleles in adaptive evolution. Nature 387:700-702.
638   16.   Torres-Barcelo C, Cabot G, Oliver A, Buckling A, Maclean RC. 2013. A trade-off between
639   oxidative stress resistance and DNA repair plays a role in the evolution of elevated mutation
640   rates in bacteria. Proc Biol Sci 280:20130007.
641   17.   Raynes Y, Sniegowski PD. 2014. Experimental evolution and the dynamics of genomic
642   mutation rate modifiers. Heredity (Edinb) 113:375-80.
643   18.   de Visser JA. 2002. The fate of microbial mutators. Microbiology 148:1247-52.
644   19.   Turrientes MC, Baquero F, Levin BR, Martinez JL, Ripoll A, Gonzalez-Alba JM, Tobes R,
645   Manrique M, Baquero MR, Rodriguez-Dominguez MJ, Canton R, Galan JC. 2013. Normal
646   mutation rate variants arise in a Mutator (Mut S) Escherichia coli population. PLoS One
647   8:e72963.
648   20.   Shaver AC, Sniegowski PD. 2003. Spontaneously arising mutL mutators in evolving
649   Escherichia coli populations are the result of changes in repeat length. J Bacteriol 185:6076-
650   82.

21.   Denamur E, Lecointre G, Darlu P, Tenaillon O, Acquaviva C, Sayada C, Sunjevaric I, Rothstein
R, Elion J, Taddei F, Radman M, Matic I. 2000. Evolutionary implications of the frequent
horizontal transfer of mismatch repair genes. Cell 103:711-21.

22.   Scott J, Thompson-Mayberry P, Lahmamsi S, King CJ, McShan WM. 2008. Phage-associated
mutator phenotype in group A streptococcus. J Bacteriol 190:6290-301.

23.   Hall LM, Henderson-Begg SK. 2006. Hypermutable bacteria isolated from humans--a critical
analysis. Microbiology 152:2505-14.

24.   Oliver A. 2010. Mutators in cystic fibrosis chronic lung infection: Prevalence, mechanisms,
and consequences for antimicrobial therapy. Int J Med Microbiol 300:563-72.

25.   Matsushima A, Takakura S, Fujihara N, Saito T, Ito I, Iinuma Y, Ichiyama S. 2010. High
prevalence of mutators among Enterobacter cloacae nosocomial isolates and their
association with antimicrobial resistance and repetitive detection. Clin Microbiol Infect
16:1488-93.

26.   Wang S, Wang Y, Shen J, Wu Y, Wu C. 2013. Polymorphic mutation frequencies in clinical
isolates of Staphylococcus aureus: the role of weak mutators in the development of
fluoroquinolone resistance. FEMS Microbiol Lett 341:13-7.

27.   Negri MC, Morosini MI, Baquero MR, del Campo R, Blazquez J, Baquero F. 2002. Very low
cefotaxime concentrations select for hypermutable Streptococcus pneumoniae populations.
Antimicrobial Agents and Chemotherapy 46:528-530.

28.   Chopra I, O'Neill AJ, Miller K. 2003. The role of mutators in the emergence of antibiotic-
resistant bacteria. Drug Resist Updat 6:137-45.

29.   Blazquez J. 2003. Hypermutation as a factor contributing to the acquisition of antimicrobial
resistance. Clinical Infectious Diseases 37:1201-1209.

30.   Mena A, Smith EE, Burns JL, Speert DP, Moskowitz SM, Perez JL, Oliver A. 2008. Genetic
Adaptation of Pseudomonas aeruginosa to the Airways of Cystic Fibrosis Patients Is
Catalyzed by Hypermutation. Journal of Bacteriology 190:7910-7917.

31.   Gutierrez O, Juan C, Perez JL, Oliver A. 2004. Lack of association between hypermutation and
antibiotic resistance development in Pseudomonas aeruginosa isolates from intensive care
unit patients. Antimicrobial Agents and Chemotherapy 48:3573-3575.

32.   Giraud A, Matic I, Tenaillon O, Clara A, Radman M, Fons M, Taddei F. 2001. Costs and
benefits of high mutation rates: Adaptive evolution of bacteria in the mouse gut. Science
291:2606-2608.

33.   Nilsson AI, Kugelberg E, Berg OG, Andersson DI. 2004. Experimental adaptation of
Salmonella typhimurium to mice. Genetics 168:1119-30.

34.   Healey KR, Zhao Y, Perez WB, Lockhart SR, Sobel JD, Farmakiotis D, Kontoyiannis DP,
Sanglard D, Taj-Aldeen SJ, Alexander BD, Jimenez-Ortigosa C, Shor E, Perlin DS. 2016.
Prevalent mutator genotype identified in fungal pathogen Candida glabrata promotes multi-
drug resistance. Nat Commun 7:11128.

35.   Labat F, Pradillon O, Garry L, Peuchmaur M, Fantin B, Denamur E. 2005. Mutator phenotype
confers advantage in Escherichia coli chronic urinary tract infection pathogenesis. FEMS
Immunol Med Microbiol 44:317-21.

36.   Mena A, Macia MD, Borrell N, Moya B, de Francisco T, Perez JL, Oliver A. 2007. Inactivation
of the mismatch repair system in Pseudomonas aeruginosa attenuates virulence but favors
persistence of oropharyngeal colonization in cystic fibrosis mice. J Bacteriol 189:3665-8.

37.   Oliver A, Mena A. 2010. Bacterial hypermutation in cystic fibrosis, not only for antibiotic
resistance. Clinical Microbiology and Infection 16:798-808.

38.   Aquino T, Nunes A. 2016. Host immunity and pathogen diversity: A computational study.
Virulence 7:121-8.

39.   Cobey S. 2014. Pathogen evolution and the immunological niche. Ann N Y Acad Sci 1320:1-
15.

40. Lukacisinova M, Novak S, Paixao T. 2017. Stress-induced mutagenesis: Stress diversity
facilitates the persistence of mutator genes. PLoS Comput Biol 13:e1005609.
41. Abel S, Abel zur Wiesch P, Davis BM, Waldor MK. 2015. Analysis of Bottlenecks in
Experimental Models of Infection. PLoS Pathog 11:e1004823.
42. Metzgar D, Wills C. 2000. Evidence for the adaptive evolution of mutation rates. Cell
101:581-4.
43. Fogle CA, Nagle JL, Desai MM. 2008. Clonal interference, multiple mutations and adaptation
in large asexual populations. Genetics 180:2163-73.
44. Methot PO, Alizon S. 2014. What is a pathogen? Toward a process view of host-parasite
interactions. Virulence 5:775-85.
45. Mackinnon MJ, Read AF. 2004. Virulence in malaria: an evolutionary viewpoint. Philos Trans
R Soc Lond B Biol Sci 359:965-86.
46. Deitsch KW, Moxon ER, Wellems TE. 1997. Shared themes of antigenic variation and
virulence in bacterial, protozoal, and fungal infections. Microbiol Mol Biol Rev 61:281-93.
47. Shrestha S, Bjornstad ON, King AA. 2014. Evolution of acuteness in pathogen
metapopulations: conflicts between "classical" and invasion-persistence trade-offs. Theor
Ecol 7:299-311.
48. Maurelli AT. 2007. Black holes, antivirulence genes, and gene inactivation in the evolution of
bacterial pathogens. FEMS Microbiology Letters 267:1-8.
49. Merino D, Reglier-Poupet H, Berche P, Charbit A, European Listeria Genome C. 2002. A
hypermutator phenotype attenuates the virulence of Listeria monocytogenes in a mouse
model. Mol Microbiol 44:877-87.
50. Gonzalez K, Faustoferri RC, Quivey RG, Jr. 2012. Role of DNA base excision repair in the
mutability and virulence of Streptococcus mutans. Mol Microbiol 85:361-77.
51. Picard B, Duriez P, Gouriou S, Matic I, Denamur E, Taddei H. 2001. Mutator natural
Escherichia coli isolates have an unusual virulence phenotype. Infection and Immunity 69:9-
14.
52. Smania AM, Segura I, Pezza RJ, Becerra C, Albesa I, Argarana CE. 2004. Emergence of
phenotypic variants upon mismatch repair disruption in Pseudomonas aeruginosa.
Microbiology-Sgm 150:1327-1338.
53. Canfield GS, Schwingel JM, Foley MH, Vore KL, Boonanantanasarn K, Gill AL, Sutton MD, Gill
SR. 2013. Evolution in fast forward: a potential role for mutators in accelerating
Staphylococcus aureus pathoadaptation. J Bacteriol 195:615-28.
54. Fukui K. 2010. DNA mismatch repair in eukaryotes and bacteria. J Nucleic Acids 2010.
55. Dai L, Muraoka WT, Wu ZW, Sahin O, Zhang QJ. 2015. A single nucleotide change in mutY
increases the emergence of antibiotic-resistant Campylobacter jejuni mutants. Journal of
Antimicrobial Chemotherapy 70:2739-2748.
56. Rajanna C, Ouellette G, Rashid M, Zemla A, Karavis M, Zhou C, Revazishvili T, Redmond B,
McNew L, Bakanidze L, Imnadze P, Rivers B, Skowronski EW, O'Connell KP, Sulakvelidze A,
Gibbons HS. 2013. A strain of Yersinia pestis with a mutator phenotype from the Republic of
Georgia. FEMS Microbiol Lett 343:113-20.
57. Ambur OH, Davidsen T, Frye SA, Balasingham SV, Lagesen K, Rognes T, Tonjum T. 2009.
Genome dynamics in major bacterial pathogens. Fems Microbiology Reviews 33:453-470.
58. Chen F, Liu WQ, Eisenstark A, Johnston RN, Liu GR, Liu SL. 2010. Multiple genetic switches
spontaneously modulating bacterial mutability. BMC Evol Biol 10:277.
59. Li GM. 2008. Mechanisms and functions of DNA mismatch repair. Cell Research 18:85-98.
60. Tham KC, Hermans N, Winterwerp HHK, Cox MM, Wyman C, Kanaar R, Lebbink JHG. 2013.
Mismatch Repair Inhibits Homeologous Recombination via Coordinated Directional
Unwinding of Trapped DNA Structures. Molecular Cell 51:326-337.
61. Lenhart JS, Pillon MC, Guarne A, Biteen JS, Simmons LA. 2016. Mismatch repair in Gram-
positive bacteria. Res Microbiol 167:4-12.

62. David SS, O'Shea VL, Kundu S. 2007. Base-excision repair of oxidative DNA damage. Nature
447:941-950.
63. Lu AL, Li X, Gu Y, Wright PM, Chang DY. 2001. Repair of oxidative DNA damage: mechanisms
and functions. Cell Biochem Biophys 35:141-70.
64. Agnew W, Barnes AC. 2007. Streptococcus iniae: an aquatic pathogen of global veterinary
significance and a challenging candidate for reliable vaccination. Vet Microbiol 122:1-15.
65. Millard CM, Baiano JC, Chan C, Yuen B, Aviles F, Landos M, Chong RS, Benedict S, Barnes AC.
2012. Evolution of the capsular operon of Streptococcus iniae in response to vaccination.
Appl Environ Microbiol 78:8219-26.
66. Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, Bentley SD, Hanage
WP, Lipsitch M. 2013. Population genomics of post-vaccine changes in pneumococcal
epidemiology. Nat Genet 45:656-63.
67. Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, McGee L, von
Gottberg A, Song JH, Ko KS, Pichon B, Baker S, Parry CM, Lambertsen LM, Shahinas D, Pillai
DR, Mitchell TJ, Dougan G, Tomasz A, Klugman KP, Parkhill J, Hanage WP, Bentley SD. 2011.
Rapid Pneumococcal Evolution in Response to Clinical Interventions. Science 331:430-434.
68. Maiden MCJ, van Rensburg MJJ, Bray JE, Earle SG, Ford SA, Jolley KA, McCarthy ND. 2013.
MLST revisited: the gene-by-gene approach to bacterial genomics. Nat Rev Micro 11:728-
736.
69. Shabalina SA, Spiridonov NA, Kashina A. 2013. Sounds of silence: synonymous nucleotides as
a key to biological regulation and complexity. Nucleic Acids Res 41:2073-94.
70. Symonds MRE, Blomberg SP. 2014. A Primer on Phylogenetic Generalised Least Squares, p
105-130. In Garamszegi LZ (ed), Modern Phylogenetic Comparative Methods and Their
Application in Evolutionary Biology: Concepts and Practice doi:10.1007/978-3-662-43550-
2_5. Springer Berlin Heidelberg, Berlin, Heidelberg.
71. Zheng Q. 2015. Methods for comparing mutation rates using fluctuation assay data. Mutat
Res 777:20-2.
72. Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B-
Methodological 57:289-300.
73. Gould CV, Sniegowski PD, Shchepetov M, Metlay JP, Weiser JN. 2007. Identifying mutator
phenotypes among fluoroquinolone-resistant strains of Streptococcus pneumoniae using
fluctuation analysis. Antimicrob Agents Chemother 51:3225-9.
74. Sprouffske K, Aguilar-Rodriguez J, Sniegowski P, Wagner A. 2018. High mutation rates limit
evolutionary adaptation in Escherichia coli. PLoS Genet 14:e1007324.
75. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR.
2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome
sequences using Gubbins. Nucleic Acids Res 43:e15.
76. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang YJ, Wishart DS. 2016. PHASTER: a better,
faster version of the PHAST phage search tool. Nucleic Acids Research 44:W16-W21.
77. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference
centre for bacterial insertion sequences. Nucleic Acids Research 34:D32-D36.
78. Chaguza C, Andam CP, Harris SR, Cornick JE, Yang M, Bricio-Moreno L, Kamng'ona AW,
Parkhill J, French N, Heyderman RS, Kadioglu A, Everett DB, Bentley SD, Hanage WP. 2016.
Recombination in Streptococcus pneumoniae Lineages Increase with Carriage Duration and
Size of the Polysaccharide Capsule. mBio 7:e01053-16.
79. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. 2016. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evolution 2.
80. Bedford T, Cobey S, Pascual M. 2011. Strength and tempo of selection revealed in viral gene
genealogies. BMC Evol Biol 11:220.

81. Neher RA, Hallatschek O. 2013. Genealogies of rapidly adapting populations. Proceedings of
the National Academy of Sciences of the United States of America 110:437-442.
82. Dupes NM, Walsh BW, Klocko AD, Lenhart JS, Peterson HL, Gessert DA, Pavlick CE, Simmons
LA. 2010. Mutations in the Bacillus subtilis beta clamp that separate its roles in DNA
replication from mismatch repair. J Bacteriol 192:3452-63.
83. Lenhart JS, Sharma A, Hingorani MM, Simmons LA. 2013. DnaN clamp zones provide a
platform for spatiotemporal coupling of mismatch detection to DNA replication. Mol
Microbiol 87:553-68.
84. Yang H, Miller JH. 2008. Deletion of dnaN1 generates a mutator phenotype in Bacillus
anthracis. DNA Repair (Amst) 7:507-14.
85. Korkmaz G, Holm M, Wiens T, Sanyal S. 2014. Comprehensive analysis of stop codon usage in
bacteria and its correlation with release factor abundance. J Biol Chem 289:30334-42.
86. Facklam R, Elliott J, Shewmaker L, Reingold A. 2005. Identification and characterization of
sporadic isolates of Streptococcus iniae isolated from humans. Journal of Clinical
Microbiology 43:933-937.
87. Weinstein MR, Litt M, Kertesz DA, Wyper P, Rose D, Coulter M, McGeer A, Facklam R, Ostach
C, Willey BM, Borczyk A, Low DE. 1997. Invasive infections due to a fish pathogen,
Streptococcus iniae. New England Journal of Medicine 337:589-594.
88. Weinberger DM, Malley R, Lipsitch M. 2011. Serotype replacement in disease after
pneumococcal vaccination. Lancet 378:1962-73.
89. Losordo T, DeLong D, Guerdat T. 2009. Advances in technology and practice for land-based
aquaculture systems: tank-based recirculating systems for finfish production. New
Technologies in Aquaculture doi:Doi 10.1533/9781845696474.6.945:945-983.
90. Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of Streptococcus pneumoniae
virulence factors in host respiratory colonization and disease. Nat Rev Microbiol 6:288-301.
91. Locke JB, Colvin KM, Datta AK, Patel SK, Naidu NN, Neely MN, Nizet V, Buchanan JT. 2007.
Streptococcus iniae capsule impairs phagocytic clearance and contributes to virulence in
fish. J Bacteriol 189:1279-87.
92. Locke JB, Colvin KM, Varki N, Vicknair MR, Nizet V, Buchanan JT. 2007. Streptococcus iniae
beta-hemolysin streptolysin S is a virulence factor in fish infection. Diseases of Aquatic
Organisms 76:17-26.
93. Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, Konradsen HB, Nahm MH.
2015. Pneumococcal Capsules and Their Types: Past, Present, and Future. Clin Microbiol Rev
28:871-99.
94. Rodriguez JL, Dalia AB, Weiser JN. 2012. Increased chain length promotes pneumococcal
adherence and colonization. Infect Immun 80:3454-9.
95. Andisi VF, Hinojosa CA, de Jong A, Kuipers OP, Orihuela CJ, Bijlsma JJ. 2012. Pneumococcal
gene complex involved in resistance to extracellular oxidative stress. Infect Immun 80:1037-
49.
96. Bjarnsholt T. 2013. The role of bacterial biofilms in chronic infections. APMIS Suppl
doi:10.1111/apm.12099:1-51.
97. Pier GB, Madin SH. 1976. Streptococcus iniae sp nov, a beta-hemolytic Streptococcus
isolated from an amazon freshwater dolphin, Inia geoffrensis. . International Journal of
Systematic Bacteriology 26:545-553.
98. Pier GB, Madin SH, Alnakeeb S. 1978. Isolation and characterisation of a 2nd isolate of
Streptococcus iniae. International Journal of Systematic Bacteriology 28:311-314.
99. Baumler A, Fang FC. 2013. Host specificity of bacterial pathogens. Cold Spring Harb Perspect
Med 3:a010041.
100. Hildebrand F, Meyer A, Eyre-Walker A. 2010. Evidence of selection upon genomic GC-
content in bacteria. PLoS Genet 6:e1001107.

101. Couce A, Caudwell LV, Feinauer C, Hindre T, Feugeas JP, Weigt M, Lenski RE, Schneider D,
Tenaillon O. 2017. Mutator genomes decay, despite sustained fitness gains, in a long-term
experiment with bacteria. Proc Natl Acad Sci U S A 114:E9026-E9035.
102. Croucher NJ, Kagedan L, Thompson CM, Parkhill J, Bentley SD, Finkelstein JA, Lipsitch M,
Hanage WP. 2015. Selective and genetic constraints on pneumococcal serotype switching.
PLoS Genet 11:e1005095.
103. Rosche WA, Foster PL. 2000. Determining mutation rates in bacterial populations. Methods
20:4-17.
104. Hedges AJ. 2002. Estimating the precision of serial dilutions and viable bacterial counts. Int J
Food Microbiol 76:207-14.
105. Kawasaki M, Delamare-Deboutteville J, Bowater RO, Walker MJ, Beatson S, Ben Zakour NL,
Barnes AC. 2018. Microevolution of aquatic Streptococcus agalactiae ST-261 from Australia
indicates dissemination via imported tilapia and ongoing adaptation to marine hosts or
environment. Appl Environ Microbiol doi:10.1128/AEM.00859-18.
106. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner
PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. J Comput Biol 19:455-77.
107. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 29:1072-5.
108. Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of
draft genomes using the Mauve aligner. Bioinformatics 25:2071-3.
109. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068-9.
110. Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-
genome alignment and visualization of thousands of intraspecific microbial genomes.
Genome Biology 15:524.
111. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30:1312-3.
112. Leache AD, Banbury BL, Felsenstein J, de Oca AN, Stamatakis A. 2015. Short Tree, Long Tree,
Right Tree, Wrong Tree: New Acquisition Bias Corrections for Inferring SNP Phylogenies.
Systematic Biology 64:1032-47.
113. Huson DH, Richter DC, Rausch C, Dezulian T, Franz M, Rupp R. 2007. Dendroscope: An
interactive viewer for large phylogenetic trees. BMC Bioinformatics 8:460.
114. He Z, Zhang H, Gao S, Lercher MJ, Chen WH, Hu S. 2016. Evolview v2: an online visualization
and management tool for customized and annotated phylogenetic trees. Nucleic Acids Res
44:W236-41.
115. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic:
An integrated and extendable desktop software platform for the organization and analysis of
sequence data. Bioinformatics 28:1647-1649.
116. Ben Zakour NL, Alsheikh-Hussain AS, Ashcroft MM, Khanh Nhu NT, Roberts LW, Stanton-
Cook M, Schembri MA, Beatson SA. 2016. Sequential Acquisition of Virulence and
Fluoroquinolone Resistance Has Shaped the Evolution of Escherichia coli ST131. MBio
7:e00347-16.
117. Umarov RK, Solovyev VV. 2017. Recognition of prokaryotic and eukaryotic promoters using
convolutional deep learning neural networks. Plos One 12.
118. Letunic I, Bork P. 2018. 20 years of the SMART protein domain annotation resource. Nucleic
Acids Res 46:D493-D496.
119. Choi Y, Chan AP. 2015. PROVEAN web server: a tool to predict the functional effect of amino
acid substitutions and indels. Bioinformatics 31:2745-7.

120.   Petkau A, Stuart-Edwards M, Stothard P, Van Domselaar G. 2010. Interactive microbial
genome visualization with GView. Bioinformatics 26:3125-6.
121.   Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image Generator (BRIG):
simple prokaryote genome comparisons. BMC Genomics 12:402.
122.   Rajagopal L. 2009. Understanding the regulation of Group B Streptococcal virulence factors.
Future Microbiol 4:201-21.
123.   Andrews JM. 2001. Determination of minimum inhibitory concentrations. J Antimicrob
Chemother 48 Suppl 1:5-16.
124.   Verotta D, Haagensen J, Spormann AM, Yang K. 2017. Mathematical Modeling of Biofilm
Structures Using COMSTAT Data. Comput Math Methods Med 2017:7246286.
125.   Hall BM, Ma CX, Liang P, Singh KK. 2009. Fluctuation analysis CalculatOR: a web tool for the
determination of mutation rate using Luria-Delbruck fluctuation analysis. Bioinformatics
25:1564-5.
126.   Zheng Q. 2017. rSalvador: An R Package for the Fluctuation Experiment. G3 (Bethesda)
7:3849-3856.

 Supporting Information Legends

Supplementary Figure 1. *Streptococcus iniae* pangenome derived from 80 strains, coloured by clade
(clade A-G innermost - outer) showing positions of MMR and OG genes relative to putative
phage/prophage and insertion sequence elements. Phage positions were predicted using Phaster and
are indicated as arcs in the outermost ring (green, complete phage; blue, incomplete phage; red,
uncertain). ISE were located with ISfinder and are indicated directionally in the outermost ring as black
arrows.

Supplementary Figure 2. Root-to-tip regression analysis (TempEst, (79)of branch lengths from best-fit-
rooted tree against time (year of isolation). The tree was derived from alignment of non-recombinant
core-genome SNPs corrected for ascertainment bias in RAxML.

Supplementary Figure 3: Alignment of the sequences found upstream from *mutY* gene in QMA0084
(representative of clade B) and QMA0248 (representative of all other analysed *S. iniae* strains) and
predicted changes in transcription pattern produced by 56 bp deletion upstream of *mutY* gene
identified in clade B strains.

Supplementary Table 1. 80-by-80 matrix of *p* values resulting from pair-wise comparisons of
fluctuation analysis data by Maximum Ratio Test corrected for multiple comparisons by controlling
for the False Discovery Rate (Microsoft Excel spreadsheet available as download).

Supplementary Table 2. Results of fluctuation analysis (Microsoft Excel spreadsheet available as
download).

Supplementary Table 3. Extended metadata for 80 *Streptococcus iniae* isolates sequenced in the
present study

Supplementary Table 4. Assembly statistics and quality data for 80 *de novo* genome assemblies
prepared in this study.

Supplementary Table 5. Recombination analysis (Gubbins) statistics for 83 *S. iniae* strains, ordered by
Clade.

Figures and tables

Figure 1: Unrooted summary phylogram of *Streptococcus iniae* strains based on alignment of core
genome SNPs filtered to remove recombination. Genetic distances were inferred by maximum
likelihood in RAXML and node support is indicated as percentage of 1000 bootstrap replicates.
Geographic origin, time range (year) of isolation, and mutation rate were added manually *post hoc*. All
nodes with bootstrap support below 75% were collapsed. The phylogram is drawn with scale

proportionate to genetic distance except the dashed branch supporting QMA0141, which is depicted at

10% scale. The inset shows the phylogram structure with QMA0141 branch drawn to the same scale.

Figure 2: Cladogram of *Streptococcus iniae* strains derived from the same dataset as Figure 1. Clade,

host species, variants in MMR and OG DNA repair genes, mutation rate phenotype, and phenotypic

variants associated with virulence were annotated with Evolview. Mutation rate (μ) phenotypes,

determined by fluctuation analysis, are depicted as box blots representing upper and lower 95%

confidence limits, with the interface representing mean values. Node support is indicated as percentage

of 1000 bootstrap replicates.

Figure 3: Phenotypic variants associated with virulence among *Streptococcus iniae* strains. A) Buoyant

density assay for capsule (CPS) presence, A1: CPS+, A2 CPS- A3: CPS+/-. B) Haemolysis of sheep red

blood cells by agar diffusion B1: +, B2:-. C) Streptococcal chain length C1: normal, C2: Long, C3: short. D)

Biofilm formation in chamber slides determined by confocal microscopy, D1: normal, D2: denser, D3:

thicker.

Figure 4: Proportions of single nucleotide polymorphism type across the core genomes of *Streptococcus*

iniae clades. Numbers to the right of the plot indicate the total numbers of SNPs that are unique to

each clade.

Table 1. *Streptococcus iniae* strains used in this study. Includes origin details (host species, time, site
of isolation), phylogenetic affiliation, virulence-associated phenotypes, and mutation rate. Atypical
places of isolation (hosts, tissues) and deviant phenotypes are in bold.

Strain	Clade	Host	Year	Location	Capsule	Hemolysis	Chains	Oxidation resistance	Biofilm formation	Mutation rate (10^{-7})
QMA0071	A	*Lates calcarifer*	2000	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2041
QMA0074	C2	*Lates calcarifer*	1998	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.6579
QMA0077	C2	*Lates calcarifer*	1995	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.7067
QMA0078	A	*Lates calcarifer*	2001	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2153
QMA0080	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1966
QMA0082	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1761
QMA0083	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1894
QMA0084	B	***Pteropus alecto***	2001	WA	wildtype	**weak**	wildtype	wildtype	wildtype	0.7293
QMA0087	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1925
QMA0130	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.468
QMA0131	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4488
QMA0133	E1	***Homo sapiens***	2001	USA	**absent**	wildtype	wildtype	wildtype	**increased**	0.5539
QMA0134	E1	***Homo sapiens***	2001	USA	**differential**	wildtype	wildtype	wildtype	**increased**	0.5457
QMA0135	E1	***Homo sapiens***	2002	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.631
QMA0137	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5495
QMA0138	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5989
QMA0139	F	*Fish sp.*	1996	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4311
QMA0140	-	***Inia geoffrensis***	1976	USA	**absent**	wildtype	wildtype	**increased**	**increased**	0.0901
QMA0141	-	***Inia geoffrensis***	1978	USA	**absent**	wildtype	wildtype	**increased**	**increased**	1.0884
QMA0142	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2044
QMA0150	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1952
QMA0155	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1493
QMA0156	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1666
QMA0157	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1545
QMA0158	A	*Lates calcarifer*	2006	SA RAS	**absent**	wildtype	**long**	wildtype	wildtype	0.1678
QMA0159	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2054
QMA0160	A	*Lates calcarifer*	1999	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1481
QMA0161	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1742
QMA0162	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0163	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1523
QMA0164	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7102
QMA0165	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6606
QMA0177	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0180	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2011
QMA0186	D	*Oncorhynchus mykiss*	2000	Israel	wildtype	**weak**	wildtype	wildtype	**increased**	0.5135
QMA0187	-	*Channa striata*	1983	Thailand	**absent**	**weak**	wildtype	wildtype	wildtype	0.4635
QMA0188	D	*Oncorhynchus mykiss*	1998	Israel	wildtype	**weak**	wildtype	wildtype	**increased**	0.5042
QMA0189	D	*Oncorhynchus mykiss*	1996	Reunion	wildtype	**weak**	wildtype	wildtype	wildtype	0.4907

QMA0190	F	*Channa striata*	1988	Thailand	**absent**	wildtype	wildtype	wildtype	wildtype	0.6413
QMA0191	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1883
QMA0207	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1486
QMA0216	A	*Lates calcarifer*	2007	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.1949
QMA0218	C2	*Lates calcarifer*	2007	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7093
QMA0220	A	*Lates calcarifer*	2006	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2086
QMA0221	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1814
QMA0222	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1829
QMA0233	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.362
QMA0234	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.354
QMA0235	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.3496
QMA0236	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.3413
QMA0244	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2101
QMA0245	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1988
QMA0246	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0247	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2059
QMA0248	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1893
QMA0249	F	*Lates calcarifer, bone*	2009	SA RAS	**differential**	**weak**	**long**	wildtype	**increased**	0.3817
QMA0250	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.195
QMA0251	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1927
QMA0252	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2006
QMA0253	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**increased**	0.3555
QMA0254	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**increased**	0.3712
QMA0258	A	*Lates calcarifer*	2008	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.1876
QMA0371	A	*Scortum barcoo*	2011	NSW	wildtype	wildtype	wildtype	wildtype	wildtype	0.1889
QMA0373	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6908
QMA0374	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.729
QMA0445	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0446	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.1687
QMA0447	E1	*Hybrid striped bass*	1996	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5902
QMA0448	E1	*Hybrid striped bass*	1998	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5529
QMA0457	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1682
QMA0458	A	*Epalzeorhynchos bicolor*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1778
QMA0462	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.4242
QMA0463	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.414
QMA0466	E2	*Oreochromis sp.*	-	USA	wildtype	wildtype	**short**	wildtype	wildtype	0.4602
QMA0467	A	*Epalzeothynchos frenatum*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1695
QMA0468	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1689
QMA0490	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1812

QMA0491	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.2
QMA0492	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1849
QMA0493	A	*Oreochromis sp.*	2016	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1987

972 Table 2: Variants in MMR and OG genes found among *S. iniae* isolates. CDS – protein coding

973 sequence, BS – predicted binding sites of transcription factors, PROVEAN score – predicted effect of

974 amino acid substitutions on the protein function; scores lower that -2 are considered to indicate a

975 deleterious effect.

Gene	Clade	Strains	Location	Mutation type	Nucleotide change	Amino acid change	Protein functional domain	PROVEAN score
mutS	F	from bone	CDS	SNP	G556T	Q -> K	Pfam MutS_II	-0.635
	E	all	CDS	SNP	T594C	-	Pfam MutS_II	n/a
mutL	F	from bone	CDS	SNP	C837A	-	DNA mis_repair	n/a
	F	all	CDS	SNP	T955G	S -> R	DNA mis_repair	-2.050
	B	all	CDS	SNP	C1117T	V -> I	-	-0.271
	F	QMA0190	CDS	SNP	A1626T	-	MutL_C	n/a
dnaN	F	QMA0139	promoter, BS of DnaA	SNP	C->T 116 bp upstream	n/a	n/a	n/a
	G	QMA0187	promoter, BS of FNR	SNP	T->C 135 bp upstream	n/a	n/a	n/a
	C2	all strains	CDS	SNP	C977T	T -> I	Pol3Bc	-2,216
	C1	all strains	stop codon	SNP	T1135C	STOP>Q	n/a	-0.005
rnhC	-	QMA0187	CDS	SNP	G778A	D ->N	Pfam: RNase_HII	-4.593
recD2	-	QMA0140	RBS	SNP	A -> G 15 bp upstream	n/a	n/a	n/a
	E1	all strains	RBS	SNP	T -> A 9 bp upstream	n/a	n/a	n/a
	B	all strains	CDS	SNP	C1605T	-	-	n/a
	C2	all strains	CDS	SNP	C1653T	-	-	n/a
mutY	B	all strains	promoter, -35 box, 1st bp of -10 box, 2 BSs of RpoD 17	large deletion (56 bp)	atcataagatttcctcctat aaatctgtgtcatatgtcat ccttctttatagtttc 247-187 bp upstream	n/a	n/a	n/a
	F	all strains	CDS	SNP	T935G	I -> T	Pfam: NUDIX_4	-2.544
mutM	D	all strains	CDS	SNP	T24G	E -> D	Fapy_DNAglyco	-2.990
mutX	E2	all strains	CDS	SNP	A193G	M -> I	Pfam: NUDIX	-0.539