Article

Mutators as drivers of adaptation in pathogenic bacteria and a risk factor for host jumps and vaccine

escape

Oleksandra Silayeva, Jan Engelstaedter, and Andrew C Barnes[*]

The University of Queensland, School of Biological Sciences, St Lucia Campus, Brisbane, Queensland

4072, Australia

Key words: mutators; DNA repair; pathogen evolution; immune escape, vaccination; Streptococcus;

aquaculture; host jumps, zoonoses

*Correspondence:

a.barnes@uq.edu.au

Abstract

Heritable hypermutable strains deficient in DNA repair genes (mutators) facilitate microbial adaptation

as they may rapidly generate beneficial mutations. Bacterial mutators deficient in mismatch (MMR) and

oxidised guanine (OG) repair are abundant in clinical samples and show increased adaptive potential in

experimental models of infection in mice. However, their role in pathoadaptation is poorly understood.

Here we investigate the role of mutators in epidemiology and evolution of the rapidly evolving aquatic

pathogen, *Streptococcus iniae*, employing a collection of 80 strains isolated globally over 40 years. We

determine phylogenetic relationship among *S. iniae* using 10,267 non-recombinant core genome single

nucleotide polymorphisms (SNPs), estimate their mutation rate by fluctuation analysis, and detect

variation in major MMR (*mutS*, *mutL*, *dnaN*, *recD2*, *rnhC*) and OG (*mutY*, *mutM*, *mutT*) genes. We find

that *S. iniae* mutation rate phenotype and genotype are strongly associated with phylogenetic

diversification and variation in major streptococcal virulence determinants (capsular polysaccharide,

hemolysin, cell chain length, resistance to oxidation, and biofilm formation). Furthermore, profound

changes in virulence determinants observed in mammalian isolates (atypical host) and vaccine-escape

isolates found in bone (atypical tissue) of vaccinated barramundi are linked to multiple MMR and OG

variants and unique mutation rates. This implies that adaptation to new host taxa, new host tissue, and

to immunity of a vaccinated host is promoted by mutator strains. Our findings support the importance of

mutation rate dynamics in evolution of pathogenic bacteria, in particular adaptation to a drastically

different immunological setting that occurs during host jump and vaccine escape events.

Introduction

Mutation rates in microbial populations are determined by a trade-off between maintenance of genetic integrity and evolvability (Stich, et al. 2010; Ferenci 2015). High fidelity of DNA repair is advantageous in adapted populations since most mutations lead to deviation from the evolved phenotype (Stich, et al. 2010; Sniegowski and Raynes 2013). Conversely, lower fidelity of DNA repair can be advantageous under stressful conditions since increased mutation supply promotes diversity and may thereby accelerate adaptation (Stich, et al. 2010; Wielgoss, et al. 2013). Bacteria can efficiently fluctuate between high and low mutation regimes: While extremely low mutation rates are maintained in favourable conditions (Wielgoss, et al. 2011), elevation of mutation rates known as stress-induced mutagenesis (SIM) occurs in adapting populations (Galhardo, et al. 2007; Sundin and Weigand 2007). One aspect of SIM is temporary hypermutability associated with changes in expression of multiple genes including error-prone DNA polymerases, recombination-preventing enzymes and enzymes involved in movement of mobile elements (Foster 2007). The most extensively studied example of transient hypermutability is the RecA-mediated general stress response, SOS, associated with up-regulation of error-prone polymerases that can by-pass DNA lesions during replication (Foster 2007). However, it is recognised that increased fitness during SOS response stems from immediate growth advantage rather than from long-term evolution of resistance (Wrande, et al. 2008; Andersson and Hughes 2010; Torres-Barcelo, et al. 2015). Another aspect of SIM, and the focus of the present study, is the evolution of heritable mutator phenotypes, also referred to as constitutive, permanent, or true mutators, that contain mutator alleles and replicate over multiple generations with a high amount of error (Galhardo, et al. 2007; Sundin and Weigand 2007). Heritably hypermutable strains, which periodically arise within a population via mutations in DNA repair genes (Miller 1996; Boe, et al. 2000), are selected under stress indirectly via 'hitchhiking' of mutator alleles with stress-resistance mutations (Mao, et al. 1997; Taddei, et al. 1997), or potentially yet-to-be established pleiotropic effects conferred by inactivation of DNA repair (Torres-Barcelo, et al. 2013; Raynes and Sniegowski 2014). There is much stronger theoretical and experimental evidence demonstrating that heritable mutators facilitate adaptive evolution (Raynes and Sniegowski 2014). However, even heritable mutators are transient phenomena in natural populations as non-mutator variants are selected within already adapted mutator clones (de Visser 2002; Turrientes, et al. 2013). Restoration of mutation rate can occur via copy-number variation leading to full reversion of the mutator allele (Shaver and Sniegowski 2003), disruption of the linkage between mutator alleles and beneficial mutations via recombination (Denamur, et al. 2000), anti-mutator compensatory mutations (Turrientes, et al. 2013; Wielgoss, et al. 2013), or prophage excision when its integration confers increase in mutation rate (Scott, et al. 2008).

Mutator strains are abundant among bacterial clinical isolates, especially in chronic cases (Hall and Henderson-Begg 2006; Oliver 2010). Multiple studies have attributed this to antibiotic selection (Negri,

et al. 2002; Matsushima, et al. 2010; Wang, et al. 2013), since mutators are a well-established risk factor for the development of antimicrobial resistance (Blazquez 2003; Chopra, et al. 2003). However, the association between frequency of mutators in diagnostic samples and antibiotic treatments is weak (Gutierrez, et al. 2004; Mena, et al. 2008), and isogenic knockout mutator strains have increased ability to colonize an animal host in challenge models devoid of antibiotic exposure (Giraud, et al. 2001; Nilsson, et al. 2004; Healey, et al. 2016). Thus, the prevalence of mutators in clinical isolates most likely results from the selective process during adaptation to the host (Labat, et al. 2005; Mena, et al. 2007; Mena, et al. 2008; Oliver and Mena 2010), where immunity acts as the ultimate selective pressure shaping the diversity of pathogenic strains (Aquino and Nunes 2016).

The 'immunological niche' concept states that, due to the diversity and complexity of immune responses, every individual host represents a unique habitat that requires adaptation (Cobey 2014). Considering that the pathogen needs to adapt to every new host, the host population taken as a whole represents an extremely heterogeneous environment where adaptive processes are ongoing and persistence of mutator alleles may be favoured (Lukacisinova, et al. 2017). Further, as a small number of infection units may enter a host, pathogen adaptation is intrinsically associated with population bottlenecks during which the rate at which beneficial variants arise might be crucial (Metzgar and Wills 2000; Abel, et al. 2015). This supports the contention that bacterial evolution is more likely to occur via multiple mutations rather than a single beneficial variant (Fogle, et al. 2008) The fitness of a pathogenic strain in the host hinges on the concept of virulence (the degree of damage to the host), which is a dynamic, complex trait determined by a multitude of virulence factors and host responses (Methot and Alizon 2014). Optimal virulence is a trade-off between acuteness and persistence with higher virulence facilitating extraction of resources and transmission, but too much damage can kill the host and stop transmission altogether (Mackinnon and Read 2004). Additionally, virulence factors are often antigens so reduced virulence may be a means of immune evasion (Deitsch, et al. 1997). Consequently, both loss and gain of virulence may be advantageous, and indicate adaptive shifts towards acute or chronic pathogenesis, respectively (Maurelli 2007; Shrestha, et al. 2014). Exacerbation and attenuation of virulence factors are repeatedly observed in both knockout and naturally occurring mutators (Picard, et al. 2001; Merino, et al. 2002; Smania, et al. 2004; Mena, et al. 2007; Gonzalez, et al. 2012; Canfield, et al. 2013). Thus, it appears that mutator strains/alleles may help pathogenic bacteria to evolve towards the optimal level of virulence.

Most bacterial mutators isolated from human clinical samples are deficient in genes from mismatch (MMR) and oxidised guanine (OG) DNA repair systems (Hall and Henderson-Begg 2006). MMR and OG genes are conserved across all domains of life (Fukui 2010), and single nucleotide polymorphisms in their sequence may produce drastic alterations in mutation rate (Rajanna, et al. 2013; Dai, et al. 2015). Nonetheless, these genes appear to be variable in major bacterial pathogens, confirming a potential role

of mutation rate dynamics in the evolution of virulence (Ambur, et al. 2009; Chen, et al. 2010). MMR is

coupled to replication and removes mispaired nucleotides as well as short insertion/deletion loops, and

prevents recombination between divergent sequences (Li 2008; Tham, et al. 2013).The MMR pathway

has been extensively characterised in *Escherichia coli,* where the MutS dimer protein interacts with the

replication processivity clamp DnaN (b-clamp), binds to the mismatch, recruits MutL, and the MutS:MutL

complex then activates the MutH endonuclease (Fukui 2010). MutH cuts the unmethylated nascent

strand at hemimethylated d(GATC) sites. MutL loads the UvrD helicase that unwinds the nicked DNA

containing the error, which is followed by excision by one of the single-strand nucleases (Fukui 2010).

This MMR organization, however, appears to be limited to *E. coli* and closely related

Gammaproteobacteria (Fukui 2010; Lenhart, et al. 2016). In most bacteria and eukaryotes, MutH is

absent and MutL has a conserved endonuclease site, dam methylase (acting at GATC sites) is absent and

the strand discrimination signal is unknown (Fukui 2010; Lenhart, et al. 2016). Studies in *Bacillus subtilis,*

a model species for gram-positive bacteria, suggest that nicks created by ribonuclease *rnhC* during

removal of misincorporated ribonucleosides direct MutL endonuclease towards the newly synthethised

strand, and that recD2 acts as a MMR helicase instead of uvrD (Lenhart, et al. 2016). The OG system

repairs oxidative DNA damage throughout the cell cycle (Lu, et al. 2001; David, et al. 2007). Among all

DNA bases, guanine is most sensitive to reactive oxygen species and its oxidised form, 8-oxo-dG or OG,

is highly mutagenic as it preferentially pairs with adenine which, if not corrected, leads to G:C to A:T

transversion on a second round of replication (Lu, et al. 2001; David, et al. 2007). OG system organization

is conserved in most bacteria and eukaryotes and generally MutT homologs remove 8-oxo-dG from the

nucleotide pool, MutM homologs excise 8-oxo-dG incorporated into DNA, and MutY homologs remove

adenine from OG:A mispairings (Lu, et al. 2001; David, et al. 2007).

The present study investigates the role of mutators and mutation rate dynamics in the epidemiology and

evolution of pathogenic bacteria using the broad host-range pathogen *Streptococcus iniae*. *S. iniae* is a

highly adaptable species, evidenced by its global distribution, ability to cross divergent host taxa

boundaries and repeated re-infection of previously immunised hosts during disease outbreaks (Agnew

and Barnes 2007; Millard, et al. 2012). Moreover, Streptococcal pathogens in general are difficult

candidates for immunisation in humans and animals due to high strain diversity evidently driven by

immunisation, yet there is little information on the role of mutators in this evolutionary process

(Croucher, et al. 2011; Croucher, et al. 2013). We determine variation in mutation rate and identify

mutations in MMR (*mutS*, *mutL*, *dnaN*, *recD2*, *rnhC*) and OG (*mutY*, *mutM*, *mutT*) genes among 80 diverse

strains of *S. iniae.* We discover that both mutation rates and mutations in MMR and OG genes correlate

with phylogenetic diversification but are highly conserved within phylogenetic clades. To determine

whether this adaptive process might be driven by the host immune response we identify variation in

major phenotypic traits that determine virulence in streptococci (capsular polysaccharide, hemolysin, length of cell chains) and bacteria in general (resistance to reactive oxygen species (ROS), and biofilm formation). *S. iniae* polysaccharide capsule and the beta-hemolysin streptolysin S both increase survival under host immune response via reduction of phagocytic killing and other mechanisms (Locke, Colvin, Datta, et al. 2007; Locke, Colvin, Varki, et al. 2007; Kadioglu, et al. 2008; Rajagopal 2009). However, both are antigens that induce antibody production and development of immune memory (Kadioglu, et al. 2008; Rajagopal 2009). In addition, both capsule production and hemolysin activity have many implications that are not fully understood (Kadioglu, et al. 2008; Geno, et al. 2015). For example, non-encapsulated strains exhibit greater ability to adhere to and colonize an epithelium (Hammerschmidt, et al. 2005). Length of cell-chains is also linked to virulence in streptococci: shorter chains are associated with acute infection and increased survival in the blood as they are less likely to activate the complement, whilst longer chains promote adherence and colonization (Rodriguez, et al. 2012). As pathogenic bacteria are continuously exposed to reactive oxygen released by immune cells, resistance to oxidative compounds contributes to virulence in pathogens allowing them to avoid immune clearance (Hegde, et al. 2008; Andisi, et al. 2012). Biofilm formation is also generally associated with virulence in bacteria; detached cells are associated with acute infections whereas biofilm aggregates are a hallmark of chronic conditions (Bjarnsholt 2013). The expression of these traits among *S. iniae* isolates is highly consistent with phylogenetic affiliation, which supports that strain diversity is primarily driven by adaptation to the host achieved via adjustment of virulence. Furthermore, we find that occurrence of abnormal virulence determinants is strongly associated with variation in MMR and OG genes and occurrence of infection in unusual immunological landscapes such as mammals and bone tissue of immunised fish. These data support the contention that mutators play a critical role in adaptation to the host and evolution of virulence, and suggest that these strains may facilitate the reinfection of immunised animals and transmission between divergent host species.

Results and Discussion

Our work investigates the link between mutation rate genotype and phenotype and phylogenetic diversification among *Streptococcus iniae* strains isolated globally between 1976 and 2016 from different host taxa. Maximum likelihood phylogenetic analysis of 80 *S. iniae* isolates (Table 1) based on non-recombinant core genome SNPs derived from whole genome data resolved 6 major clades (A-F), one lineage with two strains (clade G), and three lineages with a single strain (Figs. 1,2). Clades C-E are endemic; and clades A, B, and F contain strains from diverse geographic regions. Principally, variation in MMR and GO genes occurs only between the lineages, and they are highly conserved within clades (Fig. 2, Table 2). In fact, sequences of the major MMR and OG genes are conserved even within phylogenetic groups not readily definable by geographic origin, time, or host species, and these genes appear to be good candidates for Multilocus Sequence Typing (MLST), i.e. identification of phylogenetic position based on selected genes as opposed to a whole genome analysis (Maiden, et al. 2013). Although even synonymous SNPs in any gene locus can affect mRNA and have pronounced phenotypic consequences (Shabalina, et al. 2013), most SNPs in MMR and OG genes were identified in protein functional domains, and most amino acid substitutions were predicted to have a deleterious effect on protein function (Table 2). Further, the number of MMR and OG variants correlates significantly with mutation rate (p = 0.0056) and with the number of atypical phenotypic traits contributing to virulence (p = 0.0331), as predicted by a Phylogenetic Generalised Least Squares (PGLS) regression model that accounts for autocorrelation occurring between closely related strains (Symonds and Blomberg 2014). In turn, mutation rate is highly consistent with phylogenetic affiliation and, according to multiple pair-wise strain comparisons by Maximum Likelihood Ratio test (Zheng 2015), with probabilities corrected for multiple comparison using False Discovery Rate (Benjamini and Hochberg 1995), mutation rate difference is insignificant within but significant between major phylogenetic lineages (Suppl. Table 1).

Although mutation rate variation between divergent strains is significant, the magnitude of the differences is modest (largely around 5-fold) (Figure 1, 2, Suppl. Table 2), which is consistent with mutator phenotype transience and restoration of low mutation rate in adapted populations. On the other hand, only one order of magnitude or lower differences in mutation rate distinguishes mutator from non-mutator phenotypes in streptococci (around 10^{-7} and 10^{-8} mutation rates respectively) (Hall and Henderson-Begg 2006; Gould, et al. 2007), in contrast to other common pathogens where 2-3 orders of magnitude differences are observed (Hall and Henderson-Begg 2006). Heritably hypermutable phenotypes are costly due to inevitable accumulation of deleterious changes. Once adaptation is gained the mutator strain is predicted to evolve back towards a lower mutation rate (de Visser 2002; Raynes and Sniegowski 2014; Sprouffske, et al. 2018). This may occur via compensatory, 'anti-mutator' mutations in DNA repair genes (Turrientes, et al. 2013; Wielgoss, et al. 2013), or via recombination (Denamur, et al.

2000). Our data favour the former mechanism for several reasons. Firstly, we used a maximum likelihood

inference method to exclude areas of recombination from our phylogenetic analyses (Croucher, Page, et

al. 2015), and the MMR and OG gene regions were not identified as recombinant regions in this analysis.

Secondly, we created a pangenome for the 80 strains in our analysis and mapped on putative phage and

prophage signatures (Arndt, et al. 2016), and insertion sequence elements (Siguier, et al. 2006). None of

the MMR or OG genes, which are all present within the core genome of *S. iniae*, occurred in proximity to

phage or ISE (Suppl. Fig. 1). Whilst we cannot rule out switch by recombination, these data suggest that

accrual of SNPs via random mutation is a more likely explanation.

To investigate whether mutation rate phenotype and inferred phylogenetic association with MMR/OG

genotype was also associated with virulence phenotype, we investigated key streptococcal virulence

traits amongst the bacterial isolates in the analysis. We observe significant correlation (p = 0.0331, PGLS)

of the number of MMR and OG variants with number of atypical phenotypes for traits contributing to

virulence in streptococci (capsular polysaccharide, hemolysin, length of cell chains) and bacteria in

general (resistance to ROS, and biofilm formation). This suggests that fluctuations in mutation rate may

be linked to shifts in virulence, potentially allowing major adaptive events and leading to phylogenetic

diversification (Fig. 2, Table 3). Moreover, infections in atypical hosts (mammals) and atypical tissue

(bone) within apparently immune vaccinated hosts correlates strongly with number of atypical virulence-

associated phenotypes (p = 0 .000) and the number of variants MMR and OG variants (p = 0.002). This

suggests that mutator strains may facilitate multiple genetic changes leading to profound shifts in

virulence and antigenicity during adaptation to a drastically different immunological setting.

Of the clades identified within our collection of *S. iniae* isolates, Clade A is apparently a dominant

circulating clade as it has persisted globally for almost two decades (Clade A contains strains isolated

from USA, Honduras, and Australia between 1999 to 2016), primarily infecting closely related Perciform

fish (barramundi (*Lates calcarifer*), tilapia (*Oreochromis sp*), jade perch (*Scortum barcoo*)). Mutation rates

in all isolates among this ancestral lineage (35 strains analysed) fall within a 1.5 - 2 x 10^{-8} range (Figs. 1, 2;

Clade A), which is similar to the base mutation rate estimated for other non-mutator gram positive

bacteria (Hall and Henderson-Begg 2006; Gould, et al. 2007). All differences in mutation rates within this

clade are insignificant except three pairwise comparisons that verge on p <0.05 (Suppl. Table 1). MMR

and OG genes in these strains are identical, and only minor deviations from wildtype traits contributing

to virulence were observed among the strains (absence of capsule production and longer cell chains in

QMA0158, 216) (Figs. 1-3, Table 1). All of the other lineages (with the exceptions of clade C1 and strains

QMA0445-46 discussed below) express significantly different mutation rates to clade A, contain unique

and often multiple SNPs in *mut* genes, and show peculiar phenotypic variants related to virulence (Figs.

1-3, Tables 1,2, Suppl. Table 1). Although phylogenetic relatedness between lineages is fully resolved (Fig.

2), they originate almost simultaneously and nested lineages evolve independently sharing little phylogenetic history (Figs. 1,2). Lack of correlation between branch length and sampling date derived by root-to-tip regression analysis with time is good evidence of non-neutral evolution amongst this collection of *S. iniae* isolates (R^2=0.0936, Correlation coefficient 0.306, best-fit root; Suppl. Fig. 2, (Rambaut, et al. 2016)). Deviation from a neutral model is also supported by highly skewed branching in the tree, with some branches comprising single isolates whilst others comprise many epidemiologically unrelated isolates (Figure 1, 2). This is suggestive of frequent strong selection, presumably imposed by heterogeneity of the immune landscape encountered during transfer between host individuals, and resembles genealogical tree topology derived from viral evolution over similar timespans (Bedford, et al. 2011; Neher and Hallatschek 2013) although we acknowledge tree topology may also be affected by sampling bias.

Clade B contains QMA0084, a strain isolated from a black flying fox, *Pteropus alecto*, in Western Australia (WA) and two strains isolated from ornamental fish (clown loach or tiger botia, *Chromobotia macracanthus*) in USA. These isolates share a valine to isoleucine substitution in MutL predicted as neutral (PROVEAN score -0.271), a synonymous SNP in *recD2*, and a 56 bp deletion affecting the predicted promoter of *mutY* (Table 2). The latter mutation encompasses the -35 box, two binding sites of the RpoD17 general transcription factor, and the 1^{st} bp of the -10 box, and was predicted to abolish the original promoter (Suppl. Fig. 3). However, *mutY* might still be transcribed at some level since two other potential promoters were identified in a nearby sequence (Suppl. Fig. 3). It appears that altered *mutY* expression is also affected by the rest of the genetic background as isolates exhibit significant differences (Suppl. Table 1) in mutation rate phenotype, 7×10^{-8} in QMA0084 and 4×10^{-8} in clown loach strains (Figs. 1-2). Phenotypically, clade B has shifted towards decreased haemolytic activity (Fig. 3, B2).

Clade C shares a common ancestor with clade B and is comprised of strains obtained from barramundi farmed in WA, Northern Territory (NT) and north QLD fish farms (nested subclades C1 and C2 respectively). While a core mutation rate of 1.5-2×10^{-8} is observed in subclade C1, a significantly higher mutation rate of 7×10^{-8} has been maintained in subclade C2 from north QLD for almost two decades (1995 to 2012) (Figs. 1,2 Suppl. Table 1). We speculate that the latter might be attributable to difference in meteorological conditions: periodic heavy rainfall present in tropical north QLD is likely to bring fluctuations in water salinity and temperature, creating an unstable environment where adaptive processes are ongoing and high mutation rate is advantageous. Minor variation in virulence-related phenotypes (absence of capsule production and longer cell chains in QMA0074, 77)(Fig. 2, Table 1), are supportive that mutation rate evolution in this lineage might be driven by outside-the-host factors. The significant difference in mutation rate between nested clades C1 and C2 is potentially attributable to variation in *dnaN*. In *Bacillus subtilis*, 90% of mismatch repair is dependent on targeting MutS to nascent

DNA via the ß sliding clamp (DnaN) zone (Dupes, et al. 2010; Lenhart, et al. 2013). Inactivation of *dnaN1*

in *B. anthracis* results in a mutator phenotype with a mutation rate equivalent to a *mutS* mismatch repair-

defective strain (Yang and Miller 2008). The tyrosine to isoleucine substitution in one of the critical

residues of the loader binding interface of the ß-clamp, predicted as deleterious (PROVEAN score -2,216),

is present in all isolates of from clade C (Table 2), which may explain the high mutation rate phenotype

in clade C2 strains (Fig. 2, Table 2). In contrast, only subclade C1 harbours T1135C nucleotide substitution

in the *dnaN* that changes stop codon TAA into CAA coding for glutamine predicted as neutral (PROVEAN

score -0.005), with the next stop codon TAG found immediately downstream. Putatively, T1135C and/or

change of the *dnaN* stop codon (Korkmaz, et al. 2014) may act as compensatory to the T326I substitution,

reducing the mutation rate to a value not significantly different from the core rate found in clade A (Fig.

1, 2) via protein elongation or by affecting translation via stop codon usage bias (Korkmaz, et al. 2014).

In addition, a synonymous variant in *recD2* that is only present in subclade C2 and might affect protein

expression, may contribute to the difference in mutation rate between C1 and C2 clades.

Clade D consists of isolates from trout (*Oncorhynchus mykiss*) from Réunion and Israel. These strains have

an estimated mutation rate of 5×10^{-8}, contain a glutamate to aspartate substitution in *mutM* predicted

to be deleterious to protein function, a synonymous SNP in *RecD2*, and exhibit impeded haemolytic

activity (Fig 1,2, Table 2). Also, isolates from Israel form thicker and more dense biofilm structures (Fig.

3, D2).

Isolates from humans and fish are found in clade E where multiple phenotypic variants contributing to

virulence are observed (Figs. 2-3, Table 1). This clade contains two nested subclades that share a SNP in

mutS, but have other unique SNPs (Fig. 2, Table 2) and exhibit mutation rates that are differently

significant in most pairwise comparisons (Suppl. Table 1). Subclade E1 contains USA isolates from humans

(QMA0133-35, 37-38) and hybrid striped bass (QMA0447-48), which have a substitution in the Shine-

Dalgarno sequence of *recD2* and a mutation rate of 6.5×10^{-8} (Fig 1, Table 2). Fish strains and three human

strains (QMA0135, 37-38) form shorter cell chains and higher resistance to oxidative stress. Two human

strains QMA0133-34 produce thicker and more dense biofilm structures (Fig. 3, D2). QMA0133 is non-

encapsulated (Fig 3, A2), and QMA0134 appears to differentially express the capsular polysaccharide in

culture (Fig 3, A3). The close phylogenetic relationship presented here implicates a likely transfer from

farmed hybrid bass to humans (Facklam, et al. 2005), and this host jump may have been facilitated by the

mutator phenotype. Subclade E2 contains two human strains from Canada (QMA0130-31) and a tilapia

strain from USA (QMA0466). These strains have unique methionine to isoleucine substitution at the N-

terminus of MutX, and a mutation rate of 4.5×10^{-8} (Fig 1, Table 2). Both human strains are non-

encapsulated (Fig. 2). The tilapia strain expresses the capsule but forms short cell chains in common with

most strains from the subclade (Fig 2, Fig 3, C3). The close phylogenetic relationship again strongly

implicates transfer from tilapia farmed in USA to human patients in Canada via import into the Toronto

fish market (Weinstein, et al. 1997) and again, this host jump may have been facilitated by the mutator

state.

Clade F is not readily definable by location, time of isolation, or host species and comprises QMA0139

from unknown fish species obtained in Canada, QMA0190 isolated from snakehead murrel in Thailand

(both of which appear on long branches), and a nested terminal clade comprised of barramundi isolates

farmed in Recirculating Aquaculture Systems (RAS) in New South Wales and South Australia (Fig. 2 1,2).

The latter are of a particular interest as these mutators were sampled from barramundi bone lesions

during a disease outbreak in vaccinated fish where infection manifested itself as a slowly progressing

osteomyelitis instead of typical acute septicaemia and meningitis, implicating mutators in adaptation to

new tissue in response to host immunity (Agnew and Barnes 2007; Millard, et al. 2012). Since they are

distant to strains used in autogenous vaccines in clade A (QMA0155-57, 160, QMA0250-52), the atypical

outbreak might be classified as a case of vaccine-induced serotype replacement (VISR) – spread of co-

existing pathogenic strain/s after elimination of the dominant strain/s targeted by the vaccine

(Weinberger, et al. 2011). In contrast to natural populations where the investigation of VISR is

confounded by ecological and sampling biases, and randomised vaccine trials that lack power to detect

the population-wide effect of mass vaccination (Weinberger, et al. 2011), barramundi RAS represent a

relatively controlled and isolated natural environment where vaccination against *S. iniae* fails occasionally

but recurrently (Millard, et al. 2012). RAS are self-contained facilities with minimal water exchange and

high standards of water treatment (Losordo, et al. 2009). This ensures that gene flow and, consequently,

the probability of new serotype introduction after vaccination is reduced to a minimum. A comparatively

uniform immunized host population is exposed to dominant strains against which it was vaccinated along

with potentially co-existing lineages in the RAS. In the atypical outbreaks in RAS in Australia, vaccine-

induced immunity was partially effective in the infected fish, evident by absence of proliferation of the

new strain in typical niduses (blood, brain, and pronephros) and cross-reactivity of vaccine-induced

antibodies against bone isolates (Millard, et al. 2012). This is in contrast with the classical model of

serotype replacement where a vaccine is ineffective against the replacing co-existing strain. It has been

postulated that VISR can occur even if the vaccine is equally efficient against all circulating strains via

other trade-off mechanisms (Weinberger, et al. 2011). Apparently, evasion of immune clearance and

adaptation to osseous tissue was permitted by shift towards attenuated virulence indicated by multiple

phenotypic changes in bone isolates, including absence of capsule, impeded hemolytic activity, increased

cell-chain length, and denser biofilms (Fig. 3). All of these changes have previously been associated with

chronic infection and increased potential for colonization (Locke, Colvin, Datta, et al. 2007; Locke, Colvin,

Varki, et al. 2007; Kadioglu, et al. 2008; Andisi, et al. 2012; Rodriguez, et al. 2012; Bjarnsholt 2013; Geno,

et al. 2015), and only absence of polysaccharide capsule is shared with related QMA139 and QMA190, as
well as QMA0140-41, QMA0187, and QMA0445-46, which implicates a non-encapsulated ancestor for
these strains. The latter acapsular strain potentially also gave rise to clade E, assuming capsular
production was regained in encapsulated strains found in this branch. Multiple SNPs in repair genes (Fig.
2, Table 2) and the unique mutation rate (Figs. 1-2, Table 1) in bone isolates suggest that shift in virulence
allowing persistence in the bone tissue might have occurred via mutator phenotype. Serine to arginine
substitution in MutL and tyrosine to isoleucine substitution in MutY occur in the ancestor of QMA0139,
QMA0190, and bone strains. Both substitutions are predicted to have a deleterious effect on protein
function (PROVEAN scores -2.050 and -2.544 respectively), but their effect on mutation rate is combined
with unique SNPs found in each branch: QMA0139 has a SNP in the binding site for the DnaA transcription
factor within the *dnaN* promoter, and a mutation rate of 4.5×10^{-8}, QMA0190 has synonymous SNP in
mutL and a mutation rate of 6.5×10^{-8}, and bone strains have glutamine to glycine substitution in MutS
predicted as neutral (PROVEAN score -0.635), a synonymous SNP in *mutL*, and a mutation rate of $3-4 \times$
10^{-8}. Considering that the mutation rate of QMA0190 is significantly higher compared to rates expressed
by QMA0139 and bone isolates (Suppl. Table 1), it is likely that variants unique to the latter strains
compensate for deleterious variants shared by the isolates (Wielgoss, et al. 2013).

Clade G contains two strains from tilapia isolated in USA (QMA0445-46). Notably, despite being
phylogenetically distant from clade A, these strains have the same MMR and OG genotype and have
retained the core mutation rate and the wildtype virulence phenotype (Fig. 2, Table 1), with the exception
of capsule absence, perhaps attributable to a common non-encapsulated ancestor shared with clade F,
QMA0140-41, QMA0187, and clade E.

Three isolates of *S. iniae* were identified as independent phylogenetic lineages, with unique mutation
rate genotypes and phenotypes and combinations of phenotypes associated with virulence (Figs. 1-2,
Tables 1-2). The first long branch with a single isolate contains a strain from snakehead murrel strain
(*Channa striata*) from Thailand. This isolate is non-encapsulated and weakly haemolytic and has a
mutation rate of 4.5×10^{-8}, potentially attributable to a deleterious aspartate to asparagine substitution
in rnhC and SNP in the binding site of *fnr* transcription factor within the *dnaN* promoter. A second long
branch with a single strain contains the oldest among the analysed strains, QMA0140 isolated in 1976
from dolphin (*Inia geoffrensis*) (Pier and Madin 1976). This strain exhibits a mutation rate of 1×10^{-8},
which is unique among the strains and significantly lower than the core mutation rate, presumably
associated with increased translation rate of recD2 helicase produced by a SNP in the ribosome binding
site (Fig 2, Table 2). The longest branch on the tree contains a second dolphin strain, QMA0141, isolated
two years later in 1978 (Pier, et al. 1978). This isolate is highly divergent from the rest of strains with
around 20 kB of non-recombinant SNPs in pair-wise comparisons with other strains, accounting for

around 1% genomic difference. In contrast to QMA140, it mutates at a rate of 1x 10^{-7}, the highest

mutation rate phenotype determined among the isolates and significantly different to the rest of the

values. Multiple SNPs are observed in all MMR and OG genes: 3 in *dnaN* and *mutX*, 4 in *mutM*, 9 in *mutL*

and *rnhC*, 15 in *recD2*, 9 in *mutL* and *rnhC*, and 68 in *mutS*. Both dolphin isolates are non-encapsulated,

show increased ability to withstand oxidative stress, and form denser and thicker biofilms (Fig 2, 3).

The conservation of *S. iniae* DNA repair genotypes and phenotype within phylogenetic clades and

variation between the lineages is consistent with the idea of adaptive fluctuations in mutation rate being

a driver of pathogen evolution, and their association with variation in virulence traits supports the

contention that adaptive processes and diversification of pathogens are largely driven by host immunity.

A number of genetic mechanisms have been proposed to potentially underlie the flux of mutator alleles

and phenotypes, including random mutations (Wielgoss, et al. 2013), tandem repeat copy-number

variation (Shaver and Sniegowski 2003; Chen, et al. 2010), non-homologous recombination (Denamur, et

al. 2000), and prophage integration/excision (Scott, et al. 2008). In *S. iniae,* mutation rate dynamics

appear to be primarily associated with random mutations and compensatory evolution. Tandem-repeat

copy-number variants were not identified in *S. iniae* MMR or OG genes, supporting the contention that

occurrence of mutator phenotypes and their full reversion via this mutational type might be limited to *E.*

coli and closely related species (Shaver and Sniegowski 2003; Chen, et al. 2010). Indeed, the presence of

tandem repeats has only been detected in *mutL* in *Proteobacteria* (Shaver and Sniegowski 2003; Chen, et

al. 2010), and it is well recognized that *mutL* is structurally and functionally different in gram-positive

bacteria and eukaryotes (Fukui 2010; Lenhart, et al. 2016). With the exception of the large deletion

affecting the *mutY* promoter in clade B, all identified mutations are single nucleotide substitutions. These

SNPs are not likely to result from gene transfer, as they were not identified amongst regions of

recombination, nor within proximity to phage or ISE (Suppl. Fig. 1). These MMR and OG nucleotide

substitutions most likely result from random mutations which can be sufficient to induce profound

changes in mutation rate (Rajanna, et al. 2013; Dai, et al. 2015). Since, in most cases, SNPs occur in

multiple genes (Fig. 2) it is likely that compensatory evolution of mutation rate prevails in *S. iniae* where

some variants increase and some decrease the background mutation rate (Wielgoss, et al. 2013).

The association of MMR and OG variation with changes in virulence traits (Fig. 2) implies that adaptive

processes in *S. iniae* population driven by host immune response may be facilitated by the emergence

and decline of mutator strains. Furthermore, most profound changes in virulence-associated traits and

MMR and OG variants are observed in atypical hosts (mammals) and atypical tissue (bone) within

apparently immune vaccinated fish hosts (Fig. 2), which has major epidemiological implications. First,

mutators may promote multiple adaptive changes required for pathogen transition among divergent host

taxa known as 'host jumps', such as *Osteichthyes* and *Mammalia* discussed in the present study (Baumler

and Fang 2013). Second, mutators might present a risk factor to immune escape after vaccination and
serotype replacement. Moreover, as evidenced by *S. iniae* colonization of bone tissue of vaccinated
barramundi described previously (Millard, et al. 2012), vaccine escape leading to serotype replacement
might occur without major disruption of immune recognition when multiple changes attenuating
virulence and antigenicity allow persistence in a tissue with lower immune surveillance.

We acknowledge that our results do not exclude the possibility that variation in DNA repair genotype and
phenotype may potentially represent neutral variation coinciding with shifts in virulence occurring in each
phylogenetic lineage. However, complete conservation of MMR and OG genes and mutation rate within
S. iniae lineages (Fig. 2) and the fact that most SNPs occur in functionally relevant coding and regulatory
regions and/or change amino acid sequence (Table 2), supports that the occurrence of mutator strains is
likely to precede and facilitate shifts in virulence leading to major diversification events and
epidemiologically relevant incidents such as vaccine escape outbreaks and host jumps. Our results
indicate that high mutation rate might promote transition to a new immunological setting requiring
multiple mutations and phenotypic changes, such as adaptation to a novel host, new host tissue, or a host
with vaccine-induced adaptive immune response. Extension to other species such as *Streptococcus*
pneumoniae may be particularly interesting as serotype evolution following vaccination programmes is
well documented, but the role of mutators in this process is not yet known (Croucher, et al. 2011;
Croucher, et al. 2013; Croucher, Kagedan, et al. 2015). Future studies should determine how generalisable
the role of mutators is in the epidemiology of bacterial pathogens, evolution of virulence and antigenicity,
and emergence of new pathogenic strains.

Materials and methods
Strains and growth conditions

Eighty isolates of *S. iniae* collected in Australia, USA, Canada, Israel, Honduras, and Thailand between
1976 and 2016 from eight fish species (*Lates calcarifer*, *Scortum barcoo*, *Epalzeorhynchos frenatum,*
Epalzeorhynchos bicolor, Oreochromis sp., Channa striata, Chromobotia macracanthus, Oncorhynchus
mykiss) and three mammalian species (*Homo sapiens, Inia geoffrensis, Pteropus alecto*) were used in this
study (Table 1). Strains were received from culture collections, veterinarians, or directly from fish farms
(Suppl. Table 1) and stored as master seed stocks without further subculture at -80°C in Todd-Hewitt
Broth (THB) supplemented with 20% glycerol. Bacteria were routinely recovered from -80°C stocks on
Columbia agar supplemented with 5% defibrinated sheep blood (Oxoid, Australia), and cultured at 28°C
on Todd-Hewitt agar or in Todd-Hewitt broth (Oxoid) with agitation 200 rpm in a shaking incubator unless
otherwise specified.

Estimation of mutation rate phenotype by fluctuation analysis

To estimate mutation rates of *S. iniae* isolates, a fluctuation analysis assay for spontaneous occurrence

of rifampicin resistance was optimized according to Rosche and Foster (Rosche and Foster 2000). A single

broth culture was initiated from five separate colonies, recovered on Columbia blood agar from stock

cultures stored at -80°C, and grown overnight to late-exponential phase in Todd-Hewitt broth (THB).

Cultures were adjusted to OD_{600} = 1 corresponding to 10^8 viable CFU/mL, diluted 1:100, and distributed

in 200 μl aliquots into 8 wells of a sterile U-bottom 96-well plate (Greiner). Prior to dilution of 100 μl of

overnight OD_{600}-adjusted cultures were spread onto THA supplemented with 0.5 μg/mL rifampicin to

confirm absence of rifampicin-resistant mutants. Although comparatively large initial inoculum (1-2 x 10^5

CFU per culture, confirmed by Miles and Misra CFU count (Hedges 2002) of OD-adjusted cultures) was

used, this reduced to the minimum variation in final CFU number (N_t) as shown by the preliminary

experiments. This approach was undertaken since final population size is critical for estimation of the

mutation rate and measuring N_t for such a large number of strains and replicated cultures was technically

prohibiting. Also, invariability in N_t allowed statistical comparison of mutation rates by the Maximum

Likelihood Ratio test (Zheng 2015). To infer the N_t in final cultures and monitor their growth by optical

density, replicate plates containing two cultures per strain were prepared and incubated in a BMG

FLUOstar OPTIMA microplate reader. When these representative cultures were reaching an early

stationary phase, CFU counts were performed by Miles and Misra method (Hedges 2002) and invariably

estimated as 1-2 x 10^8 CFU in each well. Immediately after, whole 200 μL cultures were plated on Todd-

Hewitt agar containing 0.5 μg/mL rifampicin for selection of mutants, dried in a laminar flow hood, and

incubated until rifampicin resistant colonies appeared. The assay was repeated four times for each strain

in blocks of eight cultures using 20 isolates haphazardly chosen from different phylogenetic lineages in

each measurement. Processing of all 80 strains at the same time was not only time-consuming, but also

compromised the uniformity of initial inoculum size due to growth of OD_{600}-adjusted cultures after the

addition of the fresh THB. In some cases, e. g. a single strain representing an independent phylogenetic

lineage or with significantly differences in mutation rates detected in closely related strains, assays were

repeated 1-2 more times to increase a statistical power. The rifampicin-resistant mutant counts were

pooled into single data sets representing 32 to 48 cultures per strain (Suppl. Table 2).

DNA extraction, preparation and sequencing

Genomic DNA was extracted from cells collected from 10 mL late-exponential phase culture in Todd-

Hewitt broth with the DNeasy Blood & Tissue kit (Qiagen) using a modified protocol with an additional

lysis step as described previously (Kawasaki, et al. 2018). gDNA was analysed on agarose gel, quantified

by Qubit fluorimetry (Invitrogen), and 16S rRNA gene sequenced using universal 27F and 1492R primers

(Amann, et al. 1995), to confirm integrity, identity, and purity. When the required concentration for

sequencing was not achieved in samples they were dried by vacuum centrifugation (SpeedVac) at room

temperature. Sequencing was performed on the Illumina HiSeq2000 platform from Nextera XT pair-end
libraries at Australian Genome Research Facility, Melbourne. A reference genome from strain QMA0248
was constructed using both long reads derived from a single Smrt Cell using the PacBio RS II system with
P4C2 chemistry and short reads from Illumina HiSeq2000 derived from Nextera XT paired-end libraries
as reported elsewhere (NCBI accession no: GCA_002220115.1). All sequence data is deposited at NCBI
under Bioproject number PRJNA417543, SRA accession SRP145425. Sample numbers, accession numbers
and extended metadata are provided in Suppl. Table 3. Assembly statistics are provided in Suppl. Table
4.

Phylogenetic analysis

Phylogeny was constructed based on core genome SNPs from *de novo* genome assemblies filtered to
remove recombination breakpoints. Paired-end reads from Illumina were trimmed with Nesoni clip tool
version 0.132 (http://www.vicbioinformatics.com/software.nesoni.shtml), with minimum read length
50, and the first 15 bp of each read removed as quality deterioration in this region was observed when
assessed with FASTQC version 0.11.5. Assembly was performed using the SPAdes assembler version 3.7.1
(Bankevich, et al. 2012), with minimum read coverage cutoff set to 10. Quality of assemblies was assessed
with QUAST 3.2 (Gurevich, et al. 2013). Contigs were ordered by alignment to QMA248 reference genome
(CP022392.1) with Mauve Contig Mover 2.4.0 (Rissman, et al. 2009). Genome annotation was performed
using Prokka 1.11 (Seemann 2014). Rapid alignment of core genomes was carried out using parsnp in the
Harvest Tools suite version 1.2 (Treangen, et al. 2014), and the resulting alignment provided as an input
to Gubbins 1.4.7 (Croucher, Page, et al. 2015) for detection and exclusion of variants produced by
recombination. Phylogenies were then inferred from post-filtered core genome polymorphic sites by
maximum likelihood using RAxML 8.2.8 (Stamatakis 2014) with the general time reversible nucleotide
substitution model GTRGAMMA and bootstrap support from 1000 iterations. Effect of ascertainment bias
associated with using only polymorphic sites on branch length during ML inference was corrected using
Felsenstein's correction implemented in RAxML 8 (Leache, et al. 2015). The resulting phylogenetic tree
was visualized using Dendroscope v 3.5.7 (Huson, et al. 2007) with bootstrap node support value cut-off
75. For the phylogram figure, tip labels were hidden for clarity and the edge containing QMA0141 was
re-scaled as dotted line representing 100-fold decrease in length (Fig. 1). A cladogram based on the
inferred phylogeny, showing all tip labels and bootstrap support for each node, was annotated with
metadata using Evolview V2 (Fig. 2) (He, et al. 2016). To determine whether there was a strong temporal
signal in the phylogenetic data, a root-to-tip regression of branch length against time since isolation was
performed in TempEst (Rambaut, et al. 2016), using genetic distances, corrected for ascertainment bias
in an unrooted tree estimated by maximum likelihood from the alignment of non-recombinant core-
genome SNPs in RAxML 8.2.8.

Variation in DNA repair genes and their regulatory regions

SNP analysis of MMR and GO genes was performed by read-mapping with Geneious version 9.1 (Kearse, et al. 2012), using default settings unless otherwise specified. Paired-end reads from each genome were trimmed, merged into a single file with expected distances between the reads set to 250 bp, and mapped to the curated reference genome of strain QMA0248. Mapped reads were used to detect SNPs with minimum coverage set to 10 and frequency to 0.9. A consensus "pseudogenome" was generated for every strain based on the reference sequence incorporating any detected variants (Ben Zakour, et al. 2016). Multiple alignment of these pseudogenomes was carried out with *Geneious* aligning tool. Sequences of MMR genes (*dnaN,* mut*S, muTL, recD2, rnhC*) and OG genes (*mutY* , *mutM*, *mutX*) annotated in the QMA0248 reference genome were identified in each strain genome sequence. Promoter regions of repair genes were determined with BPROM (Umarov and Solovyev 2017), protein functional domains by SMART genomic (Letunic and Bork 2018), and effect of amino acid substitutions on protein function by PROVEAN Protein with a sensitivity cut-off of 1.3 (Choi and Chan 2015). To detect large variants, alignment of these regions extracted from *de novo* genome assemblies was carried out, and a 56 bp deletion detected in *mutY* promoter of clade B was confirmed by PCR with CAGAAGGAAGAAACAGAC_F/ACCTCTATTGTAGCAAAG_R primers.

Detecting possible association between variation in DNA repair genes and mobilome

A pangenome for the 80 *S. iniae* isolates was constructed with GView server (Petkau, et al. 2010) using the QMA0248 reference genbank (.gbk) file as seed by sequentially adding 79 genome assemblies by clade in the order derived from the cladogram (Fig. 2). Phage positions in the resulting pangenome were then determined by BLAST using Phaster (Arndt, et al. 2016) while IS positions were identified using ISFinder (Siguier, et al. 2006) with a BLAST e-value cutoff of 1×10^{-30}. Positions of MMR and OG genes in the pangenome were determined via manual search of the .gbk file. An image was created in BRIG (Alikhan, et al. 2011) using the pangenome as reference then aligning the 80 genome assemblies by BLAST with an e-value cutoff of 1×10^{-30}. The resulting image was annotated in BRIG from a spreadsheet derived from the positions determined above, and coloured according to clade (Suppl. Fig. 1).

Phenotypic variation related to virulence and antigenicity

A multitude of intracellular and secreted enzymes contribute to virulence in the *Streptococcus* genus (Kadioglu, et al. 2008; Rajagopal 2009). Considering the large number of strains, we limited analysis to major, easily observable, and distinct phenotypes strongly associated with virulence in streptococci

533   (capsule, hemolysis, cell chains) and bacteria in general (oxidation resistance, biofilms). All assays were

534   performed at least in triplicate per strain.

Buoyant density assay for presence of polysaccharide capsule

Presence/absence of polysaccharide capsule was estimated by Percoll buoyant density assay. Isotonic

stock Percoll (ISP) was prepared by mixing nine parts of Percoll with one part of 1.5 M NaCl. Then, 6 parts

of ISP was diluted with 4 parts of 0.15 M NaCl to make final 50 % Percoll solution, which was distributed

by 3 mL into flow cytometry tubes. 10 mL of THB cultures grown to late-exponential phase was adjusted

at OD_{600} 1 (10^8 CFU/mL), centrifuged at 3220 x g for 5min, resuspended in 0.5 mL of 0.15 M NaCl, and

layered onto the Percoll solution. Tubes were centrifuged at 4°C in a swinging bucket rotor at 4000 x g

for 3 h with low acceleration and no brake. In this assay, encapsulated cells form a clear compact band in

the Percoll gradient (Fig. 3, A1), non-encapsulated cells form a pellet in the bottom of the tube (Figure 3,

A2) and, occasionally, strains show differential expression of capsular polysaccharide evidenced by band

and a pellet (Fig. 3, A3).

Haemolytic activity assay on sheep blood agar

Rapid high-throughput detection of impaired haemolytic activity was achieved by blood-agar clearance

zone assay. Briefly, 5 mm wells were made in Columbia agar supplemented with 5% defibrinated sheep

blood (Oxoid, Australia). Bacterial cultures grown to late-exponential phase (~10^8 CFU) were diluted

1:1000 and 50 µL of diluted cultures were pipetted into punctures in the agar (initial inoculum ~$5x10^4$

CFU in each puncture). A 3 mm wide clearance zone was generally produced by bacterial lysis of sheep

erythrocytes during 24 h incubation (Fig. 3, B1). Where haemolysis was absent or fragmentary, impeded

haemolytic activity was recorded (Fig. 3, B2).

Chain formation microscopy

To assess chain formation, *S. iniae* cultures in THB were grown stationarily in 96-well plates at 28 °C for

24 h, mixed and 5 µL wet mounts prepared on a glass microscope slide. Slides were observed by bright

field microscopy under 40x objective with an Olympus BX40 microscope and captured using an Olympus

DP28 digital camera using CellSens software (Olympus Optical Co, Japan). Specimens were observed for

at least 3 min and 2-20 cell chains were generally observed (Fig. 3, C1). Where over 20 cells in a chain

were repeatedly detected (Fig. 3, C2) increased chain formation was recorded, and when more than 10

cells in a chain were not detected by similar observation and the culture was mainly composed of

detached cells (Fig. 3, C3) impeded chain formation was recorded. For the figure, 50 µL of cultures were

dried onto slides at RT, fixed with methanol, Gram stained, and images captured under 100x objective.

Oxidation resistance assay

Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) (Andrews 2001)

of hydrogen peroxide were measured to assess oxidative stress resistance among *S. iniae* strains. THB

cultures grown to late exponential phase were adjusted to OD_{600} = 1 (10^8 CFU/mL), diluted 1000-fold, and

distributed by 100 µL (~10^4 CFU) into wells of a U-bottom 96-well plate (Greiner). THB (100 µL) with 2-

fold excess concentration of hydrogen peroxide was added to the wells and serially diluted twofold,

resulting in a range from 0 to 10 mM final peroxide concentrations. Plates were incubated stationarily

for 24h and examined for presence of observable growth to determine the MIC. Viable cell counts of

cultures without visible growth were performed to determine MBC. MIC of hydrogen peroxide was

estimated as 3 mM for all strains, and MBC as 4 mM for the majority of strains. MBC elevated to 5 mM

in dolphin and human isolates from USA was classified deviation from a wildtype phenotype.

Biofilm formation and visualisation assay

S. iniae biofilms were grown for 4 days in 8-well Lab-Tek® II Chamber Slide™ Systems. To prepare an initial

inoculum of 5 x 10^5 CFU, THB cultures grown to late exponential phase were adjusted to OD_{600} = 1 (10^8

CFU/mL), diluted 100-fold, and 0.5 mL of diluted cultures were placed into Chamber slide wells. After 24

h incubation, THB was removed and replaced with fresh THB every 10-14 h for 3 days. For visualisation,

biofilms were washed in PBS, stained for 15 min with 1 µM fluorescent *Bac*Light Red bacterial stain,

washed in PBS, fixed for 30 min with 10% formalin, and washed twice PBS. Z-stacks were collected by

ZEISS LSM 710 Inverted Laser Scanning Confocal Microscope at 20x objective and visualised using

ZEN2012. Quantification of biofilms was performed using COMSTAT (Verotta, et al. 2017). Structures of

biomass larger than 3.5 and average thickness over 5 were classified as increased biofilm forming activity

(Fig. 3 D).

.

Identifying wildtype virulence-related phenotypes

We consider prevalent phenotypes as a wildtype, namely: presence of polysaccharide capsule (71.25 %

of strains; Figure 3, A1) and haemolytic activity (85% of strains; Figure 3, B1), up to 20 cells in a chain

(77.5 % of strains; Figure 3, A1), 3 mM MIC and 4 mM MBC of hydrogen peroxide (91.25 % of strains),

and 8 -12 µm thick biofilms covering up to 30% of representative image (86.25 % of strains, Fig. 4A). Other

phenotypes are regarded as deviant and discussed.

Statistical analysis

Analysis of co-variance between number of MMR and OG variants, mutation rates, number of atypical

phenotype associated with virulence, and atypical places of isolation were tested with a Phylogenetic

Generalised Least Squares model that accounts for relatedness among strains under Pagel's λ of 0.6

implemented in R (Symonds and Blomberg 2014). Estimation of mutation rates was carried out using the Ma-Sandri-Sarkar Maximum Likelihood Estimation (MSS-MLE) method as implemented in FALCOR fluctuation analysis calculator (Hall, et al. 2009). Differences in mutation rates were compared by Likelihood Ratio Test (LRT) using the rSalvador R package (Zheng 2015, 2017). LRT estimates the overlap between confidence intervals calculated for two fluctuation experiment data sets and allows to compare fluctuation assay data with different number of cultures and resolve very small differences in mutation rate (Zheng 2015, 2017). The *Compare.LD* function was applied pair-wise to all strains, and obtained *p* values were corrected for multiple comparisons by controlling the False Discovery Rate (Benjamini and Hochberg 1995) using p.adjust in R. Adjusted *p* values less than 0.05 were considered to indicate significant difference in mutation rates.

Acknowledgements

This work was supported by The Australian Research Council Discovery Project Grant DP120102755 awarded to ACB. We thank the University of Queensland School of Biological Sciences for funding a tuition fee waiver for OS. For provision of strains into our collection over 15 years we thank: Judy Forbes-Faulkner, Oonoonba Veterinary Laboratory, QLD, Australia; Lynn Shewmaker, CDC, Atlanta, USA; Christian Michel, INRA, France; Suresh Benedict, Berrimah Veterinary Laboratories, NT, Australia; Nicky Buller, WA Fisheries, WA, Australia; Mark White, Tréidlia Biovet Pty (formerly Allied Biotechnology), NSW, Australia; Matt Landos, Future Fisheries Veterinary Services, NSW, Australia; Spencer Russell, Novartis Animal Health, Canada; Ruben Arturo Lopez Crespo, Regal Springs Tilapia, Honduras. We thank Dr Simone Blomberg for advice on statistical analyses and Mr Nazar Rudenko for assistance with writing software pipelines and for use of his Unix server.

References

Abel S, Abel zur Wiesch P, Davis BM, Waldor MK. 2015. Analysis of Bottlenecks in Experimental
Models of Infection. PLoS Pathog 11:e1004823.
Agnew W, Barnes AC. 2007. Streptococcus iniae: an aquatic pathogen of global veterinary
significance and a challenging candidate for reliable vaccination. Vet Microbiol 122:1-15.
Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image Generator (BRIG): simple
prokaryote genome comparisons. BMC Genomics 12:402.
Amann RI, Ludwig W, Schleifer KH. 1995. Phylogenetic identification and in situ detection of
individual microbial cells without cultivation. Microbiology Reviews 59:143-169.
Ambur OH, Davidsen T, Frye SA, Balasingham SV, Lagesen K, Rognes T, Tonjum T. 2009. Genome
dynamics in major bacterial pathogens. FEMS Microbiol Rev 33:453-470.
Andersson DI, Hughes D. 2010. Antibiotic resistance and its cost: is it possible to reverse resistance?
Nat Rev Microbiol 8:260-271.
Andisi VF, Hinojosa CA, de Jong A, Kuipers OP, Orihuela CJ, Bijlsma JJ. 2012. Pneumococcal gene
complex involved in resistance to extracellular oxidative stress. Infection and Immunity 80:1037-
1049.
Andrews JM. 2001. Determination of minimum inhibitory concentrations. J Antimicrob Chemother
48 Suppl 1:5-16.
Aquino T, Nunes A. 2016. Host immunity and pathogen diversity: A computational study. Virulence
7:121-128.
Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang YJ, Wishart DS. 2016. PHASTER: a better, faster
version of the PHAST phage search tool. Nucleic Acids Research 44:W16-W21.
Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S,
Prjibelski AD, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-
cell sequencing. J Comput Biol 19:455-477.
Baumler A, Fang FC. 2013. Host specificity of bacterial pathogens. Cold Spring Harb Perspect Med
3:a010041.
Bedford T, Cobey S, Pascual M. 2011. Strength and tempo of selection revealed in viral gene
genealogies. BMC Evol Biol 11:220.
Ben Zakour NL, Alsheikh-Hussain AS, Ashcroft MM, Khanh Nhu NT, Roberts LW, Stanton-Cook M,
Schembri MA, Beatson SA. 2016. Sequential Acquisition of Virulence and Fluoroquinolone Resistance
Has Shaped the Evolution of Escherichia coli ST131. MBio 7:e00347-00316.
Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate - a Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society Series B-Methodological
57:289-300.
Bjarnsholt T. 2013. The role of bacterial biofilms in chronic infections. APMIS Suppl:1-51.
Blazquez J. 2003. Hypermutation as a factor contributing to the acquisition of antimicrobial
resistance. Clinical Infectious Diseases 37:1201-1209.
Boe L, Danielsen M, Knudsen S, Petersen JB, Maymann J, Jensen PR. 2000. The frequency of
mutators in populations of Escherichia coli. Mutat Res 448:47-55.
Canfield GS, Schwingel JM, Foley MH, Vore KL, Boonanantanasarn K, Gill AL, Sutton MD, Gill SR.
2013. Evolution in fast forward: a potential role for mutators in accelerating Staphylococcus aureus
pathoadaptation. J Bacteriol 195:615-628.
Chen F, Liu WQ, Eisenstark A, Johnston RN, Liu GR, Liu SL. 2010. Multiple genetic switches
spontaneously modulating bacterial mutability. Bmc Evolutionary Biology 10:277.
Choi Y, Chan AP. 2015. PROVEAN web server: a tool to predict the functional effect of amino acid
substitutions and indels. Bioinformatics 31:2745-2747.
Chopra I, O'Neill AJ, Miller K. 2003. The role of mutators in the emergence of antibiotic-resistant
bacteria. Drug Resist Updat 6:137-145.
Cobey S. 2014. Pathogen evolution and the immunological niche. Ann N Y Acad Sci 1320:1-15.

Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, Bentley SD, Hanage WP,
Lipsitch M. 2013. Population genomics of post-vaccine changes in pneumococcal epidemiology. Nat
Genet 45:656-663.
Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, McGee L, von Gottberg A,
Song JH, Ko KS, et al. 2011. Rapid pneumococcal evolution in response to clinical interventions.
Science 331:430-434.
Croucher NJ, Kagedan L, Thompson CM, Parkhill J, Bentley SD, Finkelstein JA, Lipsitch M, Hanage WP.
2015. Selective and genetic constraints on pneumococcal serotype switching. PLoS Genet
11:e1005095.
Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR. 2015. Rapid
phylogenetic analysis of large samples of recombinant bacterial whole genome sequences using
Gubbins. Nucleic Acids Res 43:e15.
Dai L, Muraoka WT, Wu ZW, Sahin O, Zhang QJ. 2015. A single nucleotide change in mutY increases
the emergence of antibiotic-resistant Campylobacter jejuni mutants. Journal of Antimicrobial
Chemotherapy 70:2739-2748.
David SS, O'Shea VL, Kundu S. 2007. Base-excision repair of oxidative DNA damage. Nature 447:941-
950.
de Visser JA. 2002. The fate of microbial mutators. Microbiology 148:1247-1252.
Deitsch KW, Moxon ER, Wellems TE. 1997. Shared themes of antigenic variation and virulence in
bacterial, protozoal, and fungal infections. Microbiol Mol Biol Rev 61:281-293.
Denamur E, Lecointre G, Darlu P, Tenaillon O, Acquaviva C, Sayada C, Sunjevaric I, Rothstein R, Elion
J, Taddei F, et al. 2000. Evolutionary implications of the frequent horizontal transfer of mismatch
repair genes. Cell 103:711-721.
Dupes NM, Walsh BW, Klocko AD, Lenhart JS, Peterson HL, Gessert DA, Pavlick CE, Simmons LA.
2010. Mutations in the Bacillus subtilis beta clamp that separate its roles in DNA replication from
mismatch repair. J Bacteriol 192:3452-3463.
Facklam R, Elliott J, Shewmaker L, Reingold A. 2005. Identification and characterization of sporadic
isolates of Streptococcus iniae isolated from humans. Journal of Clinical Microbiology 43:933-937.
Ferenci T. 2015. Trade-off Mechanisms Shaping the Diversity of Bacteria. Trends Microbiol.
Fogle CA, Nagle JL, Desai MM. 2008. Clonal interference, multiple mutations and adaptation in large
asexual populations. Genetics 180:2163-2173.
Foster PL. 2007. Stress-induced mutagenesis in bacteria. Crit Rev Biochem Mol Biol 42:373-397.
Fukui K. 2010. DNA mismatch repair in eukaryotes and bacteria. J Nucleic Acids 2010.
Galhardo RS, Hastings PJ, Rosenberg SM. 2007. Mutation as a stress response and the regulation of
evolvability. Crit Rev Biochem Mol Biol 42:399-435.
Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, Konradsen HB, Nahm MH. 2015.
Pneumococcal Capsules and Their Types: Past, Present, and Future. Clin Microbiol Rev 28:871-899.
Giraud A, Matic I, Tenaillon O, Clara A, Radman M, Fons M, Taddei F. 2001. Costs and benefits of high
mutation rates: Adaptive evolution of bacteria in the mouse gut. Science 291:2606-2608.
Gonzalez K, Faustoferri RC, Quivey RG, Jr. 2012. Role of DNA base excision repair in the mutability
and virulence of Streptococcus mutans. Molecular Microbiology 85:361-377.
Gould CV, Sniegowski PD, Shchepetov M, Metlay JP, Weiser JN. 2007. Identifying mutator
phenotypes among fluoroquinolone-resistant strains of Streptococcus pneumoniae using fluctuation
analysis. Antimicrob Agents Chemother 51:3225-3229.
Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 29:1072-1075.
Gutierrez O, Juan C, Perez JL, Oliver A. 2004. Lack of association between hypermutation and
antibiotic resistance development in Pseudomonas aeruginosa isolates from intensive care unit
patients. Antimicrob Agents Chemother 48:3573-3575.

Hall BM, Ma CX, Liang P, Singh KK. 2009. Fluctuation analysis CalculatOR: a web tool for the
determination of mutation rate using Luria-Delbruck fluctuation analysis. Bioinformatics 25:1564-
1565.
Hall LM, Henderson-Begg SK. 2006. Hypermutable bacteria isolated from humans--a critical analysis.
Microbiology 152:2505-2514.
Hammerschmidt S, Wolff S, Hocke A, Rosseau S, Muller E, Rohde M. 2005. Illustration of
pneumococcal polysaccharide capsule during adherence and invasion of epithelial cells. Infection
and Immunity 73:4653-4667.
He Z, Zhang H, Gao S, Lercher MJ, Chen WH, Hu S. 2016. Evolview v2: an online visualization and
management tool for customized and annotated phylogenetic trees. Nucleic Acids Res 44:W236-241.
Healey KR, Zhao Y, Perez WB, Lockhart SR, Sobel JD, Farmakiotis D, Kontoyiannis DP, Sanglard D, Taj-
Aldeen SJ, Alexander BD, et al. 2016. Prevalent mutator genotype identified in fungal pathogen
Candida glabrata promotes multi-drug resistance. Nat Commun 7:11128.
Hedges AJ. 2002. Estimating the precision of serial dilutions and viable bacterial counts. Int J Food
Microbiol 76:207-214.
Hegde A, Bhat GK, Mallya S. 2008. Effect of exposure to hydrogen peroxide on the virulence of
Escherichia coli. Indian J Med Microbiol 26:25-28.
Huson DH, Richter DC, Rausch C, Dezulian T, Franz M, Rupp R. 2007. Dendroscope: An interactive
viewer for large phylogenetic trees. BMC Bioinformatics 8:460.
Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of Streptococcus pneumoniae virulence
factors in host respiratory colonization and disease. Nat Rev Microbiol 6:288-301.
Kawasaki M, Delamare-Deboutteville J, Bowater RO, Walker MJ, Beatson S, Ben Zakour NL, Barnes
AC. 2018. Microevolution of aquatic Streptococcus agalactiae ST-261 from Australia indicates
dissemination via imported tilapia and ongoing adaptation to marine hosts or environment. Appl
Environ Microbiol.
Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S,
Duran C, et al. 2012. Geneious Basic: An integrated and extendable desktop software platform for
the organization and analysis of sequence data. Bioinformatics 28:1647-1649.
Korkmaz G, Holm M, Wiens T, Sanyal S. 2014. Comprehensive analysis of stop codon usage in
bacteria and its correlation with release factor abundance. Journal of Biological Chemistry
289:30334-30342.
Labat F, Pradillon O, Garry L, Peuchmaur M, Fantin B, Denamur E. 2005. Mutator phenotype confers
advantage in Escherichia coli chronic urinary tract infection pathogenesis. FEMS Immunol Med
Microbiol 44:317-321.
Leache AD, Banbury BL, Felsenstein J, de Oca AN, Stamatakis A. 2015. Short Tree, Long Tree, Right
Tree, Wrong Tree: New Acquisition Bias Corrections for Inferring SNP Phylogenies. Systematic
Biology 64:1032-1047.
Lenhart JS, Pillon MC, Guarne A, Biteen JS, Simmons LA. 2016. Mismatch repair in Gram-positive
bacteria. Res Microbiol 167:4-12.
Lenhart JS, Sharma A, Hingorani MM, Simmons LA. 2013. DnaN clamp zones provide a platform for
spatiotemporal coupling of mismatch detection to DNA replication. Mol Microbiol 87:553-568.
Letunic I, Bork P. 2018. 20 years of the SMART protein domain annotation resource. Nucleic Acids
Res 46:D493-D496.
Li GM. 2008. Mechanisms and functions of DNA mismatch repair. Cell Research 18:85-98.
Locke JB, Colvin KM, Datta AK, Patel SK, Naidu NN, Neely MN, Nizet V, Buchanan JT. 2007.
Streptococcus iniae capsule impairs phagocytic clearance and contributes to virulence in fish. J
Bacteriol 189:1279-1287.
Locke JB, Colvin KM, Varki N, Vicknair MR, Nizet V, Buchanan JT. 2007. Streptococcus iniae beta-
hemolysin streptolysin S is a virulence factor in fish infection. Diseases of Aquatic Organisms 76:17-
26.

Losordo T, DeLong D, Guerdat T. 2009. Advances in technology and practice for land-based
aquaculture systems: tank-based recirculating systems for finfish production. New Technologies in
Aquaculture:945-983.
Lu AL, Li X, Gu Y, Wright PM, Chang DY. 2001. Repair of oxidative DNA damage: mechanisms and
functions. Cell Biochem Biophys 35:141-170.
Lukacisinova M, Novak S, Paixao T. 2017. Stress-induced mutagenesis: Stress diversity facilitates the
persistence of mutator genes. PLoS Comput Biol 13:e1005609.
Mackinnon MJ, Read AF. 2004. Virulence in malaria: an evolutionary viewpoint. Philos Trans R Soc
Lond B Biol Sci 359:965-986.
Maiden MCJ, van Rensburg MJJ, Bray JE, Earle SG, Ford SA, Jolley KA, McCarthy ND. 2013. MLST
revisited: the gene-by-gene approach to bacterial genomics. Nat Rev Micro 11:728-736.
Mao EF, Lane L, Lee J, Miller JH. 1997. Proliferation of mutators in A cell population. J Bacteriol
179:417-422.
Matsushima A, Takakura S, Fujihara N, Saito T, Ito I, Iinuma Y, Ichiyama S. 2010. High prevalence of
mutators among Enterobacter cloacae nosocomial isolates and their association with antimicrobial
resistance and repetitive detection. Clin Microbiol Infect 16:1488-1493.
Maurelli AT. 2007. Black holes, antivirulence genes, and gene inactivation in the evolution of
bacterial pathogens. FEMS Microbiol Lett 267:1-8.
Mena A, Macia MD, Borrell N, Moya B, de Francisco T, Perez JL, Oliver A. 2007. Inactivation of the
mismatch repair system in Pseudomonas aeruginosa attenuates virulence but favors persistence of
oropharyngeal colonization in cystic fibrosis mice. J Bacteriol 189:3665-3668.
Mena A, Smith EE, Burns JL, Speert DP, Moskowitz SM, Perez JL, Oliver A. 2008. Genetic Adaptation
of Pseudomonas aeruginosa to the Airways of Cystic Fibrosis Patients Is Catalyzed by Hypermutation.
J Bacteriol 190:7910-7917.
Merino D, Reglier-Poupet H, Berche P, Charbit A, European Listeria Genome C. 2002. A hypermutator
phenotype attenuates the virulence of Listeria monocytogenes in a mouse model. Molecular
Microbiology 44:877-887.
Methot PO, Alizon S. 2014. What is a pathogen? Toward a process view of host-parasite interactions.
Virulence 5:775-785.
Metzgar D, Wills C. 2000. Evidence for the adaptive evolution of mutation rates. Cell 101:581-584.
Millard CM, Baiano JC, Chan C, Yuen B, Aviles F, Landos M, Chong RS, Benedict S, Barnes AC. 2012.
Evolution of the capsular operon of Streptococcus iniae in response to vaccination. Appl Environ
Microbiol 78:8219-8226.
Miller JH. 1996. Spontaneous mutators in bacteria: insights into pathways of mutagenesis and repair.
Annu Rev Microbiol 50:625-643.
Negri MC, Morosini MI, Baquero MR, del Campo R, Blazquez J, Baquero F. 2002. Very low cefotaxime
concentrations select for hypermutable Streptococcus pneumoniae populations. Antimicrob Agents
Chemother 46:528-530.
Neher RA, Hallatschek O. 2013. Genealogies of rapidly adapting populations. Proc Natl Acad Sci U S A
110:437-442.
Nilsson AI, Kugelberg E, Berg OG, Andersson DI. 2004. Experimental adaptation of Salmonella
typhimurium to mice. Genetics 168:1119-1130.
Oliver A. 2010. Mutators in cystic fibrosis chronic lung infection: Prevalence, mechanisms, and
consequences for antimicrobial therapy. Int J Med Microbiol 300:563-572.
Oliver A, Mena A. 2010. Bacterial hypermutation in cystic fibrosis, not only for antibiotic resistance.
Clinical Microbiology and Infection 16:798-808.
Petkau A, Stuart-Edwards M, Stothard P, Van Domselaar G. 2010. Interactive microbial genome
visualization with GView. Bioinformatics 26:3125-3126.
Picard B, Duriez P, Gouriou S, Matic I, Denamur E, Taddei H. 2001. Mutator natural Escherichia coli
isolates have an unusual virulence phenotype. Infection and Immunity 69:9-14.

Pier GB, Madin SH. 1976. Streptococcus iniae sp nov, a beta-hemolytic Streptococcus isolated from
an amazon freshwater dolphin, Inia geoffrensis. . International Journal of Systematic Bacteriology
26:545-553.
Pier GB, Madin SH, Alnakeeb S. 1978. Isolation and characterisation of a 2nd isolate of Streptococcus
iniae. International Journal of Systematic Bacteriology 28:311-314.
Rajagopal L. 2009. Understanding the regulation of Group B Streptococcal virulence factors. Future
Microbiol 4:201-221.
Rajanna C, Ouellette G, Rashid M, Zemla A, Karavis M, Zhou C, Revazishvili T, Redmond B, McNew L,
Bakanidze L, et al. 2013. A strain of Yersinia pestis with a mutator phenotype from the Republic of
Georgia. FEMS Microbiol Lett 343:113-120.
Rambaut A, Lam TT, Max Carvalho L, Pybus OG. 2016. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evolution 2.
Raynes Y, Sniegowski PD. 2014. Experimental evolution and the dynamics of genomic mutation rate
modifiers. Heredity (Edinb) 113:375-380.
Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of draft
genomes using the Mauve Aligner. Bioinformatics 25:2071-2073.
Rodriguez JL, Dalia AB, Weiser JN. 2012. Increased chain length promotes pneumococcal adherence
and colonization. Infection and Immunity 80:3454-3459.
Rosche WA, Foster PL. 2000. Determining mutation rates in bacterial populations. Methods 20:4-17.
Scott J, Thompson-Mayberry P, Lahmamsi S, King CJ, McShan WM. 2008. Phage-associated mutator
phenotype in group A streptococcus. J Bacteriol 190:6290-6301.
Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068-2069.
Shabalina SA, Spiridonov NA, Kashina A. 2013. Sounds of silence: synonymous nucleotides as a key to
biological regulation and complexity. Nucleic Acids Res 41:2073-2094.
Shaver AC, Sniegowski PD. 2003. Spontaneously arising mutL mutators in evolving Escherichia coli
populations are the result of changes in repeat length. J Bacteriol 185:6076-6082.
Shrestha S, Bjornstad ON, King AA. 2014. Evolution of acuteness in pathogen metapopulations:
conflicts between "classical" and invasion-persistence trade-offs. Theor Ecol 7:299-311.
Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference centre for
bacterial insertion sequences. Nucleic Acids Research 34:D32-D36.
Smania AM, Segura I, Pezza RJ, Becerra C, Albesa I, Argarana CE. 2004. Emergence of phenotypic
variants upon mismatch repair disruption in Pseudomonas aeruginosa. Microbiology-Sgm 150:1327-
1338.
Sniegowski P, Raynes Y. 2013. Mutation Rates: How Low Can You Go? Current Biology 23:R147-R149.
Sprouffske K, Aguilar-Rodriguez J, Sniegowski P, Wagner A. 2018. High mutation rates limit
evolutionary adaptation in Escherichia coli. Plos Genetics 14:e1007324.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30:1312-1313.
Stich M, Manrubia SC, Lazaro E. 2010. Variable Mutation Rates as an Adaptive Strategy in Replicator
Populations. PLoS One 5.
Sundin GW, Weigand MR. 2007. The microbiology of mutability. FEMS Microbiol Lett 277:11-20.
Symonds MRE, Blomberg SP. 2014. A Primer on Phylogenetic Generalised Least Squares. In:
Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in
Evolutionary Biology: Concepts and Practice. Berlin, Heidelberg: Springer Berlin Heidelberg. p. 105-
130.
Taddei F, Radman M, MaynardSmith J, Toupance B, Gouyon PH, Godelle B. 1997. Role of mutator
alleles in adaptive evolution. Nature 387:700-702.
Tham KC, Hermans N, Winterwerp HHK, Cox MM, Wyman C, Kanaar R, Lebbink JHG. 2013. Mismatch
Repair Inhibits Homeologous Recombination via Coordinated Directional Unwinding of Trapped DNA
Structures. Mol Cell 51:326-337.

Torres-Barcelo C, Cabot G, Oliver A, Buckling A, Maclean RC. 2013. A trade-off between oxidative
stress resistance and DNA repair plays a role in the evolution of elevated mutation rates in bacteria.
Proc Biol Sci 280:20130007.
Torres-Barcelo C, Kojadinovic M, Moxon R, MacLean RC. 2015. The SOS response increases bacterial
fitness, but not evolvability, under a sublethal dose of antibiotic. Proc Biol Sci 282:20150885.
Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-genome
alignment and visualization of thousands of intraspecific microbial genomes. Genome Biology
15:524.
Turrientes MC, Baquero F, Levin BR, Martinez JL, Ripoll A, Gonzalez-Alba JM, Tobes R, Manrique M,
Baquero MR, Rodriguez-Dominguez MJ, et al. 2013. Normal mutation rate variants arise in a Mutator
(Mut S) Escherichia coli population. PLoS One 8:e72963.
Umarov RK, Solovyev VV. 2017. Recognition of prokaryotic and eukaryotic promoters using
convolutional deep learning neural networks. PLoS ONE 12.
V. Solovyev AS. 2011. Automatic Annotation of Microbial Genomes and Metagenomic Sequences. In.
Metagenomics and its Applications in Agriculture, Biomedicine and Environmental: Nova Science
Publishers. p. 61-78.
Verotta D, Haagensen J, Spormann AM, Yang K. 2017. Mathematical Modeling of Biofilm Structures
Using COMSTAT Data. Comput Math Methods Med 2017:7246286.
Wang S, Wang Y, Shen J, Wu Y, Wu C. 2013. Polymorphic mutation frequencies in clinical isolates of
Staphylococcus aureus: the role of weak mutators in the development of fluoroquinolone resistance.
FEMS Microbiol Lett 341:13-17.
Weinberger DM, Malley R, Lipsitch M. 2011. Serotype replacement in disease after pneumococcal
vaccination. Lancet 378:1962-1973.
Weinstein MR, Litt M, Kertesz DA, Wyper P, Rose D, Coulter M, McGeer A, Facklam R, Ostach C,
Willey BM, et al. 1997. Invasive infections due to a fish pathogen, Streptococcus iniae. New England
Journal of Medicine 337:589-594.
Wielgoss S, Barrick JE, Tenaillon O, Cruveiller S, Chane-Woon-Ming B, Medigue C, Lenski RE,
Schneider D. 2011. Mutation Rate Inferred From Synonymous Substitutions in a Long-Term Evolution
Experiment With Escherichia coli. G3 (Bethesda) 1:183-186.
Wielgoss S, Barrick JE, Tenaillon O, Wiser MJ, Dittmar WJ, Cruveiller S, Chane-Woon-Ming B,
Medigue C, Lenski RE, Schneider D. 2013. Mutation rate dynamics in a bacterial population reflect
tension between adaptation and genetic load. Proc Natl Acad Sci U S A 110:222-227.
Wrande M, Roth JR, Hughes D. 2008. Accumulation of mutants in "aging" bacterial colonies is due to
growth under selection, not stress-induced mutagenesis. Proc Natl Acad Sci U S A 105:11863-11868.
Yang H, Miller JH. 2008. Deletion of dnaN1 generates a mutator phenotype in Bacillus anthracis. DNA
Repair (Amst) 7:507-514.
Zheng Q. 2015. Methods for comparing mutation rates using fluctuation assay data. Mutat Res
777:20-22.
Zheng Q. 2017. rSalvador: An R Package for the Fluctuation Experiment. G3 (Bethesda) 7:3849-3856.

Supporting Information Legends

Supplementary Figure 1. *Streptococcus iniae* pangenome derived from 80 strains, coloured by clade (clade A-G innermost - outer) showing positions of MMR and OG genes relative to putative phage/prophage and insertion sequence elements. Phage positions were predicted using Phaster and are indicated as arcs in the outermost ring (green, complete phage; blue, incomplete phage; red, uncertain). ISE were located with ISfinder and are indicated directionally in the outermost ring as black arrows.

Supplementary Figure 2. Root-to-tip regression analysis (TempEst, (Rambaut, et al. 2016)of branch lengths from best-fit-rooted tree against time (year of isolation). The tree was derived from alignment of non-recombinant core-genome SNPs corrected for ascertainment bias in RAxML.

Supplementary Figure 3: Alignment of the sequences found upstream from *mutY* gene in QMA0084 (representative of clade B) and QMA0248 (representative of all other analysed *S. iniae* strains) and predicted changes in transcription pattern produced by 56 bp deletion upstream of *mutY* gene identified in clade B strains.

Supplementary Table 1. 80-by-80 matrix of *p* values resulting from pair-wise comparisons of fluctuation analysis data by Maximum Ratio Test corrected for multiple comparisons by controlling for the False Discovery Rate (Microsoft Excel spreadsheet available as download).

Supplementary Table 2. Results of fluctuation analysis (Microsoft Excel spreadsheet available as download).

Supplementary Table 3. Extended metadata for 80 *Streptococcus iniae* isolates sequenced in the present study (Microsoft Excel spreadsheet)

Supplementary Table 4. Assembly statistics and quality data for 80 *de novo* genome assemblies prepared in this study (Microsoft Excel spreadsheet).

Figures and tables

Figure 1: Unrooted summary phylogram of *Streptococcus iniae* strains based on alignment of core genome SNPs filtered to remove recombination. Genetic distances were inferred by maximum likelihood in RAXML and node support is indicated as percentage of 1000 bootstrap replicates. Geographic origin, time range (year) of isolation, and mutation rate were added manually *post hoc*. All nodes with bootstrap support below 75% were collapsed. The phylogram is drawn with scale proportionate to genetic distance except the dashed branch supporting QMA0141, which is depicted at 10% scale. The inset shows the phylogram structure with QMA0141 branch drawn to the same scale.

Figure 2: Cladogram of *Streptococcus iniae* strains derived from the same dataset as Figure 1. Clade, host

species, variants in MMR and OG DNA repair genes, mutation rate phenotype, and phenotypic variants

associated with virulence were annotated with Evolview. Mutation rate (μ) phenotypes, determined by

fluctuation analysis, are depicted as box blots representing upper and lower 95% confidence limits, with

the interface representing mean values. Node support is indicated as percentage of 1000 bootstrap

replicates.

Figure 3: Phenotypic variants associated with virulence among *Streptococcus iniae* strains. A) Buoyant

density assay for capsule (CPS) presence, A1: CPS+, A2 CPS- A3: CPS+/-. B) Haemolysis of sheep red blood

cells by agar diffusion B1: +, B2:-. C) Streptococcal chain length C1: normal, C2: Long, C3: short. D) Biofilm

formation in chamber slides determined by confocal microscopy, D1: normal, D2: denser, D3: thicker.

Table 1. *Streptococcus iniae* strains used in this study. Includes origin details (host species, time, site

of isolation), phylogenetic affiliation, virulence-associated phenotypes, and mutation rate. Atypical

places of isolation (hosts, tissues) and deviant phenotypes are in bold.

Strain	Clade	Host	Year	Location	Capsule	Hemolysis	Chains	Oxidation resistance	Biofilm formation	Mutation rate (10^-7)
QMA0071	A	*Lates calcarifer*	2000	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2041
QMA0074	C2	*Lates calcarifer*	1998	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.6579
QMA0077	C2	*Lates calcarifer*	1995	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.7067
QMA0078	A	*Lates calcarifer*	2001	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2153
QMA0080	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1966
QMA0082	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1761
QMA0083	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1894
QMA0084	B	***Pteropus alecto***	2001	WA	wildtype	**weak**	wildtype	wildtype	wildtype	0.7293
QMA0087	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1925
QMA0130	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.468
QMA0131	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4488
QMA0133	E1	***Homo sapiens***	2001	USA	**absent**	wildtype	wildtype	wildtype	**increased**	0.5539
QMA0134	E1	***Homo sapiens***	2001	USA	**differential**	wildtype	wildtype	wildtype	**increased**	0.5457
QMA0135	E1	***Homo sapiens***	2002	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.631
QMA0137	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5495
QMA0138	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5989
QMA0139	F	*Fish sp.*	1996	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4311
QMA0140	-	***Inia geoffrensis***	1976	USA	**absent**	wildtype	wildtype	**increased**	**increased**	0.0901
QMA0141	-	***Inia geoffrensis***	1978	USA	**absent**	wildtype	wildtype	**increased**	**increased**	1.0884
QMA0142	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2044
QMA0150	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1952
QMA0155	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1493
QMA0156	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1666
QMA0157	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1545
QMA0158	A	*Lates calcarifer*	2006	SA RAS	**absent**	wildtype	**long**	wildtype	wildtype	0.1678
QMA0159	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2054
QMA0160	A	*Lates calcarifer*	1999	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1481
QMA0161	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1742
QMA0162	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0163	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1523
QMA0164	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7102
QMA0165	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6606
QMA0177	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0180	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2011
QMA0186	D	*Oncorhynchus mykiss*	2000	Israel	wildtype	**weak**	wildtype	wildtype	**increased**	0.5135
QMA0187	-	*Channa striata*	1983	Thailand	**absent**	**weak**	wildtype	wildtype	wildtype	0.4635
QMA0188	D	*Oncorhynchus mykiss*	1998	Israel	wildtype	**weak**	wildtype	wildtype	**increased**	0.5042
QMA0189	D	*Oncorhynchus mykiss*	1996	Reunion	wildtype	**weak**	wildtype	wildtype	wildtype	0.4907

QMA0190	F	*Channa striata*	1988	Thailand	**absent**	wildtype	wildtype	wildtype	wildtype	0.6413
QMA0191	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1883
QMA0207	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1486
QMA0216	A	*Lates calcarifer*	2007	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.1949
QMA0218	C2	*Lates calcarifer*	2007	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7093
QMA0220	A	*Lates calcarifer*	2006	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2086
QMA0221	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1814
QMA0222	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1829
QMA0233	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.362
QMA0234	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.354
QMA0235	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.3496
QMA0236	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**increased**	0.3413
QMA0244	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2101
QMA0245	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1988
QMA0246	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0247	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2059
QMA0248	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1893
QMA0249	F	*Lates calcarifer, bone*	2009	SA RAS	**differential**	**weak**	**long**	wildtype	**increased**	0.3817
QMA0250	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.195
QMA0251	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1927
QMA0252	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2006
QMA0253	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**increased**	0.3555
QMA0254	F	*Lates calcarifer, bone*	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**increased**	0.3712
QMA0258	A	*Lates calcarifer*	2008	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.1876
QMA0371	A	*Scortum barcoo*	2011	NSW	wildtype	wildtype	wildtype	wildtype	wildtype	0.1889
QMA0373	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6908
QMA0374	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.729
QMA0445	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0446	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.1687
QMA0447	E1	*Hybrid striped bass*	1996	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5902
QMA0448	E1	*Hybrid striped bass*	1998	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5529
QMA0457	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1682
QMA0458	A	*Epalzeorhynchos bicolor*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1778
QMA0462	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.4242
QMA0463	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.414
QMA0466	E2	*Oreochromis sp.*	-	USA	wildtype	wildtype	**short**	wildtype	wildtype	0.4602
QMA0467	A	*Epalzeothynchos frenatum*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1695
QMA0468	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1689
QMA0490	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1812

QMA0491	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.2
QMA0492	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1849
QMA0493	A	*Oreochromis sp.*	2016	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1987

955 Table 2: Variants in MMR and OG genes found among *S. iniae* isolates. CDS – protein coding sequence,

956 BS – predicted binding sites of transcription factors, PROVEAN score – predicted effect of amino acid

957 substitutions on the protein function; scores lower that -2 are considered to indicate a deleterious

958 effect.

Gene	Clade	Strains	Location	Mutation type	Nucleotide change	Amino acid change	Protein functional domain	PROVEAN score
mutS	F	from bone	CDS	SNP	G556T	Q -> K	Pfam MutS_II	-0.635
	E	all	CDS	SNP	T594C	-	Pfam MutS_II	n/a
mutL	F	from bone	CDS	SNP	C837A	-	DNA mis_repair	n/a
	F	all	CDS	SNP	T955G	S -> R	DNA mis_repair	-2.050
	B	all	CDS	SNP	C1117T	V -> I	-	-0.271
	F	QMA0190	CDS	SNP	A1626T	-	MutL_C	n/a
dnaN	F	QMA0139	promoter, BS of DnaA	SNP	C->T 116 bp upstream	n/a	n/a	n/a
	G	QMA0187	promoter, BS of FNR	SNP	T->C 135 bp upstream	n/a	n/a	n/a
	C2	all strains	CDS	SNP	C977T	T -> I	Pol3Bc	-2,216
	C1	all strains	stop codon	SNP	T1135C	TAA->Q	n/a	-0.005
rnhC	-	QMA0187	CDS	SNP	G778A	D ->N	Pfam: RNase_HII	-4.593
recD2	-	QMA0140	RBS	SNP	A -> G 15 bp upstream	n/a	n/a	n/a
	E1	all strains	RBS	SNP	T -> A 9 bp upstream	n/a	n/a	n/a
	B	all strains	CDS	SNP	C1605T	-	-	n/a
	C2	all strains	CDS	SNP	C1653T	-	-	n/a
mutY	B	all strains	promoter, -35 box, 1st bp of -10 box, 2 BSs of RpoD17	large deletion (56 bp)	atcataagatttcctcctat aaatctgtgtcatatgtcat ccttctttatagtttc 247-187 bp upstream	n/a	n/a	n/a
	F	all strains	CDS	SNP	T935G	I -> T	Pfam: NUDIX_4	-2.544
mutM	D	all strains	CDS	SNP	T24G	E -> D	Fapy_DNAglyco	-2.990
mutX	E2	all strains	CDS	SNP	A193G	M -> I	Pfam: NUDIX	-0.539

969   Supplementary Figures

```
QMA248_500     cagattatgaataactaatctctggaaaaattagtctccattcttgtaacgtcctttatt     60
QMA0084_444    cagattatgaataactaatctctggaaaaattagtctccattcttgtaacgtcctttatt     60
               ************************************************************

QMA248_500     gaatttaagcatatactatacctttcaatattggagattttccaaaacttatactgtaaa     120
QMA0084_444    gaatttaagcatatactatacctttcaatattggagattttccaaaacttatactgtaaa     120
               ************************************************************

QMA248_500     gagcgctcctttggccaaataacgcatacttcaataagttcaacaatgtgatcactttgt     180
QMA0084_444    gagcgctcctttggccaaataacgcatacttcaataagttcaacaatgtgatcactttgt     180
               ************************************************************

QMA248_500     tagttgtaaaagtcagtttgcctaccgttgttggagctacttagtctccgttcttgtacc     240
QMA0084_444    tagttgtaaaagtcagtttgcctaccgttgttggagctacttagtctccgttcttgtacc     240
               ************************************************************

QMA248_500     gtgcgacatgtagtttcatcataagatttcctcctataaatctgtgtcatatgtcatcct     300
QMA0084_444    gtgcgacatgtagtttc-------------------------------------------     257
               *****************

QMA248_500     tctttatagtttcattataatgtaagcgtttgcttttgtcaagaaaaaatatttccctat     360
QMA0084_444    -------------attataatgtaagcgtttgcttttgtcaagaaaaaatatttccctat     304
                            ***********************************************

QMA248_500     gacattttccacttgcctaataaaaagataagaaaaaaaagctgttatcattgttttttg     420
QMA0084_444    gacattttccacttgcctaataaaaagataagaaaaaaaagctgttatcattgttttttg     364
               ************************************************************

QMA248_500     aaaactctgatagaagctattattttgactttctagcctagccttttcgcatattgaaaa     480
QMA0084_444    aaaactctgatagaagctattattttgactttctagcctagccttttcgcatattgaaaa     424
               ************************************************************

QMA248_500     aactttgctacaatagaggt     500
QMA0084_444    aactttgctacaatagaggt     444
               ********************
```

BROM prediction:

> *QMA248_500*

Length of sequence-   500
Threshold for promoters - 0.20

Number of predicted promoters -   1

Promoter Pos: 328 LDF- 7.23
10 box at pos.   313 cattataat   Score   75
35 box at pos.   291 atgtca   Score   23

Oligonucleotides from known TF binding sites:

For promoter at   328:
rpoD17: TTCCTCCT at position 268   Score - 8
rpoD17: GTGTCATA at position 284   Score - 7
rpoD17: TAATGTAA at position 318   Score - 12
galR: TGTAAGCG at position 321   Score - 8
rpoD18: TTGCTTTT at position 330   Score - 6
rpoD16: TGTCAAGA at position 337   Score - 19

⊳ *QMA0084_444*

Length of sequence-   444
Threshold for promoters - 0.20

Number of predicted promoters -   2

Promoter Pos: 362 LDF- 6.79
10 box at pos.   347 tgttatcat   Score   75
35 box at pos.   324 ataaaa   Score   7

Promoter Pos: 62 LDF- 3.55
10 box at pos.   44 ttgtaacgt   Score   38
35 box at pos.   23 tggaaa   Score   23

Oligonucleotides from known TF binding sites:

For promoter at   362:
arcA: AATAAAAA at position 323   Score - 12
purR: ATAAAAAG at position 324   Score - 11
cpxR: TAAAAAGA at position 325   Score - 9
rpoD18: ATCATTGT at position 351   Score - 5
rpoD19: ATTGTTTT at position 354   Score - 7
fnr: TTTTTTGA at position 358   Score - 9

For promoter at   62:
lrp: ATAACTAA at position 11   Score - 14
tus: TAACTAAT at position 12   Score - 17
tus: TTGTAACG at position 44   Score - 7
rpoD17: ATACTATA at position 73   Score - 11

38