Long title:

Mutators as drivers of adaptation in pathogenic bacteria and a risk factor for host jumps and vaccine

escape: insights into evolutionary epidemiology of the global aquatic pathogen *Streptococcus iniae*

Short title:

Mutators as a risk factor for host jumps and immune escape in pathogenic bacteria

Oleksandra Silayeva[1,2], Jan Engelstaedter[1], and Andrew C Barnes[1,2*]

[1]The University of Queensland, School of Biological Sciences and [2]Centre for Marine Science, St Lucia

Campus, Brisbane, Queensland 4072, Australia

Key words: Mutators; DNA repair; pathogen evolution; immune escape, vaccination; Streptococcus;

aquaculture; host jumps, zoonoses

*Correspondence:

a.barnes@uq.edu.au

Abstract

Hypermutable strains (mutators) occur via mutations in DNA repair genes and facilitate microbial adaptation as they may rapidly generate stress-resistance mutations. Bacterial mutators deficient in mismatch repair (MMR) and oxidised guanine repair (OG) are abundant in clinical settings, however their role in epidemiology and evolution of virulence remains poorly understood. Here we determine phylogenetic relationship among 80 strains of *Streptococcus iniae* isolated globally over 40 years based using non-recombinant core genome single nucleotide polymorphisms (SNPs), estimate their mutation rate by fluctuation analysis, and identify variation in major MMR (*mutS*, *mutL*, *dnaN*, *recD2*, *rnhC*) and OG (*yfhQ*, *mutM*, *mutT*) genes. We find that *S. iniae* mutation rate phenotype and genotype are strongly associated with phylogenetic diversification and variation in major streptococcal virulence determinants (capsular polysaccharide, hemolysin, cell chain length, resistance to oxidation, and biofilm formation). Furthermore, we find that profound changes in virulence determinants observed in mammalian isolates (atypical host) and vaccine-escape isolates found in bone (atypical tissue) of vaccinated barramundi are linked to multiple MMR and OG variants and unique mutation rates. This implies that adaptation to new host taxa, to new host tissue, and to immunity of a vaccinated host may be facilitated by mutator strains. Our findings highlight the importance of mutators and mutation rate dynamics in evolution of pathogenic bacteria, in particular adaptation to a drastically different immunological setting that occurs during host jump and vaccine escape events.

Author Summary

Host immune response is a powerful selective pressure that drives diversification of pathogenic microorganisms and, ultimately, evolution of new strains. Major adaptive events in pathogen evolution, such as transmission to a new host species or infection of vaccinated hosts, require adaptation to a drastically different immune landscape. Such adaptation may be favoured by hypermutable strains (or mutators) that are defective in normal DNA repair and consequently capable of generating multiple potentially beneficial and compensatory mutations. This permits rapid adjustment of virulence and antigenicity in a new immunological setting. Here we show that mutators, through mutations in DNA repair genes and corresponding shifts in mutation rate, are associated with major diversification events and virulence evolution in the broad host-range pathogen *Streptococcus iniae*. We show that mutators underpin infection of vaccinated hosts, transmission to new host species and the evolution of new strains.

Introduction

Optimal mutation rates in microbial populations are determined by a trade-off between maintenance of genetic integrity and evolvability [1, 2]. High fidelity of DNA repair is advantageous in adapted populations since most mutations lead to deviation from the evolved phenotype [2, 3]. Conversely, lower fidelity of DNA repair can be advantageous under stressful conditions since increased mutation supply promotes diversity and may thereby accelerate adaptation [2, 4]. Bacteria can efficiently fluctuate between high and low mutation regimes: While extremely low mutation rates are maintained in favourable conditions [5], elevation of mutation rates known as stress-induced mutagenesis (SIM) occurs in adapting populations [6, 7]. SIM is associated with temporary hypermutability during SOS response and increased frequency of heritable hypermutable strains (mutator strains, or simply mutators) [6, 7]. Mutators periodically arise within a population via mutations in DNA repair genes [8, 9], and become prevalent under stress as, in a clonal population, mutator alleles 'hitchhike' with beneficial mutations that arise at increased rate in the mutator genetic backgrounds [10, 11]. When adaptation is gained low mutation rate can be restored via anti-mutator compensatory mutations [4].

Mutator strains are abundant among bacterial clinical isolates, especially in chronic cases [12, 13]. Multiple studies have attributed this to antibiotic selection [14-16], since mutators are a well-established risk factor for the development of antimicrobial resistance [17, 18]. However, the association between frequency of mutators in diagnostic samples and antibiotic treatments appears to be weak [19, 20], and isogenic knockout mutator strains show an increased ability to colonize an animal host in challenge models devoid of antibiotic exposure [21-23]. Thus, the prevalence of mutators in clinical isolates most likely results from the selective process during adaptation to the host [19, 24-26], where immunity acts as the ultimate selective pressure shaping the diversity of pathogenic strains [27]. The 'Immunological niche' concept states that, due to the diversity and complexity of immune responses, every individual host represents a unique habitat that requires adaptation [28]. Increased mutation supply in a mutator strain could allow rapid adjustment to the particular immunological environment found within each host. Considering that the pathogen needs to adapt to every new host, the host population taken as a whole represents an extremely heterogeneous environment where adaptive processes are ongoing and persistence of mutator alleles is favoured [29]. The fitness of a pathogenic strain in the host hinges on the concept of virulence (the degree of damage to the host), which is a dynamic, complex trait determined by a multitude of virulence factors and host responses [30]. Optimal virulence appears to be a trade-off between acuteness and persistence with higher virulence facilitating extraction of resources and transmission, but too much damage can kill the host and stop transmission altogether [31]. Additionally, virulence factors are often antigens activating the immune response, so lower levels of virulence may be a means of immune evasion [32]. Consequently, both loss and gain of virulence may be

advantageous, and indicate adaptive shifts towards acute or chronic pathogenesis, respectively [33, 34].

Exacerbation and attenuation of virulence factors are repeatedly observed in both knockout and naturally

occurring mutators [25, 35-39]. Thus, it appears that mutator strains/alleles may help pathogenic bacteria

to evolve towards the optimal level of virulence.

Most bacterial mutators isolated from human clinical samples are deficient in genes from mismatch

(MMR) and oxidised guanine (OG) DNA repair systems [12]. MMR and OG genes are conserved across all

domains of life [40], and single nucleotide polymorphisms in their sequence may produce drastic

alterations in mutation rate [41, 42]. Nonetheless, these genes appear to be variable in major bacterial

pathogens, confirming a potential role of mutation rate dynamics in the evolution of virulence [43, 44].

MMR is coupled to replication and removes mispaired nucleotides as well as short insertion/deletion

loops, and prevents recombination between divergent sequences [45, 46].The MMR pathway has been

extensively characterised in *Escherichia coli,* where the MutS dimer protein interacts with the replication

processivity clamp DnaN (b-clamp), binds to the mismatch, recruits MutL, and the MutS:MutL complex

then activates the MutH endonuclease [40]. MutH cuts the unmethylated nascent strand at

hemimethylated d(GATC) sites. MutL loads the UvrD helicase that unwinds the nicked DNA containing

the error, which is followed by excision by one of the single-strand nucleases [40]. This MMR organization,

however, appears to be limited to *E. coli* and closely related Gammaproteobacteria [40, 47]. In most

bacteria and eukaryotes, MutH is absent and MutL has a conserved endonuclease site, dam methylase

(acting at GATC sites) is absent and the strand discrimination signal is unknown [40, 47]. Studies in *Bacillus*

subtilis, a model species for gram-positive bacteria, suggest that nicks created by ribonuclease *rnhC*

during removal of misincorporated ribonucleosides direct MutL endonuclease towards the newly

synthethised strand, and that recD2 acts as a MMR helicase instead of uvrD [47]. The OG system repairs

oxidative DNA damage throughout the cell cycle [48, 49]. Among all DNA bases, guanine is most sensitive

to reactive oxygen species and its oxidised form, 8-oxo-dG or OG, is highly mutagenic as it preferentially

pairs with adenine which, if not corrected, leads to G:C to A:T transversion on a second round of

replication [48, 49]. OG system organization is conserved in most bacteria and eukaryotes and generally

MutT homologs remove 8-oxo-dG from the nucleotide pool, MutM homologs excise 8-oxo-dG

incorporated into DNA, and MutY homologs remove adenine from OG:A mispairings [48, 49].

The present study investigates the role of mutators and mutation rate dynamics in the epidemiology and

evolution of pathogenic bacteria using the broad host-range pathogen *Streptococcus iniae*. *S. iniae* is a

model highly adaptable species, evidenced by its global distribution, ability to cross divergent host taxa

boundaries and repeated re-infection of previously immunised hosts during disease outbreaks in

vaccinated fish in aquaculture [50, 51]. Moreover, Streptococcal pathogens in general are difficult

candidates for immunisation in humans and animals due to high strain diversity evidently driven by

immunisation, yet there is little information on the role of mutators in this evolutionary process [52, 53]. We determine variation in mutation rate and identify mutations in MMR (*mutS*, *mutL*, *dnaN*, *recD2*, *rnhC*) and OG (*yfhQ*, *mutM*, *mutT*) genes among 80 diverse strains of *S. iniae* and discover that they correlate with phylogenetic diversification but are highly conserved within phylogenetic clades. To identify whether adaptive process might be driven by the host immune response we identify variation in major phenotypic traits that determine level of virulence in streptococci (capsular polysaccharide, hemolysin, length of cell chains) and bacteria in general (resistance to reactive oxygen species (ROS), and biofilm formation). Two major streptococcal virulence factors in *S. iniae* infection are the polysaccharide capsule and the beta-hemolysin streptolysin S [54-57]. Both increase survival under native immune response via reduction of phagocytic killing and other mechanisms, however, both are antigens that induce antibody production and development of immune memory [54, 55]. In addition, both capsule production and hemolysin activity have many implications that are not fully understood [55, 58]. For example, non-encapsulated strains exhibit greater ability to adhere to and colonize an epithelium [59]. Length of cell-chains is another phenotype linked to virulence in streptococci: shorter chains are associated with acute infection and increased survival in the blood as they are less likely to activate the complement, whilst longer chains promote adherence and colonization [60]. As pathogenic bacteria are continuously exposed to reactive oxygen released by immune cells (a phenomenon known as 'oxidative burst'), resistance to oxidative compounds contributes to virulence in pathogens allowing them to avoid immune clearance [61, 62]. Another feature generally associated with virulence in bacteria is biofilm forming ability: detached cells are associated with acute infections whereas biofilm aggregates are a hallmark of chronic conditions [63]. The expression of these traits among *S. iniae* isolates is highly consistent with phylogenetic affiliation, which confirms that strain diversity is primarily driven by adaptation to the host achieved via adjustment of virulence. Furthermore, we find that occurrence of abnormal virulence determinants is strongly associated with number of mutations in MMR and OG genes and occurrence of infection in unusual immunological landscapes such as mammals and bone tissue of immunised fish. These data support the critical role of mutators in adaptation to the host and evolution of virulence. Mutators also facilitate the reinfection of immunised animals and transmission between divergent host species.

Results and Discussion

High mutation rates are costly due to inevitable accumulation of deleterious changes. Once adaptation

is gained the mutator strain is predicted to evolve back towards a low mutation rate [64], most likely via

compensatory, 'anti-mutator' mutations in DNA repair genes [4, 65]. Considering this volatility of mutator

phenotypes, their involvement in adaptation cannot be accurately assessed by *de facto* presence among

sporadic isolates, as the latter might have already (partially or fully) adapted and restored a lower

mutation rate. However, mutator and anti-mutator mutations that cause adaptive fluctuations in the

fidelity of DNA repair could be identified in the genotype of an adapted strain. Also, after loss and gain of

functional DNA repair alleles, the mutation rate of an evolved strain is not likely to be exactly the same

as before adaptation. Consequently, variation in DNA repair genes and mutation rate determined

phenotypically by fluctuation analysis might indicate adaptation via a mutator state when placed in a

phylogenetic context.

Our work investigates the link between mutation rate genotype and phenotype and phylogenetic

diversification among *Streptococcus iniae* strains isolated globally between 1976 and 2016 from different

host taxa. Maximum likelihood phylogenetic analysis of 80 *S. iniae* isolates (Table 1) based on non-

recombinant core genome SNPs derived from whole genome data resolved 6 major clades (A-F), one

lineage with two strains (clade G), and three lineages with a single strain (Figure 1). Principally, variation

in MMR and GO genes occurs only between the lineages, and they are highly conserved within clades

(Figure 2, Table 2). In fact, sequence of the major MMR and GO genes is conserved even within

phylogenetic groups not readily definable by geographic origin, time, or host species, and these genes

appear to be good candidates for Multilocus Sequence Typing (MLST, identification of phylogenetic

position based on selected genes as opposed to a whole genome analysis). Although even synonymous

SNPs in any gene locus could affect mRNA and have pronounced phenotypic consequences [66], most

SNPs were identified in protein functional domains, and most amino acid substitutions were predicted to

have a deleterious effect on protein function (Table 2). Further, MMR and GO variation correlates

significantly with mutation rate (p = 0.0056) and deviations from wildtype traits contributing to virulence

(p = 0.0331). In turn, mutation rate is highly consistent with phylogenetic affiliation, and mutation rate

difference is classified as insignificant in pair-wise strain comparisons within major phylogenetic lineages

but significant between (Figure 1, 2).

Isolate QMA0248 (Clade A, used as a reference genome for contig re-ordering in *de novo* genome

assembly and read-mapping for SNP detection) a phenotype mutation rate of 1.9×10^{-8}, which is

consistent with a base mutation rate estimated for other non-mutator gram positive bacteria [12, 67].

Similar mutation rates (within $1.5 - 2 \times 10^{-8}$) are maintained by all isolates (35 strains) along the same

ancestral lineage as QMA0248 (Fig 1, 2; clade A), and all the differences are insignificant except three

pairwise comparisons that verge on commonly accepted significance level. This lineage appears to have

persisted globally in a range of hosts for at least two decades, as it contains strains isolated from 1999 to

2016 in USA, Honduras, and Australia from barramundi (*Lates calcarifer*), tilapia (*Oreochromis sp*), jade

perch (*Scortum barcoo*), and ornamental fish (*Epalzeorhynchos frenatum, Epalzeorhynchos bicolor*).

MMR and GO genes in these strains are identical to QMA0248, and only minor deviations from wildtype

traits contributing to virulence were observed among the strains (absence of capsule production and

longer cell chains in QMA0158, 216) (Figs. 1-3). The rest of the lineages express significantly different

mutation rates (except strains from clade C1 and QMA045-46 discussed below), contain unique and often

multiple SNPs in *mut* genes, and show peculiar phenotypic variants related to virulence (Figures 2, 3).

Although phylogenetic relatedness between lineages is fully resolved, they originate almost

simultaneously and nested lineages evolve independently sharing little phylogenetic history (Figure 1,2).

Clade B contains QMA0084, a strain isolated from a black flying fox, *Pteropus alecto*, in Western Australia

(WA) and two strains isolated from ornamental fish species (clown loach or tiger botia, *Chromobotia*

macracanthus) in USA. These isolates share valine to isoleucine substitution in MutL, a synonymous SNP

in *recD2*, and 56 bp deletion within the promoter of *yfhQ*. The latter mutation is predicted to drastically

changes *yfhQ* transcription pattern, affecting transcription factor binding sites of the original promoter

and introducing a second promoter downstream. It appears that transcription is also affected by the rest

of the genetic background as isolates exhibit significant difference in mutation rate phenotype, 7×10^{-8}

in QMA0084 and 4×10^{-8} in clown loach strains. Phenotypically, clade B has shifted towards decreased

haemolytic activity. Clade B shares a common ancestor with clade C comprised of strains obtained from

barramundi farmed in WA, Northern Territory (NT) (nested clade C1), and north QLD fish farms (nested

clade C2). While a core mutation rate of $1.5\text{-}2 \times 10^{-8}$ is observed in clade C1, a significantly higher mutation

rate of 7×10^{-8} has been maintained in sister clade C2 from north QLD for almost two decades (1995 to

2012) (Fig 1). We speculate that the latter might be attributable to difference in meteorological

conditions: periodic heavy rainfall present in tropical north QLD is likely to bring fluctuations in water

salinity and temperature, creating an unstable environment where adaptive processes are ongoing and

high mutation rate is advantageous. Minor variation in virulence-related phenotypes (absence of capsule

production and longer cell chains in QMA0074, 77)(Fig 2), are supportive that mutation rate evolution in

this lineage might be driven by outside-the-host factors. The significant difference in mutation rate

between nested clades C1 and C2 is potentially attributable to variation in *dnaN*: A predicted deleterious

tyrosine to isoleucine substitution (Table 2, PROVEAN score -2.216) is present in all isolates of C2, which

may explain the high mutation rate phenotype in clade C2 strains (which also have a synonymous variant

in *recD2*)(Fig. 2, Table 2). Indeed, deletion of *dnaN1* in *Bacillus anthracis* results in a mutator phenotype

with equivalent mutation rate to a *mutS* mismatch repair-defective strain, supportive of the connection

between deleterious mutation in DnaN and mutator phenotype found here [68]. Moreover, the T326I

substitution in *S. iniae* clade C2 DnaN occurs in one of the critical residues of the loader binding interface

of the ß-clamp, which may impede interaction with MutS. While MutS can perform mismatch repair

independently of DnaN in *B. subtilis*, 90% of mismatch repair is dependent on targeting MutS to nascent

DNA via the DnaN clamp zone [69, 70]. In contrast the T1135C nucleotide substitution in the *dnaN* stop

codon is unique to clade C1 strains and changes stop codon TAA into CAA coding for glutamine, with the

next stop codon TAG found immediately downstream. This mutation does not appear to affect mutation

and was not predicted to have any affect on the encoded protein function (Table 2) via single amino acid

protein elongation [71] producing a mutation rate that is insignificantly different from the core rate found

in clade A (Fig 2). Clade D consists of isolates from trout (*Oncorhynchus mykiss*) from Réunion and Israel.

These strains have an estimated mutation rate of 5×10^{-8}, contain a glutamate to aspartate substitution

in *mutM* predicted to be deleterious to protein function, a synonymous SNP in *RecD2*, and exhibit

impeded haemolytic activity (Fig 1,2, Table 2).

Isolates from humans and related fish strains are found in clade E where multiple phenotypic variants

contributing to virulence are observed. This clade contains two nested subclades that share a SNP in

mutS, but have other unique SNPs and different mutation rates, which is significant in most pairwise

comparisons. Subclade E1 contains USA isolates from human (QMA0133-35, 37-38) and hybrid striped

bass (QMA0447-48), which have a substitution in the Shine-Dalgarno sequence of *recD2* and a mutation

rate of 6.5×10^{-8} (Fig 1, Table 2). Fish strains and three human strains (QMA0135, 37-38) show a decreased

ability to form cell chains and an elevated ability to withstand oxidative stress. Two human strains

QMA0133-34 form somewhat thicker and considerably denser biofilm structures. QMA0133 is non-

encapsulated, and QMA0134 appears to differentially express the capsular polysaccharide in a culture.

The close phylogenetic relationship presented here implicates a likely transfer from farmed hybrid bass

to humans [72], and this host jump may have been facilitated by the mutator phenotype. Subclade E2

contains two human strains from Canada (QMA0130-31) and tilapia strain from USA (QMA0466). These

strains have unique methionine to isoleucine substitution at the N-terminus of the MutX, and mutation

rate of 4.5×10^{-8} (Fig 1, Table 2). Both human strains are non-encapsulated (Fig 2). The tilapia strain

expresses the capsule but forms short cell chains in common with most strains from the subclade (Fig 2).

The close phylogenetic relationship again strongly implicates transfer from tilapia farmed in USA to

human patients in Canada via import into the Toronto fish market [73] and again, this host jump may

have been facilitated by the mutator state. Clade F is not readily definable by location, time of isolation,

or host species and comprises QMA0139 from unknown fish species obtained in Canada, QMA0190

isolated from snakehead murrel in Thailand (both of which appear on long branches), and a nested

terminal clade comprised of barramundi isolates farmed in Recirculating Aquaculture Systems (RAS) in New South Wales and South Australia. The latter are of a particular interest as they were sampled from barramundi bone lesions during a disease outbreak in vaccinated fish where infection manifested itself as a slowly progressing osteomyelitis instead of typical acute septicaemia and meningitis [50, 51]. Since they are distant to strains used in autogenous vaccines that are found in clade A (QMA0155-57, 160, QMA0250-52), the atypical outbreak might be classified as a case of vaccine-induced serotype replacement (VISR) – spread of co-existing pathogenic strain/s after elimination of the dominant strain/s targeted by the vaccine [74]. In contrast to natural populations where the investigation of VIRS is confounded by ecological and sampling biases, and randomised vaccine trials that lack power to detect the population-wide effect of mass vaccination [74], barramundi RAS represent a relatively controlled and isolated natural environment where vaccination against *S. iniae* fails occasionally but recurrently [51]. RAS are self-contained facilities with minimal water exchange and high standards of water treatment [75]. This ensures that gene flow and, consequently, the probability of new serotype introduction after vaccine implementation is reduced to a minimum. A comparatively uniform immunized host population is exposed to dominant strains against which it was vaccinated along with potentially co-existing lineages in the RAS. In the atypical outbreaks in RAS in Australia, vaccine-induced immunity was partially effective in the infected fish, evident by absence of proliferation of the new strain in typical niduses (blood, brain, and pronephros) and cross-reactivity of vaccine-induced antibodies against bone isolates [51]. This is in contrast with the classical model of serotype replacement where a vaccine is ineffective against the replacing co-existing strain. It has been postulated that VISR can occur even if the vaccine is equally efficient against all circulating strains via other trade-off mechanisms [74]. Apparently, evasion from immune clearance and adaptation to osseous tissue was allowed by shift towards attenuated virulence as suggested by multiple phenotypic changes in bone isolates: absence of capsular production, impeded hemolytic activity, increased cell-chain length, and denser biofilms. All of these changes have previously been associated with chronic infection and increased potential for colonization [55-58, 60, 62, 63], and only absence of polysaccharide capsule is shared with related QMA139 and QMA190. Multiple SNPs in repair genes and the high mutation rate in bone isolates suggest that shift in virulence allowing persistence in the bone tissue might have occurred via mutator phenotype. Serine to arginine substitution in MutL and tyrosine to isoleucine substitution in MutY occur in the ancestor of QMA0139, QMA0190, and bone strains. Both substitutions are predicted to have a deleterious effect on protein function, but their effect on mutation rate is combined with unique SNPs found in each branch: QMA0139 has a SNP in the binding site of the DnaA transcription factor within the *dnaN* promoter, and a mutation rate of 4.5×10^{-8}, QMA0190 has synonymous SNP in *mutL* and mutation rate of 6.5×10^{-8}, and bone strains have glutamine to glycine substitution in MutS and synonymous SNP in *mutL*, and a mutation

rate of 3-4 x 10^{-8}. Considering that the mutation rate of QMA0190 is significantly higher compared to

rates expressed by QMA0139 and bone isolates, it is likely that variants unique to the latter strains

compensate for deleterious variants shared by the isolates [4].

Clade G contains two strains from tilapia isolated in USA (QMA0445-46). Notably, despite being

phylogenetically distant from clade A, these strains have the same MMR and GO genotype and have

retained the core mutation rate and the wildtype virulence phenotype. The first long branch with a single

isolate contains a strain from snakehead murrel strain (*Channa striata*) from Thailand. This isolate is non-

encapsulated and weakly haemolytic and has a mutation rate of 4.5 x 10^{-8}, potentially attributable to

deleterious aspartate to asparagine substitution in rnhC and SNP in the binding site of *fnr* transcription

factor within the *dnaN* promoter. A second long branch with a single strain contains the oldest among

the analysed strains, QMA0140 isolated in 1976 from dolphin (*Inia geoffrensis*)[76]. This strain exhibits a

mutation rate of 1 x 10^{-8}, which is unique among the strains and significantly lower than the core mutation

rate, presumably associated with increased translation rate of recD2 helicase produced by a SNP in the

ribosome binding site (Fig 2, Table 2). The longest branch on the tree contains a second dolphin strain,

QMA0141, isolated two years later in 1978 [77]. This isolate is highly divergent from the rest of strains

with around 20 kB of non-recombinant SNPs in pair-wise comparisons with other strains, accounting for

around 1% genomic difference. In contrast to QMA140, it mutates at a rate of 1x 10^{-7}, the highest

mutation rate phenotype determined among the isolates and significantly different to the rest of the

values. Multiple SNPs are observed in all MMR and GO genes: 3 in *dnaN* and *mutX*, 4 in *mutM*, 9 in *mutL*

and *rnhC*, 15 in *recD2*, 9 in *mutL* and *rnhC*, and 68 in *mutS*. Both dolphin isolates are non-encapsulated,

show increased ability to withstand oxidative stress, and form denser and thicker biofilms (Fig 2, 3).

The conservation of *S. iniae* DNA repair genotypes and phenotype within phylogenetic clades and

variation between the lineages is consistent with the idea of adaptive fluctuations in mutation rate being

a driver of pathogen evolution, and their association with variation in virulence traits supports the

contention that adaptive processes and diversification of pathogens are largely driven by host immunity.

Moreover, infections in atypical hosts (mammals) and atypical tissues (bone) within apparently immune

animals correlates strongly with the degree of deviation from the wildtype virulence phenotype (p=0.000)

and the number of variants in DNA repair genes (p=0.002). This indicates that adaptation to a drastically

different immunological landscape might be facilitated by elevated mutation supply permitting rapid

adjustment of virulence and antigenicity, which has major epidemiological implications. First, mutators

might facilitate pathogen transition among divergent host species known as 'host jumps', including higher

host taxa (eg. *Osteichthyes* and *Mammalia*) [78] . Second, mutators might present a risk factor to immune

escape after vaccination and serotype replacement. Moreover, as evidenced by *S. iniae* colonization of

vaccinated barrumundi bone described previously [51], vaccine escape leading to serotype replacement

might occur without major disruption of immune recognition when multiple changes attenuating virulence and antigenicity allow persistence in a tissue with lower immune surveillance .

To summarize, we estimate mutation rates among the strains by fluctuation analysis, identify the potential genetic bases for the observed differences by comparing MMR and GO genes and their regulatory regions, highlight correlating phenotypic changes associated with virulence, and interpret the data in a context of the phylogeny and the case histories of strain isolation. Our results highlight the role of mutators and mutation rate dynamics in evolution and epidemiology of pathogenic bacteria. In particular, mutators associate with major adaptive events leading to colonization of a new host tissue, infection of novel host taxa, and survival in vaccinated host with specific adaptive immune response. Further experimental studies are required to determine how generalizable these findings are to other pathogens and the evolution of novel strains. It would be of particular interest to conduct similar studies in the Pneumococcus where serotype evolution following vaccination programmes is well documented, but the role of mutators in this process is not yet known [52, 53, 79].

Materials and methods

Strains and growth conditions

Eighty isolates of *S. iniae* collected in Australia, USA, Canada, Israel, Honduras, and Thailand between 1976 and 2016 from eight fish species (*Lates calcarifer*, *Scortum barcoo*, *Epalzeorhynchos frenatum, Epalzeorhynchos bicolor, Oreochromis sp., Channa striata, Chromobotia macracanthus, Oncorhynchus mykiss*) and three mammalian species (*Homo sapiens, Inia geoffrensis, Pteropus alecto*) were used in this study (Table 1). Strains were received from culture collections, veterinarians, or directly from fish farms (Supplementary Table 1) and stored as master seed stocks without further subculture at -80°C in Todd-Hewitt Broth (THB) supplemented with 20% glycerol. Bacteria were routinely recovered from -80°C stocks on Columbia agar supplemented with 5% defibrinated sheep blood (Oxoid, Australia), and cultured at 28°C on Todd-Hewitt agar or in Todd-Hewitt broth (Oxoid) with agitation 200 rpm in a shaking incubator unless otherwise specified.

Estimation of mutation rate phenotype by fluctuation analysis

To estimate mutation rates of *S. iniae* isolates, a fluctuation analysis assay for spontaneous occurrence of rifampicin resistance was optimized according to Rosche and Foster [80]. A single broth culture was initiated from five separate colonies, recovered on Columbia blood agar from stock cultures stored at -80°C, and grown overnight to late-exponential phase in Todd-Hewitt broth. Cultures were adjusted to $OD_{600} = 1$ corresponding to ~10^8 viable CFU, diluted 1:100, and distributed in 200 µl aliquots into 8 wells

of a sterile U-bottom 96-well plate (Greiner) (initial number of CFU (N_0) ~10^5 per well; lower inocula

produced variability in final CFU number, N_t). A replicate plate containing at least two replicate cultures

per strain was also prepared in order to monitor the culture growth by optical density with a BMG

FLUOstar OPTIMA microplate reader. The 96-well plates were incubated without agitation to early

stationary phase. Viable CFU counts were performed for two representative cultures for each strain by

Miles and Misra method [81], and estimated as 1-2 x 10^8 CFU per culture. For selection of mutants, whole

200 µL cultures were plated on Todd-Hewitt agar containing 0.5 µg/mL rifampicin, dried in a laminar flow

hood, and incubated until rifampicin resistant colonies appeared. The assay was repeated three or more

times for each strain, and results were pooled into single data sets representing 32 to 120 cultures per

strain. Estimation of mutation rate (µ - probability of mutation per cell per generation), was carried out

using the Ma-Sandri-Sarkar Maximum Likelihood Estimation (MSS-MLE) method as implemented in

FALCOR fluctuation analysis calculator [82] .

DNA extraction, preparation and sequencing

Genomic DNA was extracted with the DNeasy Blood & Tissue kit (Qiagen) using a modified protocol with

an additional lysis step. Briefly, 10 mL of late-exponential phase culture was centrifuged and the pellet

resuspended in 200 µL lysis buffer comprising 141 µL MilliQ water, 40 µL lysozyme (200 mg/mL), 8 uL of

0.5 M EDTA, 5 µL Triton X100, 4 µL RNAase A (100 mg/mL), and 2 µL of 1M Tris-HCl, and incubated at

37°C, 700 rpm for 1 h in an Eppendorf thermomixer. Then 200 µL of lysis buffer AL (Qiagen) and 20 µL of

proteinase K were added to the sample, mixed by inversion and incubated in the Eppendorf thermomixer

at 56°C, 700 rpm for 1 h. Finally, 200 µL of molecular grade absolute ethanol was added, samples

vortexed, loaded onto the purification column, left to stand for 2 min, washed with AW1 and AW2 buffers

and dried for 3 min. Elution was performed with 80 µL of nuclease-free water pre-heated to 70°C in two

steps: first with 50 µL, then with 30 µL. Eluate (70 µL) was transferred into a new tube and stored at

-80°C until submission for sequencing. The remaining 10 µL was used for gDNA visualization on agarose

gel, quantification by Qubit fluorimetry (Invitrogen), and sequencing 16S rRNA gene using universal 27F

and 1492R primers [83], to confirm integrity, identity, and purity of gDNA. When the required

concentration for sequencing was not achieved in samples they were dried by vacuum centrifugation

(SpeedVac) at room temperature. Sequencing was performed on the Illumina HiSeq2000 platform from

Nextera XT pair-end libraries at Australian Genome Research Facility, Melbourne. A reference genome

from strain QMA0248 was constructed using both long reads derived from a single Smrt Cell using the

PacBio RS II system with P4C2 chemistry and short reads from Illumina HiSeq2000 derived from Nextera

XT paired-end libraries as reported elsewhere (NCBI accession no: GCA_002220115.1). All sequence data

is deposited at NCBI under Bioproject number PRJNA417543, SRA accession SRP145425. Sample

numbers, accession numbers and extended metadata are provided in the supplementary table S1.

Assembly statistics are provided in Supplementary table S2

Phylogenetic analysis

Phylogeny was constructed based on core genome SNPs from *de novo* genome assemblies filtered to

remove recombination breakpoints. Paired-end reads from Illumina were trimmed with Nesoni clip tool

version 0.132 (http://www.vicbioinformatics.com/software.nesoni.shtml), with minimum read length

50, and the first 15 bp of each read removed as quality deterioration in this region was observed when

assessed with FASTQC version 0.11.5. Assembly was performed using the SPAdes assembler version 3.7.1

[84], with minimum read coverage cutoff set to 10. Quality of assemblies was assessed with QUAST 3.2

[85]. Contigs were ordered by alignment to QMA248 reference genome (CP022392.1) with Mauve Contig

Mover 2.4.0 [86]. Genome annotation was performed using Prokka 1.11 [87]. Rapid alignment of core

genomes was carried out using parsnp in the Harvest Tools suite version 1.2 [88], and the resulting

alignment provided as an input to Gubbins 1.4.7 [89] for detection and exclusion of variants produced by

recombination. Phylogenies were then inferred from post-filtered core genome polymorphic sites by

maximum likelihood using RAxML 7.2.8 [90] with the general time reversible nucleotide substitution

model GTRGAMMA and bootstrap support from 1000 iterations. Ascertainment bias associated with

using only polymorphic sites was accounted for using Felsenstein's correction in RAXML. The resulting

phylogenetic tree was visualized using Dendroscope v 3.5.7 [91] with bootstrap node support value cut-

off 75. For the phylogram figure, tip labels were hidden for clarity and the edge containing QMA0141 was

re-scaled as dotted line representing 100-fold decrease in length (Figure 1). A cladogram based on the

inferred phylogeny, showing all tip labels and bootstrap support for each node, was annotated with

metadata using Evolview V2 (Figure 2) [92].

Variation in DNA repair genes and their regulatory regions

SNP analysis of MMR and GO genes was performed by read-mapping with Geneious version 9.1 [93],

using default settings unless otherwise specified. Paired-end reads from each genome were trimmed,

merged into a single file with expected distances between the reads set to 250 bp, and mapped to

reference genome of strain QMA0248. Mapped reads were used to detect SNPs with minimum coverage

set to 10 and frequency to 0.9. A consensus pseudogenome was generated for every strain based on the

reference sequence including any detected variants. Multiple alignment of pseudogenomes was carried

out with *Geneious* aligning tool. Sequences of MMR genes (*dnaN,* mut*S, muTL, recD2, rnhC*) and OG genes

(*yfhQ*, *mutM*, *mutX*) annotated in the QMA0248 reference genome were identified in each strain genome

sequence. Promoter regions of repair genes were determined with BPROM [94], protein functional

domains by SMART genomic [95], and effect of amino acid substitutions on protein function by PROVEAN

Protein with a sensitivity cut-off of 1.3 [96]. To detect large variants, alignment of these regions extracted

from *de novo* genome assemblies was carried out, and a 56 bp deletion detected in *yfhQ* promoter of

clade B was confirmed by PCR using CAGAAGGAAGAAACAGAC_F/ACCTCTATTGTAGCAAAG_R primers

followed by Sanger sequencing.

Phenotypic variation related to virulence and antigenicity

A multitude of intracellular and secreted enzymes contribute to virulence in the *Streptococcus* genus [54,

55]. Considering the large number of strains, we limited analysis to major, easily observable, and distinct

phenotypes strongly associated with virulence in streptococci (capsule, hemolysis, cell chains) and

bacteria in general (oxidation resistance, biofilms). All assays were performed at least in triplicate per

strain.

Buoyant density assay for presence of polysaccharide capsule

Presence/absence of polysaccharide capsule was estimated by Percoll buoyant density assay. Isotonic

stock Percoll (ISP) was prepared by mixing nine parts of Percoll with one part of 1.5 M NaCl. Then, 6 parts

of ISP was diluted with 4 parts of 0.15 M NaCl to make final 50 % Percoll solution, which was distributed

by 3 mL into flow cytometry tubes. 10 mL of THB cultures grown to late-exponential phase was adjusted

at OD_{600} 1 (10^8 CFU/mL), centrifuged at 3220 x g for 5min, resuspended in 0.5 mL of 0.15 M NaCl, and

layered onto the Percoll solution. Tubes were centrifuged at 4°C in a swinging bucket rotor at 4000 x g

for 3 h with low acceleration and no brake. In this assay, encapsulated cells form a clear compact band in

the Percoll gradient (Figure 3, A1), non-encapsulated cells form a pellet in the bottom of the tube (Figure

3, A2) and, occasionally, strains show differential expression of capsular polysaccharide evidenced by

band and a pellet (Figure 3, A3).

Haemolytic activity assay on sheep blood agar

Rapid high-throughput detection of impaired haemolytic activity was achieved by blood-agar clearance

zone assay. Briefly, 5 mm wells were made in Columbia agar supplemented with 5% defibrinated sheep

blood (Oxoid, Australia). Bacterial cultures grown to late-exponential phase (~10^8 CFU) were diluted

1:1000 and 50 µL of diluted cultures were pipetted into punctures in the agar (initial inoculum ~$5x10^4$

CFU in each puncture). A 3 mm wide clearance zone was generally produced by bacterial lysis of sheep

erythrocytes during 24 h incubation (Figure 3, B1). Where haemolysis was absent or fragmentary,

impeded haemolytic activity was recorded (Figure 3, B2).

Chain formation microscopy

To assess chain formation, *S. iniae* cultures in THB were grown stationarily in 96-well plates at 28 °C for

24 h, mixed and 5 µL wet mounts prepared on a glass microscope slide. Slides were observed by bright

field microscopy under 40x objective with an Olympus BX40 microscope and captured using an Olympus

DP28 digital camera using CellSens software (Olympus Optical Co, Japan). Specimens were observed for

at least 3 min and 2-20 cell chains were generally observed (Figure 3, C1). Where over 20 cells in a chain

were repeatedly detected (Figure 3, C2) increased chain formation was recorded, and when more than

10 cells in a chain were not detected by similar observation and the culture was mainly composed of

detached cells (Figure 3, C3) impeded chain formation was recorded. For the figure, 50 µL of cultures

were dried onto slides at RT, fixed with methanol, Gram stained, and images captured under 100x

objective.

Oxidation resistance assay

Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) [97] of

hydrogen peroxide were measured to assess oxidative stress resistance among *S. iniae* strains. THB

cultures grown to late exponential phase were adjusted to $OD_{600} = 1$ (10^8 CFU/mL), diluted 1000-fold, and

distributed by 100 µL (~10^4 CFU) into wells of a U-bottom 96-well plate (Greiner). THB (100 µL) with 2-

fold excess concentration of hydrogen peroxide was added to the wells and serially diluted twofold,

resulting in a range from 0 to 10 mM final peroxide concentrations. Plates were incubated stationarily

for 24h and examined for presence of observable growth to determine the MIC. Viable cell counts of

cultures without visible growth were performed to determine MBC. MIC of hydrogen peroxide was

estimated as 3 mM for all strains, and MBC as 4 mM for the majority of strains. MBC elevated to 5 mM

in dolphin and human isolates from USA was classified deviation from a wildtype phenotype.

Biofilm formation and visualisation assay

S. iniae biofilms were grown for 4 days in 8-well Lab-Tek® II Chamber Slide™ Systems. To prepare an initial

inoculum of 5 x 10^5 CFU, THB cultures grown to late exponential phase were adjusted to $OD_{600} = 1$ (10^8

CFU/mL), diluted 100-fold, and 0.5 mL of diluted cultures were placed into Chamber slide wells. After 24

h incubation, THB was removed and replaced with fresh THB every 10-14 h for 3 days. For visualisation,

biofilms were washed in PBS, stained for 15 min with 1 µM fluorescent *Bac*Light Red bacterial stain,

washed in PBS, fixed for 30 min with 10% formalin, and washed twice PBS. Z-stacks were collected by

ZEISS LSM 710 Inverted Laser Scanning Confocal Microscope at 20x objective and visualised using

ZEN2012. 8 -12 µm thick biofilms covering up to 30% of representative 200 x 200 µm surface produced

by most strains were classified as typical (Figure 3, D1), denser (Figure 3, D2) or/and thicker (Figure 3, D3)

structures were recorded as deviant.

Identifying wildtype virulence-related phenotypes

We consider prevalent phenotypes as a wildtype, namely: presence of polysaccharide capsule (71.25 % of strains; Figure 3, A1) and haemolytic activity (85% of strains; Figure 3, B1), up to 20 cells in a chain (77.5 % of strains; Figure 3, A1), 3 mM MIC and 4 mM MBC of hydrogen peroxide (91.25 % of strains), and 8 -12 µm thick biofilms covering up to 30% of representative image (86.25 % of strains, Fig. 4A). Other phenotypes are regarded as deviant and discussed.

Statistical analysis

Estimation of mutation rates was performed by Ma-Sandri-Sarkar-Maximum Likelihood Estimator (MSS-MLE) method. Fluctuation assay results were adjusted to equal number of mutants in the final culture and allowed comparison of mutation rates by Likelihood Ratio test [98] using rSalvador R package [99]. *Compare.LD* function was applied pair-wise to all strains. Pairwise comparisons resulting in p values less than 0.05 were considered significantly different. Associations between variation in DNA repair genes, mutation rates, deviations from the wildtype virulence phenotype, and typical places of isolation were tested with a Phylogenetic Generalised Least Squares model that accounts for relatedness among strains under Pagel's λ of 0.6 implemented in R [100].

Acknowledgements

For provision of strains into our collection over 15 years we thank: Judy Forbes-Faulkner, Oonoonba Veterinary Laboratory, QLD, Australia; Lynn Shewmaker, CDC, Atlanta, USA; Christian Michel, INRA, France; Suresh Benedict, Berrimah Veterinary Laboratories, NT, Australia; Nicky Buller, WA Fisheries, WA, Australia; Mark White, Tréidlia Biovet Pty (formerly Allied Biotechnology), NSW, Australia; Matt Landos, Future Fisheries Veterinary Services, NSW, Australia; Spencer Russel, Novartis Animal Health, Canada; Ruben Arturo Lopez Crespo, Regal Springs Tilapia, Honduras. ACB and OS extend their thanks to Mr Nazar Rudenko for assistance with writing software pipelines and for use of his unix server.

References

1.   Ferenci T. Trade-off Mechanisms Shaping the Diversity of Bacteria. Trends in microbiology.
2015. doi: 10.1016/j.tim.2015.11.009. PubMed PMID: 26705697.
2.   Stich M, Manrubia SC, Lazaro E. Variable Mutation Rates as an Adaptive Strategy in
Replicator Populations. PloS one. 2010;5(6). doi: ARTN e11186

10.1371/journal.pone.0011186. PubMed PMID: WOS:000278886300028.
3.   Sniegowski P, Raynes Y. Mutation Rates: How Low Can You Go? Current Biology.
2013;23(4):R147-R9. doi: Doi 10.1016/J.Cub.2013.01.018. PubMed PMID: WOS:000315178400007.
4.   Wielgoss S, Barrick JE, Tenaillon O, Wiser MJ, Dittmar WJ, Cruveiller S, et al. Mutation rate
dynamics in a bacterial population reflect tension between adaptation and genetic load. Proc Natl
Acad Sci U S A. 2013;110(1):222-7. doi: 10.1073/pnas.1219574110. PubMed PMID: 23248287;
PubMed Central PMCID: PMCPMC3538217.
5.   Wielgoss S, Barrick JE, Tenaillon O, Cruveiller S, Chane-Woon-Ming B, Medigue C, et al.
Mutation Rate Inferred From Synonymous Substitutions in a Long-Term Evolution Experiment With
Escherichia coli. G3. 2011;1(3):183-6. doi: 10.1534/g3.111.000406. PubMed PMID: 22207905;
PubMed Central PMCID: PMC3246271.
6.   Galhardo RS, Hastings PJ, Rosenberg SM. Mutation as a stress response and the regulation of
evolvability. Critical reviews in biochemistry and molecular biology. 2007;42(5):399-435. doi:
10.1080/10409230701648502. PubMed PMID: 17917874; PubMed Central PMCID: PMC3319127.
7.   Sundin GW, Weigand MR. The microbiology of mutability. Fems Microbiol Lett.
2007;277(1):11-20. doi: Doi 10.1111/J.1574-6968.2007.00901.X. PubMed PMID:
WOS:000250619200002.
8.   Boe L, Danielsen M, Knudsen S, Petersen JB, Maymann J, Jensen PR. The frequency of
mutators in populations of Escherichia coli. Mutation research. 2000;448(1):47-55. PubMed PMID:
10751622.
9.   Miller JH. Spontaneous mutators in bacteria: insights into pathways of mutagenesis and
repair. Annual review of microbiology. 1996;50:625-43. doi: 10.1146/annurev.micro.50.1.625.
PubMed PMID: 8905093.
10.   Mao EF, Lane L, Lee J, Miller JH. Proliferation of mutators in A cell population. Journal of
bacteriology. 1997;179(2):417-22. PubMed PMID: 8990293; PubMed Central PMCID: PMC178711.
11.   Taddei F, Radman M, MaynardSmith J, Toupance B, Gouyon PH, Godelle B. Role of mutator
alleles in adaptive evolution. Nature. 1997;387(6634):700-2. doi: Doi 10.1038/42696. PubMed PMID:
WOS:A1997XD86900052.
12.   Hall LM, Henderson-Begg SK. Hypermutable bacteria isolated from humans--a critical
analysis. Microbiology. 2006;152(Pt 9):2505-14. doi: 10.1099/mic.0.29079-0. PubMed PMID:
16946246.
13.   Oliver A. Mutators in cystic fibrosis chronic lung infection: Prevalence, mechanisms, and
consequences for antimicrobial therapy. International journal of medical microbiology : IJMM.
2010;300(8):563-72. doi: 10.1016/j.ijmm.2010.08.009. PubMed PMID: 20837399.
14.   Matsushima A, Takakura S, Fujihara N, Saito T, Ito I, Iinuma Y, et al. High prevalence of
mutators among Enterobacter cloacae nosocomial isolates and their association with antimicrobial
resistance and repetitive detection. Clinical microbiology and infection : the official publication of
the European Society of Clinical Microbiology and Infectious Diseases. 2010;16(9):1488-93. doi:
10.1111/j.1469-0691.2009.03145.x. PubMed PMID: 20041896.
15.   Wang S, Wang Y, Shen J, Wu Y, Wu C. Polymorphic mutation frequencies in clinical isolates
of Staphylococcus aureus: the role of weak mutators in the development of fluoroquinolone
resistance. Fems Microbiol Lett. 2013;341(1):13-7. doi: 10.1111/1574-6968.12085. PubMed PMID:
23330696.

16.   Negri MC, Morosini MI, Baquero MR, del Campo R, Blazquez J, Baquero F. Very low
cefotaxime concentrations select for hypermutable Streptococcus pneumoniae populations.
Antimicrobial agents and chemotherapy. 2002;46(2):528-30. doi: Doi 10.1128/Aac.46.2.528-
530.2002. PubMed PMID: WOS:000173427900042.
17.   Chopra I, O'Neill AJ, Miller K. The role of mutators in the emergence of antibiotic-resistant
bacteria. Drug resistance updates : reviews and commentaries in antimicrobial and anticancer
chemotherapy. 2003;6(3):137-45. PubMed PMID: 12860461.
18.   Blazquez J. Hypermutation as a factor contributing to the acquisition of antimicrobial
resistance. Clin Infect Dis. 2003;37(9):1201-9. PubMed PMID: WOS:000185916200010.
19.   Mena A, Smith EE, Burns JL, Speert DP, Moskowitz SM, Perez JL, et al. Genetic Adaptation of
Pseudomonas aeruginosa to the Airways of Cystic Fibrosis Patients Is Catalyzed by Hypermutation.
Journal of bacteriology. 2008;190(24):7910-7. doi: Doi 10.1128/Jb.01147-08. PubMed PMID:
WOS:000261217900007.
20.   Gutierrez O, Juan C, Perez JL, Oliver A. Lack of association between hypermutation and
antibiotic resistance development in Pseudomonas aeruginosa isolates from intensive care unit
patients. Antimicrobial agents and chemotherapy. 2004;48(9):3573-5. doi: Doi
10.1128/Aac.48.9.3573-3575.2004. PubMed PMID: WOS:000223625800055.
21.   Giraud A, Matic I, Tenaillon O, Clara A, Radman M, Fons M, et al. Costs and benefits of high
mutation rates: Adaptive evolution of bacteria in the mouse gut. Science. 2001;291(5513):2606-8.
doi: Doi 10.1126/Science.1056421. PubMed PMID: WOS:000167861000043.
22.   Nilsson AI, Kugelberg E, Berg OG, Andersson DI. Experimental adaptation of Salmonella
typhimurium to mice. Genetics. 2004;168(3):1119-30. doi: 10.1534/genetics.104.030304. PubMed
PMID: 15579674; PubMed Central PMCID: PMC1448776.
23.   Healey KR, Zhao Y, Perez WB, Lockhart SR, Sobel JD, Farmakiotis D, et al. Prevalent mutator
genotype identified in fungal pathogen Candida glabrata promotes multi-drug resistance. Nature
communications. 2016;7:11128. doi: 10.1038/ncomms11128. PubMed PMID: 27020939.
24.   Labat F, Pradillon O, Garry L, Peuchmaur M, Fantin B, Denamur E. Mutator phenotype
confers advantage in Escherichia coli chronic urinary tract infection pathogenesis. FEMS Immunol
Med Microbiol. 2005;44(3):317-21. doi: 10.1016/j.femsim.2005.01.003. PubMed PMID: 15907455.
25.   Mena A, Macia MD, Borrell N, Moya B, de Francisco T, Perez JL, et al. Inactivation of the
mismatch repair system in Pseudomonas aeruginosa attenuates virulence but favors persistence of
oropharyngeal colonization in cystic fibrosis mice. Journal of bacteriology. 2007;189(9):3665-8. doi:
10.1128/JB.00120-07. PubMed PMID: 17307847; PubMed Central PMCID: PMC1855878.
26.   Oliver A, Mena A. Bacterial hypermutation in cystic fibrosis, not only for antibiotic
resistance. Clin Microbiol Infec. 2010;16(7):798-808. doi: 10.1111/j.1469-0691.2010.03250.x.
PubMed PMID: WOS:000278645700002.
27.   Aquino T, Nunes A. Host immunity and pathogen diversity: A computational study.
Virulence. 2016;7(2):121-8. doi: 10.1080/21505594.2016.1149284. PubMed PMID: 26836775;
PubMed Central PMCID: PMC4994829.
28.   Cobey S. Pathogen evolution and the immunological niche. Annals of the New York Academy
of Sciences. 2014;1320:1-15. doi: 10.1111/nyas.12493. PubMed PMID: 25040161; PubMed Central
PMCID: PMC4141700.
29.   Lukacisinova M, Novak S, Paixao T. Stress-induced mutagenesis: Stress diversity facilitates
the persistence of mutator genes. PLoS computational biology. 2017;13(7):e1005609. doi:
10.1371/journal.pcbi.1005609. PubMed PMID: 28719607; PubMed Central PMCID: PMC5538753.
30.   Methot PO, Alizon S. What is a pathogen? Toward a process view of host-parasite
interactions. Virulence. 2014;5(8):775-85. doi: 10.4161/21505594.2014.960726. PubMed PMID:
25483864.
31.   Mackinnon MJ, Read AF. Virulence in malaria: an evolutionary viewpoint. Philosophical
transactions of the Royal Society of London Series B, Biological sciences. 2004;359(1446):965-86.
doi: 10.1098/rstb.2003.1414. PubMed PMID: 15306410; PubMed Central PMCID: PMC1693375.

32.   Deitsch KW, Moxon ER, Wellems TE. Shared themes of antigenic variation and virulence in
bacterial, protozoal, and fungal infections. Microbiology and molecular biology reviews : MMBR.
1997;61(3):281-93. PubMed PMID: 9293182; PubMed Central PMCID: PMC232611.

33.   Shrestha S, Bjornstad ON, King AA. Evolution of acuteness in pathogen metapopulations:
conflicts between "classical" and invasion-persistence trade-offs. Theoretical ecology. 2014;7(3):299-
311. doi: 10.1007/s12080-014-0219-7. PubMed PMID: 25214895; PubMed Central PMCID:
PMC4157682.

34.   Maurelli AT. Black holes, antivirulence genes, and gene inactivation in the evolution of
bacterial pathogens. Fems Microbiol Lett. 2007;267(1):1-8. doi: 10.1111/j.1574-6968.2006.00526.x.

35.   Merino D, Reglier-Poupet H, Berche P, Charbit A, European Listeria Genome C. A
hypermutator phenotype attenuates the virulence of Listeria monocytogenes in a mouse model.
Molecular microbiology. 2002;44(3):877-87. PubMed PMID: 11994166.

36.   Gonzalez K, Faustoferri RC, Quivey RG, Jr. Role of DNA base excision repair in the mutability
and virulence of Streptococcus mutans. Molecular microbiology. 2012;85(2):361-77. doi:
10.1111/j.1365-2958.2012.08116.x. PubMed PMID: 22651851; PubMed Central PMCID:
PMC3391326.

37.   Picard B, Duriez P, Gouriou S, Matic I, Denamur E, Taddei H. Mutator natural Escherichia coli
isolates have an unusual virulence phenotype. Infection and immunity. 2001;69(1):9-14. doi: Doi
10.1128/Iai.69.1.9-14.2001. PubMed PMID: WOS:000165943500002.

38.   Smania AM, Segura I, Pezza RJ, Becerra C, Albesa I, Argarana CE. Emergence of phenotypic
variants upon mismatch repair disruption in Pseudomonas aeruginosa. Microbiol-Sgm.
2004;150:1327-38. doi: Doi 10.1099/Mic.0.26751-0. PubMed PMID: WOS:000221538000030.

39.   Canfield GS, Schwingel JM, Foley MH, Vore KL, Boonanantanasarn K, Gill AL, et al. Evolution
in fast forward: a potential role for mutators in accelerating Staphylococcus aureus pathoadaptation.
Journal of bacteriology. 2013;195(3):615-28. doi: 10.1128/JB.00733-12. PubMed PMID: 23204459;
PubMed Central PMCID: PMC3554003.

40.   Fukui K. DNA mismatch repair in eukaryotes and bacteria. Journal of nucleic acids.
2010;2010. doi: 10.4061/2010/260512. PubMed PMID: 20725617; PubMed Central PMCID:
PMC2915661.

41.   Dai L, Muraoka WT, Wu ZW, Sahin O, Zhang QJ. A single nucleotide change in mutY increases
the emergence of antibiotic-resistant Campylobacter jejuni mutants. J Antimicrob Chemoth.
2015;70(10):2739-48. doi: 10.1093/jac/dkv190. PubMed PMID: WOS:000363249800009.

42.   Rajanna C, Ouellette G, Rashid M, Zemla A, Karavis M, Zhou C, et al. A strain of Yersinia
pestis with a mutator phenotype from the Republic of Georgia. Fems Microbiol Lett.
2013;343(2):113-20. doi: 10.1111/1574-6968.12137. PubMed PMID: 23521061.

43.   Ambur OH, Davidsen T, Frye SA, Balasingham SV, Lagesen K, Rognes T, et al. Genome
dynamics in major bacterial pathogens. FEMS microbiology reviews. 2009;33(3):453-70. doi:
10.1111/j.1574-6976.2009.00173.x. PubMed PMID: WOS:000264956900002.

44.   Chen F, Liu WQ, Eisenstark A, Johnston RN, Liu GR, Liu SL. Multiple genetic switches
spontaneously modulating bacterial mutability. BMC evolutionary biology. 2010;10:277. doi:
10.1186/1471-2148-10-277. PubMed PMID: 20836863; PubMed Central PMCID: PMC2955026.

45.   Li GM. Mechanisms and functions of DNA mismatch repair. Cell Res. 2008;18(1):85-98. doi:
10.1038/cr.2007.115. PubMed PMID: WOS:000252920000009.

46.   Tham KC, Hermans N, Winterwerp HHK, Cox MM, Wyman C, Kanaar R, et al. Mismatch
Repair Inhibits Homeologous Recombination via Coordinated Directional Unwinding of Trapped DNA
Structures. Mol Cell. 2013;51(3):326-37. doi: Doi 10.1016/J.Molcel.2013.07.008. PubMed PMID:
WOS:000323086400008.

47.   Lenhart JS, Pillon MC, Guarne A, Biteen JS, Simmons LA. Mismatch repair in Gram-positive
bacteria. Research in microbiology. 2016;167(1):4-12. doi: 10.1016/j.resmic.2015.08.006. PubMed
PMID: 26343983.

48.   David SS, O'Shea VL, Kundu S. Base-excision repair of oxidative DNA damage. Nature.
2007;447(7147):941-50. doi: 10.1038/nature05978. PubMed PMID: WOS:000247373100036.
49.   Lu AL, Li X, Gu Y, Wright PM, Chang DY. Repair of oxidative DNA damage: mechanisms and
functions. Cell biochemistry and biophysics. 2001;35(2):141-70. doi: 10.1385/CBB:35:2:141. PubMed
PMID: 11892789.
50.   Agnew W, Barnes AC. Streptococcus iniae: an aquatic pathogen of global veterinary
significance and a challenging candidate for reliable vaccination. Veterinary microbiology.
2007;122(1-2):1-15. doi: 10.1016/j.vetmic.2007.03.002. PubMed PMID: 17418985.
51.   Millard CM, Baiano JC, Chan C, Yuen B, Aviles F, Landos M, et al. Evolution of the capsular
operon of Streptococcus iniae in response to vaccination. Appl Environ Microbiol. 2012;78(23):8219-
26. Epub 2012/09/25. doi: AEM.02216-12 [pii]

10.1128/AEM.02216-12. PubMed PMID: 23001668; PubMed Central PMCID: PMC3497394.
52.   Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, et al. Population
genomics of post-vaccine changes in pneumococcal epidemiology. Nat Genet. 2013;45(6):656-63.
Epub 2013/05/07. doi: 10.1038/ng.2625. PubMed PMID: 23644493; PubMed Central PMCID:
PMCPMC3725542.
53.   Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, et al. Rapid
Pneumococcal Evolution in Response to Clinical Interventions. Science. 2011;331(6016):430-4. doi:
10.1126/science.1198545. PubMed PMID: WOS:000286635900044.
54.   Rajagopal L. Understanding the regulation of Group B Streptococcal virulence factors. Future
microbiology. 2009;4(2):201-21. doi: 10.2217/17460913.4.2.201. PubMed PMID: 19257847; PubMed
Central PMCID: PMC2691590.
55.   Kadioglu A, Weiser JN, Paton JC, Andrew PW. The role of Streptococcus pneumoniae
virulence factors in host respiratory colonization and disease. Nat Rev Microbiol. 2008;6(4):288-301.
doi: 10.1038/nrmicro1871. PubMed PMID: 18340341.
56.   Locke JB, Colvin KM, Datta AK, Patel SK, Naidu NN, Neely MN, et al. Streptococcus iniae
capsule impairs phagocytic clearance and contributes to virulence in fish. J Bacteriol.
2007;189(4):1279-87. doi: 10.1128/JB.01175-06. PubMed PMID: 17098893; PubMed Central PMCID:
PMC1797360.
57.   Locke JB, Colvin KM, Varki N, Vicknair MR, Nizet V, Buchanan JT. Streptococcus iniae beta-
hemolysin streptolysin S is a virulence factor in fish infection. Diseases of aquatic organisms.
2007;76(1):17-26. doi: Doi 10.3354/Dao076017. PubMed PMID: WOS:000247715900003.
58.   Geno KA, Gilbert GL, Song JY, Skovsted IC, Klugman KP, Jones C, et al. Pneumococcal
Capsules and Their Types: Past, Present, and Future. Clin Microbiol Rev. 2015;28(3):871-99. Epub
2015/06/19. doi: 10.1128/cmr.00024-15. PubMed PMID: 26085553; PubMed Central PMCID:
PMCPmc4475641.
59.   Hammerschmidt S, Wolff S, Hocke A, Rosseau S, Muller E, Rohde M. Illustration of
pneumococcal polysaccharide capsule during adherence and invasion of epithelial cells. Infection
and immunity. 2005;73(8):4653-67. doi: 10.1128/IAI.73.8.4653-4667.2005. PubMed PMID:
16040978; PubMed Central PMCID: PMC1201225.
60.   Rodriguez JL, Dalia AB, Weiser JN. Increased chain length promotes pneumococcal
adherence and colonization. Infection and immunity. 2012;80(10):3454-9. doi: 10.1128/IAI.00587-
12. PubMed PMID: 22825449; PubMed Central PMCID: PMC3457561.
61.   Hegde A, Bhat GK, Mallya S. Effect of exposure to hydrogen peroxide on the virulence of
Escherichia coli. Indian journal of medical microbiology. 2008;26(1):25-8. PubMed PMID: 18227593.
62.   Andisi VF, Hinojosa CA, de Jong A, Kuipers OP, Orihuela CJ, Bijlsma JJ. Pneumococcal gene
complex involved in resistance to extracellular oxidative stress. Infection and immunity.
2012;80(3):1037-49. doi: 10.1128/IAI.05563-11. PubMed PMID: 22215735; PubMed Central PMCID:
PMC3294666.
63.   Bjarnsholt T. The role of bacterial biofilms in chronic infections. APMIS Supplementum.
2013;(136):1-51. doi: 10.1111/apm.12099. PubMed PMID: 23635385.

64.   de Visser JA. The fate of microbial mutators. Microbiology. 2002;148(Pt 5):1247-52. PubMed
PMID: 11988499.
65.   Turrientes MC, Baquero F, Levin BR, Martinez JL, Ripoll A, Gonzalez-Alba JM, et al. Normal
mutation rate variants arise in a Mutator (Mut S) Escherichia coli population. PloS one.
2013;8(9):e72963. doi: 10.1371/journal.pone.0072963. PubMed PMID: 24069167; PubMed Central
PMCID: PMC3771984.
66.   Shabalina SA, Spiridonov NA, Kashina A. Sounds of silence: synonymous nucleotides as a key
to biological regulation and complexity. Nucleic Acids Res. 2013;41(4):2073-94. doi:
10.1093/nar/gks1205. PubMed PMID: 23293005; PubMed Central PMCID: PMC3575835.
67.   Gould CV, Sniegowski PD, Shchepetov M, Metlay JP, Weiser JN. Identifying mutator
phenotypes among fluoroquinolone-resistant strains of Streptococcus pneumoniae using fluctuation
analysis. Antimicrob Agents Chemother. 2007;51(9):3225-9. Epub 2007/08/01. doi: AAC.00336-07
[pii]

10.1128/AAC.00336-07 [doi]. PubMed PMID: 17664329.
68.   Yang H, Miller JH. Deletion of dnaN1 generates a mutator phenotype in Bacillus anthracis.
DNA Repair (Amst). 2008;7(3):507-14. Epub 2008/02/05. doi: 10.1016/j.dnarep.2007.10.003.
PubMed PMID: 18242150; PubMed Central PMCID: PMCPMC2367095.
69.   Dupes NM, Walsh BW, Klocko AD, Lenhart JS, Peterson HL, Gessert DA, et al. Mutations in
the Bacillus subtilis beta clamp that separate its roles in DNA replication from mismatch repair. J
Bacteriol. 2010;192(13):3452-63. Epub 2010/05/11. doi: 10.1128/JB.01435-09. PubMed PMID:
20453097; PubMed Central PMCID: PMCPMC2897676.
70.   Lenhart JS, Sharma A, Hingorani MM, Simmons LA. DnaN clamp zones provide a platform for
spatiotemporal coupling of mismatch detection to DNA replication. Mol Microbiol. 2013;87(3):553-
68. Epub 2012/12/12. doi: 10.1111/mmi.12115. PubMed PMID: 23228104; PubMed Central PMCID:
PMCPMC5938748.
71.   Korkmaz G, Holm M, Wiens T, Sanyal S. Comprehensive analysis of stop codon usage in
bacteria and its correlation with release factor abundance. The Journal of biological chemistry.
2014;289(44):30334-42. doi: 10.1074/jbc.M114.606632. PubMed PMID: 25217634; PubMed Central
PMCID: PMC4215218.
72.   Facklam R, Elliott J, Shewmaker L, Reingold A. Identification and characterization of sporadic
isolates of Streptococcus iniae isolated from humans. Journal of Clinical Microbiology.
2005;43(2):933-7. PubMed PMID: ISI:000227045600071.
73.   Weinstein MR, Litt M, Kertesz DA, Wyper P, Rose D, Coulter M, et al. Invasive infections due
to a fish pathogen, Streptococcus iniae. N Engl J Med. 1997;337(9):589-94. PubMed PMID:
ISI:A1997XT39400002.
74.   Weinberger DM, Malley R, Lipsitch M. Serotype replacement in disease after pneumococcal
vaccination. Lancet. 2011;378(9807):1962-73. doi: 10.1016/S0140-6736(10)62225-8. PubMed PMID:
21492929; PubMed Central PMCID: PMC3256741.
75.   Losordo T, DeLong D, Guerdat T. Advances in technology and practice for land-based
aquaculture systems: tank-based recirculating systems for finfish production. Basics. 2009:945-83.
doi: Doi 10.1533/9781845696474.6.945. PubMed PMID: WOS:000276209900032.
76.   Pier GB, Madin SH. Streptococcus iniae sp nov, a beta-hemolytic Streptococcus isolated from
an amazon freshwater dolphin, Inia geoffrensis. . International Journal of Systematic Bacteriology.
1976;26(4):545-53. PubMed PMID: ISI:A1976CT37500020.
77.   Pier GB, Madin SH, Alnakeeb S. Isolation and characterisation of a 2nd isolate of
Streptococcus iniae. International Journal of Systematic Bacteriology. 1978;28(2):311-4. PubMed
PMID: ISI:A1978EW90500021.
78.   Baumler A, Fang FC. Host specificity of bacterial pathogens. Cold Spring Harbor perspectives
in medicine. 2013;3(12):a010041. Epub 2013/12/04. doi: 10.1101/cshperspect.a010041. PubMed
PMID: 24296346; PubMed Central PMCID: PMCPmc3839602.

79.   Croucher NJ, Kagedan L, Thompson CM, Parkhill J, Bentley SD, Finkelstein JA, et al. Selective
and genetic constraints on pneumococcal serotype switching. PLoS Genet. 2015;11(3):e1005095.
Epub 2015/04/01. doi: 10.1371/journal.pgen.1005095. PubMed PMID: 25826208; PubMed Central
PMCID: PMCPmc4380333.
80.   Rosche WA, Foster PL. Determining mutation rates in bacterial populations. Methods.
2000;20(1):4-17. doi: 10.1006/meth.1999.0901. PubMed PMID: 10610800; PubMed Central PMCID:
PMC2932672.
81.   Hedges AJ. Estimating the precision of serial dilutions and viable bacterial counts.
International journal of food microbiology. 2002;76(3):207-14. PubMed PMID: 12051477.
82.   Hall BM, Ma CX, Liang P, Singh KK. Fluctuation analysis CalculatOR: a web tool for the
determination of mutation rate using Luria-Delbruck fluctuation analysis. Bioinformatics.
2009;25(12):1564-5. doi: 10.1093/bioinformatics/btp253. PubMed PMID: 19369502; PubMed
Central PMCID: PMC2687991.
83.   Amann RI, Ludwig W, Schleifer KH. Phylogenetic identification and in situ detection of
individual microbial cells without cultivation. Microbiology Reviews. 1995;59(1):143-69.
84.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new
genome assembly algorithm and its applications to single-cell sequencing. Journal of computational
biology : a journal of computational molecular cell biology. 2012;19(5):455-77. doi:
10.1089/cmb.2012.0021. PubMed PMID: 22506599; PubMed Central PMCID: PMC3342519.
85.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics. 2013;29(8):1072-5. doi: 10.1093/bioinformatics/btt086. PubMed PMID:
23422339; PubMed Central PMCID: PMC3624806.
86.   Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. Reordering contigs of draft
genomes using the Mauve aligner. Bioinformatics. 2009;25(16):2071-3. doi:
10.1093/bioinformatics/btp356. PubMed PMID: 19515959; PubMed Central PMCID:
PMCPMC2723005.
87.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068-9. doi: 10.1093/bioinformatics/btu153. PubMed PMID: 24642063.
88.   Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome
alignment and visualization of thousands of intraspecific microbial genomes. Genome Biology.
2014;15(11):524. doi: 10.1186/PREACCEPT-2573980311437212. PubMed PMID: 25410596.
89.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid phylogenetic
analysis of large samples of recombinant bacterial whole genome sequences using Gubbins. Nucleic
Acids Res. 2015;43(3):e15. doi: 10.1093/nar/gku1196. PubMed PMID: 25414349; PubMed Central
PMCID: PMCPMC4330336.
90.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014;30(9):1312-3. doi: 10.1093/bioinformatics/btu033. PubMed PMID:
24451623; PubMed Central PMCID: PMCPMC3998144.
91.   Huson DH, Richter DC, Rausch C, Dezulian T, Franz M, Rupp R. Dendroscope: An interactive
viewer for large phylogenetic trees. BMC bioinformatics. 2007;8:460. doi: 10.1186/1471-2105-8-460.
PubMed PMID: 18034891; PubMed Central PMCID: PMC2216043.
92.   He Z, Zhang H, Gao S, Lercher MJ, Chen WH, Hu S. Evolview v2: an online visualization and
management tool for customized and annotated phylogenetic trees. Nucleic Acids Res.
2016;44(W1):W236-41. doi: 10.1093/nar/gkw370. PubMed PMID: 27131786; PubMed Central
PMCID: PMCPMC4987921.
93.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: An
integrated and extendable desktop software platform for the organization and analysis of sequence
data. Bioinformatics. 2012;28(12):1647-9. doi: 10.1093/bioinformatics/bts199. PubMed PMID:
WOS:000305419800052.

94.   V. Solovyev AS. Automatic Annotation of Microbial Genomes and Metagenomic Sequences.
Metagenomics and its Applications in Agriculture, Biomedicine and Environmental Nova Science
Publishers; 2011. p. 61-78.
95.   Letunic I, Bork P. 20 years of the SMART protein domain annotation resource. Nucleic Acids
Res. 2018;46(D1):D493-D6. doi: 10.1093/nar/gkx922. PubMed PMID: 29040681; PubMed Central
PMCID: PMC5753352.
96.   Choi Y, Chan AP. PROVEAN web server: a tool to predict the functional effect of amino acid
substitutions and indels. Bioinformatics. 2015;31(16):2745-7. doi: 10.1093/bioinformatics/btv195.
PubMed PMID: 25851949; PubMed Central PMCID: PMCPMC4528627.
97.   Andrews JM. Determination of minimum inhibitory concentrations. The Journal of
antimicrobial chemotherapy. 2001;48 Suppl 1:5-16. PubMed PMID: 11420333.
98.   Zheng Q. Methods for comparing mutation rates using fluctuation assay data. Mutation
research. 2015;777:20-2. doi: 10.1016/j.mrfmmm.2015.04.002. PubMed PMID: 25912079.
99.   Zheng Q. rSalvador: An R Package for the Fluctuation Experiment. G3. 2017;7(12):3849-56.
doi: 10.1534/g3.117.300120. PubMed PMID: 29084818; PubMed Central PMCID: PMC5714482.
100.   Symonds MRE, Blomberg SP. A Primer on Phylogenetic Generalised Least Squares. In:
Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in
Evolutionary Biology: Concepts and Practice. Berlin, Heidelberg: Springer Berlin Heidelberg; 2014. p.
105-30.

Supporting Information Legends

Supplementary Table 1. Extended metadata for 80 *Streptococcus iniae* isolates sequenced in the present study (Microsoft Excel spreadsheet

Supplementary Table 2. Assembly statistics and quality data for 80 *de novo* genome assemblies prepared in this study (Microsoft Excel spreadsheet).

Figures and tables

Figure 1: Summary phylogram of *Streptococcus iniae* strains based on alignment of core genome SNPs filtered to remove recombination. Genetic distances were inferred by maximum likelihood in RAXML and node support is indicated as percentage of 1000 bootstrap replicates. Geographic origin, time range (year) of isolation, and mutation rate were added manually *post hoc*. All nodes with bootstrap support below 75% were collapsed. The phylogram is drawn with scale proportionate to genetic distance except the dashed branch supporting QMA0141, which is depicted at 10% scale. The inset shows the phylogram structure with QMA0141 branch drawn to the same scale.

Figure 2: Cladogram of *Streptococcus iniae* strains derived from the same dataset as Figure 1. Clade, host species, variants in MMR and GO DNA repair genes, mutation rate phenotype, and phenotypic variants associated with virulence were annotated with Evolview. Node support is indicated as percentage of 1000 bootstrap replicates.

Figure 3: Phenotypic variants associated with virulence among *Streptococcus iniae* strains. A) Buoyant density assay for capsule (CPS) presence, A1: CPS+, A2 CPS- A3: CPS+/-. B) Haemolysis of sheep red blood cells by agar diffusion B1: +, B2:-. C) Streptococcal chain length C1: normal, C2: Long, C3: short. D) Biofilm formation in chamber slides determined by confocal microscopy, D1: normal, D2: Denser, D3: thicker.

841   Table 1. *Streptococcus iniae* strains used in this study. Includes origin details (host species, time, site

842   of isolation), phylogenetic affiliation, virulence-associated phenotypes, and mutation rate. Atypical

843   places of isolation (hosts, tissues) and deviant phenotypes are in bold.

Strain	Clade	Host	Year	Location	Capsule	Hemolysis	Chains	Oxidation resistance	Biofilms	Mutation rate (10^{-7})
QMA0071	A	*Lates calcarifer*	2000	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2041
QMA0074	C2	*Lates calcarifer*	1998	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.6579
QMA0077	C2	*Lates calcarifer*	1995	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.7067
QMA0078	A	*Lates calcarifer*	2001	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.2153
QMA0080	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1966
QMA0082	C1	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1761
QMA0083	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1894
QMA0084	B	***Pteropus alecto***	2001	WA	wildtype	**weak**	wildtype	wildtype	wildtype	0.7293
QMA0087	A	*Lates calcarifer*	2004	WA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1925
QMA0130	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.468
QMA0131	E2	***Homo sapiens***	1995	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4488
QMA0133	E1	***Homo sapiens***	2001	USA	**absent**	wildtype	wildtype	wildtype	**denser**	0.5539
QMA0134	E1	***Homo sapiens***	2001	USA	**differential**	wildtype	wildtype	wildtype	**denser**	0.5457
QMA0135	E1	***Homo sapiens***	2002	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.631
QMA0137	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5495
QMA0138	E1	***Homo sapiens***	2004	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5989
QMA0139	F	*Fish sp.*	1996	Canada	**absent**	wildtype	wildtype	wildtype	wildtype	0.4311
QMA0140	G	***Inia geoffrensis***	1976	USA	**absent**	wildtype	wildtype	**increased**	**denser, thicker**	0.0901
QMA0141		***Inia geoffrensis***	1978	USA	**absent**	wildtype	wildtype	**increased**	**denser, thicker**	1.0884
QMA0142	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2044
QMA0150	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1952
QMA0155	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1493
QMA0156	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1666
QMA0157	A	*Lates calcarifer*	2005	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1545
QMA0158	A	*Lates calcarifer*	2006	SA RAS	**absent**	wildtype	**long**	wildtype	wildtype	0.1678
QMA0159	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2054
QMA0160	A	*Lates calcarifer*	1999	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1481
QMA0161	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1742
QMA0162	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0163	A	*Lates calcarifer*	2000	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1523
QMA0164	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7102
QMA0165	C2	*Lates calcarifer*	2006	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6606
QMA0177	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1718
QMA0180	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.2011
QMA0186	D	*Oncorhynchus mykiss*	2000	Israel	wildtype	**weak**	wildtype	wildtype	wildtype	0.5135
QMA0187	G	*Channa striata*	1983	Thailand	**absent**	**weak**	wildtype	wildtype	wildtype	0.4635
QMA0188	D	*Oncorhynchus mykiss*	1998	Israel	wildtype	**weak**	wildtype	wildtype	wildtype	0.5042

QMA0189	D	*Oncorhynchus mykiss*	1996	Reunion	wildtype	**weak**	wildtype	wildtype	wildtype	0.4907
QMA0190	F	*Channa striata*	1988	Thailand	**absent**	wildtype	wildtype	wildtype	wildtype	0.6413
QMA0191	C1	*Lates calcarifer*	2005	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1883
QMA0207	C1	*Lates calcarifer*	2006	NT	wildtype	wildtype	wildtype	wildtype	wildtype	0.1486
QMA0216	A	*Lates calcarifer*	2007	QLD	**absent**	wildtype	**long**	wildtype	wildtype	0.1949
QMA0218	C2	*Lates calcarifer*	2007	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.7093
QMA0220	A	*Lates calcarifer*	2006	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2086
QMA0221	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1814
QMA0222	A	*Lates calcarifer*	2006	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1829
QMA0233	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**denser**	0.362
QMA0234	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**denser**	0.354
QMA0235	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**denser**	0.3496
QMA0236	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	**weak**	**long**	wildtype	**denser**	0.3413
QMA0244	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2101
QMA0245	A	*Lates calcarifer*	2008	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1988
QMA0246	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0247	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2059
QMA0248	A	*Lates calcarifer*	2009	SA RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1893
QMA0249	F	*Lates calcarifer,* ***bone***	2009	SA RAS	**differential**	**weak**	**long**	wildtype	**denser**	0.3817
QMA0250	A	*Lates calcarifer*	2007	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.195
QMA0251	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.1927
QMA0252	A	*Lates calcarifer*	2008	NSW RAS	wildtype	wildtype	wildtype	wildtype	wildtype	0.2006
QMA0253	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**denser**	0.3555
QMA0254	F	*Lates calcarifer,* ***bone***	2009	NSW RAS	**absent**	wildtype	**long**	wildtype	**denser**	0.3712
QMA0258	A	*Lates calcarifer*	2008	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.1876
QMA0371	A	*Scortum barcoo*	2011	NSW	wildtype	wildtype	wildtype	wildtype	wildtype	0.1889
QMA0373	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.6908
QMA0374	C2	*Lates calcarifer*	2012	QLD	wildtype	wildtype	wildtype	wildtype	wildtype	0.729
QMA0445	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.2149
QMA0446	G	*Oreochromis sp.*	1998	USA	**absent**	wildtype	wildtype	wildtype	wildtype	0.1687
QMA0447	E1	*Hybrid striped bass*	1996	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5902
QMA0448	E1	*Hybrid striped bass*	1998	USA	wildtype	wildtype	**short**	**increased**	wildtype	0.5529
QMA0457	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1682
QMA0458	A	*Epalzeorhynchos bicolor*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1778
QMA0462	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.4242
QMA0463	B	*Chromobotia macracanthus*	2005	USA	wildtype	**weak**	wildtype	wildtype	wildtype	0.414
QMA0466	E2	*Oreochromis sp.*	-	USA	wildtype	wildtype	**short**	wildtype	wildtype	0.4602
QMA0467	A	*Epalzeothynchos frenatum*	2004	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1695

QMA0468	A	*Oreochromis sp.*	2005	USA	wildtype	wildtype	wildtype	wildtype	wildtype	0.1689
QMA0490	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1812
QMA0491	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.2
QMA0492	A	*Oreochromis sp.*	2015	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1849
QMA0493	A	*Oreochromis sp.*	2016	Honduras	wildtype	wildtype	wildtype	wildtype	wildtype	0.1987

849   Table 2: Variants in MMR and GO genes found among *S. iniae* isolates. D letter next to PROVEAN
850   output value indicates predicted deleterious effect of amino acid change on protein function.

Gene	Clade	Isolates	Mutation	Codon/Amino acid	Locus	PROVEAN score
mutS	G	bone strains	G556T	Q186K	Pfam MutS_II	-0.635
	E1,2	all strains	T594C	198	Pfam MutS_II	n/a
mutL	G	bone strains	C837A	279	DNA mis_repair	n/a
	F	all strains	T955G	S319R	DNA mis_repair	-2.050, D
	B	all strains	C1117T	V393I	-	-0.271
	F	QMA0190	A1626T	542	MutL_C	n/a
dnaN	F	QMA0139	C->T	n/a	promoter, *dnaA* BS 116 bp upstream	n/a
	G	QMA0187	T->C	n/a	promoter, *fnr* BS 135 bp upstream	n/a
	C2	all strains	C977T	T326I	Pol3Bc	-2,216, D
	C1	all strains	T1135C	TAA->Q	Stop codon	n/a
rnhC	G	QMA0187	G778A	D260N	Pfam: RNase_HII	-4.593, D
recD2	G	QMA0140	A -> G	n/a	RBS 15 b. p. upstream	n/a
	E1	all strains	T -> A	n/a	RBS 9 b. p. upstream	n/a
	B	all strains	C1605T	535	-	n/a
	C2	all strains	C1653T	551	-	n/a
yfhQ	B	all strains	56 b.p. deletion	n/a	promoter 187 b. p. upstream	n/a
	F	all strains	T935G	I312T	Pfam: NUDIX_4	-2.544, D
mutM	D	all strains	T24G	E8D	Fapy_DNAglyco	-2.990, D
mutX	E2	all strains	A193G	M65I	Pfam: NUDIX	-0.539

Flying fox
WA 2001
$\mu = 7 \times 10^{-8}$

Ornamental
USA 2005
$\mu = 4 \times 10^{-8}$

Clade C1
$\mu = 1.5\text{-}2 \times 10^{-8}$

Barramundi
WA & NT
2005-2006

Clade C2
$\mu = 6.5\text{-}7.5 \times 10^{-8}$

Barramundi
QLD
1995-2012

Clade D
$\mu = 5 \times 10^{-8}$

Trout
Israel 1998-2000

Clade B

Clade E1
$\mu = 5\text{-}6.5 \times 10^{-8}$

Trout
Réunion
1996

Human & tilapia
USA 1996-2004

Ornamental
& tilapia
USA 2005

Human
Canada 1995

Jade perch
NSW 2011

Tilapia
USA 1998

Clade E2

Barramundi
SA (inc QMA0248),
NSW & WA
1999-2008

Barramundi
QLD 2000-1

Dolphin
USA 1976
$\mu = 0.9 \times 10^{-7}$

Ornamental
& tilapia
USA 2004

Barramundi
QLD
2007-8

Snakehead murrel
Thailand 1983
$\mu = 0.9 \times 10^{-7}$

Clade A
$\mu = 1.5 \text{-} \times 10^{-8}$

Tilapia
Honduras
2015

Tilapia
USA 1998
$\mu = 1.5\text{-}2 \times 10^{-8}$

Clade G

Fish sp.
Canada 1996
$\mu = 4.5 \times 10^{-8}$

QMA0141
Dolphin
USA 1978
$\mu = 1 \times 10^{-7}$

Snakehead murrel
Thailand 1988
$\mu = 6.5 \times 10^{-8}$

Clade F

Barramundi (bone)
SA & NSW 2008-2009
$\mu = 3.5\text{-}4 \times 10^{-8}$

100 100 100 100 100 100 100 100 100 100 100 100 100 100 100 100 100

QMA0141

0.2