Title:

Comparative genomics analyses indicate differential methylated amine

utilisation trait within the member of the genus *Gemmobacter*

Running title:

Methylated amine utilisation trait in Gemmobacter

Kröber, E.[1], Spurgin L.[2], Wischer, D., Kumaresan, D.[3#]

[1] Microbial Biogeochemistry, RA Landscape Functioning, ZALF Leibniz Centre for Landscape

Research, Müncheberg, Germany

[2] School of Environmental Sciences, University of East Anglia, Norwich, UK

[4] School of Biological Sciences & Institute for Global Food Security, Queen's University

Belfast, Belfast, Northern Ireland, UK

[#]Correspondence to

Dr D Kumaresan, School of Biological Sciences, Queen's University Belfast, Belfast, Northern

Ireland, United Kingdom, d.kumaresan@qub.ac.uk

Abstract

Methylated amines are ubiquitous in the environment and play a role in regulating the earth's climate via a set of complex biological and chemical reactions. Microbial degradation of these compounds is thought to be a major sink. Recently we isolated a facultative methylotroph, *Gemmobacter* sp. LW-1, an isolate from the unique environment Movile Cave, Romania, which is capable of methylated amine utilisation as a carbon source. Here, using a comparative genomics approach, we investigate how widespread methylated amine utilisation trait is within the member of the bacterial genus *Gemmobacter*. Five genomes of different *Gemmobacter* species isolated from diverse environments, such as activated sludge, fresh water, sulphuric cave waters (Movile Cave) and the marine environment were available from the public repositories and used for the analysis. Our results indicate that some members of the genus *Gemmobacter*, namely *G. aquatilis*, *G. caeni* and *G.* sp. LW-1 have the genetic potential of methylated amine utilisation while others (*G. megaterium* and *G. nectariphilus*) have not. Ancestral state reconstruction analysis also suggested that methylated amine utilisation trait might not be ancestral to members of *Gemmobacter* and has been gained. Based on our analysis, we suggest that the trait of methylated amine utilisation within the members of the genus *Gemmobacter* might be independent of their habitat and more randomly distributed.

Introduction

Methylated amines (MAs) are ubiquitous in the environment with a variety of natural and anthropogenic sources including the oceans, vegetation, sediments and organic-rich soils, animal husbandry, food industry, pesticides, sewage, and automobiles, to mention only a few (Schade and Crutzen, 1995;Latypova et al., 2010;Ge et al., 2011).

Methylated amines are also known to influence earth's climate, via a series of

complex biological and chemical interactions (Carpenter et al., 2012). Some of the

most abundant methylated amines found in the atmosphere are trimethylamine

(TMA), dimethylamine (DMA) and monomethylamine (MMA) (Ge et al., 2011).

Microbial metabolism of methylated amines involves both aerobic and anaerobic

microorganisms, e.g. some methanogenic archaea such as *Methanosarcina* and

Methanomicrobium can use MAs to produce methane (Burke et al., 1998;Liu and

Whitman, 2008;Lyimo et al., 2009) while Gram-positive and Gram-negative

methylotrophic bacteria can use MAs as carbon and nitrogen source (Chen et al.,

2010b). Previously, MAs were typically associated with marine ecosystems as they

are by-products of degradation of osmolytic chemicals such as glycine betaine,

carnitine, choline and trimethylamine N-oxide (Chen et al., 2010b). However, recent

studies have reported the detection and activity of aerobic methylotrophic bacteria

that utilise MAs in a variety of natural and engineered environments (Chen et al.,

2009;Chistoserdova et al., 2009;Chistoserdova, 2011;Ge et al., 2011;Wischer et al.,

2015) and could play a major role in global C and N budgets.

Aerobic methylotrophs are a polyphyletic group of microorganisms capable of

utilising one-carbon (C_1) compounds such as methane, methanol or methylated

amines as their sole source of carbon and energy (Anthony, 1982;Lidstrom,

2006;Chistoserdova et al., 2009). Methylotrophs can degrade TMA to DMA by using

the enzymes TMA dehydrogenase, TMA monooxygenase or TMA methyltransferase,

encoded by the genes *tdm*, *tmm* and *mtt*, respectively (Paul et al., 2000;Chen,

2012;Lidbury et al., 2014). The enzymes DMA dehydrogenase (*dmd*) or DMA

monooxygenase (*dmmDABC*) modulate the conversion of DMA to MMA (Lidstrom,

2006;Chen, 2012). Two distinct pathways have been characterised for the oxidation of

MMA (Chistoserdova, 2011). The direct MMA-oxidation pathway mediated by a

single enzyme (MMA dehydrogenase in Gram-negative bacteria and MMA oxidase in

Gram-positive bacteria) converts MMA to formaldehyde and releases ammonium

(McIntire et al., 1991;Chistoserdova et al., 1994). The alternate pathway, referred to

as the *N*-methylglutamate (NMG) pathway or indirect MMA-oxidation pathway, is

mediated by three individual enzymes via the oxidation of MMA to gamma-

glutamylmethylamide (GMA) and its further degradation to *N*-methylglutamate

(NMG) and 5,10-methylenetetrahydrofolate ($CH_2 = H_4F$) (Latypova et al.,

2010;Chistoserdova, 2011). A stepwise conversion of MMA in the NMG pathway is

modulated by the enzymes GMA synthetase (*gmaS*), NMG synthase (*mgsABC*) and

NMG dehydrogenase (*mgdABCD*) (Chen et al., 2010b;Latypova et al., 2010). The

capability to use MMA not only as a source for carbon but also for nitrogen is

widespread in bacteria. Notably, the NMG pathway is not only restricted to

methylotrophs but also present in non-methylotrophic bacteria that use MMA as a

nitrogen but not as a carbon source (Chen et al., 2010a;Chen, 2012;Taubert et al.,

2017).

In a recent study, we isolated an alphaproteobacterial facultative methylotrophic

bacterium, *Gemmobacter* sp. LW-1 (recently renamed from *Catellibacterium* (Chen et

al., 2013)) from the Movile Cave ecosystem (Kumaresan et al., 2014) (Mangalia,

Romania) that can use methylated amines as both carbon and nitrogen source

(Wischer et al., 2015) and subsequently obtained its genome sequence (Kumaresan et

al., 2015). Using a ^{13}C-MMA DNA based stable-isotope probing (SIP) experiment we

also showed that *Gemmobacter* sp. LW-1 was indeed an active MMA utiliser in

microbial mats from this environment (Wischer et al., 2015). This was the first report of methylated amine utilisation in a member of the bacterial genus *Gemmobacter*. However, growth on C_1 compounds (methanol and formate) has been reported for the genus *Gemmobacter*, e.g. in *G. caeni* (Zheng et al., 2011).

The genus *Gemmobacter* (family *Rhodobacteraceae*) currently comprises ten validated species: *Gemmobacter megaterium* (Liu et al., 2014), *G. nectariphilum* (Tanaka et al., 2004;Chen et al., 2013), *G. aquatilis* (Rothe et al., 1987), *G. caeni* (Zheng et al., 2011;Chen et al., 2013), *G. aquaticus* (Liu et al., 2010;Chen et al., 2013), *G. nanjingense* (Zhang et al., 2012;Chen et al., 2013), *G. intermedius* (Kämpfer et al., 2015), *G. lanyuensis* (Sheu et al., 2013b), *G. tilapiae* (Sheu et al., 2013a) and *G. fontiphilus* (Chen et al., 2013). These species were isolated from a wide range of environments including fresh water environments (freshwater pond (Rothe et al., 1987;Sheu et al., 2013a), freshwater spring (Chen et al., 2013;Sheu et al., 2013b)), coastal planktonic seaweed (Liu et al., 2014), white stork nestling (Kämpfer et al., 2015), waste water and activated sludge (Tanaka et al., 2004;Zheng et al., 2011;Zhang et al., 2012), suggesting that members of the genus *Gemmobacter* are widely distributed in engineered and natural environments.

Here, using a comparative genomics approach we study how widespread methylated amine utilisation trait is within the members of the genus *Gemmobacter*. We used five isolate genomes for members within the genus *Gemmobacter* (*G*. sp. LW-1, *G. caeni*, *G. aquatilis*, *G. nectariphilus* and *G. megaterium)* available from public repositories (Accessed June 2018). These genomes were used for comparative genomics and

ancestral state reconstruction analyses to understand patterns of gain/loss of

methylated amine utilisation across the *Gemmobacter* phylogeny.

Materials and Methods

Genome data acquisition

Five *Gemmobacter* genomes (*G. caeni*, *G. aquatilis*, *G. nectariphilus*, *G. megaterium*,

Gemmobacter sp. LW-1) available through the Integrated Microbial Genomes (IMG)

database (https://img.jgi.doe.gov/) were used for comparative genome analysis

(Markowitz et al., 2013). Accession numbers and genome characteristics are listed in

Supplementary Table S1.

Phylogenetic analysis

Phylogenetic relatedness between the different members of the genus *Gemmobacter*

was determined using phylogenetic trees constructed from 16S rRNA gene sequences

(nucleotide) and metabolic gene sequences (*gmaS* and *mauA*; amino acids) involved

in MMA utilisation. RNAmmer (Lagesen et al., 2007) was used to retrieve 16S rRNA

gene sequences from the genome sequences. Multiple sequence alignment of 16S

rRNA gene sequences was performed using the SINA alignment service via SILVA

(Pruesse et al., 2007;Pruesse et al., 2012) and subsequently imported into MEGA7

(Kumar et al., 2016) to construct a maximum-likelihood nucleotide-based

phylogenetic tree (Saitou and Nei, 1987). Bootstrap analysis was performed with

1000 replicates to provide confidence estimates for phylogenetic tree topologies

(Felsenstein, 1985).

To determine phylogenetic affiliations for the protein encoding genes *gmaS* and *mauA*, gene sequences retrieved from the genome sequences were aligned to homologous sequences retrieved from the NCBI Genbank database using Basic Local Alignment Search Tool (BLAST, blastx) (Altschul et al., 1990) and curated *gmaS* sequences used for primer design in our previous study (Wischer et al., 2015). Amino acid sequences were aligned in MEGA7 (Kumar et al., 2016) using ClustalW (Thompson et al., 1994) and the alignment was subsequently used to construct maximum likelihood phylogenetic trees based on the JTT matrix-based model (Jones et al., 1992). Bootstrap analysis was performed with 1000 replicates to provide confidence estimates for phylogenetic tree topologies (Felsenstein, 1985).

Comparative genomic analyses

CGView Comparison Tool (CCT) was used to visually compare the genomes within the genus *Gemmobacter* (Grant et al., 2012). CCT utilises BLAST to compare the genomes and the BLAST results are presented in a DNA-based graphical map (Grant et al., 2012). Average Nucleotide Identity (ANI) (Rodriguez-R and Konstantinidis, 2016) between different genomes was estimated using one-way ANI (best hit) and two-way ANI (reciprocal best hit) based on Goris *et al*. (Goris et al., 2007). In addition the whole-genome based average nucleotide identity (gANI) and the $p_r^{intra\text{-}species}$ value were determined for *G*. sp. LW-1 and *G. caeni* (these two genomes revealed the closest ANI) based on Konstantinidis and Tiedje (Konstantinidis and Tiedje, 2005) via the Joint Genome Institute (JGI) platform (https://ani.jgi-psf.org/html/home.php; Version 0.3, April 2014). In order to determine if two genomes belong to the same species, the computation of empirical probabilities ($p_r^{intra\text{-}species}$) can be calculated as follows,

$$p_r^{intra-species}[AF = a, ANI = b] = p_r^{intra-species}[AF = a] * p_r^{intra-species}[ANI = b|AF = a]$$

AF represents alignment fraction.

Pan-genome analysis including average amino acid identity (AAI) analysis, pan-

genome tree construction and determination of core and dispensable genes and

singletons (unique genes) was carried out using the Efficient Database framework for

comparative Genome Analyses using BLAST score Ratios (EDGAR) platform (Blom

et al., 2016).

In order to compare the genetic potential for methylated amine utilisation within the

available *Gemmobacter* genomes, known protein sequences involved in methylated

amine utilisation pathways (Latypova et al., 2010;Chen, 2012) were used as query

sequences through the BLAST (blastp) program (Altschul et al., 1990) available

within the Rapid Annotation using Subsystem Technology (RAST) server (Aziz et al.,

2008). The list of protein queries used is given in Supplementary Table S2.

We examined patterns of gain/loss of methylated amine utilisation along the above

16S rRNA gene phylogeny, by performing ancestral state reconstruction analysis

using the phytools package in R (Revell, 2012). We used stochastic character

mapping (Nielsen, 2002) to map presence/absence of methylated amine utilisation,

assigning a prior probability of one to species known to utilise methylated amines,

zero to those known not to do so, and 0.5 to those where the trait value was unknown.

We used MCMC (with the function *make.simmap)* to simulate 1000 stochastic maps,

and then we used the function *densityMap* (Revell, 2013) to visualise the aggregate

result from the stochastic mapping analysis.

Results and discussion

Phylogenetic relatedness based on the 16S rRNA and metabolic gene

The phylogenetic relatedness of the five members within the genus *Gemmobacter* (*G.*

sp. LW-1, *G. caeni*, *G. aquatilis*, *G. nectariphilus* and *G. megaterium*) was resolved

based on 16S rRNA gene sequences (Figure 1). Three members of the genus

Gemmobacter (*G.* sp. LW-1, *G. caeni*, and *G. aquatilis)* clustered together with

several other related *Gemmobacter* and *Rhodobacter* 16S rRNA gene sequences

retrieved from fresh water, soil and sediment and activated sludge environments

(Figure 1). *G. nectariphilus* and *G. megaterium* sequences clustered together with

Paracoccus kawasakiensis and other related *Gemmobacter* sequences from marine,

fresh water and activated sludge environments (Figure 1). Based on the 16S rRNA

gene sequences retrieved from public database, we observed that the members of the

genus *Gemmobacter* are widely distributed in engineered (such as activated sludge

and clinical environments) and natural environments i.e. fresh water, soil and

sediment, and marine environments (Figure 1). GMA synthetase, a key enzyme in the

NMG pathway, is encoded by the gene *gmaS*. *gmaS* sequences retrieved from the

isolate genomes along with other ratified *gmaS* sequences were used to construct an

amino acid-based phylogenetic tree (Figure 2). *gmaS* gene sequences retrieved from

genomes of *G.* sp. LW-1, *G. caeni* and *G. aquatilis* clustered within Group I of

alphaproteobacterial *gmaS* sequences containing sequences from marine and non-

marine bacteria within the orders *Rhodobacterales* and *Rhizobiales* as described in

Wischer et al. (Wischer et al., 2015) and were closely related to *Paracoccus yeei*, *P.*

sp. 1W-5 and *Rhodobacter* sp. 1W-5 (Figure 2).

Whilst *gmaS* gene sequences were detected in three of the five investigated

Gemmobacter genomes, *mauA* gene sequences were identified only in the genomes of

G. caeni and *G.* sp. LW-1 (Supplementary Figure S1). It has been suggested that the

NMG pathway for MMA utilisation is more universally distributed and more

abundant across proteobacterial methylotrophs than the direct MMA oxidation

pathway (Nayak and Marx, 2015). However, it should be noted that genes encoding

for the enzymes within the NMG pathway (*gmaS*) can not only be detected in

methylotrophs but also in non-methylotrophic bacteria that use MMA as a nitrogen

source, but not as a carbon source (Chen, 2012;Wischer et al., 2015).

A comparative genome analysis of members within the genus Gemmobacter

At the time of the analysis, five *Gemmobacter* genomes obtained from isolates from

different environments were available (Figure 1 and Table 2). *Gemmobacter* genome

sizes range from ~3.96 Mb to ~5.14 Mb with GC contents between 64.71% to 66.19%

(Table 2). Analysis of sequence annotations revealed that on average 91.13% of the

genomes consist of coding sequences.

The genomes were compared using the CGView comparison tool (Grant et al., 2012)

(Figure 3). *Gemmobacter* sp. LW-1, isolated from the Movile Cave ecosystem was

used as the reference genome and the results of the BLAST comparison with other

Gemmobacter genomes are represented as a BLAST ring for each genome (Figure 3).

Similarities between segments of the reference genome sequence and the other

genome sequences are shown by a coloured arc beneath the region of similarity

indicating the percentage of similarity as a colour code. Our analysis (Figure 3)

revealed low identity levels (mostly <88%) between *Gemmobacter* sp. LW-1 and *G.*

aquatilis, *G. nectariphilus* and *G. megaterium* across the genomes. Moreover, the

analysis suggested several sites of potential insertion/deletion events in the genome of

Gemmobacter sp. LW-1. Possible insertion/deletion regions can be identified as those

gaps in the map where no homology is detected. For example the region between

2200-2300 kbp (Figure 3) where a gap can be found in the otherwise contiguous

homologous regions between the reference genome *G.* sp. LW-1 and the first of the

query genomes (*G. caeni*). This might likely be due to a lack of hits or hits with low

identity that can be spurious matches. Since it covers a large region we could possibly

rule out that it is not an artefact arising from a lack of sensitivity in the BLAST

analysis. Even though the genomes of *G.* sp. LW-1 and *G. caeni* are closely related,

our analysis demonstrates that their genomes are not completely identical. Despite the

fact that the majority of their genomes indicate very high identity levels (mostly >96-

98% as shown by the dominance of dark red colours of the circle representing the

BLAST hit identity between *G.* sp. LW-1 and *G. caeni*), many segments appear to be

exclusive to *G.* sp. LW-1.

In order to further resolve the similarity between these genomes we calculated the

average nucleotide identity (ANI) (Rodriguez-R and Konstantinidis, 2016)

(Supplementary Table S3 and Supplementary Figure S2A-D) and the average amino

acid identity,(AAI; Supplementary Figure S2E). It is generally accepted that an ANI

value of >95-96% can be used for species delineation (Richter and Rossello-Mora,

2009;Kim et al., 2014). Our analysis revealed that *Gemmobacter* sp. LW-1 and

Gemmobacter caeni share an ANI value of 98.62 (Supplementary Table S3) implying

that both are in fact the same species. The genome-based average nucleotide identity

(gANI) between *G.* sp. LW-1 and *G. caeni* was calculated as 98.70. The AF was

calculated to be 0.91, which would result in a computed probability of 0.98

suggesting that both genomes might belong to the same species. However, it should

be noted that these are draft genomes and a more in depth characterization of their

physiology and phenotype is required to delineate these organisms at the level of

strain.

Pan-genome analysis, carried out using the EDGAR platform (Blom et al., 2016),

identified metabolic genes present in all *Gemmobacter* species (core genes), two or

more *Gemmobacter* species (accessory or dispensable genes), and unique

Gemmobacter species (singleton genes). A pan-genome tree was constructed (Figure

4A) based on the pan-genome dataset and neighbor-joining method (Saitou and Nei,

1987). As with the 16S-rRNA gene based phylogenetic tree (Figure 1), the five

Gemmobacter species formed two main clusters in the pan-genome tree analysis

(Figure 4A). The pan-genome tree also confirmed the phylogenetic closeness between

Gemmobacter caeni and *Gemmobacter* sp. LW-1 (Figure 4A). According to pan-

genome analysis of the five *Gemmobacter* genomes, a total of 9,286 genes were

identified, consisting of 1,806 core genes, 3085 dispensable genes and 305, 1,072,

896, 1,165 and 957 singletons for *G.* sp. LW-1, *G. caeni*, *G. aquatilis*, *G.*

nectariphilus and *G. megaterium*, respectively (Figure 4B).

Methylated amine utilisation, N assimilation and C$_1$ oxidation

Investigation of the methylated amine utilisation pathways in five *Gemmobacter*

species revealed the presence of the genes encoding enzymes TMA dehydrogenase

(*tmd*), TMA monooxygenase (*tmm*), TMAO demethylase (*tdm*) and DMA

monooxygenase in genomes of *G.* sp. LW-1, *G. caeni* and *G. aquatilis* while none of

these genes were detected in *G. nectariphilus* or *G. megaterium* (Figure 5). These

findings are supported by results from a previous study which showed growth of *G.*

sp. LW-1 on TMA as a carbon and nitrogen source (Wischer et al., 2015). *G.* sp. LW-

1, *G. caeni* and *G. aquatilis* could potentially use the TMA oxidation pathway to

convert TMA to DMA. Based on the genome sequences, it can be suggested that

these three species could use the enzyme DMA monooxygenase (*dmmDABC*) to

oxidize DMA to MMA but not the DMA dehydrogenase since the corresponding

protein encoding gene (*dmd*) was not found (Figure 5).

We also compared the distribution of the direct MMA-oxidation and the NMG

pathways in the genomes of five *Gemmobacter* species (Figure 5). The direct MMA-

oxidation pathway (*mauA*-dependent) is so far only known to be present in

methylotrophic bacteria that can use MMA as a carbon source. Whereas the NMG

pathway (*gmaS*-dependent) has been shown to be present in non-methylotrophic

bacteria that can use MMA as a nitrogen source (Chen et al., 2010a;Nayak and Marx,

2015;Wischer et al., 2015;Nayak et al., 2016). Analysis of the genome sequences

revealed that both *G.* sp. LW-1 and *G. caeni* possess genes for both MMA oxidation

pathways (Figure 5). We have previously shown that *Gemmobacter* sp. LW-1 can use

MMA and TMA as both a carbon and nitrogen source (Wischer et al., 2015). Genome

sequence of *G. aquatilis* indicated the presence of genes involved only in the NMG

pathway. In the facultative methylotroph *Methylobacterium extorquens* AM1 it has

been shown that the NMG pathway is advantageous compared to the direct MMA-

oxidation pathway (Nayak et al., 2016). NMG pathway enables facultative

methylotrophic bacteria to switch between using MMA as a nitrogen source or as a

carbon and energy source whereas the direct MMA oxidation pathway allows for

334 rapid growth on MMA only as the primary energy and carbon source (Nayak et al.,

335 2016). This could suggest that *G. aquatilis* might use the NMG pathway for utilising

336 MMA as both nitrogen and carbon source. However, growth assays are required to

337 confirm whether *G. aquatilis* can use MMA as a carbon source. We did not detect

338 genes for either MMA oxidation pathways in the genome sequences of *G.*

339 *nectariphilus* and *G. megaterium* suggesting the lack of genetic potential of these

340 organisms to use MMA as either C or N source.

342 The C_1 units derived from methylated amines need to be further oxidized when the

343 nitrogen is sequestered without assimilation of the carbon from the methylated

344 amines. Genome analysis confirmed that all five *Gemmobacter* species possess the

345 genetic capability for C_1 oxidation and also indicate that tetrahydrofolate (H_4F) is the

346 C_1 carrier (Figure 5). The bifunctional enzyme 5,10-methylene-tetrahydrofolate

347 dehydrogenase/ cyclohydrolase, encoded by the gene *folD*, was detected in all the

348 *Gemmobacter* genomes (Figure 5/ Table 1). Genes encoding key enzymes in the C_1

349 oxidation pathway via tetrahydromethanopterin (H_4MPT) were not detected.

350 (Chistoserdova, 2011). The formate-tetrahydrofolate ligase, encoded by the gene *fhs*

351 (Figure 5), provides C_1 units for biosynthetic pathways (Chen, 2012). However, the

352 oxidation of formyl-H_4F (CHO-H_4F) can also be facilitated by *purU*, the gene

353 encoding for the formyl-H_4F deformylase. The formate dehydrogenase (*fdh*) mediates

354 the last step of the C_1 oxidation pathway, the oxidation of formate to CO_2. The genes

355 for the C_1 oxidation pathway via H_4F were detected in all five *Gemmobacter*

356 genomes.

The *fae* gene, encoding the formaldehyde-activating enzyme that catalyses the reduction of formaldehyde with H_4MPT was not detected in any of the five *Gemmobacter* genomes confirming that these members of the genus *Gemmobacter* lack the H_4MPT pathway for formaldehyde oxidation (Figure 5/ Table 1). Investigation of the nitrogen assimilation pathway revealed the presence of the genes encoding glutamine synthetase (GS; *gluL*) and glutamine synthase (GOGAT; *glxB*) in all five *Gemmobacter* genomes. In bacteria this pathway is essential for glutamate synthesis at low ammonium concentrations (Chen, 2012).

Using comparative genome analysis we provide genome-based evidence that the two *Gemmobacter* isolates *G*. sp. LW-1 and *G. caeni* are capable of generating energy from complete oxidation of methylated amines via the H_4F-dependent pathway using either the NMG pathway or the direct MMA oxidation pathway. *Gemmobacter aquatilis* is genetically capable of methylated amine degradation to yield formaldehyde and only encodes the genes for the NMG pathway, which indicates that *G. aquatilis* could use this pathway to use MMA as a nitrogen source. Both *G. nectariphilus* and *G. megaterium* genomes indicate the lack of potential to use methylated amines (Figure 5/Table 1). Therefore, the question arises if the genes for methylated amine utilisation have been acquired or lost.

Stochastic character mapping along the 16S rRNA gene phylogeny suggested that the ability to use methylated amines has either been gained or lost multiple times (Supplementary Figure S3). Most likely is that methylated amine utilisation is not ancestral in *Gemmobacter,* and has evolved three times, once in the clade containing *G*. sp. LW-1, *G. caeni* and *G. aquatilis,* once in *Haematobacter* and once in

Paracoccus (Supplementary Figure S3)*.* An alternative, but less plausible, scenario is

that methylated amine utilisation is an ancestral trait in *Gemmobacter,* and has been

lost and regained multiple times across the phylogeny.

Gemmobacter sp. LW-1 was isolated from the Movile Cave ecosystem (Wischer et

al., 2015). Microbial mats and lake water within the cave have been shown to harbor a

wide diversity of methylated amine-utilising bacteria (Wischer et al.,

2015;Kumaresan et al., 2018). Whilst the mechanism of MAs production within the

system has to be elucidated, it can be speculated that degradation of floating microbial

mats (i.e. organic matter) could result in MAs (Wischer et al., 2015). Similarly, *G.*

caeni isolated from activated sludge (Zheng et al., 2011) could possibly use the MAs

generated from organic matter degradation. Interestingly, whilst *G. megaterium* was

isolated from a marine environment (seaweed (Liu et al., 2014)) possibly

encountering MAs from the degradation of osmolytes such as glycine betaine (*N,N,N-*

trimethylglycine) we did not detect metabolic genes involved in methylated amine

utilisation. Our analyses suggest that the trait for methylated amine utilisation could

be independent of the habitat.

Conclusions

In summary, three of the five investigated *Gemmobacter* genomes (*G.* sp. LW-1, *G.*

caeni and *G. aquatilis*) indicated metabolic potential to utilise methylated amines, of

which only two (*G.* sp. LW-1 and *G. caeni*) possess the genes for both MMA

oxidation pathways, the NMG pathway and the direct MMA oxidation pathway. *G.*

sp. LW-1 and *G. caeni* are facultative methylotrophs which could potentially use

these pathways to utilise MMA as both a carbon and nitrogen source, while

potentially *G. aquatilis* could only use the NMG pathway as a nitrogen source.

Furthermore, the genomes of *G.* sp. LW-1 and *G. caeni* showed a high similarity to

each other (>98%) suggesting that both belong to the same species. *G. megaterium*

and *G. nectariphilus* genomes indicated no metabolic potential to utilise MAs.

Phylogenetic, pan-genome and ANI analyses revealed that *G.* sp. LW-1 and *G. caeni*

are closely related, although they were isolated from different environments. Whilst

G. caeni and *G. nectariphilus* were isolated from a similar environment (activated

sludge) it revealed a high amount of evolutionary change from the common ancestor.

Overall, these results suggest that the trait for methylated amine utilisation could be

independent from the habitat and localised factors or selection pressures could

influence the ability of these organisms to use methylated amines. Access to

Gemmobacter isolates with or without the genetic potential for methylated amine

utilisation trait will allow us to perform physiological experiments in future to test

how this trait can affect fitness of closely related organisms. Ancestral state

reconstruction analysis confirms that across *Gemmobacter* and related genera,

methylated amine utilisation has either evolved or been lost multiple times over the

evolutionary history of this group. The adaptive or non-adaptive processes behind this

pattern remain to be investigated.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank Dr Felipe Hernandes Coutinho (Federal University of Rio de Janeiro) for a

fruitful discussion on genomic regions of insertion and deletion in *Gemmobacter* sp.

LW-1. The EDGAR platform is financially supported by the BMBF grant FKZ

031A533 within the de.NBI network.

458   REFERENCES

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local
alignment search tool. *Journal of Molecular Biology* 215**,** 403-410.
Anthony, C. (1982). *The biochemistry of methylotrophs.* London: Academic Press.
Aziz, R.K., Bartels, D., Best, A.A., Dejongh, M., Disz, T., Edwards, R.A., Formsma, K.,
Gerdes, S., Glass, E.M., and Kubal, M. (2008). The RAST Server: rapid
annotations using subsystems technology. *BMC Genomics* 9**,** 75.
Blom, J., Kreis, J., Spanig, S., Juhre, T., Bertelli, C., Ernst, C., and Goesmann, A.
(2016). EDGAR 2.0: an enhanced software platform for comparative gene
content analyses. *Nucleic Acids Research* 44**,** W22-W28.
Burke, S.A., Lo, S.L., and Krzycki, J.A. (1998). Clustered genes encoding the
methyltransferases of methanogenesis from monomethylamine. *Journal*
of Bacteriology 180**,** 3432-3440.
Carpenter, L.J., Archer, S.D., and Beale, R. (2012). Ocean-atmosphere trace gas
exchange. *Chemical Society Reviews* 41**,** 6473-6506.
Chen, W.M., Cho, N.T., Huang, W.C., Young, C.C., and Sheu, S.Y. (2013). Description
of *Gemmobacter fontiphilus* sp nov., isolated from a freshwater spring,
reclassification of *Catellibacterium nectariphilum* as *Gemmobacter*
nectariphilus comb. nov., *Catellibacterium changlense* as *Gemmobacter*
changlensis comb. nov., *Catellibacterium aquatile* as *Gemmobacter*
aquaticus nom. nov., *Catellibacterium caeni* as *Gemmobacter caeni* comb.
nov., *Catellibacterium nanjingense* as *Gemmobacter nanjingensis* comb.
nov., and emended description of the genus *Gemmobacter* and of
Gemmobacter aquatilis. International Journal of Systematic and
Evolutionary Microbiology 63**,** 470-478.
Chen, Y. (2012). Comparative genomics of methylated amine utilization by
marine *Roseobacter* clade bacteria and development of functional gene
markers (*tmm, gmaS*). *Environmental Microbiology* 14**,** 2308-2322.
Chen, Y., Mcaleer, K.L., and Murrell, J.C. (2010a). Monomethylamine as a nitrogen
source for a nonmethylotrophic bacterium, *Agrobacterium tumefaciens*.
Applied and Environmental Microbiology 76**,** 4102-4104.
Chen, Y., Scanlan, J., Song, L., Crombie, A., Rahman, M.T., Schäfer, H., and Murrell,
J.C. (2010b). γ-Glutamylmethylamide is an essential intermediate in the
metabolism of methylamine by *Methylocella silvestris. Applied and*
Environmental Microbiology 76**,** 4530-4537.
Chen, Y., Wu, L., Boden, R., Hillebrand, A., Kumaresan, D., Moussard, H., Baciu, M.,
Lu, Y., and Colin Murrell, J. (2009). Life without light: microbial diversity
and evidence of sulfur- and ammonium-based chemolithotrophy in
Movile Cave. *ISME Journal* 3**,** 1093-1104.
Chistoserdova, A.Y., Christoserdova, L.V., Mcintire, W.S., and Lidstrom, M.E.
(1994). Genetic organization of the mau gene cluster in *Methylobacterium*
extorquens AM1: complete nucleotide sequence and generation and
characteristics of mau mutants. *Journal of Bacteriology* 176**,** 4052-4065.
Chistoserdova, L. (2011). Modularity of methylotrophy, revisited. *Environmental*
Microbiology 13**,** 2603-2622.
Chistoserdova, L., Kalyuzhnaya, M.G., and Lidstrom, M.E. (2009). The expanding
world of methylotrophic metabolism. *Annual Review of Microbiology* 63**,**
477-499.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the
508   bootstrap. *Evolution* 39**,** 783-791.
509   Ge, X., Wexler, A.S., and Clegg, S.L. (2011). Atmospheric amines – Part I. A review.
510   *Atmospheric Environment* 45**,** 524-546.
511   Goris, J., Konstantinidis, K.T., Klappenbach, J.A., Coenye, T., Vandamme, P., and
512   Tiedje, J.M. (2007). DNA–DNA hybridization values and their relationship
513   to whole-genome sequence similarities. *International Journal of*
514   *Systematic and Evolutionary Microbiology* 57**,** 81-91.
515   Grant, J.R., Arantes, A.S., and Stothard, P. (2012). Comparing thousands of
516   circular genomes using the CGView Comparison Tool. *BMC genomics* 13**,**
517   202.
518   Jones, D.T., Taylor, W.R., and Thornton, J.M. (1992). The rapid generation of
519   mutation data matrices from protein sequences. *Bioinformatics* 8**,** 275-
520   282.
521   Kämpfer, P., Jerzak, L., Wilharm, G., Golke, J., Busse, H.-J., and Glaeser, S.P. (2015).
522   *Gemmobacter intermedius* sp. nov., isolated from a white stork (Ciconia
523   ciconia). *International Journal of Systematic and Evolutionary Microbiology*
524   65**,** 778-783.
525   Kim, M., Oh, H.-S., Park, S.-C., and Chun, J. (2014). Towards a taxonomic
526   coherence between average nucleotide identity and 16S gene sequence
527   similarity for species demarcation of prokaryotes. *International Journal of*
528   *Systematic and Evolutionary Microbiology* 64**,** 346-351.
529   Konstantinidis, K.T., and Tiedje, J.M. (2005). Genomic insights that advance the
530   species definition for prokaryotes. *Proceedings of the National Academy of*
531   *Sciences of the United States of America* 102**,** 2567-2572.
532   Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary
533   Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and*
534   *Evolution* 33**,** 1870-1874.
535   Kumaresan, D., Stephenson, J., Doxey, A.C., Bandukwala, H., Brooks, E.,
536   Hillebrand-Voiculescu, A., Whiteley, A.S., and Murrell, J.C. (2018). Aerobic
537   proteobacterial methylotrophs in Movile Cave: genomic and metagenomic
538   analyses. *Microbiome* 6.
539   Kumaresan, D., Wischer, D., Hillebrand-Voiculescu, A.M., and Murrell, J.C. (2015).
540   Draft genome sequences of facultative methylotrophs, *Gemmobacter* sp.
541   strain LW1 and *Mesorhizobium* sp. strain 1M-11, isolated from Movile
542   Cave, Romania. *Genome Announcements* 3.
543   Kumaresan, D., Wischer, D., Stephenson, J., Hillebrand-Voiculescu, A., and
544   Murrell, J.C. (2014). Microbiology of Movile Cave - A
545   chemolithoautotrophic ecosystem. *Geomicrobiology Journal* 31**,** 186-193.
546   Lagesen, K., Hallin, P., Rødland, E.A., Stærfeldt, H.-H., Rognes, T., and Ussery, D.W.
547   (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA
548   genes. *Nucleic acids research* 35**,** 3100-3108.
549   Latypova, E., Yang, S., Wang, Y.S., Wang, T., Chavkin, T.A., Hackett, M., Schäfer, H.,
550   and Kalyuzhnaya, M.G. (2010). Genetics of the glutamate‐mediated
551   methylamine utilization pathway in the facultative methylotrophic beta‐
552   proteobacterium *Methyloversatilis universalis* FAM5. *Molecular*
553   *Microbiology* 75**,** 426-439.

Lidbury, I., Murrell, J.C., and Chen, Y. (2014). Trimethylamine N-oxide
metabolism by abundant marine heterotrophic bacteria. *Proceedings of*
the National Academy of Sciences 111**,** 2710-2715.
Lidstrom, M.E. (2006). "Aerobic Methylotrophic Prokaryotes," in *The*
Prokaryotes: Volume 2: Ecophysiology and Biochemistry, eds. M. Dworkin,
S. Falkow, E. Rosenberg, K.-H. Schleifer & E. Stackebrandt. (New York, NY:
Springer New York), 618-634.
Liu, J.-J., Zhang, X.-Q., Chi, F.-T., Pan, J., Sun, C., and Wu, M. (2014). *Gemmobacter*
megaterium sp. nov., isolated from coastal planktonic seaweeds.
International Journal of Systematic and Evolutionary Microbiology 64**,** 66-
71.
Liu, Y., and Whitman, W.B. (2008). Metabolic, phylogenetic, and ecological
diversity of the methanogenic archaea. *Annals of the New York Academy of*
Sciences 1125**,** 171-189.
Liu, Y., Xu, C.-J., Jiang, J.-T., Liu, Y.-H., Song, X.-F., Li, H., and Liu, Z.-P. (2010).
Catellibacterium aquatile sp. nov., isolated from fresh water, and emended
description of the genus *Catellibacterium* Tanaka et al. 2004. *International*
Journal of Systematic and Evolutionary Microbiology 60**,** 2027-2031.
Lyimo, T.J., Pol, A., Jetten, M.S.M., and Op Den Camp, H.J.M. (2009). Diversity of
methanogenic archaea in a mangrove sediment and isolation of a new
Methanococcoides strain. *FEMS Microbiology Letters* 291**,** 247-253.
Markowitz, V.M., Chen, I.-M.A., Palaniappan, K., Chu, K., Szeto, E., Pillay, M.,
Ratner, A., Huang, J., Woyke, T., and Huntemann, M. (2013). IMG 4 version
of the integrated microbial genomes comparative analysis system. *Nucleic*
acids research 42**,** D560-D567.
Mcintire, W.S., Wemmer, D.E., Chistoserdova, A., and Lidstrom, M.E. (1991). A
new cofactor in a prokaryotic enzyme: tryptophan tryptophylquinone as
the redox prosthetic group in methylamine dehydrogenase. *Science* 252**,**
817-824.
Nayak, D.D., Agashe, D., Lee, M.-C., and Marx, C.J. (2016). Selection maintains
apparentlydegenerate metabolic pathwaysdueto tradeoffsin using
methylamine for carbon versus nitrogen. *Current Biology* 26**,** 1-11.
Nayak, D.D., and Marx, C.J. (2015). Experimental horizontal gene transfer of
methylamine dehydrogenase mimics prevalent exchange in nature and
overcomes the methylamine growth constraints posed by the sub-optimal
N-Methylglutamate pathway. *Microorganisms* 3**,** 60-79.
Nielsen, R. (2002). Mapping mutations on phylogenies. *Systematic Biology* 51**,**
729-739.
Paul, L., Ferguson, D.J., and Krzycki, J.A. (2000). The trimethylamine
methyltransferase gene and multiple dimethylamine methyltransferase
genes of *Methanosarcina barkeri* contain in-frame and read-through
amber codons. *Journal of Bacteriology* 182**,** 2520-2529.
Pruesse, E., Peplies, J., and Glöckner, F.O. (2012). SINA: accurate high-throughput
multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* 28**,**
1823-1829.
Pruesse, E., Quast, C., Knittel, K., Fuchs, B.M., Ludwig, W., Peplies, J., and Glöckner,
F.O. (2007). SILVA: a comprehensive online resource for quality checked
and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic*
Acids Research 35**,** 7188-7196.

Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology
(and other things). *Methods in Ecology and Evolution* 3**,** 217-223.
Revell, L.J. (2013). Two new graphical methods for mapping trait evolution on
phylogenies. *Methods in Ecology and Evolution* 4**,** 754-759.
Richter, M., and Rossello-Mora, R. (2009). Shifting the genomic gold standard for
the prokaryotic species definition. *Proceedings of the National Academy of*
Sciences 106**,** 19126-19131.
Rodriguez-R, L.M., and Konstantinidis, K.T. (2016). The enveomics collection: a
toolbox for specialized analyses of microbial genomes and metagenomes.
PeerJ Preprints.
Rothe, B., Fischer, A., Hirsch, P., Sittig, M., and Stackebrandt, E. (1987). The
phylogenetic position of the budding bacteria *Blastobacter aggregatus*
and *Gemmobacter aquatilis* gen., nov. sp. nov. *Archives of Microbiology*
147**,** 92-99.
Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for
reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4**,** 406-
425.
Schade, G.W., and Crutzen, P.J. (1995). Emission of aliphatic-amines from animal
husbandry and their reactions - potential source of N_2O and HC_n. *Journal*
of Atmospheric Chemistry 22**,** 319-346.
Sheu, S.-Y., Sheu, D.-S., Sheu, F.-S., and Chen, W.-M. (2013a). *Gemmobacter tilapiae*
sp. nov., a poly-β-hydroxybutyrate-accumulating bacterium isolated from
a freshwater pond. *International Journal of Systematic and Evolutionary*
Microbiology 63**,** 1550-1556.
Sheu, S.-Y., Shiau, Y.-W., Wei, Y.-T., and Chen, W.-M. (2013b). *Gemmobacter*
lanyuensis sp. nov., isolated from a freshwater spring. *International*
Journal of Systematic and Evolutionary Microbiology 63**,** 4039-4045.
Tanaka, Y., Hanada, S., Manome, A., Tsuchida, T., Kurane, R., Nakamura, K., and
Kamagata, Y. (2004). *Catellibacterium nectariphilum* gen. nov., sp. nov.,
which requires a diffusible compound from a strain related to the genus
Sphingomonas for vigorous growth. *International Journal of Systematic*
and Evolutionary Microbiology 54**,** 955-959.
Taubert, M., Grob, C., Howat, A.M., Burns, O.J., Pratscher, J., Jehmlich, N., Von
Bergen, M., Richnow, H.H., Chen, Y., and Murrell, J.C. (2017). Methylamine
as a nitrogen source for microorganisms from a coastal marine
environment. *Environmental Microbiology* 19**,** 2246-2257.
Thompson, J.D., Higgins, D.G., and Gibson, T.J. (1994). CLUSTAL W: improving the
sensitivity of progressive multiple sequence alignment through sequence
weighting, position-specific gap penalties and weight matrix choice.
Nucleic Acids Research 22**,** 4673-4680.
Wischer, D., Kumaresan, D., Johnston, A., El Khawand, M., Stephenson, J.,
Hillebrand-Voiculescu, A.M., Chen, Y., and Colin Murrell, J. (2015).
Bacterial metabolism of methylated amines and identification of novel
methylotrophs in Movile Cave. *ISME Journal* 9**,** 195-206.
Zhang, J., Chen, S.-A., Zheng, J.-W., Cai, S., Hang, B.-J., He, J., and Li, S.-P. (2012).
Catellibacterium nanjingense sp. nov., a propanil-degrading bacterium
isolated from activated sludge, and emended description of the genus
Catellibacterium. International Journal of Systematic and Evolutionary
Microbiology 62**,** 495-499.

Zheng, J.-W., Chen, Y.-G., Zhang, J., Ni, Y.-Y., Li, W.-J., He, J., and Li, S.-P. (2011).
Description of *Catellibacterium caeni* sp. now., reclassification of
Rhodobacter changlensis Anil Kumar et al. 2007 as *Catellibacterium*
changlense comb. nov. and emended description of the genus
Catellibacterium. International Journal of Systematic and Evolutionary
Microbiology 61**,** 1921-1926.

Figures and Tables

Figure 1. Phylogenetic tree based on 16S rRNA gene sequences. The tree was constructed using the maximum likelihood method for clustering and the Tamura-Nei model for computing evolutionary distances. Numbers at branches are bootstrap percentages >50% of 1000 replicates. Star represents the *Gemmobacter* species used for comparative genome analysis. Coloured boxes represent the habitat where the sequence was retrieved: blue (fresh water), orange (soil and sediment), green (activated sludge), grey (marine), purple (clinical source). Triangles represent sequences that are listed as *Catellibacterium* in the NCBI database, which have been recently reclassified to *Gemmobacter* (Chen et al., 2013). Scale bar: 0.02 substitutions per nucleotide position.

Figure 2. Maximum-likelihood phylogenetic tree based on *gmaS* sequences. The tree was constructed using amino acid sequences (GmaS) using the maximum-likelihood method based on the JTT matrix-based model. Members of the genus *Gemmobacter* used for genome comparison are represented with a star. Numbers at branches are bootstrap percentages >50% of 1000 replicates. Amino acid sequences of the glutamine synthetase type III (GlnA) were used as out-group. Scale bar: 0.1 substitutions per amino acid position. MRC, marine *Roseobacter* clade.

Figure 3. DNA BLAST map of *Gemmobacter* genomes. *Gemmobacter* sp. LW-1 was used as a reference genome against *Gemmobacter megaterium* (inner ring), *Gemmobacter aquatilis* (second inner ring), *Gemmobacter nectariphilus* (third ring), and *Gemmobacter caeni* (fourth ring). The fifth and sixth ring (outer rings) represent the CDS (blue), tRNA (maroon), and rRNA (purple) on the reverse and forward

strand, respectively. The colour scale (inset) shows the level of sequence identity with

the respective sequences from *G. megaterium*, *G. aquatilis*, *G. nectariphilus* and *G.*

caeni. The locations of genes involved in methylotrophy are indicated at the outside

of the map.

Figure 4. Pan-genome analysis. (A) Pan-genome tree consisting of five

Gemmobacter species was constructed using the neighbour-joining method within the

EDGAR platform. **(B)** Number of core, dispensable, and specific genes (singletons)

of each *Gemmobacter* species.

Figure 5. Metabolic pathways involved in methylated amine utilisation and one-

carbon utilisation annotated with presence/absence of specific genes in the

genomes of *Gemmobacter*. The analysis was based on a five-way comparison among

Gemmobacter sp. LW-1 (L), *Gemmobacter caeni* (C), *Gemmobacter aquatilis* (A),

Gemmobacter nectariphilus (N) and *Gemmobacter megaterium* (M). The color-coded

boxes next to the genes indicate the presence (green) or absence (orange) of a gene in

each genome.

Table 1. Comparative genomic analysis of methylated amine-utilising genes in

genomes-sequenced *Gemmobacters* in comparison to selected marine *Roseobacter*

clade bacteria. Shown is the presence (+) or absence (-) of specific genes in the

genome sequences.

Supplementary Figure S1. Maximum-likelihood phylogenetic tree (JTT matrix-

based) of *mauA* sequences. Sequences from the genus *Gemmobacter* are marked

with a star. Amino acid sequences (MauA) were aligned using the ClustalW

algorithm. Numbers at branches are bootstrap percentages >50% of 1000 replicates.

Scale bar: 0.1 substitutions per amino acid. Coloured boxes indicate

*Alphaproteobacteria (*yellow), *Gammaproteobacteria (*red) and *Betaproteobacteria*

*(*black).

Supplementary Figure S2. (A-D) Average nucleotide identity (ANI) analysis of

***Gemmobacter* sp. LW-1 and *Gemmobacter caeni*, *Gemmobacter aquatilis*,**

***Gemmobacter nectariphilus* and *Gemmobacter megaterium* and (E) AAI analysis**

between those species

Supplementary Figure S3. Ancestral state reconstruction of methylated amine

utilisation along the 16S rRNA gene phylogeny, using stochastic mapping. Branch

colour represents the posterior probability (computed as the relative frequency across

stochastic maps) of methylated amine utilisation through the phylogeny. Red indicates

a high posterior probability of methylated amine utilisation.

Supplementary Table S1. Genome characteristics of the five *Gemmobacter* isolate

genomes used in this study.

Supplementary Table S2. List of protein queries used for the genome

comparison with their accession number.

Supplementary Table S3. Average nucleotide identity (ANI) values between

Gemmobacter sp. LW-1 and *Gemmobacter caeni*, *Gemmobacter aquatilis*,

Gemmobacter nectariphilus, Gemmobacter megaterium, Rhodobacter sphaeroides

and *Paracoccus denitrificans*

Figure 1

Figure 2

Figure 3

Protein	Corresponding gene(s)
TMA monooxygenase	tmm
TMA methyltransferase	mttB1, mttB2, mttB3, mttB4
MMA dehydrogenase	mauA, mauD, mauE, mauG
NMG synthase	mgsA, mgsB, mgsC
NMG dehydrogenase	mgdA, mgdB, mgdC, mgdD
HCHO-activating enzyme	gfa1, gfa2, gfa3, gfa4
GSH-dependent HCHO dehydrogenase	adhI
S-formyl-GSH hydrolase	fghA
Glutamine synthetase (GS)	gluL
Glutamate synthase (GOGAT)	glxB, glxC
Methylene-H_4F-dehydro-genase/cyclohydrolase	folD
Formate-H_4F ligase	fhs
Formyl-H_4F deformylase	purU
Formate dehydrogenase	fdh
Carbon monoxide dehydrogenase	coxD, coxE, coxF,coxG, coxL, coxM, coxS
Methanol dehydrogenase	xoxF
N-acetylglutamate synthase	nagS
Gamma-glutamylcysteine synthetase	gcl
Methylenetetrahydrofolate reductase	metF
Glutamate dehydrogenase	gdh2
Ammonium transporter	amtB

Figure 4A

Tree scale: 0.01

Gemmobacter aquatilis
Gemmobacter sp. LW-1
Gemmobacter caeni
Gemmobacter megaterium
Gemmobacter nectariphilus

Figure 4B

G. sp. LW-1

G. aquatilis

G. caeni

1072

Dispensable

Core
1806

3085

G. nectariphilus

1165

G. megaterium

Figure 5

Table 1. Comparative genomic analysis of methylamine-utilizing genes in genome-sequenced Gemmobacters in comparison to selected marine *Roseobacter* clade bacteria. Shown is the presence (+) or absence (-) of the corresponding genes in the genome sequences.

Organism	gmaS	tmm	xoxF	fae	foldD	purU	fhs (ftfL)	gfa	adhI (flhA)	fghA	fdh	amtB	mtdA/ mtdB	mchA	ftr	fmdA /B/C	mauA	cox
Citreicella sp. SE45	+	+	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	+
Roseovarius sp. TM1035	+	+	+	-	+	-	+	-	+	+	+	+	-/-	-	-	-	-	+
Paracoccus denitrificans PD1222	-	-	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	-
Rhodobacter sphaeroides 241	-	+	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter caeni	+	+	+	-	+	+	+	+a	+	+	+	+	-/-	-	-	-	+	+
Gemmobacter sp. LW-1	+	+	+	-	+	+	+	+a	+	+	+	+	-/-	-	-	-	+	+
Gemmobacter aquatilis	+	+	+	-	+	+	+	-	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter nectariphilus DSM-15620	-	-	-	-	+	+	+	+a	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter megaterium DSM-26375	-	-	-	-	+	+	+	+a	+	+	+	+	-/-	-	-	-	-	+

a gfa-like protein-encoding gene