Title:

Comparative genomics analyses indicate differential methylated amine

utilisation trait within members of the genus *Gemmobacter*

Running title:

Methylated amine utilisation trait in *Gemmobacter*

Kröber, E.[1], Peixoto, J[2]., Cunningham, MR[4]., Spurgin L.[3], Wischer, D[3]., Kruger, R[2].,

Kumaresan, D.[4#]

[1] Microbial Biogeochemistry, RA Landscape Functioning, ZALF Leibniz Centre for

Landscape Research, Mu☐ncheberg, Germany

[2]Department of Cellular Biology, University of Brasília, Brasília, Brazil

[3] School of Environmental Sciences, University of East Anglia, Norwich, UK

[4] School of Biological Sciences & Institute for Global Food Security, Queen's

University Belfast, Belfast, Northern Ireland, UK

[#]Correspondence to

Dr D Kumaresan, School of Biological Sciences, Queen's University Belfast, Belfast,

Northern Ireland, United Kingdom, d.kumaresan@qub.ac.uk

Abstract

Methylated amines are ubiquitous in the environment and play a role in regulating the earth's climate via a set of complex biological and chemical reactions. Microbial degradation of these compounds is thought to be a major sink. Recently we isolated a facultative methylotroph, *Gemmobacter* sp. LW-1, an isolate from the unique environment Movile Cave, Romania, which is capable of methylated amine utilisation as a carbon source. Here, using a comparative genomics approach, we investigate how widespread methylated amine utilisation trait is within the member of the bacterial genus *Gemmobacter*. Five genomes of different *Gemmobacter* species isolated from diverse environments, such as activated sludge, fresh water, sulphuric cave waters (Movile Cave) and the marine environment were available from the public repositories and used for the analysis. Our results indicate that some members of the genus *Gemmobacter*, namely *G. aquatilis*, *G. caeni* and *G.* sp. LW-1 have the genetic potential of methylated amine utilisation while others (*G. megaterium* and *G. nectariphilus*) have not.

Introduction

Methylated amines (MAs) are ubiquitous in the environment with a variety of natural and anthropogenic sources including the oceans, vegetation, sediments and organic-rich soils, animal husbandry, food industry, pesticides, sewage, and automobiles, to mention only a few [1-3]. Methylated amines are also known to influence earth's climate, via a series of complex biological and chemical interactions [4]. Some of the most abundant methylated amines found in the atmosphere are trimethylamine (TMA), dimethylamine (DMA) and monomethylamine (MMA) [1]. Microbial

metabolism of methylated amines involves both aerobic and anaerobic

microorganisms, e.g. some methanogenic archaea such as *Methanosarcina* and

Methanomicrobium can use MAs to produce methane [5-7] while Gram-positive and

Gram-negative methylotrophic bacteria can use MAs as carbon and nitrogen source

[8]. Previously, MAs were typically associated with marine ecosystems as they are

by-products of degradation of osmolytic chemicals such as glycine betaine, carnitine,

choline and trimethylamine N-oxide [8]. However, recent studies have reported the

detection and activity of aerobic methylotrophic bacteria that utilise MAs in a variety

of natural and engineered environments [1, 9-12] and could play a major role in

global C and N budgets.

Aerobic methylotrophs are a polyphyletic group of microorganisms capable of

utilising one-carbon (C_1) compounds such as methane, methanol or methylated

amines as their sole source of carbon and energy [9, 13, 14]. Methylotrophs can

degrade TMA to DMA by using the enzymes TMA dehydrogenase, TMA

monooxygenase or TMA methyltransferase (under anaerobic conditions by

methylotrophic methanogens), encoded by the genes *tdm*, *tmm* and *mtt*, respectively

[15-18]. The enzymes DMA dehydrogenase (*dmd*) or DMA monooxygenase

(*dmmDABC*) modulate the conversion of DMA to MMA [14, 15]. Two distinct

pathways have been characterised for the oxidation of MMA [10]. The direct MMA-

oxidation pathway mediated by a single enzyme (MMA dehydrogenase in Gram-

negative bacteria and MMA oxidase in Gram-positive bacteria) converts MMA to

formaldehyde and releases ammonium [19, 20]. The alternate pathway, referred to as

the *N*-methylglutamate (NMG) pathway or indirect MMA-oxidation pathway, is

mediated by three individual enzymes via the oxidation of MMA to gamma-

glutamylmethylamide (GMA) and its further degradation to N-methylglutamate

(NMG) and 5,10-methylenetetrahydrofolate ($CH_2 = H_4F$) [3, 10]. A stepwise

conversion of MMA in the NMG pathway is modulated by the enzymes GMA

synthetase (*gmaS*), NMG synthase (*mgsABC*) and NMG dehydrogenase (*mgdABCD*)

[3, 8]. The capability to use MMA not only as a source for carbon but also for

nitrogen is widespread in bacteria. Notably, the NMG pathway is not only restricted

to methylotrophs but also present in non-methylotrophic bacteria that use MMA as a

nitrogen but not as a carbon source [15, 21, 22].

In a recent study, we isolated an alphaproteobacterial facultative methylotrophic

bacterium, *Gemmobacter* sp. LW-1 (recently renamed from *Catellibacterium* [23])

from the Movile Cave ecosystem (Mangalia, Romania) [24] that can use methylated

amines as both carbon and nitrogen source [12] and subsequently obtained its genome

sequence [25]. Using a ^{13}C-MMA DNA based stable-isotope probing (SIP)

experiment we also showed that *Gemmobacter* sp. LW-1 was indeed an active MMA

utiliser in microbial mats from this environment [12]. This was the first report of

methylated amine utilisation in a member of the bacterial genus *Gemmobacter*.

However, growth on C_1 compounds (methanol and formate) has been reported for the

genus *Gemmobacter*, e.g. in *G. caeni* [26]. The genus *Gemmobacter* (family

Rhodobacteraceae) currently comprises ten validated species: *Gemmobacter*

megaterium [27], *G. nectariphilum* [23, 28], *G. aquatilis* [29], *G. caeni* [23, 26], *G.*

aquaticus [23, 30], *G. nanjingense* [23, 31], *G. intermedius* [32], *G. lanyuensis* [33],

G. tilapiae [34] and *G. fontiphilus* [23]. These species were isolated from a wide

range of environments including fresh water environments (freshwater pond [29, 34],

freshwater spring [23, 33]), coastal planktonic seaweed [27], white stork nestling [32],

waste water and activated sludge [26, 28, 31], suggesting that members of the genus

Gemmobacter are widely distributed in engineered and natural environments.

Here, using a comparative genomics approach we study how widespread methylated

amine utilisation trait (i.e. metabolic potential) is within the members of the genus

Gemmobacter. We used five isolate genomes for members within the genus

Gemmobacter (*G*. sp. LW-1, *G. caeni*, *G. aquatilis*, *G. nectariphilus* and *G.*

megaterium) alongside genomes of four closely related organisms within the family

Rhodobacteraceae to show that the methylated amine utilisation trait is not common

within the members of the genus *Gemmobacter.*

Materials and Methods

Genome data acquisition

Five *Gemmobacter* genomes (*G. caeni*, *G. aquatilis*, *G. nectariphilus*, *G. megaterium*,

Gemmobacter sp. LW-1) available through the Integrated Microbial Genomes (IMG)

database (https://img.jgi.doe.gov/) were used for comparative genome analysis [35].

Accession numbers and genome characteristics are listed in Supplementary Table S1.

Phylogenetic analysis

Phylogenetic relatedness between the different members of the genus *Gemmobacter*

was determined using phylogenetic trees constructed from 16S rRNA gene sequences

(nucleotide) and metabolic gene sequences (*gmaS* and *mauA*; amino acids) involved

in MMA utilisation. RNAmmer [36] was used to retrieve 16S rRNA gene sequences

from the genome sequences. Multiple sequence alignment of 16S rRNA gene

sequences was performed using the SINA alignment service via SILVA [37, 38] and

subsequently imported into MEGA7 [39] to construct a maximum-likelihood

nucleotide-based phylogenetic tree [40]. Bootstrap analysis was performed with 1000

replicates to provide confidence estimates for phylogenetic tree topologies [41].

To determine phylogenetic affiliations for the protein encoding genes *gmaS* and

mauA, gene sequences retrieved from the genome sequences were aligned to

homologous sequences retrieved from the NCBI Genbank database using Basic Local

Alignment Search Tool (BLAST, blastx) [42] and curated *gmaS* sequences used for

primer design in our previous study [12]. Amino acid sequences were aligned in

MEGA7 [39] using ClustalW [43] and the alignment was subsequently used to

construct maximum likelihood phylogenetic trees based on the JTT matrix-based

model [44]. Bootstrap analysis was performed with 1000 replicates to provide

confidence estimates for phylogenetic tree topologies [41].

We investigated the phylogeny of nine bacterial species – *Gemmobacter aquatilis*,

Gemmobacter caeni, *Gemmobacter sp. LW-1*, *Gemmobacter megabacterium* DSM-

26375, *Gemmobacter nectariphilus* DSM-15620, *Paracoccus denitrificans* PD1222,

Citreicella sp. SE45, *Roseovarius* sp. TM1035 and *Rhodobacter sphaeroides* 241 –

using the protein sequences of 831 orthologs present in single copy in all species

identified using OrthoFinder [45] with default settings. For each set of orthologous

proteins, multiple alignments were produced using MAFFT [46] with the --auto

settings. Subsequently, conserved alignment blocks were identified using trimal [47]

with the option -automated1. The phylogenetic reconstruction analysis was based in a

final concatenated alignment consisting of 265,536 amino acid positions, which was

the input for RAxML [48] with the option PROTCATLGF. The selection of best

amino acid substitution model was performed after implementing the script

ProteinModelSelection.pl provided by the RAxML package. In order to assess the

environmental distribution of the genus *Gemmobacter*, we used MAPseq [49]

(https://beta.microbeatlas.org) to survey the relative abundance based on 16S rRNA

gene sequences (query sequence: *Gemmobacter aquatilis* DSM3857 (NR_104740.1 at

97% cut-off).

Comparative genomic analyses

CGView Comparison Tool (CCT) was used to visually compare the genomes within

the genus *Gemmobacter* [50]. CCT utilises BLAST to compare the genomes and the

BLAST results are presented in a DNA-based graphical map [50]. Average

Nucleotide Identity (ANI) [51] between different genomes was estimated using one-

way ANI (best hit) and two-way ANI (reciprocal best hit) based on Goris *et al*. [52].

In addition the whole-genome based average nucleotide identity (gANI) and the $p_r^{\text{intra-}}$

$^{\text{species}}$ value were determined for *G.* sp. LW-1 and *G. caeni* (these two genomes

revealed the closest ANI) based on Konstantinidis and Tiedje [53] via the Joint

Genome Institute (JGI) platform (https://ani.jgi-psf.org/html/home.php; Version 0.3,

April 2014). In order to determine if two genomes belong to the same species, the

computation of empirical probabilities ($p_r^{\text{intra-species}}$) can be calculated as follows,

$$p_r^{intra-species}[AF = a, ANI = b] = p_r^{intra-species}[AF = a] * p_r^{intra-species}[ANI = b | AF = a]$$

AF represents alignment fraction.

Pan-genome analysis including average amino acid identity (AAI) analysis, pan-

genome tree construction and determination of core and dispensable genes and

singletons (unique genes) was carried out using the Efficient Database framework for

comparative Genome Analyses using BLAST score Ratios (EDGAR) platform [54].

Estimation of genomic completeness and contamination was carried out using the

CheckM program [55].

In order to compare the genetic potential for methylated amine utilisation within the

available *Gemmobacter* genomes, known protein sequences involved in methylated

amine utilisation pathways [3, 15] were used as query sequences through the BLAST

(blastp) program [42] available within the Rapid Annotation using Subsystem

Technology (RAST) server [56]. The list of protein queries used is given in

Supplementary Table S2.

Results and discussion

Analysis of phylogenetic relatedness

The phylogenetic relatedness of the five members within the genus *Gemmobacter* (*G.*

sp. LW-1, *G. caeni*, *G. aquatilis*, *G. nectariphilus* and *G. megaterium*) was resolved

based on 16S rRNA gene sequences (Figure 1). Three members of the genus

Gemmobacter (*G.* sp. LW-1, *G. caeni*, and *G. aquatilis)* clustered together with

several other related *Gemmobacter* and *Rhodobacter* 16S rRNA gene sequences

retrieved from fresh water, soil and sediment and activated sludge environments

(Figure 1). *G. nectariphilus* and *G. megaterium* sequences clustered together with

Paracoccus kawasakiensis and other related *Gemmobacter* sequences from marine,

fresh water and activated sludge environments (Figure 1). Based on the 16S rRNA

gene sequences retrieved from public database and MAPseq analyses, we observed

that the members of the genus *Gemmobacter* are widely distributed in engineered

(such as activated sludge and clinical environments) and natural environments i.e.

fresh water, soil and sediment, and marine environments (Figure 1 and Supplementary

Figure S3). Phylogenomic analysis based on single copy marker genes

(Supplementary Figure S4) revealed that *G.* sp LW1, *G. caeni and G. aquatilis* were

closely related to *Rhodobacter sphaeroides* 241 whereas *G. megabacterium and G.*

nectariphilus to *Paracoccus denitrificans.*

GMA synthetase, a key enzyme in the NMG pathway, is encoded by the gene *gmaS*.

gmaS sequences retrieved from the isolate genomes along with other ratified *gmaS*

sequences were used to construct an amino acid-based phylogenetic tree (Figure 2).

gmaS gene sequences retrieved from genomes of *G.* sp. LW-1, *G. caeni* and *G.*

aquatilis clustered within Group I of alphaproteobacterial *gmaS* sequences containing

sequences from marine and non-marine bacteria within the orders *Rhodobacterales*

and *Rhizobiales* as described in Wischer et al. [12] and were closely related to

Paracoccus yeei, *P.* sp. 1W-5 and *Rhodobacter* sp. 1W-5 (Figure 2). Whilst *gmaS*

gene sequences were detected in three of the five investigated *Gemmobacter*

genomes, *mauA* gene sequences were identified only in the genomes of *G. caeni* and

G. sp. LW-1 (Supplementary Figure S1). It has been suggested that the NMG

pathway for MMA utilisation is more universally distributed and more abundant

across proteobacterial methylotrophs than the direct MMA oxidation pathway [57].

However, it should be noted that genes encoding for the enzymes within the NMG

pathway (*gmaS*) can not only be detected in methylotrophs but also in non-

methylotrophic bacteria that use MMA as a nitrogen source, but not as a carbon

source [12, 15].

A comparative genome analysis of members within the genus Gemmobacter

At the time of the analysis, five *Gemmobacter* genomes obtained from isolates from

different environments were available (Figure 1 and Table 2). *Gemmobacter* genome

sizes range from ~3.96 Mb to ~5.14 Mb with GC contents between 64.71% to 66.19%

and genome completeness between 98.61% to 99.69% (Table 2; Supplementary Table

1). Analysis of sequence annotations revealed that on average 91.13% of the genomes

consist of coding sequences.

The genomes were compared using the CGView comparison tool [50] (Figure 3).

Gemmobacter sp. LW-1, isolated from the Movile Cave ecosystem was used as the

reference genome and the results of the BLAST comparison with other *Gemmobacter*

genomes are represented as a BLAST ring for each genome (Figure 3). Similarities

between segments of the reference genome sequence and the other genome sequences

are shown by a coloured arc beneath the region of similarity indicating the percentage

of similarity as a colour code. Our analysis (Figure 3) revealed low identity levels

(mostly <88%) between *Gemmobacter* sp. LW-1 and *G. aquatilis*, *G. nectariphilus*

and *G. megaterium* across the genomes. Moreover, the analysis suggested several

sites of potential insertion/deletion events in the genome of *Gemmobacter* sp. LW-1.

Possible insertion/deletion regions can be identified as those gaps in the map where

no homology is detected. For example, the region between 2200-2300 kbp (Figure 3)

where a gap can be found in the otherwise contiguous homologous regions between

the reference genome *G.* sp. LW-1 and the first of the query genomes (*G. caeni*).

This might likely be due to a lack of hits or hits with low identity that can be spurious

matches. Since it covers a large region we could possibly rule out that it is not an

artefact arising from a lack of sensitivity in the BLAST analysis. Even though the

genomes of *G.* sp. LW-1 and *G. caeni* are closely related, our analysis demonstrates

that their genomes are not completely identical. Despite the fact that the majority of

their genomes indicate very high identity levels (mostly >96-98% as shown by the

dominance of dark red colours of the circle representing the BLAST hit identity

between *G.* sp. LW-1 and *G. caeni*), many segments appear to be exclusive to *G.* sp.

LW-1.

In order to further resolve the similarity between these genomes we calculated the

average nucleotide identity (ANI) [51] (Supplementary Table S3 and Supplementary

Figure S2A-D) and the average amino acid identity,(AAI; Supplementary Figure

S2E). It is generally accepted that an ANI value of >95-96% can be used for species

delineation [58, 59]. Our analysis revealed that *Gemmobacter* sp. LW-1 and

Gemmobacter caeni share an ANI value of 98.62 (Supplementary Table S3) implying

that both are in fact the same species. The genome-based average nucleotide identity

(gANI) between *G.* sp. LW-1 and *G. caeni* was calculated as 98.70. The AF was

calculated to be 0.91, which would result in a computed probability of 0.98

suggesting that both genomes might belong to the same species. However, it should

be noted that these are draft genomes and a more in depth characterization of their

physiology and phenotype is required to delineate these organisms at the level of

strain.

Pan-genome analysis, carried out using the EDGAR platform [54], identified

metabolic genes present in all *Gemmobacter* species (core genes), two or more

Gemmobacter species (accessory or dispensable genes), and unique *Gemmobacter*

species (singleton genes). A pan-genome tree was constructed (Figure 4A) based on

the pan-genome dataset and neighbor-joining method [40]. As with the 16S-rRNA

gene based phylogenetic tree (Figure 1), the five *Gemmobacter* species formed two

main clusters in the pan-genome tree analysis (Figure 4A). The pan-genome tree also

confirmed the phylogenetic closeness between *Gemmobacter caeni* and *Gemmobacter*

sp. LW-1 (Figure 4A). According to pan-genome analysis of the five *Gemmobacter*

genomes, a total of 9,286 genes were identified, consisting of 1,806 core genes, 3085

dispensable genes and 305, 1,072, 896, 1,165 and 957 singletons for *G.* sp. LW-1, *G.*

caeni, *G. aquatilis*, *G. nectariphilus* and *G. megaterium*, respectively (Figure 4B).

Methylated amine utilisation, N assimilation and C₁ oxidation

Investigation of the methylated amine utilisation pathways in five *Gemmobacter*

species revealed the presence of the genes encoding enzymes TMA dehydrogenase

(*tmd*), TMA monooxygenase (*tmm*), TMAO demethylase (*tdm*) and DMA

monooxygenase in genomes of *G.* sp. LW-1, *G. caeni* and *G. aquatilis* while none of

these genes were detected in *G. nectariphilus* or *G. megaterium* (Figure 5). These

findings are supported by results from a previous study which showed growth of *G.*

sp. LW-1 on TMA as a carbon and nitrogen source [12]. *G.* sp. LW-1, *G. caeni* and

G. aquatilis could potentially use the TMA oxidation pathway to convert TMA to

DMA. Based on the genome sequences, it can be suggested that these three species

could use the enzyme DMA monooxygenase (*dmmDABC*) to oxidize DMA to MMA

but not the DMA dehydrogenase since the corresponding protein encoding gene

(*dmd*) was not found (Figure 5).

We also compared the distribution of the direct MMA-oxidation and the NMG

pathways in the genomes of five *Gemmobacter* species (Figure 5) and gene

arrangement (Supplementary Figure S5). The direct MMA-oxidation pathway (*mauA-*

dependent) is so far only known to be present in methylotrophic bacteria that can use

MMA as a carbon source. Whereas the NMG pathway (*gmaS*-dependent) has been

shown to be present in non-methylotrophic bacteria that can use MMA as a nitrogen

source [12, 21, 57, 60]. Analysis of the genome sequences revealed that both *G*. sp.

LW-1 and *G. caeni* possess genes for both MMA oxidation pathways (Figure 5). We

have previously shown that *Gemmobacter* sp. LW-1 can use MMA and TMA as both

a carbon and nitrogen source [12]. Genome sequence of *G. aquatilis* indicated the

presence of genes involved only in the NMG pathway. In the facultative methylotroph

Methylobacterium extorquens AM1 it has been shown that the NMG pathway is

advantageous compared to the direct MMA-oxidation pathway [60]. NMG pathway

enables facultative methylotrophic bacteria to switch between using MMA as a

nitrogen source or as a carbon and energy source whereas the direct MMA oxidation

pathway allows for rapid growth on MMA only as the primary energy and carbon

source [60]. This could suggest that *G. aquatilis* might use the NMG pathway for

utilising MMA as both nitrogen and carbon source. However, growth assays are

required to confirm whether *G. aquatilis* can use MMA as a carbon source. We did

not detect genes for either MMA oxidation pathways in the genome sequences of *G.*

nectariphilus and *G. megaterium* suggesting the lack of genetic potential of these

organisms to use MMA as either C or N source.

The C_1 units derived from methylated amines need to be further oxidized when the

nitrogen is sequestered without assimilation of the carbon from the methylated

amines. Genome analysis confirmed that all five *Gemmobacter* species possess the

genetic capability for C_1 oxidation and also indicate that tetrahydrofolate (H_4F) is the

C_1 carrier (Figure 5). The bifunctional enzyme 5,10-methylene-tetrahydrofolate

dehydrogenase/ cyclohydrolase, encoded by the gene *folD*, was detected in all the

Gemmobacter genomes (Figure 5/ Table 1). Genes encoding key enzymes in the C_1

oxidation pathway via tetrahydromethanopterin (H_4MPT) were not detected. [10]. The

formate-tetrahydrofolate ligase, encoded by the gene *fhs* (Figure 5), provides C_1 units

for biosynthetic pathways [15]. However, the oxidation of formyl-H_4F (CHO-H_4F)

can also be facilitated by *purU*, the gene encoding for the formyl-H_4F deformylase.

The formate dehydrogenase (*fdh*) mediates the last step of the C_1 oxidation pathway,

the oxidation of formate to CO_2. The genes for the C_1 oxidation pathway via H_4F

were detected in all five *Gemmobacter* genomes.

The *fae* gene, encoding the formaldehyde-activating enzyme that catalyses the

reduction of formaldehyde with H_4MPT was not detected in any of the five

Gemmobacter genomes confirming that these members of the genus *Gemmobacter*

lack the H_4MPT pathway for formaldehyde oxidation (Figure 5/ Table 1).

Investigation of the nitrogen assimilation pathway revealed the presence of the genes

encoding glutamine synthetase (GS; *gluL)* and glutamine synthase (GOGAT; *glxB*) in

all five *Gemmobacter* genomes. In bacteria this pathway is essential for glutamate

synthesis at low ammonium concentrations [15].

Using comparative genome analysis we provide genome-based evidence that the two

Gemmobacter isolates *G.* sp. LW-1 and *G. caeni* are capable of generating energy

from complete oxidation of methylated amines via the H_4F-dependent pathway using

either the NMG pathway or the direct MMA oxidation pathway. *Gemmobacter*

aquatilis is genetically capable of methylated amine degradation to yield

formaldehyde and only encodes the genes for the NMG pathway, which indicates that

G. aquatilis could use this pathway to use MMA as a nitrogen source. Both *G.*

nectariphilus and *G. megaterium* genomes indicate the lack of potential to use

methylated amines (Figure 5/Table 1).

Gemmobacter sp. LW-1 was isolated from the Movile Cave ecosystem [12].

Microbial mats and lake water within the cave have been shown to harbor a wide

diversity of methylated amine-utilising bacteria [12, 61]. Whilst the mechanism of

MAs production within the system has to be elucidated, it can be speculated that

degradation of floating microbial mats (i.e. organic matter) could result in MAs [12].

Similarly, *G. caeni* isolated from activated sludge [26] could possibly use the MAs

generated from organic matter degradation. Interestingly, whilst *G. megaterium* was

isolated from a marine environment (seaweed [27]) possibly encountering MAs from

the degradation of osmolytes such as glycine betaine (*N,N,N*-trimethylglycine) we did

not detect metabolic genes involved in methylated amine utilisation. Our genome-

based analyses suggest that the potential for methylated amine utilisation is

differential within the members of the genus *Gemmobacter* that requires confirmation

using growth/activity assays.

Conclusions

In summary, three of the five investigated *Gemmobacter* genomes (*G.* sp. LW-1, *G.*

caeni and *G. aquatilis*) indicated metabolic potential to utilise methylated amines, of

which only two (*G.* sp. LW-1 and *G. caeni*) possess the genes for both MMA

oxidation pathways, the NMG pathway and the direct MMA oxidation pathway. *G.*

sp. LW-1 and *G. caeni* are facultative methylotrophs which could potentially use

these pathways to utilise MMA as both a carbon and nitrogen source, while

potentially *G. aquatilis* could only use the NMG pathway as a nitrogen source.

Furthermore, the genomes of *G.* sp. LW-1 and *G. caeni* showed a high similarity to

each other (>98%) suggesting that both belong to the same species. *G. megaterium*

and *G. nectariphilus* genomes indicated no metabolic potential to utilise MAs.

Phylogenetic, pan-genome and ANI analyses revealed that *G.* sp. LW-1 and *G. caeni*

are closely related, although they were isolated from different environments. Overall,

our results suggest that the trait for methylated amine utilisation could be independent

from the habitat and localised factors or selection pressures could influence the ability

of these organisms to use methylated amines. Access to *Gemmobacter* isolates with or

without the genetic potential for methylated amine utilisation trait will allow us to

perform physiological experiments in future to test how this trait can affect fitness of

closely related organisms.

Conflict of Interest

400   The authors declare no conflict of interest.

Acknowledgements

403   We thank Dr Felipe Hernandes Coutinho (Federal University of Rio de Janeiro) for a

404   fruitful discussion on genomic regions of insertion and deletion in *Gemmobacter* sp.

405   LW-1. The EDGAR platform is financially supported by the BMBF grant FKZ

406   031A533 within the de.NBI network. This manuscript has been published as preprint

407   at the preprint server for biology (bioRxiv.org, Kröber et al., 2018).

REFERENCES

1. Ge X, Wexler AS, Clegg SL (2011) Atmospheric amines – Part I. A review. Atmospheric Environment 45: 524-546. doi: http://dx.doi.org/10.1016/j.atmosenv.2010.10.012

2. Schade GW, Crutzen PJ (1995) Emission of Aliphatic-Amines from Animal Husbandry and Their Reactions - Potential Source of N2o and Hcn. J Atmos Chem 22: 319-346. doi: Doi 10.1007/Bf00696641

3. Latypova E, Yang S, Wang YS, Wang T, Chavkin TA, Hackett M, Schäfer H, Kalyuzhnaya MG (2010) Genetics of the glutamate‐mediated methylamine utilization pathway in the facultative methylotrophic beta‐proteobacterium Methyloversatilis universalis FAM5. Molecular microbiology 75: 426-439.

4. Carpenter LJ, Archer SD, Beale R (2012) Ocean-atmosphere trace gas exchange. Chemical Society Reviews 41: 6473-6506. doi: 10.1039/c2cs35121h

5. Lyimo TJ, Pol A, Jetten MSM, Op den Camp HJM (2009) Diversity of methanogenic archaea in a mangrove sediment and isolation of a new Methanococcoides strain. Fems Microbiol Lett 291: 247-253. doi: 10.1111/j.1574-6968.2008.01464.x

6. Burke SA, Lo SL, Krzycki JA (1998) Clustered genes encoding the methyltransferases of methanogenesis from monomethylamine. J Bacteriol 180: 3432-3440.

7. Liu Y, Whitman WB (2008) Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. Ann N Y Acad Sci 1125: 171-189. doi: 10.1196/annals.1419.019

8. Chen Y, Scanlan J, Song L, Crombie A, Rahman MT, Schäfer H, Murrell JC (2010) γ-Glutamylmethylamide is an essential intermediate in the metabolism of methylamine by Methylocella silvestris. Applied and environmental microbiology 76: 4530-4537.

9. Chistoserdova L, Kalyuzhnaya MG, Lidstrom ME (2009) The Expanding World of Methylotrophic Metabolism. Annual Review of Microbiology 63: 477-499. doi: 10.1146/annurev.micro.091208.073600

10. Chistoserdova L (2011) Modularity of methylotrophy, revisited. Environmental Microbiology 13: 2603-2622.

11. Chen Y, Wu L, Boden R, Hillebrand A, Kumaresan D, Moussard H, Baciu M, Lu Y, Colin Murrell J (2009) Life without light: microbial diversity and evidence of sulfur- and ammonium-based chemolithotrophy in Movile Cave. ISME J 3: 1093-1104.

12. Wischer D, Kumaresan D, Johnston A, El Khawand M, Stephenson J, Hillebrand-Voiculescu AM, Chen Y, Colin Murrell J (2015) Bacterial

456		metabolism of methylated amines and identification of novel
457		methylotrophs in Movile Cave. ISME J 9: 195-206. doi:
458		10.1038/ismej.2014.102
459	13.	Anthony C (1982) The biochemistry of methylotrophs. Academic Press,
460		London
461	14.	Lidstrom ME (2006) Aerobic Methylotrophic Prokaryotes. In: Dworkin, M,
462		Falkow, S, Rosenberg, E, Schleifer, K-H, Stackebrandt, E (eds.) The
463		Prokaryotes: Volume 2: Ecophysiology and Biochemistry. Springer New
464		York, New York, NY, pp. 618-634
465	15.	Chen Y (2012) Comparative genomics of methylated amine utilization by
466		marine *Roseobacter* clade bacteria and development of functional gene
467		markers (tmm, gmaS). Environmental Microbiology 14: 2308-2322.
468	16.	Lidbury I, Murrell JC, Chen Y (2014) Trimethylamine N-oxide metabolism
469		by abundant marine heterotrophic bacteria. Proceedings of the National
470		Academy of Sciences 111: 2710-2715.
471	17.	Paul L, Ferguson DJ, Krzycki JA (2000) The trimethylamine
472		methyltransferase gene and multiple dimethylamine methyltransferase
473		genes of Methanosarcina barkeri contain in-frame and read-through
474		amber codons. J Bacteriol 182: 2520-2529.
475	18.	Lidbury I, Mausz MA, Scanlan DJ, Chen Y (2017) Identification of
476		dimethylamine monooxygenase in marine bacteria reveals a metabolic
477		bottleneck in the methylated amine degradation pathway. ISME J 11:
478		1592-1601. doi: 10.1038/ismej.2017.31
479	19.	McIntire WS, Wemmer DE, Chistoserdova A, Lidstrom ME (1991) A new
480		cofactor in a prokaryotic enzyme: tryptophan tryptophylquinone as the
481		redox prosthetic group in methylamine dehydrogenase. Science 252: 817-
482		824.
483	20.	Chistoserdova AY, Christoserdova LV, McIntire WS, Lidstrom ME (1994)
484		Genetic organization of the mau gene cluster in Methylobacterium
485		extorquens AM1: complete nucleotide sequence and generation and
486		characteristics of mau mutants. J Bacteriol 176: 4052-4065.
487	21.	Chen Y, McAleer KL, Murrell JC (2010) Monomethylamine as a nitrogen
488		source for a nonmethylotrophic bacterium, Agrobacterium tumefaciens.
489		Applied and environmental microbiology 76: 4102-4104.
490	22.	Taubert M, Grob C, Howat AM, Burns OJ, Pratscher J, Jehmlich N, von
491		Bergen M, Richnow HH, Chen Y, Murrell JC (2017) Methylamine as a
492		nitrogen source for microorganisms from a coastal marine environment.
493		Environmental Microbiology 19: 2246-2257. doi: 10.1111/1462-
494		2920.13709
495	23.	Chen WM, Cho NT, Huang WC, Young CC, Sheu SY (2013) Description of
496		Gemmobacter fontiphilus sp nov., isolated from a freshwater spring,
497		reclassification of Catellibacterium nectariphilum as Gemmobacter
498		nectariphilus comb. nov., Catellibacterium changlense as Gemmobacter
499		changlensis comb. nov., Catellibacterium aquatile as Gemmobacter
500		aquaticus nom. nov., Catellibacterium caeni as Gemmobacter caeni comb.
501		nov., Catellibacterium nanjingense as Gemmobacter nanjingensis comb.
502		nov., and emended description of the genus Gemmobacter and of
503		Gemmobacter aquatilis. International Journal of Systematic and
504		Evolutionary Microbiology 63: 470-478. doi: 10.1099/ijs.0.042051-0

24. Kumaresan D, Wischer D, Stephenson J, Hillebrand-Voiculescu A, Murrell
JC (2014) Microbiology of Movile Cave - A chemolithoautotrophic
ecosystem. Geomicrobiology Journal 31: 186-193.
25. Kumaresan D, Wischer D, Hillebrand-Voiculescu AM, Murrell JC (2015)
Draft Genome Sequences of Facultative Methylotrophs, Gemmobacter sp.
Strain LW1 and Mesorhizobium sp. Strain 1M-11, Isolated from Movile
Cave, Romania. Genome Announc 3. doi: 10.1128/genomeA.01266-15
26. Zheng J-W, Chen Y-G, Zhang J, Ni Y-Y, Li W-J, He J, Li S-P (2011)
Description of Catellibacterium caeni sp. now., reclassification of
Rhodobacter changlensis Anil Kumar et al. 2007 as Catellibacterium
changlense comb. nov. and emended description of the genus
Catellibacterium. International Journal of Systematic and Evolutionary
Microbiology 61: 1921-1926.
27. Liu J-J, Zhang X-Q, Chi F-T, Pan J, Sun C, Wu M (2014) Gemmobacter
megaterium sp. nov., isolated from coastal planktonic seaweeds.
International Journal of Systematic and Evolutionary Microbiology 64: 66-
71. doi: doi:10.1099/ijs.0.050955-0
28. Tanaka Y, Hanada S, Manome A, Tsuchida T, Kurane R, Nakamura K,
Kamagata Y (2004) Catellibacterium nectariphilum gen. nov., sp. nov.,
which requires a diffusible compound from a strain related to the genus
Sphingomonas for vigorous growth. International Journal of Systematic
and Evolutionary Microbiology 54: 955-959. doi:
doi:10.1099/ijs.0.02750-0
29. Rothe B, Fischer A, Hirsch P, Sittig M, Stackebrandt E (1987) The
phylogenetic position of the budding bacteria Blastobacter aggregatus
and Gemmobacter aquatilis gen., nov. sp. nov. Archives of Microbiology
147: 92-99. doi: 10.1007/bf00492911
30. Liu Y, Xu C-J, Jiang J-T, Liu Y-H, Song X-F, Li H, Liu Z-P (2010)
Catellibacterium aquatile sp. nov., isolated from fresh water, and
emended description of the genus Catellibacterium Tanaka et al. 2004.
International Journal of Systematic and Evolutionary Microbiology 60:
2027-2031. doi: doi:10.1099/ijs.0.017632-0
31. Zhang J, Chen S-A, Zheng J-W, Cai S, Hang B-J, He J, Li S-P (2012)
Catellibacterium nanjingense sp. nov., a propanil-degrading bacterium
isolated from activated sludge, and emended description of the genus
Catellibacterium. International Journal of Systematic and Evolutionary
Microbiology 62: 495-499. doi: doi:10.1099/ijs.0.029819-0
32. Kämpfer P, Jerzak L, Wilharm G, Golke J, Busse H-J, Glaeser SP (2015)
Gemmobacter intermedius sp. nov., isolated from a white stork (Ciconia
ciconia). International Journal of Systematic and Evolutionary
Microbiology 65: 778-783. doi: doi:10.1099/ijs.0.000012
33. Sheu S-Y, Shiau Y-W, Wei Y-T, Chen W-M (2013) Gemmobacter lanyuensis
sp. nov., isolated from a freshwater spring. International Journal of
Systematic and Evolutionary Microbiology 63: 4039-4045. doi:
doi:10.1099/ijs.0.052399-0
34. Sheu S-Y, Sheu D-S, Sheu F-S, Chen W-M (2013) Gemmobacter tilapiae sp.
nov., a poly-β-hydroxybutyrate-accumulating bacterium isolated from a
freshwater pond. International Journal of Systematic and Evolutionary
Microbiology 63: 1550-1556. doi: doi:10.1099/ijs.0.044735-0

35. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner
A, Huang J, Woyke T, Huntemann M (2013) IMG 4 version of the
integrated microbial genomes comparative analysis system. Nucleic acids
research 42: D560-D567.
36. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW
(2007) RNAmmer: consistent and rapid annotation of ribosomal RNA
genes. Nucleic acids research 35: 3100-3108.
37. Pruesse E, Peplies J, Glöckner FO (2012) SINA: accurate high-throughput
multiple sequence alignment of ribosomal RNA genes. Bioinformatics 28:
1823-1829.
38. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO
(2007) SILVA: a comprehensive online resource for quality checked and
aligned ribosomal RNA sequence data compatible with ARB. Nucleic acids
research 35: 7188-7196.
39. Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary
Genetics Analysis version 7.0 for bigger datasets. Molecular biology and
evolution 33: 1870-1874.
40. Saitou N, Nei M (1987) The neighbor-joining method: a new method for
reconstructing phylogenetic trees. Molecular biology and evolution 4:
406-425.
41. Felsenstein J (1985) Confidence limits on phylogenies: an approach using
the bootstrap. Evolution 39: 783-791.
42. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local
alignment search tool. Journal of molecular biology 215: 403-410.
43. Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the
sensitivity of progressive multiple sequence alignment through sequence
weighting, position-specific gap penalties and weight matrix choice.
Nucleic acids research 22: 4673-4680.
44. Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of
mutation data matrices from protein sequences. Bioinformatics 8: 275-
282.
45. Emms DM, Kelly S (2015) OrthoFinder: solving fundamental biases in
whole genome comparisons dramatically improves orthogroup inference
accuracy. Genome biology 16: 157. doi: 10.1186/s13059-015-0721-2
46. Nakamura T, Yamada KD, Tomii K, Katoh K (2018) Parallelization of
MAFFT for large-scale multiple sequence alignments. Bioinformatics 34:
2490-2492. doi: 10.1093/bioinformatics/bty121
47. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool
for automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics 25: 1972-1973.
48. Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis
and post-analysis of large phylogenies. Bioinformatics 30: 1312-1313.
49. Matias Rodrigues JF, Schmidt TSB, Tackmann J, von Mering C (2017)
MAPseq: highly efficient k-mer search with confidence estimates, for
rRNA sequence analysis. Bioinformatics 33: 3808-3810. doi:
10.1093/bioinformatics/btx517
50. Grant JR, Arantes AS, Stothard P (2012) Comparing thousands of circular
genomes using the CGView Comparison Tool. BMC genomics 13: 202.

51.   Rodriguez-R LM, Konstantinidis KT (2016) The enveomics collection: a
603   toolbox for specialized analyses of microbial genomes and metagenomes.
604   PeerJ Preprints.
605   52.   Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje
606   JM (2007) DNA–DNA hybridization values and their relationship to
607   whole-genome sequence similarities. International journal of systematic
608   and evolutionary microbiology 57: 81-91.
609   53.   Konstantinidis KT, Tiedje JM (2005) Genomic insights that advance the
610   species definition for prokaryotes. P Natl Acad Sci USA 102: 2567-2572.
611   doi: 10.1073/pnas.0409727102
612   54.   Blom J, Kreis J, Spanig S, Juhre T, Bertelli C, Ernst C, Goesmann A (2016)
613   EDGAR 2.0: an enhanced software platform for comparative gene content
614   analyses. Nucleic Acids Research 44: W22-W28. doi:
615   10.1093/nar/gkw255
616   55.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015)
617   CheckM: assessing the quality of microbial genomes recovered from
618   isolates, single cells, and metagenomes. Genome research: gr.
619   186072.186114.
620   56.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K,
621   Gerdes S, Glass EM, Kubal M (2008) The RAST Server: rapid annotations
622   using subsystems technology. BMC genomics 9: 75.
623   57.   Nayak DD, Marx CJ (2015) Experimental horizontal gene transfer of
624   methylamine dehydrogenase mimics prevalent exchange in nature and
625   overcomes the methylamine growth constraints posed by the sub-optimal
626   N-Methylglutamate pathway. Microorganisms 3: 60-79.
627   58.   Richter M, Rossello-Mora R (2009) Shifting the genomic gold standard for
628   the prokaryotic species definition. Proceedings of the National Academy
629   of Sciences of the United States of America 106: 19126-19131.
630   59.   Kim M, Oh H-S, Park S-C, Chun J (2014) Towards a taxonomic coherence
631   between average nucleotide identity and 16S gene sequence similarity for
632   species demarcation of prokaryotes. International Journal of Systematic
633   and Evolutionary Microbiology 64: 346-351.
634   60.   Nayak DD, Agashe D, Lee M-C, Marx CJ (2016) Selection maintains
635   apparentlydegenerate metabolic pathwaysdueto tradeoffsin using
636   methylamine for carbon versus nitrogen. Current Biology 26: 1-11.
637   61.   Kumaresan D, Stephenson J, Doxey AC, Bandukwala H, Brooks E,
638   Hillebrand-Voiculescu A, Whiteley AS, Murrell JC (2018) Aerobic
639   proteobacterial methylotrophs in Movile Cave: genomic and metagenomic
640   analyses. Microbiome 6. doi: ARTN 1
641   10.1186/s40168-017-0383-2

642
643

Figures and Tables

Figure 1. Phylogenetic tree based on 16S rRNA gene sequences. The tree was

constructed using the maximum likelihood method for clustering and the Tamura-Nei

model for computing evolutionary distances. Numbers at branches are bootstrap

percentages >50% of 1000 replicates. Star represents the *Gemmobacter* species used

for comparative genome analysis. Coloured boxes represent the habitat where the

sequence was retrieved: blue (fresh water), orange (soil and sediment), green

(activated sludge), grey (marine), purple (clinical source). Triangles represent

sequences that are listed as *Catellibacterium* in the NCBI database, which have been

recently reclassified to *Gemmobacter* [23]. Scale bar: 0.02 substitutions per

nucleotide position.

Figure 2. Maximum-likelihood phylogenetic tree based on *gmaS* sequences. The

tree was constructed using amino acid sequences (GmaS) using the maximum-

likelihood method based on the JTT matrix-based model. Members of the genus

Gemmobacter used for genome comparison are represented with a star. Numbers at

branches are bootstrap percentages >50% of 1000 replicates. Amino acid sequences

of the glutamine synthetase type III (GlnA) were used as out-group. Scale bar: 0.1

substitutions per amino acid position. MRC, marine *Roseobacter* clade.

Figure 3. DNA BLAST map of *Gemmobacter* genomes. *Gemmobacter* sp. LW-1

was used as a reference genome against *Gemmobacter megaterium* (inner ring),

Gemmobacter aquatilis (second inner ring), *Gemmobacter nectariphilus* (third ring),

and *Gemmobacter caeni* (fourth ring). The fifth and sixth ring (outer rings) represent

the CDS (blue), tRNA (maroon), and rRNA (purple) on the reverse and forward

strand, respectively. The colour scale (inset) shows the level of sequence identity with

the respective sequences from *G. megaterium, G. aquatilis, G. nectariphilus* and *G.*

caeni. The locations of genes involved in methylotrophy are indicated at the outside

of the map.

Figure 4. Pan-genome analysis. (A) Pan-genome tree consisting of five

Gemmobacter species was constructed using the neighbour-joining method within the

EDGAR platform. **(B)** Number of core, dispensable, and specific genes (singletons)

of each *Gemmobacter* species.

Figure 5. Metabolic pathways involved in methylated amine utilisation and one-

carbon utilisation annotated with presence/absence of specific genes in the

genomes of *Gemmobacter*. The analysis was based on a five-way comparison among

Gemmobacter sp. LW-1 (L), *Gemmobacter caeni* (C), *Gemmobacter aquatilis* (A),

Gemmobacter nectariphilus (N) and *Gemmobacter megaterium* (M). The color-coded

boxes next to the genes indicate the presence (green) or absence (orange) of a gene in

each genome.

Table 1. Comparative genomic analysis of methylated amine-utilising genes in

genomes-sequenced *Gemmobacters* in comparison to selected marine *Roseobacter*

clade bacteria. Shown is the presence (+) or absence (-) of specific genes in the

genome sequences.

Supplementary Figure S1. Maximum-likelihood phylogenetic tree (JTT matrix-

based) of *mauA* sequences. Sequences from the genus *Gemmobacter* are marked with

a star. Amino acid sequences (MauA) were aligned using the ClustalW algorithm.

Numbers at branches are bootstrap percentages >50% of 1000 replicates. Scale bar:

0.1 substitutions per amino acid. Coloured boxes indicate *Alphaproteobacteria*

*(*yellow), *Gammaproteobacteria (*red) and *Betaproteobacteria (*black).

Supplementary Figure S2. (A-D) Average nucleotide identity (ANI) analysis of

Gemmobacter sp. LW-1 and *Gemmobacter caeni*, *Gemmobacter aquatilis*,

Gemmobacter nectariphilus and *Gemmobacter megaterium* and (E) AAI analysis

between those species

Supplementary Figure S3. Distribution of *Gemmobacter* in different ecosystems

Supplementary Figure S4. Maximum-likelihood phylogenetic tree of concatenated

single copy marker genes from genomes of *Gemmobacter* and closely related

organisms.

Supplementary Figure S5. Arrangement of genes involved in methylated amine

utilisation

Supplementary Table S1. Genome characteristics of the five *Gemmobacter* isolate

genomes used in this study.

Supplementary Table S2. List of protein queries used for the genome comparison

with their accession number.

Supplementary Table S3. Average nucleotide identity (ANI) values between

Gemmobacter sp. LW-1 and *Gemmobacter caeni*, *Gemmobacter aquatilis*,

Gemmobacter nectariphilus, Gemmobacter megaterium, Rhodobacter sphaeroides

and *Paracoccus denitrificans*

Figure 1

Figure 2

Figure 3

Protein	Corresponding gene(s)
TMA monooxygenase	tmm
TMA methyltransferase	mttB1, mttB2, mttB3, mttB4
MMA dehydrogenase	mauA, mauD, mauE, mauG
NMG synthase	mgsA, mgsB, mgsC
NMG dehydrogenase	mgdA, mgdB, mgdC, mgdD
HCHO-activating enzyme	gfa1, gfa2, gfa3, gfa4
GSH-dependent HCHO dehydrogenase	adhI
S-formyl-GSH hydrolase	fghA
Glutamine synthetase (GS)	gluL
Glutamate synthase (GOGAT)	glxB, glxC
Methylene-H$_4$F-dehydro-genase/cyclohydrolase	folD
Formate-H$_4$F ligase	fhs
Formyl-H$_4$F deformylase	purU
Formate dehydrogenase	fdh
Carbon monoxide dehydrogenase	coxD, coxE, coxF, coxG, coxL, coxM, coxS
Methanol dehydrogenase	xoxF
N-acetylglutamate synthase	nagS
Gamma-glutamylcysteine synthetase	gcl
Methylenetetrahydrofolate reductase	metF
Glutamate dehydrogenase	gdh2
Ammonium transporter	amtB

Figure 4A

Tree scale: 0.01 ⊢—⊣
Gemmobacter aquatilis
Gemmobacter sp. LW-1
Gemmobacter caeni
Gemmobacter megaterium
Gemmobacter nectariphilus

Figure 4B

G. sp. LW-1

G. aquatilis

G. caeni

1072

Dispensable

Core
1806

3085

G. nectariphilus

1165

G. megaterium

Table 1. Comparative genomic analysis of methylamine-utilizing genes in genome-sequenced Gemmobacters in comparison to selected marine *Roseobacter* clade bacteria. Shown is the presence (+) or absence (-) of the corresponding genes in the genome sequences.

Organism	gmaS	tmm	xoxF	fae	foldD	purU	fhs (ftfL)	gfa	adhI (flhA)	fghA	fdh	amtB	mtdA/ mtdB	mchA	ftr	fmdA /B/C	mauA	cox
Citreicella sp. SE45	+	+	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	+
Roseovarius sp. TM1035	+	+	+	-	+	-	+	-	+	+	+	+	-/-	-	-	-	-	+
Paracoccus denitrificans PD1222	-	-	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	-
Rhodobacter sphaeroides 241	-	+	+	-	+	+	+	+	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter caeni	+	+	+	-	+	+	+	+[a]	+	+	+	+	-/-	-	-	-	+	+
Gemmobacter sp. LW-1	+	+	+	-	+	+	+	+[a]	+	+	+	+	-/-	-	-	-	+	+
Gemmobacter aquatilis	+	+	+	-	+	+	+	-	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter nectariphilus DSM-15620	-	-	-	-	+	+	+	+[a]	+	+	+	+	-/-	-	-	-	-	+
Gemmobacter megaterium DSM-26375	-	-	-	-	+	+	+	+[a]	+	+	+	+	-/-	-	-	-	-	+

[a] gfa-like protein-encoding gene