1 **Architectural traits constrain the evolution of unisexual flowers and**

2 **sexual segregation within inflorescences: an interspecific approach**

5 **Rubén Torices[1,2,3], Ana Afonso[4], Arne A. Anderberg[5], José M. Gómez[1] and Marcos**

6 **Méndez[3]**

7
8 [1] Departamento de Ecología Funcional y Evolutiva, Estación Experimental de Zonas Áridas (EEZA-
9 CSIC), Carretera de Sacramento s/n, 04120 La Cañada de San Urbano, Almería, Spain
10
11 [2] Department of Ecology and Evolution, University of Lausanne, Biophore Building, CH-1015 Lausanne,
12 Switzerland
13
14 [3] Área de Biodiversidad y Conservación, Universidad Rey Juan Carlos, E-28933 Móstoles-Madrid, Spain
15
16 [4] Centro de Ecologia Funcional, Departamento de Ciências da Vida, Universidade de Coimbra, Calçada
17 Martim de Freitas, 3000-456 Coimbra, Portugal
18
19 [5] Department of Botany, Swedish Museum of Natural History, P.O. Box 50007, SE-104 05 Stockholm,
20 Sweden
21
22
23
24

Contact information
Rubén Torices
Área de Biodiversidad y Conservación
Universidad Rey Juan Carlos
c/ Tulipán s/n.
E-28933. Móstoles (Madrid)
Spain
e-mail: ruben.torices@urjc.es
33

34 **Running head**: Sexual specialization of flowers

35 **Keywords:** Asteraceae, hermaphroditism, gamete packaging, gynomonoecy, monoecy,
36 sex allocation, sexual systems.

ABSTRACT

Male and female unisexual flowers have repeatedly evolved from the ancestral bisexual flowers in different lineages of flowering plants. This sex specialization in different flowers often occurs within inflorescences. We hypothesize that inflorescence architecture may impose a constraint on resource availability for late flowers, potentially leading to different optima in floral sex allocation and unisexuality. Under this hypothesis we expect that inflorescence traits increasing the difference in resource availability between early and late flowers would be phylogenetically correlated with a higher level of sexual specialization. To test this hypothesis, we performed a comparative analysis of inflorescence traits (inflorescence size, number of flowers and flower density) in the sunflower family, which displays an extraordinary variation in floral sexual specialization at the inflorescence level, i.e. hermaphroditic, gynomonoecious and monoecious species. We found that species with a complete sex separation in unisexual flowers (monoecy) had significantly denser inflorescences. Furthermore, those species arranging their flowers in denser inflorescences also showed greater differences in the size of early and late fruits, a proxy of resource variation between flowers. Our findings support the idea that floral sexual specialization and consequently sexual segregation may be the consequence of different floral sex allocation optima driven by the sequential development of flowers that results in a persistent resource decline from earlier to later flowers.

INTRODUCTION

Most angiosperms are hermaphrodite, i.e., they produce bisexual or 'perfect' flowers bearing both functional pollen grains and ovules. Combining both sexual functions in the same flower reduces direct reproduction costs, such as sharing attractive structures for pollinators (Charnov et al. 1976), and acts as an insurance against vagaries in the mating environment by allowing selfing (Wilson and Harder 2003). Thus, evolutionary biologists have wondered, since Darwin (1877), what favours non-hermaphroditic sexual systems, where flowers of different sex -male, female, or bisexual- are combined in the same or different individuals of a population. Darwin (1877) interpreted the evolution of unisexual flowers as a means of reducing selfing. Nevertheless, the simultaneous presence of self-incompatibility and unisexual flowers in many angiosperms (Bertin 1993) suggests that avoiding selfing is not the only driver of sexual specialization in flowers. Alternative explanations for the evolution of sexual specialization are the avoidance of interference between sexual functions (Charnov 1979; Willson 1979) and optimal resource allocation (Janzen 1977; Willson 1979; Lloyd 1982).

Any evolutionary explanation of sexual specialization of flowers should consider that sexes are often segregated within inflorescences due to two constraints set by inflorescence architecture. The first constraint is ontogenetic (Lee 1988; Diggle 1995, 2003): because flowers within inflorescences usually develop sequentially, early flowers usually flower and ripen fruits first and pre-empt resources for late flowers. The second constraint is strictly positional (Diggle 2003) and results in a persistent limitation of resources at certain flower positions inherent to the architecture of inflorescence axes. Both constraints provide a general proximate mechanism for sexual selection, because they can strongly influence the mating environment or resource

availability experienced by individual flowers (Brunet and Charlesworth 1995; Diggle

2003; Harder and Prusinkiewicz 2013). As resource availability may differentially

affect male and female performance (Freeman et al. 1980, Lau and Stephenson 1993),

this resource gradient across flowers within an inflorescence could also lead to

differential sexual specialization in female or male functions at different floral

positions.

On theoretical grounds, a female-biased floral allocation can be predicted in

those floral positions in the inflorescence with higher resource availability, whereas a

male-biased allocation will be expected in positions with less resources (Brunet and

Charlesworth 1995). Intraspecific observational (Diggle 2003) and experimental

evidence (Emms 1993; Diggle 1994) supports this idea, at least for plants with

elongated inflorescences. For instance, *Solanum hirtum* produces inflorescences with

bisexual flowers, but the late (distal) ones are labile, becoming male in resource-

depleted plants (Diggle 1994). Nevertheless, it remains poorly understood whether these

intraspecific plastic responses in sexual specialization and segregation within

inflorescences can actually become fixed during the evolution of a lineage (Diggle

2003; Torices and Méndez 2010), giving rise to the sexual segregation within

inflorescences observed in many angiosperm families. The exception is a pioneer

comparative study of *Solanum* that found how plastic responses in the production of

male unisexual flowers are ancestral to fixed position effects (Diggle and Miller 2013).

Here, we take this comparative approach a step further, by using Asteraceae to test at a

family level whether the degree of sexual specialization is phylogenetically correlated

with inflorescence architectural traits which may lead to a resource decline from early to

late flowers.

We hypothesize that three architectural traits of inflorescences, namely larger

size, and higher flower number and density increase resource competition among

flowers and likely constrain resource availability for late flowers, potentially leading to

male-biased floral sex allocation and unisexuality. Testing this hypothesis at an

interspecific level faces two challenges. First, inflorescences may differ impressively

across species in size, number of flowers, fruit size and flower size, even within the

same main inflorescence type (Leppik 1977; Burtt 1977; Wyatt 1982; Harder et al.

2004). Second, these traits do not necessarily covary across species, because

inflorescence size and flower number are under selection by several ecological drivers,

such as pollinators (Willson and Rathcke 1974; Thomson 1988; Andersson 1996;

Kirchner et al. 2005; Iwata et al. 2012), seed predators (Fenner et al. 2002), or altitude

and geographic ranges (Fenner et al. 2001). Consequently, the diversity of trait

combinations across species could obscure the detection of relationships between

inflorescence traits and floral sexual specialization. In particular, a negative covariation

between flower size and number (Sargent et al. 2007; Goodwillie et al. 2010;

Vasconcelos and Proença 2015) could lead to similar levels of resource competition

among flowers/fruits in inflorescences having different number of flowers, since an

increase in the number of flowers is commonly associated with a decrease in flower and

fruit size. Thus, flower density, in which flower size and inflorescence size are taken

into account, may be a more reliable indicator of the strength of resource competition

among flowers within an inflorescence than the number of flowers per inflorescence or

inflorescence size alone.

In this study, we compared species of the sunflower family showing different

levels of floral sexual specialization to test whether they show different inflorescence

traits. We then assessed how these inflorescence traits correlate with resource

differences between flowers within the same inflorescence. The sunflower family

(Asteraceae) is a suitable model for testing this hypothesis. First, all Asteraceae share

the same basic inflorescence architecture, the head or capitulum (Funk et al. 2009),

which mainly follows a centripetal pattern in floral development and blooming (Harris

1995; Pozner et al. 2012). Second, three different sexual systems are common in

Asteraceae: hermaphroditism, gynomonoecy and monoecy (Fig. 1). From the ancestral

hermaphroditism, monoecy likely evolved through a gynomonoecious intermediate

(Torices et al. 2011). Third, a clear pattern of sexual specialization is present within

inflorescences in non-hermaphroditic species. Namely, gynomonoecious species

produce female flowers at the outermost positions and bisexual ones in the inner

positions, while monoecious species show female flowers in the outermost positions

and male flowers in inner positions (Fig. 1). Finally, there is anatomical (Alkio and

Grimm 2003), physiological (Alkio et al. 2003;) and experimental evidence that

architectural constraints occur within heads, and both outer flowers and positional

effects can limit the available resources to late-blooming, inner flowers (Torices and

Méndez 2010).

Figure 1. Sexual specialization in the Asteraceae inflorescences. Three different
types of functionally hermaphroditic inflorescences and individuals can be observed in

the Asteraceae: hermaphroditic, gynomonoecious and monoecious. Allocation of
gametes to different flowers shows a gradient in floral sexual specialization from
hermaphroditic species (only bisexual flowers) to gynomonecious species (female and
bisexual flowers) and to monoecious species (female and male flowers). In Asteraceae,
sexual specialization (i.e., bi- vs. unisexuality in flowers) and sexual segregation within
inflorescences occur in concert. Lower panels show representative species from a)
hermaphroditic heads (*Tragopogon porrifolius* L.), b) gynomonoecious heads
(*Anacyclus valentinus* L.), and c) monoecious heads (*Tussilago farfara* L.).
162

In this paper, we followed a comparative approach to study the role of

architectural constraints in the evolution of floral sexual specialization and sexual

segregation within inflorescences in the Asteraceae. Specifically, we assessed how

inflorescence traits (namely, inflorescence size, number of flowers, and flower density)

were associated with different levels of floral sexual specialization and sexual

segregation represented by hermaphroditic, gynomonoecious and monoecious species

(Fig. 1). In addition, we assessed how these inflorescence traits affected variation in

fruit size within the inflorescences, as a proxy for the resource gradient between flower

positions (Lee 1988; Stephenson 1992). Finally, we tested whether female flowers in an

inflorescence produce larger fruits than bisexual ones, as would be expected if sexual

specialization 'releases' female flowers from expending resources on male structures and

compensated the lack of the male sex function by producing larger fruits than bisexual

flowers.

MATERIAL AND METHODS

Study species

The Asteraceae is the largest family of angiosperms, with over 1,500 genera and

25,000 species, and a worldwide distribution (Funk et al. 2009). All Asteraceae share

the same basic inflorescence, the head or capitulum, a dense indeterminate inflorescence

where all the flowers are sessile and attached to a common receptacle (Fig. 1). Heads

represent the basic pollinator attraction unit (Burtt 1977; Leppik 1977). Different

degrees of sexual segregation within the heads can be observed among species of the

family (Fig. 1). From hermaphroditism to monoecy, a floral sexual specialization

occurring in individual flowers. The evolutionary transition between them occurs

through a gynomonoecious intermediate that bears both female and bisexual flowers

(Torices et al. 2011).

We included a total of 91 species in our study, including 42 hermaphroditic, 28

gynomonoecious and 18 monoecious species (Supplementary Table S1). The species

were studied on material from two herbaria: the Asteraceae collection of the Swedish

Natural History Museum Herbarium (S), and the Herbarium of the University of

Coimbra (COI). Herbarium sampling allowed a comprehensive sampling of Asteraceae

diversity, since different evolutionary lineages coming from different continents and

biomes were sampled. We sampled specimens from 26 tribes and 7 subfamilies

(including the "Heliantheae alliance"). Hermaphroditism is the ancestral condition in

the family and also in most of the early diverging lineages (Supplementary Fig. S1;

Torices et al. 2011). Once gynomonoecy evolved at the origin of the subfamily

Asteroideae, it remained as the ancestral condition for most of the tribes of this large

clade (Fig. S1). This evolutionary history of sexual systems was taken into account

when selecting the species to be sampled and we put a strong emphasis in sampling

representatives of all sexual systems at the early and late diverging lineages, as well as

in sampling basal and derived hermaphroditic and gynomonoecious species. All

monoecious species were almost always derived from gynomonoecious lineages in the

subfamily Asteroideae where all monoecious species evolved in this family with very

rare exceptions.

Inflorescence and fruit traits

Three traits were measured at an inflorescence level in the 91 species: (i)

inflorescence size, measured as the head diameter in mm; (ii) the number of flowers per

inflorescence; and (iii) flower density, calculated as the ratio between the number of

flowers and the area of each head. Flower density was used as a measure of floral

aggregation and may provide a better proxy of resource competition between flowers

within the head than head size or number of flowers per head.

To minimize specimen damage, we sampled only those specimens with mature

fruits, i.e. fruiting heads, in which we measured inflorescence and fruit traits. Fruiting

heads in Asteraceae usually retain the size and structure of the inflorescence and

therefore they can be used to describe inflorescence traits such as size and number of

flowers. Otherwise, we assume that any change in size during the ripening of the

fruiting head is proportional to head size. First, we searched those specimens belonging

to the species included in the phylogenetic supertrees published for this family (Funk et

al., 2005; Funk, 2009). Second, we selected herbarium specimens with enough mature

fruiting heads and maintaining their original positions within heads. For each species,

one specimen having fruiting heads at desirable conditions was selected, and at least

one capitulum was sampled. We sampled the specimen in which fruiting head removal

caused the least damage possible to the specimen. Following this procedure, 109

herbarium specimens were initially sampled, and from 91 of them we could collect

inflorescence traits (Supplementary Table S1).

All fruiting heads were dissected to separate all fruits in their relative positions

within heads from the outermost to the innermost positions. When necessary, heads

were placed in water with a detergent for rehydration and to reduce damage to the head.

Data were collected for 70 species. Heads and fruits were measured using pictures taken

with a tripod-stabilized digital camera. Fruit area of more than 2,700 fruits were

measured as using Image J 1.54s software (Schneider et al. 2012). Although low

intraspecific sample sizes may lead to increased type I error in comparative studies, this

effect is important only when coupled with high intraspecific variation (Harmon and

Losos 2005). However, when the range of taxa studied is wide, variation across species

is usually much greater than variation within species. In our study, fruit size varied

between species investigated, the largest fruits were more than 100-fold larger than the

smallest fruits, whereas no single species showed such a degree of variation in fruit size.

Statistical analyses

Relationship between inflorescence traits and floral sexual specialization

We used phylogenetic generalized least squared (PGLS) models (Freckleton et

al. 2002; Paradis 2012) to explore the relationship between the degree of floral sexual

specialization (hermaphroditism, gynomonoecy or monoecy) and inflorescence traits.

All models were evaluated under both an adaptive model (OU, Ornstein-Uhlenbeck

model) and a neutral model of evolution (BM, Brownian motion model) (Felsenstein

2004). PGLS were fitted using the "ape" (Paradis et al. 2004) and "geiger" packages

(Harmon et al. 2008) in R. The fittest model for each combination of variables was

selected using a likelihood ratio test comparing BM and OU models. For all fitted

models, the OU model had a higher goodness of fit than the BM (results not shown).

Therefore, we present only the results fitted under an OU model. The phylogenetic

relationship between the species included in the analyses was considered by using the

phylogenetic supertree published for the Asteraceae (Funk et al. 2005), adding a

calibration to include branch lengths (Torices 2010). The root of this tree was

previously scaled to 1 for all the analyses. Specific comparisons between

hermaphroditism, gynomonoecy and monoecy were explored using the marginal means,

using the 'lsmeans' package (Lenth 2016) in R (R Development Core Team, 2015),

which can be defined as a linear combination of the estimated effects from a linear

model.

262
Relationships between fruit size, inflorescence traits and floral sexual specialization

We explored how fruit size variation within inflorescences, as a proxy of the

resource gradient between flower positions, was related to inflorescence traits and floral

sexual specialization, using the meta-analytical effect size to get a standardized measure

of the magnitude of the difference among the size of the outer and inner fruits (fruit size

difference, hereafter FSD). A random-effects meta-analysis was used. Effect sizes were

calculated using the 'meta' package in R (Schwarzer 2015).

The correlation between FSD and fruit size with the degree of floral sexual

specialization was explored using PGLS models. FSD and fruit size were the response

variables, whereas the degree of floral sexual specialization was the predictor variable.

All models were evaluated under both an OU and a BM model (see above).

In addition, we assessed the allometric relationships of inflorescence traits (head

diameter, number of flowers and flower density) with FSD and outer and inner fruit

sizes. These allometric relationships were tested by fitting PGLS models. FSD and fruit

size were the response variables, and the three inflorescence traits were the predictor

variables. For those significant correlations, we estimated phylogenetic reduced major

axis regressions and tested whether slopes were significantly different from one using

the 'phyl.RMA' function included in the 'phytools' package (Revell 2012). All

variables were log-transformed before analysis.

282
Relationship between floral sexual specialization and fruit size

We explored whether unisexual flowers produced larger fruits than bisexual

flowers. We assessed the effect of floral sexual specialization (female vs. bisexual

flowers) using only the outer fruits because strictly female flowers are found only in

these positions. In addition, given the strong effect of inflorescence traits on fruit size

(see Results), we included in the model the density of flowers, to control for the size of

the inflorescence and the number of flowers. We fitted two PGLS models, where fruit

size was the response variable and flower sex was included as a predictor categorical

variable. Flower density was included as a continuous predictor variable. The only

difference between both models was the inclusion of an interaction term between the

sex of the flower and the flower density. The model including an interaction did not

perform better and was thus dropped. Fruit size and flower density were log-

transformed. We evaluated this model under OU and BM correlation structures.

RESULTS

Inflorescence traits and floral sexual specialization

Inflorescence traits differed significantly among the degrees of floral sexual

specialization, i.e., hermaphroditism, gynomonoecy, and monoecy (Table 1).

Hermaphroditic and gynomonoecious species displayed larger inflorescences than

monoecious species (Fig. 2a). Gynomonoecious species had significantly more flowers

per inflorescence than hermaphroditic and monoecious species (Fig. 2b). Nevertheless,

flower density was correlated with the degree of floral sexual specialization, increasing

from hermaphroditic through gynomonoecious to monoecious species (Fig. 2c).

Inflorescence size was significantly correlated to the other two inflorescence

traits (Supplementary Table S2). The number of flowers increased disproportionally

with an increase in inflorescence size, whereas flower density disproportionally

decreased with an increase in inflorescence size, measured as head diameter

(Supplementary Table S3). Number of flowers and flower density were only marginally

correlated (Supplementary Table S2), and flower density increased proportionally with

an increased number of flowers (Supplementary Table S3).

Figure 2. Inflorescence and fruit traits of different sexual systems. Phylogenetically
controlled least-squares means (± 95% confidence interval) of (a) inflorescence size
(mm), (b) number of flowers, (c) flower density (no. flowers / mm^2), (d) outer fruit size,
(e) inner fruit size, and (f) Fruit Size Difference (FSD) for different levels of sexual
systems representing increasing levels of floral sexual specialization within
inflorescences: hermaphroditism, gynomonoecy and monoecy. FSD is the standardized
fruit size difference between outer and inner fruits measured as the meta-analytical
effect size. Means sharing the same superscript letter were not significantly different at
the $P < 0.05$ level.

324
325

Fruit size variation within inflorescences

Fruit size was not statistically different between hermaphroditic,

gynomonoecious and monoecious species, either for outer or inner positions (Table 1;

Fig. 2d,e). Two different relationships between fruit size and inflorescence traits were

observed, independently of position in the head. First, fruit size significantly increased

with an increase in inflorescence size (Fig. 3a). The phylogenetic RMA slopes were

significantly higher than one (outer fruits: $b = 1.34$, $t = 3.16$, d.f. = 68.9, $P = 0.002$;

inner fruits: $b = 1.40$, $t = 3.64$, d.f. = 57.4, $P < 0.001$) indicating a disproportionate

increase in fruit size with an increase in inflorescence diameter (Fig. 3a). Second, fruit

size decreased with an increase in flower density (Fig. 3c). The phylogenetic RMA

slopes were significantly < 1.0 (outer fruits: $b = -0.79$, $t = 3.31$, d.f. = 61.4, $P = 0.002$;

inner fruits: $b = -0.82$, $t = 2.78$, d.f. = 51.7, $P = 0.008$). Therefore, fruit size decreased at

a lower rate than the increase in flower density (Fig. 3c). Fruit size was not statistically

correlated with flower number (Fig. 3b).

Figure 3. Allometric relationships between fruit size and inflorescence traits.
Phylogenetic RMA regressions between fruit size and a) inflorescence size (mm), b)
number of flowers, and c) flower density (no. flowers / mm^2); and between FSD and d)
inflorescence size (mm), e) number of flowers, and f) flower density (no. flowers /
mm^2). In the upper row black dots and solid lines represent outer fruits whereas white
dots and dashed lines indicate inner fruits. Lines represent RMA slopes. Phylogenetic
standard regression slopes are showed in Supplementary Table 4.
348

FSD, the standardized fruit size difference, significantly decreased with an

increase in inflorescence size (Fig. 3d; $b = -3.20$, $t = 10.24$, d.f. = 62.2, $P < 0.0001$),

implying that smaller heads showed a higher difference between outer and inner fruits.

FSD did not show any significant relationship with the number of flowers (Fig. 3e),

although it significantly increased with flower density (Fig. 3f), revealing that an

increase in flower density was associated with a higher difference in size between outer

and inner fruits ($b = 1.85$, $t = 5.28$, d.f. $= 61.9$, $P <0.0001$). The degree of sexual

specialization did not significantly affect FSD (Table 1). However, the post-hoc

comparison between degrees of sexual specialization showed that monoecious species

had significantly larger FSD than hermaphroditic and gynomonoecious species (Fig.

2e). This difference was mainly mediated by the differences among sexual systems in

flower density, because the inclusion of flower density as a covariate removed any

statistical difference between monoecious species and the other two categories

considered (Supplementary Table S5).

Floral sexual specialization and fruit size

Fruit size decreased with increasing flower density ($F_{1,76} = 109.03$, $P < 0.0001$;

Fig. 4). Nevertheless, female flowers produced significantly larger fruits than bisexual

flowers after controlling by flower density ($F_{1,76} = 5.87$, $P = 0.018$, n = 78; Fig. 4).

Figure 4. Relationship between fruit size produced by female (filled dots and solid
line) and bisexual flowers (white dots and dashed line) and flower density. Female

flowers produced by monoecious species are shown as black dots whereas those
produced by gynomonoecious species are indicated by grey dots.
373

DISCUSSION

Our comparative study across three sexual systems in Asteraceae provided clear

support for a phylogenetic association between inflorescence traits, fruit size variation

within inflorescences, and the degrees of sexual specialization and segregation. In

particular, we observed that monoecious species bore smaller and denser inflorescences

than hermaphroditic ones, also showing the largest fruit size difference (FSD) between

outer and inner fruits. These results, together with the lack of a correlation between the

number of flowers and the degree of sexual specialization, support the idea that floral

sexual specialization and consequently sexual segregation within inflorescences of

Asteraceae might in part be the result of different sex allocation optima brought about

by architecturally mediated persistent resource limitation of the inner flowers.

Although selfing and geitonogamy avoidance has been considered an important

factor in the evolution of floral sex specialization and sexual segregation within

inflorescences (Harder et al. 2000), on its own it is unlikely to explain the pattern

observed in Asteraceae. The negative effects of geitonogamy are expected to be higher

in large inflorescences, where a higher number of flowers can lead to longer floral bouts

by pollinators (Harder and Barrett 1995). Thus, many-flowered inflorescences would be

expected to show a higher probability of exhibiting unisexual flowers than few-flowered

inflorescences. However, in our data set the number of flowers per inflorescence was

not clearly related to sexual specialization (Fig. 3b). Moreover, outer seeds should be

more outcrossed than inner ones, but evidence is scarce and inconclusive: two species

have shown higher outcrossing rate in outer flowers than inner ones (Marshall and

Abbott 1984, Cheptou et al. 2001), whereas two other studies have found no differences

(Gibson 2001, Gibson and Tomlinson 2002). Finally, the species of this family usually

show other mechanisms to avoid selfing such as self-incompatibility (Ferrer et al. 2007)

and dichogamy (both at the flower and inflorescence levels; Burtt 1977). Therefore,

whether limiting geitonogamy plays a key role in the evolution of sexual segregation in

Asteraceae requires further examination, because empirical evidence remains

contradictory.

An alternative to the selfing avoidance hypothesis is that evolutionary transitions

between sexual systems might have evolved as a way to optimize gamete packaging.

Inflorescence traits, such as the inflorescence size, inflorescence number, but also the

number of flowers per inflorescence, are key components of gamete packaging

strategies (Schoen and Dubuc 1990; Fishbein and Venable 1996). Our results suggest

that shifts in the so called 'inflorescence design', can have effects at the flower level

too. For instance, the increase in flower number is usually negatively correlated with

flower size (Sargent et al. 2007; Goodwillie et al. 2010; Vasconcelos and Proença

2015). In the Asteraceae, these shifts can have triggered shifts in floral sex expression

leading to transitions between hermaphroditism, gynomonoecy and monoecy, whereas

the pollination unit keeps their mating opportunities through the retention of both male

and female functions. Two 'inflorescence design' syndromes has been proposed related

to sexual systems for the tribe Inuleae (Torices and Anderberg 2009) which are also

supported by our results. In the first group, including hermaphroditic and

gynomonoecious species, attraction units were mostly large solitary or corymbose heads

with rather few heads together. In contrast, the attraction unit of the second group,

composed of monoecious species only, were clusters of several small heads. Moreover,

we observed that although larger inflorescences produced both more flowers and larger

fruits in species in our sample, fruit size decreased with flower density. This indicates

that resource competition can be the mechanism underlaying the interspecific negative

relationships between the number of flowers and fruit size.

Importantly, our study provides insights into how intraspecific size-number

trade-offs can translate into negative covariation between traits across species. The

consequences of this negative correlation between size and number at the inflorescence

level were not the same for all flower positions. We observed that increased flower

density led to decreases fruit size, especially at the innermost positions, resulting in a

higher FSD (fruit size difference between outer and inner fruits). Flower density might

thus amplify the effects of architectural constraints, which pervasively limit resources at

the innermost positions. Under these circumstances, theory predicts that a high resource

difference between flower positions can cause plants to allocate more resources to their

female function in flowers with more resources, and to their male function in resource-

depleted flowers (Brunet and Charlesworth 1995). This expectation agrees with the

positional pattern observed for the sex of flowers in Asteraceae inflorescences (Fig. 1),

where female unisexual flowers consistently appear at those earlier (or outer) floral

positions that generally have a higher resource supply, whereas male unisexual flowers

are displayed at the later (or inner) positions, which usually are the most resource-

limited positions (Alkio and Grimm 2003; Alkio et al. 2003; Torices and Méndez 2010).

Therefore, shifts in inflorescence traits modifying the density of flowers might

secondarily cascade to other important floral traits such as flower and fruit size.

Specially, flower density might have a role on the evolution of floral sex functions,

given its effects on fruit size and thus on the floral female performance.

Under the resource optimization hypothesis, floral sex specialization is expected

to entail an improvement in fitness compared to bisexual flowers. Previous studies

indicate that dioecious species have larger fruit set than cosexual species (Sutherland

1986) and larger seed size (Vamosi et al. 2008). While there is not a formal test on

dioecious species of the Asteraceae or in other groups where flowers are no longer the

functional unit, our results support that when the inflorescence is the main pollination

unit, architectural traits such as flower density may obscure direct comparisons between

dioecious and non-dioecious species, given its effect on fruit size. In the Asteraceae,

female unisexual flowers from both gynomonoecious and monoecious species had

significantly larger fruits than bisexual species, when the confounding effect of flower

density was factored out. Thus, our study provides support for an intrinsic advantage of

flower specialization at least in terms of female fitness.

Expectations of a negative covariation of traits across species assumes that

everything else remains equal. Vamosi et al. (2008) did not find any difference in seed

size between hermaphroditic and monoecious species. However it is unclear if

hermaphroditic and monoecious species in their dataset differed in inflorescence traits,

such as flower density, which could confound the effect of the floral sexual

specialization on fruit and seed size. Positive correlations between inflorescence parts

might be found if species differ in resource budget (the big house-big car effect, *sensu*

Reznick et al. 2000). In addition, additional selective factors on dispersal performance,

which usually occur at the fruit level, could indirectly drive shifts on inflorescence

design. Our study only revealed the evolutionary drivers in the specialization of floral

sex functions after taking into account other inflorescence traits that might otherwise

have masked those drivers. This underlies the strength of a comparative approach for

understanding the mechanistic basis of the evolution of non-hermaphroditic sexual

systems, in the line of Diggle & Miller (2013), even when considering phylogenetic

scales above the genus level.

Conclusion

Our results highlight the importance of considering architectural traits to understand phenotypic diversity in modular organisms, which can have important functional consequences (Herrera 2009 for a review). Architectural constraints may have profound consequences in modular organisms such as plants, influencing how male and female functions perform at different positions within one individual. In particular, the sequential development of inflorescences and asymmetric competition between early and late flowers lead to a gradient in the resource availability experienced by individual flowers within the inflorescences. Thus, a separation of male and female functions in different flowers might evolve not only to maximize mating patterns, but also to optimize resource allocation.

This combination of architectural constraints and selection for optimal sex allocation at the flower level could explain sexual specialization and segregation in other clades beside Asteraceae. Differential access to resources and/or architectural constraints derive from basic anatomical properties of inflorescences and therefore should be general to most plant clades. In fact, in a sample of 127 families showing andro-, gyno- or monoecy, two thirds showed sexual segregation along inflorescences (M. Méndez, unpubl. data), suggesting that inflorescence architecture is a general constraint in the evolution of these sexual systems. Particular aspects of gamete packaging could differ among clades or inflorescence architectures. For example, floral density can have a minor role in elongated inflorescences compared to inflorescence size and flower number. This is why a comparative approach to sexual specialization and segregation has to be aware that all else will rarely remain equal and consider that different solutions may exist to a common problem. Comparative studies including

inflorescence traits, sexual expression and fruit size variation across flowers offer great

promise in elucidating the generality of this mechanism in the evolution of unisexual

flowers in Angiosperms.

Acknowledgements

We are especially grateful to John Pannell for the review of a previous version of the

manuscript, and to L. DeSoto, S. Castro and F. Sales for their help in different stages of

the study. RT was partially supported by the Spanish Ministry of Education (BVA

2010-0375).

References

Andersson, S. 1996. Floral display and pollination success in *Senecio jacobaea*

(Asteraceae): interactive effects of head and corymb size. *Am. J. Bot.* 83:71–75.

Alkio, M., and E. Grimm. 2003. Vascular connections between the receptacle and

empty achenes in sunflower (*Helianthus annuus* L.). *J. Exp. Bot.* 54:345–348.

Alkio, M., A. Schubert, W. Diepenbrock, and E. Grimm. 2003. Effect of source – sink

ratio on seed set and filling in sunflower (*Helianthus annuus* L .). *Plant. Cell*

Environ. 26:1609–1619.

Bertin, R. I. 1993. Incidence of monoecy and dichogamy in relation to self-fertilization

in angiosperms. *Amer. J. Bot.* 80:557-560.

Brunet, J., and D. Charlesworth. 1995. Floral sex allocation in sequentially blooming

plants. *Evolution* 49:70–79.

Burtt, B. L. 1977. Aspects of diversification in the capitulum. Pp. 41-59 *in* V. Heywood,

J. B. Harborne, B. L. Turner, eds. The biology and chemistry of the Compositae.

Vol. 1. Academic Press, London.

Charnov, E. L. 1979. Simultaneous hermaphroditism and sexual selection. *Proc. Natl.*

Acad. Sci. USA 76: 2480-2484.

Charnov, E. L., J. Maynard Smith, J. J. Bull. 1976. Why be an hermaphrodite? *Nature*

263: 125-126.

Charnov, E. L. 1987. On sex allocation and selfing in higher plants. *Evol. Ecol*. 12:30–

36.

Cheptou, P. O., J. Lepart, and J. Escarre. 2001. Differential outcrossing rates in

dispersing and non-dispersing achenes in the heterocarpic plant *Crepis sancta*

(Asteraceae). *Evol. Ecol.* 15:1–13.

Darwin, C. 1877. The different forms of flowers on plants of the same species. John

Murray, London, U.K.

Diggle, P. K. 1994. The expression of andromonoecy in *Solanum hirtum* (Solanaceae):

phenotypic plasticity and ontogenetic contingency. *Am. J. Bot.* 81: 1354-1365.

Diggle, P. K. 1995. Architectural effects and the interpretation of patterns of fruit and

seed development. *Annu. Rev. Ecol. Syst*. 26:531–552.

Diggle, P. K. 2003. Architectural effects on floral form and function: a review. Pp. 63–

80 *in* T. Stuessy, E. Hörandl, and V. Mayer, eds. Deep morphology: toward a

renaissance of morphology in plant systematics. Ganter Verlag, Königstein,

Germany.

Diggle, P. K., and J. S. Miller. 2013. Developmental plasticity, genetic assimilation, and

the evolutionary diversification of sexual expression in *Solanum. Am. J. Bot.*

100:1050-1056.

Emms, S. K. 1993. Andromonoecy in *Zigadenus paniculatus* (Liliaceae): Spatial and

temporal patterns of sex allocation. *Am. J. Bot.* 80:914-923.

Felsenstein, J. 2004. Inferring phylogenies. Sinauer Associates, Sunderland.

Fenner, M., W. G. Lee, and E. H. Pinn. 2001. Reproductive features of *Celmisia* species

(Asteraceae) in relation to altitude and geographical range in New Zealand. *Biol. J.*

Linn. Soc. 74:51–58.

Fenner, M., J. E. Cresswell, R. A. Hurley, and T. Baldwin. 2002. Relationship between

capitulum size and pre-dispersal seed predation by insect larvae in common

Asteraceae. *Oecologia* 130:72–77.

Ferrer, M. M., and S. V Good-Avila. 2007. Macrophylogenetic analyses of the gain and

loss of self-incompatibility in the Asteraceae. *New Phytol*. 173:401–14.

Fishbein, M., and D. L. Venable. 1996. Evolution of inflorescence design: theory and

data. *Evolution* 50:2165–2177.

Freckleton, R. P., P. H. Harvey, and M. Pagel. 2002. Phylogenetic analysis and

comparative data: a test and review of evidence. *Am. Nat*. 160:712–26.

Freeman, D. C., K. T. Harper, and El. L. Charnov. 1980. Sex change in plants: old and

new observations and new hypotheses. *Oecologia* 47:222-232.

Funk, V. A., L. Watson, B. Gemeinholzer, E. Schilling, A. Susanna, and R. K. Jansen.

2005. Everywhere but Antarctica: using a supertree to understand the diversity and

distribution of the Compositae. *Biol. Skr*. 55:343–374.

Funk, V. A., A. Susanna, T. Stuessy, and R. J. Bayer, eds. 2009. Systematics, evolution,

and biogeography of Compositae. International Association for Plant Taxonomy,

Viena.

Gibson, J. 2001. Ecological and genetic comparison between ray and disc achene pools

of the heteromorphic species *Prionopsis ciliata* (Asteraceae). *Int. J. Plant Sci.*

162:137–145.

Gibson, J. P., A. D. Tomlinson, and A. S. College. 2002. Genetic diversity and mating

system comparisons between ray and disc achene seed pools of the heterocarpic

species *Heterotheca subaxillaris* (Asteraceae). *Int. J. Plant Sci.* 163:1025–1034.

Goodwillie, C., R. D. Sargent, C. G. Eckert, E. Elle, M. A. Geber, M. O. Johnston, S.

Kalisz, D. A. Moeller, R. H. Ree, M. Vallejo-Marín, and A. A. Winn. 2010.

Correlated evolution of mating system and floral display traits in flowering plants

and its implications for the distribution of mating system variation. *New Phytol.*

185:311-321.

Harder, L. D., and S. C. H. Barrett. 1995. Mating cost of large floral displays in

hermaphrodite plants. *Nature* 373:512–515.

Harder, L. D., S. C. H. Barrett, and W. W. Cole. 2000. The mating consequences of

sexual segregation within inflorescences of flowering plants. *Proc. R. Soc. B Biol.*

Sci. 267:315–20.

Harder, L. D., C. Y. Jordan, W. E. Gross, and M. B. Routley. 2004. Beyond

floricentrism: the pollination function of inflorescences. *Plant Species Biol.* 19:137–

148.

Harder, L. D., and P. Prusinkiewicz. 2013. The interplay between inflorescence

development and function as the crucible of architectural diversity. *Ann. Bot.*

112:1477–1493.

Harmon, L. J., and J. B. Losos. 2005. The effect of intraspecific sample size on type I

and type II error rates in comparative studies. *Evolution* 59:2705–2710.

Harmon L. J, J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER:

investigating evolutionary radiations. *Bioinformatics* 24:129-131.

Harris, E. M. 1995. Inflorescence and floral ontogeny in Asteraceae: a synthesis of

historical and current concepts. *Bot. Rev.* 61:93-278.

Herrera, C. M. 2009. Multiplicity in unity: plant subindividual variation and interactions

with animals. The University of Chicago Press, London.

Imbert, E., J. Escarre, and J. Lepart. 1997. Seed heteromorphism in *Crepis sancta*

(Asteraceae): performance of two morphs in different environments. *Oikos* 79:325–

332.

Iwata, T., O. Nagasaki, H. S. Ishii, and A. Ushimaru. 2012. Inflorescence architecture

affects pollinator behaviour and mating success in *Spiranthes sinensis* (Orchidaceae).

New Phytol. 193:196–203.

Kirchner, F., S. H. Luijten, E. Imbert, M. Riba, M. Mayol, S. C. Gonza, and B. Colas.

2005. Effects of local density on insect visitation and fertilization success in the

narrow-endemic *Centaurea corymbosa* (Asteraceae). *Oikos* 111:130–142.

Janzen, D. H. 1977. A note on optimal mate selection by plants. *Am. Nat.* 111: 365-371.

Lau, T.-C., and A. G. Stephenson. 1993. Effects of soil nitrogen on pollen production,

pollen grain size, and pollen performance in *Cucurbita pepo* (Cucurbitaceae). *Amer.*

J. Bot. 80:763-768.

Lee, T. D. 1988 Patterns of fruit and seed production. Pp 179-202 *in* J. Lovett-Doust

and L. Lovett-Doust. Plant reproductive ecology: patterns and strategies. Oxford

University Press, New York.

Lenth, R. V. 2016. Least-squares means: the R package lsmeans. *J Stat Soft* 69:1-33.

Leppik, E. E. 1977. The evolution of capitulum types of the Compositae in the light of

insect-flower interaction. Pp 61-89 *in* V. H. Heywood, J. B. Harborne, B. L. Turner,

eds. The biology and chemistry of the Compositae. Vol. 1. Academic Press, London,

UK.

Lloyd, D. G. 1982. Selection of combined versus separate sexes in seed plants. *Am. Nat.*

120:571-585.

Marshall, D. F., and R. J. Abbott. 1984. Polymorphism for outcrossing frequency at ray

floret locus in *Senecio vulgaris* L. III. Causes. *Heredity* 53:145–149.

Paradis, E. 2012. Analysis of phylogenetics and evolution with R (Second Edition).

Springer. New York.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of phylogenetics and

evolution in R language. *Bioinformatics* 20:289–290.

Pozner, R., C. Zanotti, and L. A. Johnson. 2012. Evolutionary origin of the Asteraceae

capitulum: Insights from Calyceraceae. *Am. J. Bot.* 99:1–13.

R Core Team. 2015. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and

other things). *Methods Ecol. Evol*. 3:217–223.

Reznick, D., :L. Nunney, and A. Tessier. 2000. Big houses, big cars, superfleas and the

costs of reproduction. *Trends Ecol. Evol.* 15:421-425.

Sargent, R. D., C. Goodwillie, S. Kalisz, and R. H. Ree. 2007. Phylogenetic evidence

for a flower size and number trade-off. *Am. J. Bot*. 94:2059–2062.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25

years of image analysis. *Nat. Methods* 9:671–5.

Schoen, D. J., and M. Dubuc. 1990. The evolution of inflorescence size and number: a

gamete-packaging strategy in plants. *Am. Nat*. 135:841–857.

Schwarzer, G. 2015. meta: general package for meta-analysis. R package version 4.1-0.

http://CRAN.R-project.org/package=meta

Stephenson, A. G. 1992. The regulation of maternal investment in plants. Pp. 151-171

in C. Marshall, J. Grace eds. Fruit and seed production: aspects of development,

environmental physiology and ecology. Cambridge University Press, Cambridge.

Sutherland, S. 1986. Patterns of fruit-set: what controls fruit-flower ratios in plants?

Evolution 40:117–128.

Thomson, J. D. 1988. Effects of variation in inflorescence size and floral rewards on the

visitation rates of traplining pollinators of *Aralia hispida*. *Evol. Ecol.* 2:65–76.

Torices, R. 2010. Adding time-calibrated branch lengths to the Asteraceae supertree. *J.*

Syst. Evol. 48:271–278.

Torices, R., and A. A. Anderberg. 2009. Phylogenetic analysis of sexual systems in

Inuleae (Asteraceae). *Am. J. Bot.* 96:1011–1019.

Torices, R., and M. Méndez. 2010. Fruit size decline from the margin to the center of

capitula is the result of resource competition and architectural constraints. *Oecologia*

164:949–58.

Torices, R., M. Méndez, and J. M. Gómez. 2011. Where do monomorphic sexual

systems fit in the evolution of dioecy? Insights from the largest family of

angiosperms. *New Phytol.* 190:234–48.

Vamosi, S. M., S. J. Mazer, and F. Cornejo. 2008. Breeding systems and seed size in a

neotropical flora: testing evolutionary hypotheses. *Ecology* 89:2461–72.

Vasconcelos, T. N. C., and C. E. B. Proença. 2015. Floral cost vs. floral display:

Insights from the megadiverse Myrtales suggest that energetically expensive floral

parts are less phylogenetically constrained. *Am. J. Bot.* 102:900–909.

Willson, M. F. 1979. Sexual selection in plants. *Am. Nat.* 113:777-790.

Willson, M. F. 1982. Sexual selection and dicliny in angiosperms. *Am. Nat.* 119:579-

583.

Willson, M. F., and B. J. Rathcke. 1974. Adaptive design of the floral display in

Asclepias syriaca L. *Am. Midl. Nat.* 92:47–57.

Wilson, W. G., and L. D. Harder. 2003. Reproductive uncertainty and the relative

competitiveness of simultaneous hermaphroditism versus dioecy. *Am. Nat.* 162:220-

241.

Wyatt, R. 1982. Inflorescence architecture: how flower number, arrangement, and

phenology affect pollination and fruit-set. *Am. J. Bot.* 69:585–594.

Tables

Table 1. Differences in inflorescence and fruit traits between inflorescences with
different degree of floral sexual specialization. *F* and *P* values were obtained after a
deviance analysis of the phylogenetic generalized linear models fitted for each
inflorescence trait, with the degree of floral sexual specialization within inflorescence
(hermaphroditism, gynomonoecy or monoecy) as the main factor. FSD is the
standardized fruit size difference between outer and inner fruits measured as the meta-
analytical effect size.
684

Inflorescence traits	*F*	d.f.	*P*
Capitulum diameter (mm)	7.27	2, 85	0.001
Number of flowers	3.65	2, 84	0.030
Flower density (no. flowers/mm^2)	6.99	2, 84	0.002
Outer fruit size (mm^2)	0.69	2, 77	0.505
Inner fruit size (mm^2)	2.01	2, 66	0.142
FSD	2.30	2, 64	0.109

685
686

Figure Legends

Figure 1. Sexual specialization in the Asteraceae inflorescences. Three different
types of functionally hermaphroditic inflorescences and individuals can be observed in
the Asteraceae: hermaphroditic, gynomonoecious and monoecious. Allocation of
gametes to different flowers shows a gradient in floral sexual specialization from
hermaphroditic species (only bisexual flowers) to gynomonecious species (female and
bisexual flowers) and to monoecious species (female and male flowers). In Asteraceae,
sexual specialization (i.e., bi- vs. unisexuality in flowers) and sexual segregation within
inflorescences occur in concert. Lower panels show representative species from a)
hermaphroditic heads (*Tragopogon porrifolius* L.), b) gynomonoecious heads
(*Anacyclus valentinus* L.), and c) monoecious heads (*Tussilago farfara* L.).
700

Figure 2. Inflorescence and fruit traits of different sexual systems. Phylogenetically
controlled least-squares means (± 95% confidence interval) of (a) inflorescence size
(mm), (b) number of flowers, (c) flower density (no. flowers / mm^2), (d) outer fruit size,
(e) inner fruit size, and (f) Fruit Size Difference (FSD) for different levels of sexual
systems representing increasing levels of floral sexual specialization within
inflorescences: hermaphroditism, gynomonoecy and monoecy. FSD is the standardized
fruit size difference between outer and inner fruits measured as the meta-analytical
effect size. Means sharing the same superscript letter were not significantly different at
the $P < 0.05$ level.
710

Figure 3. Allometric relationships between fruit size and inflorescence traits.
Phylogenetic RMA regressions between fruit size and a) inflorescence size (mm), b)
number of flowers, and c) flower density (no. flowers / mm^2); and between FSD and d)
inflorescence size (mm), e) number of flowers, and f) flower density (no. flowers /
mm^2). In the upper row black dots and solid lines represent outer fruits whereas white
dots and dashed lines indicate inner fruits. Lines represent RMA slopes. Phylogenetic
standard regression slopes are showed in Supplementary Table 4.
718

719 **Figure 4. Relationship between fruit size produced by female (filled dots and solid**
720 **line) and bisexual flowers (white dots and dashed line) and flower density**. Female
721 flowers produced by monoecious species are shown as black dots whereas those
722 produced by gynomonoecious species are indicated by grey dots.