Identification of evolutionarily conserved virulence factor by selective pressure analysis

of *Streptococcus pneumoniae*

Masaya Yamaguchi[a], Kana Goto[a, b], Yujiro Hirose[a], Yuka Yamaguchi[a], Tomoko Sumitomo[a],

Masanobu Nakata[a], Kazuhiko Nakano[b], Shigetada Kawabata[a]*

[a]Department of Oral and Molecular Microbiology, Osaka University Graduate School of

Dentistry, Suita, Osaka, Japan

[b]Department of Pediatric Dentistry, Osaka University Graduate School of Dentistry, Suita,

Osaka, Japan

*Address correspondence to Shigetada Kawabata, kawabata@dent.osaka-u.ac.jp

Running title: Identification of virulence factor by evolutionary analysis

Keywords: *Streptococcus pneumoniae*, Evolutionary analysis, Choline-binding protein,

Evolutionary conservation

Abstract

Streptococcus pneumoniae is a pathogen that poses one of the greatest threats to human

health, causing pneumonia, sepsis, and meningitis. Here, we investigated the evolutionary

selective pressures on 16 pneumococcal choline-binding cell-surface proteins. Phylogenetic

and molecular analyses revealed that the *cbpJ* and *lytA* genes had the highest codon rates

under negative selection among those examined. Although LytA is an important virulence

factor inducing pneumococcal autolysis, the role of CbpJ in pneumococcal pathogenesis is

unknown. We therefore examined whether CbpJ acts as a virulence factor *in vitro* and *in vivo*.

In neutrophil bactericidal assays, the *cbpJ* mutant showed reduced survival as compared to

wild-type (WT) *S. pneumoniae*. Mice intranasally infected with a *cbpJ* mutant strain showed

a lower bacterial burden in bronchoalveolar lavage fluid and higher survival rate than those

infected with WT cells, whereas no differences were observed between strains by intravenous

infection. Thus, molecular evolutionary analysis is a powerful tool that reveals the

importance of virulence factors in real-world infection and transmission, since calculations

are performed based on bacterial genome diversity following transmission of infection in an

uncontrolled population. In addition, evolutionarily conserved virulence factors are candidate

therapeutic targets or vaccine antigens.

Improper use of antibiotics creates evolutionary pressures that drive bacteria to acquire drug

resistance by natural mutation and/or horizontal transfer of resistance genes. This is a major

public health threat: it is estimated that drug-resistant infections cause 10 million deaths

annually and may result in economic losses reaching 100 trillion US dollars by 2050[1].

However, a target-to-hit screen typically requires 24 works-in-process and 94 million US

dollars, and the baseline total cost is 1.8 billion US dollars over 13 years to launch a new

drug[2]. In fact, the number of new antibiotics developed and approved has steadily decreased

in the past three decades, leaving fewer options for treating resistant bacteria[3].

Streptococcus pneumoniae is one of the pathogens posing the greatest threat to human

health[4,5]. *S. pneumoniae* belongs to the mitis group[6,7] and is a major cause of pneumonia,

sepsis, and meningitis[8,9]. In 2015, pneumococcal pneumonia caused over 1.5 million deaths

in individuals of all ages, and this rate increased in people over 70 years old between 2005

and 2015[10], which is especially problematic since the elderly population is growing in many

parts of the world. Although pneumococcal conjugate vaccines have considerable benefits,

non-vaccine pneumococcus serotypes have increased worldwide[11,12].

The conflict between the host immune system and pathogens leads to an evolutionary

arms races known as the "Red Queen" scenario[13,14]. Protein regions at the host–pathogen

interface are subjected to the strongest selective pressure and thus evolve under positive

selection. Adaptive evolution has been reported in genes related to the mammalian immune

system such as pattern recognition receptors[14]. In contrast, bacterial proteins are under

negative/purifying selection, which is important for the survival and/or success of the species

since non-synonymous substitution of codons can lead to lineage extinction (Fig. 1).

Phylogenetic and molecular evolutionary analyses can reveal the number of codons under

negative/purifying selection in a species, since alterations in amino acid residues in regions

under negative selective pressure are not allowed. Thus, drugs targeting these regions would

be less likely to promote the development of resistance through natural mutation.

We analysed pneumococcal choline-binding proteins (CBPs) localised on the bacterial

cell surface through interaction with choline-binding repeats and phosphoryl choline on the

cell wall. *S. pneumoniae* harbours various CBPs including *N*-acetylmuramoyl L-alanine

amidase (LytA), which induces pneumococcal-specific autolysis[15]. Although several CBPs

have been characterised, their phylogenetic relationships remain unclear and the unclassified

gene names are confusing. We first analysed the distribution of genes encoding CBPs based

on pneumococcal genome sequences. Orthologues of genes in each strain were identified by

phylogenetic analysis. We then calculated the evolutionary selective pressure on each codon

from phylogenetic trees and aligned sequences. We found that the *cbpJ* gene contains the

highest rate of codons under negative selection. CbpJ has no known functional domains

except signal sequences and choline-binding repeats, and its role in pneumococcal

pathogenesis is unclear. Functional analyses revealed that CbpJ contributes to evasion of host

neutrophil-mediated killing in pneumococcal pneumonia. Thus, evolutionary analysis

focusing on negative selection can reveal novel virulence factors.

Results

Distribution of *cbp* genes among pneumococcal strains

Genes encoding CBPs among pneumococcal strains were extracted by tBLASTn search

(Supplementary Table 1). Some genes were re-annotated since the search results showed that

certain homologous regions were not matched to annotated open reading frames (ORFs). In

strain SPNA45, the *SPNA_01670* gene contains both predicted promoter regions and intact

ORF structures of the *cbpF* and *cbpJ* genes. On the other hand, *cbpG*-homologous regions in

strains R6, D39, SPN034183, SPN994038, and SPN994039 did not contain promoters

(Supplementary Table 1 and Supplementary Table 2). Orthologous relationships of each gene

were analysed. The distribution of *cbp* genes was unrelated to capsular serotypes. Four

genes—i.e., *lytA*, *lytB*, *cbpD*, and *cbpE*—were conserved as intact ORFs in all 28

pneumococcal strains (Fig. 2A). Other *cbp* genes contained frameshift mutations in the

orthologues or were absent in some strains.

Phylogenetic relationships in pneumococcal CBPs

Phylogenetic relationships of genes encoding CBPs in pneumococcal species are confusing

since some genes in the same cluster show high similarity to each other. To clarify the

relationships, we compared common nucleotide sequences among genes encoding CBPs in

strain TIGR4. Maximum likelihood and Bayesian phylogenetic analyses revealed two

common clusters: one comprising *cbpF*, *cbpG*, *cbpJ*, *cbpK*, and *cbpC*, and the other *lytA*, *lytB*,

lytC, *cbpL*, and *cbpE* (Fig. 2B and Supplementary Fig. 1). The names of some CBP genes

were not consistent with those of phylogenetically related genes. In particular, *cbpF*, *cbpG*,

cbpJ, and *cbpK* were located close to each other in pneumococcal genomes and showed high

similarity. We thus defined orthologous genes in each pneumococcal strain based on

maximum likelihood and Bayesian phylogenetic analyses (Fig. 3 and Supplementary Fig. 2).

The gene locus tag numbers in orthologous relationships are shown in Supplementary Table 1.

The sequence similarity of *cbpF*, *cbpG*, *cbpJ*, and *cbpK* and their close proximity within

genomes indicated that a common ancestral *S. pneumoniae* acquired the genes by gene

duplication. Phylogenetic trees showed well-separated clusters of each gene. These

independent relationships indicated that horizontal gene transfer did not contribute to the

spread of *cbpF*, *cbpG*, *cbpJ*, and *cbpK* in *S. pneumoniae* species, despite their ability to take

up exogenous DNA. The genetic diversity of these genes may have been established by

accumulation of natural mutations during pneumococcal transmission.

108 **Evolutionary selective pressures on each of the CBP codons**

109 To evaluate the significance of CBPs in real-life infection and transmission, we performed

110 molecular evolutionary calculations based on bacterial genome diversity established after

111 transmission of infection in an uncontrolled population. The nucleotide sequences of each

112 CBP were aligned by codon, and conserved common codons were used for phylogenetic

113 analysis (Supplementary Fig. 3). The selective pressure on each gene was calculated based on

114 the phylogenetic trees and aligned sequences (Table 1). The rates of codons under negative

115 selection are shown in Supplementary Figure 4. Over 13% of total codons in *cbpJ* and *lytA*

116 genes were under negative selection as compared to less than 5% for other *cbp* genes,

117 indicating that these genes play an important role in the success of *S. pneumoniae* species. On

118 the other hand, the *pspA* gene encoding the genetically divergent virulence factor PspA

119 contained fewer evolutionarily conserved codons, but had the highest numbers of codons

120 under positive pressure. Additionally, there were no evolutionarily conserved codons in the

121 *cbpG*, *cbpC*, and *cbpL* genes. The latter two had no common codons since a few genes had

122 frameshift mutations. When we re-calculated selective pressure without these genes, we

123 found a low rate of codons under negative selection among CBP-encoding genes

124 (Supplementary Table 3).

CbpJ acts as a virulence factor in pneumococcal pneumonia

While CbpJ had the highest rate of codons under negative selection among pneumococcal

CBPs, it has no known functional domains except a choline-binding repeat in its amino acid

sequence; moreover, its role in pneumococcal pathogenesis is unknown. In contrast, CbpL

showed limited numbers of evolutionarily conserved codons even after the above-described

adjustment. Domain structures and codons of CbpJ and CbpL under negative selection are

shown in Figure 4A. To assess the roles of CbpJ and CbpL in pneumococcal pathogenesis, we

generated mutant strains deficient in the corresponding genes. We first performed a blood

bactericidal assay to investigate the role of CbpJ and CbpL in sepsis. The survival rates of

ΔcbpJ and *ΔcbpL* strains in mouse blood were comparable to that of the wild-type (WT)

strain (Fig. 4B). We also carried out a human neutrophil bactericidal assay without serum.

Strain *ΔcbpJ* had a lower survival rate than the WT. These results suggest that CbpJ

contributes to the evasion of neutrophil-mediated killing (Fig. 4C).

In a mouse intravenous infection model, the survival rates of *ΔcbpL*- and

ΔcbpJ-infected mice did not differ significantly from that of mice infected with WT *S.*

pneumoniae (Fig. 5A). We also found that incubation of *S. pneumoniae* in human plasma for

3 h inhibited the expression of *cbpL* and *cbpJ*, as determined by quantitative real-time

(q)PCR (Fig. 5B). On the other hand, mice intranasally infected with strain *ΔcbpJ* showed an

improved survival rate as compared to those infected with WT *S. pneumoniae*; although a

similar tendency was observed for *ΔcbpL*-infected relative to WT mice, the difference was

not statistically significant (Fig. 5C). The number of bacteria in bronchoalveolar lavage fluid

(BALF) from *ΔcbpJ*-infected mice was lower than that in BALF from *ΔcbpL*- and

WT-infected mice (Fig. 5D). We also examined whether CbpL or CbpJ contributes to the

association of *S. pneumoniae* to alveolar epithelial cells and found that WT *S. pneumoniae* as

well as *ΔcbpL* and *ΔcbpJ* mutant strains did not differ in their ability to adhere to A549

human alveolar epithelial cells (Fig. 5E). However, *S. pneumoniae* WT strain exhibited

extensive inflammatory cell infiltration and bleeding as compared to the *ΔcbpJ* strain.

Histological examination of lung tissue from intranasally-infected mice showed that *ΔcbpJ*

induced milder inflammation than the WT strain. Lung tissue from *ΔcbpL*-infected mice

showed moderate inflammation (Fig. 5F). These results indicate that CbpJ acts as a

pneumococcal virulence factor in lung infection by contributing to the evasion of

neutrophil-mediated killing, whereas CbpJ has no role in bacterial survival in blood. In

addition, *cbpL* deficiency in strain TIGR4 did not significantly attenuate pathogenesis in

mouse lung and blood infection.

Discussion

Discussion

In this study, we investigated evolutionarily conserved rates of CBP codons since these cell

surface proteins directly interact with the external environment, which induces rapid rates of

evolution in genes involved in genetic conflicts[14]. Evolutionary analysis based on

phylogenetic relationships can reveal regions in which the encoded amino acids are not

allowed to change even under selective pressure. The genetic diversity of *S. pneumoniae*

isolated from patients was the result of transmission in a real population. Thus, the

evolutionary conservation rate is a parameter that reflects the importance of the protein in

human infection. Although so-called arms races involve both the host and bacteria, most

studies of genetic diversity have focused on the former[14,16-18]. For example, evolutionary

studies based on inter-species comparisons have shown that most of the positive selection

targets in host receptors are located in regions that are responsible for direct interactions with

pathogens. Our study focused on negative selection targets in bacterial surface proteins

through an evolutionary analysis based on intra-species comparisons. This approach enabled

us to estimate the contribution of bacterial proteins to species success throughout the life

cycle, including inside the host and during the transmission phase.

We previously detected bacterial virulence factors by function prediction—e.g., by

searching for conserved motif/domains, constructing random transposon libraries, or

analysing the biochemical properties of the pathogen[19-23]. Although these laboratory-based

approaches are valuable, they are time-consuming and costly, and may not yield the expected

results. It is useful to examine the correlation between a target molecule and clinical features

as this can minimise the time and cost required for analysis. Furthermore, in basic studies of

bacterial pathogens, animal infection models are often used to determine whether a bacterial

molecule acts as a virulence factor. Although this is the best means of obtaining *in vivo*

information, it is unclear how accurately it reflects the clinical condition in humans.

Combining an evolutionary analysis and an animal model would be highly effective for

evaluating the functional significance of a putative virulence factor.

187   Genome-wide association study (GWAS) is a powerful tool for identifying the

relationship between genetic variants—mainly single nucleotide polymorphisms

(SNPs)—and phenotype, such as in diseases. As GWAS has become more prevalent, various

programs and software packages have been developed for this purpose[24,25]. On the other hand,

this approach has certain limitations, including the requirements for an appropriate control

group and detailed information regarding phenotype. In infectious diseases, it can be difficult

to quantify clinical features recorded at different medical centres. Furthermore, in the case of

most pathogens, there are no natural attenuated or avirulent strains that can serve as a control

group. Our evolutionary analysis has the advantage that it can be performed with genomic

information of pathogenic strains only by assuming the presence of pathogens as a phenotype

evading natural selection. Since synonymous and non-synonymous substitutions are

estimated to occur with equal probability under no selective pressure, a population in which

the latter has resulted in extinction by natural selection can serve as a control group. While

we have shown in the current study that evolutionary analysis with a small population has the

power to detect evolutionarily conserved proteins, a larger population would allow a

higher-resolution analysis, including the detection of conserved regions in some pathogenic

strains isolated from a specific site of infection or pathological condition. Since this analysis

involves simultaneous processing of aligned nucleotide and amino acid sequences, more

information is obtained from only SNPs extracted from nucleotide sequences. In addition,

automatised phylogenetic and evolutionary analyses are needed to analyse a large population.

Therefore, the development of software packages for meta-data is expected to aid widespread

application of this analytical approach.

There are some limitations to our evolutionary analysis. Firstly, although it can detect

evolutionarily conserved proteins, it cannot identify diverse virulence factors such as PspA

and CbpA within species[26-28]. Similarly, virulence factors recently acquired by horizontal

gene transfer have not been under selective pressure for a sufficiently long period of time to

perform this analysis. In addition, the high rate of codons under negative selection indicate

their universal importance in bacterial species. In other words, a molecule under relaxed

selective pressure could contribute to the virulence of some populations of the species.

However, these features of molecular evolutionary analysis can be advantages when

screening for therapeutic target sites or vaccine antigens with a low frequency of missense

mutations, which could reduce the virulence or survivability of the pathogen. Evolutionary

analysis could also be an effective alternative strategy for overcoming drug resistance

through antigen replacement, and could reduce costs associated with drug discovery and

development.

The *lytA* gene, which was conserved among virtually all pneumococcal strains,

showed the highest rates of codons under negative selection, except for the *cbpJ* gene that

was only present in some strains. LytA is known to induce pneumococcal-specific autolysis[29]

and contributes to pneumococcal virulence[30,31]. Our evolutionary analysis supports previous

reports that the *lytA* gene is a suitable genetic marker[32,33] due to its evolutionary conservation.

We also showed that the *pspA* and *cbpA* genes contain relatively high rates of codons under

positive selection, and both encode polymorphic virulent proteins[26,27,34] that are candidate

vaccine antigens, although the genes are not universally present within a global serotype 1

collection[28]. In addition, selective pressure by vaccines can easily cause differentiation or

deficiency of these proteins since the corresponding genes contain few codons under negative

selection. A multivalent system would be required for vaccines using these antigens.

Although the difference was not statistically significant, mice intranasally infected

with TIGR4 *ΔcbpL* strain showed a trend towards improved survival relative to the

WT-infected mice. In a previous study, a D39*lux cbpL*-deficient strain showed reduced

virulence as compared to the WT strain[35]. Since CbpL sequences in TIGR4 and D39 strains

are similar, the discrepancy between the previous study and our findings is likely due to

differences in other surface proteins in each strain. For example, the absence of CbpJ, which

contributes to the evasion of neutrophil killing, could affect the survivability of D39.

Recently, antivirulence drugs have been developed as an additional strategy to treat or

prevent bacterial infections. Drugs targeting bacterial virulence factors are expected to reduce

the selective pressure of conventional antibiotics since they would not affect the natural

survival of targeted bacteria[36]. Furthermore, the abundance of candidate targets is a major

advantage of antivirulence strategies. Effective design of vaccines and antivirulence drugs

requires a thorough understanding of virulence factors; combining our evolutionary analysis

and traditional molecular microbiological approaches can improve the detection of potential

drug targets. In this study, we identified CbpJ as a novel evolutionarily conserved virulence

factor. Thus, molecular evolutionary analysis is a powerful system that can reveal the

importance of virulence factors in real-world infections and transmission.

 Methods

Phylogenetic and evolutionary analyses

Phylogenetic and evolutionary analyses were performed as previously described[37,38], with

minor modifications. Homologues and orthologues of *cbp* genes were searched using the

tBLASTn function of NCBI BLAST. Bacterial ORFs and promoters were predicted by

FGENESB (Bacterial Operon and Gene Prediction) and BPROM, respectively[39]. To prevent

node density artefacts, sequences with 100% identity were treated as the same sequence in

Phylogears2[40,41]. The sequences were aligned using MAFFT v.7.221 with an L-INS-i

strategy,[42] and ambiguously aligned regions were removed using Jalview[43,44]. Calculated

orthologous regions were used for further phylogenetic analysis, and edited codon sequences

were re-aligned using MAFFT with an L-INS-i strategy. The best-fitting codon evolutionary

models for MrBayes and RAxML analyses were determined using Kakusan4[45]. Bayesian

Markov chain Monte Carlo analyses were performed with MrBayes v.3.2.5[46], and 2×10^6

generations were sampled after confirming that the standard deviation of split frequencies

was < 0.01 for up to 8×10^6 generations. To validate phylogenetic inferences, maximum

likelihood phylogenetic trees with bootstrap values were generated with RAxML v.8.1.20[47].

Phylogenetic trees were generated using FigTree v.1.4.2[48] based on the calculated data.

Evolutionary analyses were performed based on aligned orthologous regions of *cbp*

genes and Bayesian phylogenetic trees. Whole-gene non-synonymous/synonymous ratio

calculations as well as statistical tests for negative or positive selection of individual codons

were performed using the two-rate fixed-effects likelihood function in HyPhy software

package[49].

Bacterial strains and construction of mutant strains

Streptococcus pneumoniae strains were cultured in Todd-Hewitt broth (BD Biosciences,

Franklin Lakes, NJ, USA) supplemented with 0.2% yeast extract (BD Biosciences) (THY

medium) at 37°C. For mutant selection and maintenance, spectinomycin (Wako Pure

Chemical Industries, Osaka, Japan) was added to the medium at a concentration of 120

µg/ml.

S. pneumoniae TIGR4 isogenic *cbpJ* (*ΔcbpJ*) and *cbpL* (*ΔcbpL*) mutant strains were

generated as previously described[22]. Briefly, the upstream region of *cbpJ* or *cbpL*, an *aad9*

cassette, and the downstream region of *cbpJ* or *cbpL* were combined by PCR using primers

shown in Supplementary Table 4. The products were used to construct the mutant strains by

double-crossover recombination with synthesised CSP2[50]. All mutations were confirmed by

PCR amplification of genomic DNA isolated from the mutant strains.

Blood and neutrophil bactericidal assays

A blood bactericidal assay was performed as previously described[20,22,51]. Mouse blood was

obtained via cardiac puncture from healthy female CD-1 mice (Slc:ICR, 6 weeks old; Japan

SLC, Hamamatsu, Japan). For human neutrophil isolation, blood was collected via

venopuncture from healthy donors after obtaining written, informed consent according to a

protocol approved by the institutional review board of Osaka University Graduate School of

Dentistry (H26-E43). Neutrophils were isolated from fresh human blood by density gradient

centrifugation using Polymorphprep (Alere Technologies, Jena, Germany). Pneumococcal

cells grown to the mid-log phase were washed and resuspended in phosphate-buffered saline

(PBS). Bacterial cells (1×10^4 CFU/20 μl) were combined with fresh mouse blood (180 μl)

or human neutrophils (2×10^5 cells/180 μl), and the mixture was incubated at 37°C in 5%

CO_2 for 1, 2, and 3 h. Viable cell counts were determined by seeding diluted samples onto

THY blood agar. The percent of original inoculum was calculated as the number of CFU at

the specified time point divided by the number of CFU in the initial inoculum.

Mouse infection assays

All mouse experiments were conducted in accordance with animal protocols approved by the

Animal Care and Use Committee of Osaka University Graduate School of Dentistry

(28-002-0). Female CD-1 mice (Slc:ICR, 6 weeks old) were intranasally infected with $5 \times$

10^7 or 2×10^6 CFU of *S. pneumoniae* via the tail vein. Mouse survival was monitored for 14

days. At 24 h after intranasal infection, animals were euthanized by lethal intraperitoneal

injection of sodium pentobarbital and lung tissue or BALF samples were collected. Bacterial

counts in BALF were determined by plating serial dilutions. Lung tissue specimens were

fixed with 4% formaldehyde, embedded in paraffin, and cut into sections that were stained

with haematoxylin and eosin solution (Applied Medical Research, Osaka, Japan) and

visualized with a BZ-X710 microscope (Keyence, Osaka, Japan).

qPCR

qPCR was performed as previously described[37,38], with minor modifications. Primers are

listed in Supplementary Table 4. Total RNA of pneumococcal strains grown to the mid-log

phase ($OD_{600} = 0.5$) was isolated with an RNeasy Mini kit (Qiagen, Valencia, CA, USA) and

RQ1 RNase-Free DNase (Promega, Madison, WI, USA), and cDNA was synthesised with

SuperScript IV VILO Master Mix (Life Technologies, Carlsbad, CA, USA). qPCR analysis

was performed on a StepOnePlus Real-Time PCR system using Power SYBR Green Master

PCR mix (Thermo Fisher Scientific, Waltham, MA, USA). 16S rRNA was used as a

normalising control.

Statistical analysis

Statistical analysis of *in vitro* and *in vivo* data was performed with a non-parametric

Kruskal-Wallis test with Dunn's multiple comparisons test. Mouse survival curves were

compared with the log-rank test. $P < 0.05$ was considered statistically significant. The tests

were carried out with Prism v.7.0d software (GraphPad Inc., La Jolla, CA, USA). All

experiments were repeated at least three times. In the evolutionary analyses, $P < 0.1$ was

regarded as a significant difference with the HyPhy default setting.

Acknowledgements

This study was supported in part by Japanese Society for the Promotion of Science

KAKENHI (grant numbers 15H05012, 16H05847, 16K15787, 17H05103, 17K11666, and

17H04369); SECOM Science and Technology Foundation; Takeda Science Foundation; GSK

Japan Research Grant; Asahi Glass Foundation; Kurata Memorial Hitachi Science and

Technology Foundation; and Kobayashi International Scholarship Foundation. The funders

had no role in study design, data collection or analysis, decision to publish, or preparation of

the manuscript.

Author contributions

M.Y. and S.K. designed the study. M.Y. and Y.Y. performed bioinformatics analyses. K.G.,

M.Y., and Y.H. performed the experiments. M.Y., T.S., M.N., and S.K. contributed to the

setup of the experimentation. M.Y. wrote the manuscript. G.K., Y.H., Y.Y., T.S., M.N., K.N.,

and S.K. contributed to the writing of the manuscript.

Competing financial interests statement

The authors declare that they have no competing interests.

References

References

1   O'Neill, J. Tackling Drug-Resistant Infections Globally: final report and

recommendations. The review on antimicrobial resistance; London: HM Government

and the Wellcome Trust; 2016.

2   Paul, S.M. *et al.* How to improve R&D productivity: the pharmaceutical industry's grand

challenge. *Nat Rev Drug Discov* **9**, 203–214 (2010).

3   CDC. Antibiotic Resistance Threats in the United States (2013).

4   CDC. Biggest Threats, <https://www.cdc.gov/drugresistance/biggest_threats.html>

(2017).

5   WHO. WHO Priority Pathogens List for R&D of New Antibiotics,

<http://www.who.int/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/>

(2017).

6   Richards, V.P. *et al.* Phylogenomics and the dynamic genome evolution of the genus

Streptococcus. *Genome Biol Evol* **6**, 741–753 (2014).

7   Kawamura, Y., Hou, X.G., Sultana, F., Miura, H. & Ezaki, T. Determination of 16S

rRNA sequences of *Streptococcus mitis* and *Streptococcus gordonii* and phylogenetic

relationships among members of the genus *Streptococcus*. *Int J Syst Bacteriol* **45**, 406–

408 (1995).

8   Walker, C.L. *et al.* Global burden of childhood pneumonia and diarrhoea. *Lancet* **381**,

1405–1416 (2013).

9   Castelblanco, R.L., Lee, M. & Hasbun, R. Epidemiology of bacterial meningitis in the

USA from 1997 to 2010: a population-based observational study. *Lancet Infect Dis* **14**,

813–819 (2014).

10   Collaborators, G.L. Estimates of the global, regional, and national morbidity, mortality,

and aetiologies of lower respiratory tract infections in 195 countries: a systematic

analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis* **17**, 1133–1161

(2017).

11   Golubchik, T. *et al.* Pneumococcal genome sequencing tracks a vaccine escape variant

formed through a multi-fragment recombination event. *Nat Genet* **44**, 352–355 (2012).

12   Flasche, S. *et al.* Effect of pneumococcal conjugate vaccination on serotype-specific

carriage and invasive disease in England: a cross-sectional study. *PLoS Med* **8**, e1001017

(2011).

13   Brockhurst, M.A. *et al.* Running with the Red Queen: the role of biotic conflicts in

evolution. *Proc Biol Sci* **281**, doi:10.1098/rspb.2014.1382 (2014).

14   Sironi, M., Cagliani, R., Forni, D. & Clerici, M. Evolutionary insights into

host-pathogen interactions from mammalian sequence data. *Nat Rev Genet* **16**, 224–236
(2015).

15   Maestro, B. & Sanz, J.M. Choline binding proteins from *Streptococcus pneumoniae*: a
dual role as enzybiotics and targets for the design of new antimicrobials. *Antibiotics*
(Basel) **5**, doi:10.3390/antibiotics5020021 (2016).

16   Fumagalli, M. & Sironi, M. Human genome variability, natural selection and infectious
diseases. *Curr Opin Immunol* **30**, 9–16 (2014).

17   Karlsson, E.K., Kwiatkowski, D.P. & Sabeti, P.C. Natural selection and infectious
disease in human populations. *Nat Rev Genet* **15**, 379–393 (2014).

18   Siddle, K.J. & Quintana-Murci, L. The Red Queen's long race: human adaptation to
pathogen pressure. *Curr Opin Genet Dev* **29**, 31–38 (2014).

19   Terao, Y. *et al.* Group A streptococcal cysteine protease degrades C3 (C3b) and
contributes to evasion of innate immunity. *J Biol Chem* **283**, 6253–6260 (2008).

20   Yamaguchi, M., Terao, Y., Mori, Y., Hamada, S. & Kawabata, S. PfbA, a novel plasmin-
and fibronectin-binding protein of *Streptococcus pneumoniae*, contributes to
fibronectin-dependent adhesion and antiphagocytosis. *J Biol Chem* **283**, 36272–36279
(2008).

21   Sumitomo, T. *et al.* Streptolysin S contributes to group A streptococcal translocation
across an epithelial barrier. *J Biol Chem* **286**, 2750–2761 (2011).

22   Mori, Y. *et al.* alpha-Enolase of *Streptococcus pneumoniae* induces formation of
neutrophil extracellular traps. *J Biol Chem* **287**, 10472–10481 (2012).

23   Yamaguchi, M. *et al. Streptococcus pneumoniae* invades erythrocytes and utilizes them
to evade human innate immunity. *PLoS One* **8**, e77282, (2013).

24   Gallagher, M.D. & Chen-Plotkin, A.S. The post-GWAS era: from association to function.
Am J Hum Genet **102**, 717–730 (2018).

25   Marigorta, U.M., Rodriguez, J.A., Gibson, G. & Navarro, A. Replicability and
prediction: lessons and challenges from GWAS. *Trends Genet*
doi:10.1016/j.tig.2018.03.005 (2018).

26   Hollingshead, S.K. *et al.* Pneumococcal surface protein A (PspA) family distribution
among clinical isolates from adults over 50 years of age collected in seven countries. *J*
Med Microbiol **55**, 215–221 (2006).

27   Brooks-Walter, A., Briles, D.E. & Hollingshead, S.K. The *pspC* gene of *Streptococcus*
pneumoniae encodes a polymorphic protein, PspC, which elicits cross-reactive
antibodies to PspA and provides immunity to pneumococcal bacteremia. *Infect Immun*
67, 6533–6542 (1999).

28 Cornick, J.E. *et al.* The global distribution and diversity of protein vaccine candidate

antigens in the highly virulent *Streptococcus pneumoniae* serotype 1. *Vaccine* **35**, 972–

980 (2017).

29 Mosser, J.L. & Tomasz, A. Choline-containing teichoic acid as a structural component of

pneumococcal cell wall and its role in sensitivity to lysis by an autolytic enzyme. *J Biol*

Chem **245**, 287–298 (1970).

30 Berry, A.M., Lock, R.A., Hansman, D. & Paton, J.C. Contribution of autolysin to

virulence of *Streptococcus pneumoniae*. *Infect Immun* **57**, 2324–2330 (1989).

31 Orihuela, C.J., Gao, G., Francis, K.P., Yu, J. & Tuomanen, E.I. Tissue-specific

contributions of pneumococcal virulence factors to pathogenesis. *J Infect Dis* **190**, 1661–

1669 (2004).

32 Carvalho Mda, G. *et al.* Evaluation and improvement of real-time PCR assays targeting

lytA, *ply*, and *psaA* genes for detection of pneumococcal DNA. *J Clin Microbiol* **45**,

2460–2466 (2007).

33 Saukkoriipi, A. *et al. lytA* quantitative PCR on sputum and nasopharyngeal swab

samples for detection of pneumococcal pneumonia among the elderly. *J Clin Microbiol*

56, doi:10.1128/JCM.01231-17 (2018).

34 Hakenbeck, R., Madhour, A., Denapaite, D. & Bruckner, R. Versatility of choline

metabolism and choline-binding proteins in *Streptococcus pneumoniae* and commensal

streptococci. *FEMS Microbiol Rev* **33**, 572–586 (2009).

35 Gutierrez-Fernandez, J. *et al.* Modular architecture and unique teichoic acid recognition

features of choline-binding protein L (CbpL) contributing to pneumococcal pathogenesis.

Sci Rep **6**, 38094 (2016).

36 Dickey, S.W., Cheung, G.Y.C. & Otto, M. Different drugs for bad bugs: antivirulence

strategies in the age of antibiotic resistance. *Nat Rev Drug Discov* **16**, 457–471 (2017).

37 Yamaguchi, M. *et al.* Evolutionary inactivation of a sialidase in group B *Streptococcus*.

Sci Rep **6**, 28852 (2016).

38 Yamaguchi, M. *et al.* Zinc metalloproteinase ZmpC suppresses experimental

pneumococcal meningitis by inhibiting bacterial invasion of central nervous systems.

Virulence **8**, 1516–1524 (2017).

39 Solovyev, V. & Salamov, A. *Metagenomics and its Applications in Agriculture,*

Biomedicine and Environmental Studies (ed. Li W.R.) Ch. 4, 61–78 (Nova Science

Publishers, 2011).

40 Tanabe, A.S. Phylogears2 ver. 2.0, <http://www.fifthdimension.jp/> (2008).

41 Venditti, C., Meade, A. & Pagel, M. Detecting the node-density artifact in phylogeny

reconstruction. *Syst Biol* **55**, 637–643 (2006).

42   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772–780 (2013).

43   Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M. & Barton, G.J. Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics*
25, 1189–1191 (2009).

44   Talavera, G. & Castresana, J. Improvement of phylogenies after removing divergent and
ambiguously aligned blocks from protein sequence alignments. *Syst Biol* **56**, 564–577
(2007).

45   Tanabe, A.S. Kakusan4 and Aminosan: two programs for comparing nonpartitioned,
proportional and separate models for combined molecular phylogenetic analyses of
multilocus sequence data. *Mol Ecol Resour* **11**, 914–921 (2011).

46   Ronquist, F. *et al.* MrBayes 3.2: efficient Bayesian phylogenetic inference and model
choice across a large model space. *Syst Biol* **61**, 539–542 (2012).

47   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

48   Rambaut, A. FigTree ver.1.4.2, <http://tree.bio.ed.ac.uk/software/figtree/> (2014).

49   Pond, S.L., Frost, S.D. & Muse, S.V. HyPhy: hypothesis testing using phylogenies.
Bioinformatics **21**, 676–679 (2005).

50   Bricker, A.L. & Camilli, A. Transformation of a type 4 encapsulated strain of
Streptococcus pneumoniae. *FEMS Microbiol Lett* **172**, 131–135 (1999).

51   Yamaguchi, M. *et al.* Role of *Streptococcus sanguinis* sortase A in bacterial colonization.
Microbes Infect **8**, 2791–2796 (2006).

A. Phylogenetic relationship before natural selection

B. Real population resulting from natural selection

Figure 1. Scheme for intra-species molecular evolutionary analysis. A. Random genetic drift induces synonymous and non-synonymous mutations with equal probability. However, non-synonymous mutations in the essential region cause host selection. **B.** As a result of natural selection, synonymous substitutions are concentrated in important genes. Phylogenetic and molecular evolutionary analyses can detect significant accumulation of synonymous substitutions in codons of host proteins. Codon-based analysis yields much more information than nucleotide- or amino acid-based analyses.

Figure 1. Yamaguchi *et al.*

A

B

Figure 2. Yamaguchi *et al*.

Figure 2. Distribution of *cbp* genes and phylogenetic relationship in TIGR4. A. Distribution of genes encoding CBPs among pneumococcal strains. Blue, yellow, and gray show the presence, pseudogenisation, and absence of genes, respectively. *These genes are annotated as one gene, but our bioinformatic analysis indicates that they are independent genes. **B.** Nucleotide-based Bayesian phylogenetic tree of *cbp* genes of *S. pneumoniae* strain TIGR4. The tree is unrooted and posterior probabilities are shown near the nodes. The scale bar indicates nucleotide substitutions per site.

A

B

Figure 3. Yamaguchi *et al.*

Figure 3: Phylogenetic analyses of *cbp* genes with high similarity. A, B. Nucleotide-based Bayesian phylogenetic tree of the *lytA*, *lytB*, *lytC*, *cbpE*, and *cbpL* genes (A) and the *cbpF*, *cbpG*, *cbpJ*, and *cbpK* genes (B) in *S. pneumoniae*. The trees are unrooted although they are presented as midpoint-rooted for clarity. Strains with identical sequences are listed on the same branch. Posterior probabilities are shown near the nodes. The scale bar indicates nucleotide substitutions per site.

A

TIGR4 CbpL

MNKRLFSKMSLVTLPILALFSQSVLAEENI
HFSSCKEAWANGYSDIHEGEPGYSAKLDRD
HDGVACELKNAPKGAFKAKQSTAIQINTSS
ATTSGWVKQDGAWYYFDGNGNLVKNAWQGS
YYLKADGKMAQSEWIYDSSYQAWYYLKSDG
SYAKNAWQGAYYLKSNGKMAQGEWVYDSSY
QAWYYLKSDGSYARNAWQGNYYLKSDGKMA
KGEWVYDATYQAWYYLTSDGSYAYSTWQGN
YYLKSDGKMAVNEWVDGGRYYVGADGVWKE
VQASTASSSNDSNSEYSAALGKAKSYNSLF
HMSKKRMYRQLTSDFDKFSNDAAQYAIDHL
DD*

TIGR4 CbpJ

MKILKKTMQVGLTVFFFGLLGTSTVFADS
EGWQFVQENGRTYYKKGDLKETYWRVIDGK
YYYFDSLSGEMVVGWQYIPFPSKGSTIGPY
PNGIRLEGFPKSEWYYFDKNGVLQEFVGWK
TLEIKTKDSVGRKYGEKREDSEDKEEKRYY
TNYYFNQNHSLETGWLYDQSNWYYLAKTEI
NGENYLGGERRAGWINDDSTWYYLDPTTGI
MQTGWQYLGNKWYYLRSSGAMATGWYQDGS
TWYYLNAGNGDMKTGWFQVNGNWYYAYSSG
ALAVNTTVDGYSVNYNGEWVR*

| 🟦 Signal Sequence | 🟪 Excalibur domain | 🟩 Choline binding domain | 🟧 Lipoprotein domain |

B

C

Figure 4. Yamaguchi *et al.*

Figure 4: Blood and neutrophil bactericidal assays using TIGR4 WT, *ΔcbpJ*, and *ΔcbpL* strains. A. Amino acid sequences and domain structures of CbpL and CbpJ in strain TIGR4. Bold, black underlined, and magenta underlined characters represent comparable codons and those under purifying or positive selection, respectively. **B, C.** Growth of pneumococcal strains in mouse blood (B) and in the presence of human neutrophils (C). Bacterial cells were incubated in blood or with neutrophils for 1, 2, and 3 h at 37°C and 5% CO_2, then serially diluted and plated on TS blood agar. The number of CFUs was determined following incubation. Growth index was calculated by dividing CFU after incubation by the CFU of the original inoculum. Data are presented as the mean of six samples with standard error.

Figure 5. Yamaguchi *et al.*

Figure 5: Intranasal and intravenous infection with *S. pneumoniae* TIGR4 WT, *ΔcbpJ*, and *ΔcbpL* strains. A. Mouse sepsis model. Mice were intravenously infected with 2×10^6 CFU of *S. pneumoniae* TIGR4 WT, *ΔcbpL*, or *ΔcbpJ*, and survival was monitored for 14 days. Differences between infected mouse groups were analysed with the log-rank test. **B.** Fold transcript levels of *cbpL* and *cbpJ* in TIGR4 WT *S. pneumoniae* cells in the presence or absence of human plasma. 16S rRNA was used as an internal standard. Data were pooled and normalised from three independent experiments, each performed in quadruplicate. **C.** Mouse pneumonia model. Mice were intranasally infected with 5×10^7 CFU of *S. pneumoniae* TIGR4 WT, *ΔcbpL*, or *ΔcbpJ* strains, and survival was monitored for 14 days. **D.** Pneumococcal CFU in BALF collected 24 h after intranasal infection. The difference between groups was analysed with the Kruskal-Wallis test with Dunn's multiple comparisons test. **E.** *S. pneumoniae* TIGR4 WT, *ΔcbpL*, and *ΔcbpJ* strains were examined for their ability to associate with A549 cells. Data are presented as the mean of six samples. Differences between groups were analysed with the Kruskal-Wallis test with Dunn's multiple comparisons test. **F.** Haematoxylin and eosin staining of infected mouse lung tissue collected 24 h after intranasal infection. Scale bars, 200 μm (upper panels), 50 μm (middle panels), and 20 μm (lower panels). Data are shown as mean with standard error (B, D, and E).

Figure 5. Yamaguchi *et al.*

476   **Table 1.** Evolutionary analyses of genes encoding choline-binding proteins*

Genes	Number of sequences[1]	dN/dS	Coverage of comparable codons relative to whole protein in TIGR4	Codons evolving under positive selection	Codons evolving under purifying selection	% Of codons under purifying selection relative to total codons
pspA	24	0.857	19.060% (142/745)	6.338% (9/142)	12.676% (18/142)	2.416%
cbpF	19	0.395	60.411% (206/341)	0.485% (1/206)	3.398% (7/206)	2.053%
cbpJ	15	0.346	84.084% (280/333)	1.429% (4/280)	18.571% (52/280)	15.616%
cbpC	13	–	0% (0/93)	–	–	0.000%
cbpG	21	–	0% (0/286)	–	–	0.000%
cbpK	11	0.353	85.630% (292/341)	0.342% (1/292)	3.082% (9/292)	2.639%
cbpL	20	–	0% (0/333)	–	–	0.000%
cbpE	18	0.325	99.363% (624/628)	0.160% (1/624)	4.968% (31/624)	4.936%
lytB	22	0.185	92.868% (612/659)	0% (0/612)	4.739% (29/612)	4.401%
cbpM	10	0.642	98.462% (128/130)[2]	0% (0/128)	0% (0/128)	0.000%
lytC	23	0.400	19.348% (95/491)	0% (0/95)	5.263% (5/95)	1.018%
lytA	14	0.141	80.564% (257/319)	0% (0/257)	17.121% (44/257)	13.793%
cbpA	19	0.864	22.334% (155/694)	3.226% (5/155)	7.742% (12/155)	1.729%
pcpA	18	0.261	77.010% (479/622)	0% (0/479)	0.418% (2/479)	0.322%
cbpD	19	0.359	75.278% (338/449)	0.296% (1/338)	3.550% (12/338)	2.672%
cbpI	2	–	–	–	–	–

477   [1]Sequences with 100% identity were treated as the same sequence; [2]compared to D39.

478   *Evolutionary analysis was performed by Bayesian inference of aligned *cbp* sequences from complete genomes of *S. pneumoniae* with the two-rate fixed-effects

479   likelihood function in HyPhy software package. dN/dS is the ratio of non-synonymous to synonymous changes in overall analysed genes. Individual codons with a

480   statistically significant signature were also calculated and are expressed as a percentage of the total number of codons included in the analysis.