1

A Genome-Wide Association Study Reveals a Novel Regulator of Ovule Number

and Fertility in *Arabidopsis thaliana*

Jing Yuan[1,2] and Sharon A. Kessler[1,2*]

[1]Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana USA

[2]Purdue Center for Plant Biology, Purdue University, West Lafayette, Indiana USA

* Corresponding author
Email: sakessler@purdue.edu

Short title

Regulation of ovule number in Arabidopsis

Abstract

Ovules contain the female gametophytes which are fertilized during pollination to initiate

seed development. Thus, the number of ovules that are produced during flower

development is an important determinant of seed crop yield and plant fitness. Mutants

with pleiotropic effects on development often alter the number of ovules, but specific

regulators of ovule number have been difficult to identify in traditional mutant screens.

We used natural variation in Arabidopsis accessions to identify new genes involved in

the regulation of ovule number. The ovule numbers per flower of 189 Arabidopsis

accessions were determined and found to have broad phenotypic variation that ranged

from 39 ovules to 84 ovules per pistil. Genome-Wide Association tests revealed several

genomic regions that are associated with ovule number. T-DNA insertion lines in

candidate genes from the most significantly associated loci were screened for ovule

number phenotypes. The *NEW ENHANCER of ROOT DWARFISM (NERD1)* gene was

found to have pleiotropic effects on plant fertility that include regulation of ovule number

and both male and female gametophyte development. Overexpression of NERD1

increased ovule number per fruit in a background-dependent manner and more than

doubled the total number of flowers produced in all backgrounds tested, indicating that

manipulation of NERD1 levels can be used to increase plant productivity.

Author Summary

Ovules are the precursors of seeds in flowering plants. Each ovule contains an egg cell

and a central cell that fuse with two sperm cells during double fertilization to generate

seeds containing an embryo and endosperm. The number of ovules produced during

flower development determines the maximum number of seeds that can be produced by

a flower. In this paper, we used natural variation in *Arabidopsis thaliana* accessions to

identify regions of the genome that are associated with ovule number. Polymorphisms

in the plant-specific NERD1 gene on chromosome 3 were significantly associated with

ovule number. Mutant and overexpression analyses revealed that *NERD1* is a positive

regulator of ovule number, lateral branching, and flower number in Arabidopsis.

Manipulation of NERD1 expression levels could potentially be used to increase yield in

crop plants.

Key Words

Ovule number, NERD1, Arabidopsis, Branching, Carpel Margin Meristem, GWAS

Introduction

During plant reproduction, pollen tubes deliver two sperm cells to female gametophytes

contained within ovules. This allows double fertilization to occur in order to produce the

embryo and endosperm in the developing seed. Angiosperms with all kinds of

pollination syndromes (insect-, wind-, and self-pollinated) produce much more pollen

than ovules in order to ensure successful pollination. For example, most soybean

varieties produce only 2 ovules per flower, but more than 3,000 pollen grains, a 1,500-

fold difference(1). Wind-pollinated plants such as maize have an even more extreme

difference in pollen production vs. ovule production per plant, with more than 1 million

pollen grains versus an average of 250 ovules per plant (a 4000-fold difference(2)).

Arabidopsis thaliana, which is a self-pollinating plant, also produces an excess of pollen,

with at least 2000 pollen grains per flower compared to an average of 60 ovules per

flower(3). Since pollen is produced in excess, in self-pollinated plants the number of

ovules (i.e. female gametes) sets the maximum seed number per flower.

The ability to manipulate ovule number to increase the reproductive potential of plants

requires an understanding of the molecular pathways that control ovule initiation.

The model plant *Arabidopsis thaliana* produces flowers with four whorls of organs:

sepals, petals, stamens, and carpels. The inner whorls (3 and 4) are responsible for

sexual reproduction, with pollen (the male gametophytes) produced in the whorl 3

stamens and the female gametophytes (also known as the embryo sacs), produced in

ovules contained within the whorl 4 carpels. Specification of the 4 whorls is controlled by

the "ABC" genes, with the C-class gene AGAMOUS (AG) a major regulator of carpel

development(4).

In Arabidopsis, ovules are initiated from the carpel margin meristem (CMM) at stage 9

of floral development(5). The Arabidopsis gynoecium comprises two carpels that are

fused vertically at their margins (6). The CMM develops on the adaxial face of the

carpels (inside the fused carpel cylinder) and will give rise to the placenta, ovules,

septum, transmitting tract, style, and stigma. Once the placenta is specified, all of the

ovule primordia are initiated at the same time(7). Subsequently, each primordium will be

patterned into three different regions: the funiculus, which connects the ovule to the

septum; the chalaza, which gives rise to the integuments; and the nucellus, which gives

rise to the embryo sac. Ovule development concludes with the specification of the

megaspore mother cell within the nucellus which undergoes meiosis followed by three

rounds of mitosis to form the mature haploid embryo sac (reviewed in(8)).

In Arabidopsis, CMM development requires coordination of transcriptional regulators

involved in meristem function with hormone signaling (reviewed in(6)). Most mutants

that have been reported to affect ovule number have pleiotropic effects related to the

establishment of polarity and boundaries during gynoecial development (reviewed in

(9)). For example, the *AINTEGUMENTA* (*ANT*) transcription factor regulates organ

initiation and cell divisions during flower development(10). *ANT* acts redundantly with

the related gene *AINTEGUMENTA-LIKE6/PLETHORA3* to regulate carpel margin

development and fusion which leads to a modest reduction in ovule number. This

phenotype is exacerbated when *ant* is combined with mutations in other carpel

development transcriptional regulators, such as *SEUSS* (*SEU*), *LEUNIG* (*LUG*),

SHATTERPROOF1 and *2* (*SHP1* and *SHP2*), *CRABSCLAW* (*CRC*), *FILAMENTOUS*

FLOWER (*FIL*), and *YABBY3* (*YAB3*). Mutant combinations between *ant* and these

mutants leads to severe defects in carpel fusion coupled with severe reductions in the

marginal tissues that give rise to the CMM (summarized in(6)). An extreme example is

the double mutant *seu-3 ant-1* which results in a complete loss of ovule initiation due to

defects in CMM development(11). The organ boundary genes, *CUP-SHAPED*

COTYLEDON1 and *2* (*CUC1* and *CUC2*), are also required for CMM development and

subsequent ovule initiation. *ant cuc2* mutants with *cuc1* levels decreased specifically in

the CMM by an RNAi construct driven by the *SEEDSTICK* promoter show an 80%

reduction in ovule number, indicating that *ANT* controls cell proliferation while *CUC1/2*

are necessary to set up the boundaries that allow ovule primordia to be initiated(12).

Plant hormones are also involved in gynoecium development and can have both indirect

and direct effects on ovule number. Auxin biosynthesis, transport, signaling, and

transport mutants have varying effects on gynoecium development and patterning,

many of which lead to pleiotropic effects on tissues and organs derived from the

CMM(13). Treatment of developing flowers with the auxin polar transport inhibitor NPA

showed that an apical-basal auxin gradient in the developing gynoecium is necessary

for patterning events that lead to ovule initiation(13). Cytokinin has also been implicated

in ovule initiation and development in Arabidopsis. Notably, triple mutants in the

ARABIDOPSIS HISTIDINE KINASE (*AHK*) cytokinin receptors, *AHK2*, *AHK3*, and

AHK4/CRE1, displayed a 90% reduction in ovule number due to decreased cytokinin

signaling(14). Conversely, double mutants in the cytokinin degrading cytokinin

oxidases/dehydrogenases (CKXs) displayed higher cytokinin levels in inflorescences

and produced more than double the number of ovules as in wild-type controls(15).

Brassinosteroids (BR) may also be positive regulators of ovule initiation. Gain-of-

function mutants in the BR-induced transcription factor *BZR1* had increased ovule

number per flower while BR-deficient and insensitive mutants had decreased ovule

number compared to wild-type controls. Upregulation of the ovule regulators *ANT* and

HULLENLOS (*HUL*) was correlated with *BZR1* activity, indicating the BR signaling

positively regulates ovule development(16). Recently, gibberellins were shown to

negatively regulate ovule number independently of auxin transport and signaling. A

gain-of-function mutation in the DELLA gene, GA-INSENSITVE (GAI) and loss-of-

function mutations in GA receptors all led to increased ovule number in Arabidopsis

(17).

To date, research on the factors controlling ovule number has been dominated by the

analyses of mutants that have been identified based on pleiotropic effects on

gynoecium development. In an attempt to identify loci that regulate ovule number

without affecting other aspects of flower morphology, we took advantage of natural

variation in ovule number in Arabidopsis accessions from diverse geographical

locations. Over 7,000 natural accessions are now available with intraspecific variation,

and next generation sequencing has been used to generate data on single nucleotide

polymorphisms (SNPs) from over 1,000 of these accessions as part of the 1001

genomes project(18). 100s of different phenotypes have been analyzed in this collection

such as flowering time, leaf shape and size, the ability to resist to pathogens, etc.(19).

In this study, we identified variation in ovule number per flower in a screen of 189

Arabidopsis accessions and conducted a Genome Wide Association Study (GWAS) to

identify loci associated with the ovule number trait. Further analysis of two loci identified

in our GWAS revealed that the *NEW ENHANCER of ROOT DWARFISM* (*NERD1*) and

OVULE NUMBER ASSOCIATED 2 (*ONA2*) genes participate in the determination of

ovule number during Arabidopsis flower development. The discovery of new ovule

number regulators in Arabidopsis has the potential to provide targets for the

development of crop varieties with greater yield.

Results

Natural Variation in Ovule Number

We set out to identify new regulators of ovule number in Arabidopsis by taking

advantage of phenotypic variation in naturally occurring accessions. We obtained 189

Arabidopsis accessions from the ABRC and assayed them for variation in ovule number

(Table S1). Since ovule number can vary throughout the life cycle of the plant[20], we

determined the average number of ovules from flowers 6-10 on the main stem of plants

that were vernalized for 4 weeks and then grown in long days at 22˚C. Under these

growth conditions, accessions displayed a remarkable diversity in ovule number per

flower, with a range of 39-82 ovules per flower (Fig 1A-B). The commonly used

reference accession, Col-0, falls in the middle of the range with an average ovule

number of 63±3 ovules.

In contrast to flowering time variation which has been shown to correlate with latitude of

origin in Arabidopsis accessions(21), ovule number was not strongly correlated with

location of origin in the accessions analyzed (Fig 1C, S1). Mapping ovule number data

onto a cladogram of the accessions used in our study revealed a cluster of low ovule

number accessions in one specific clade, indicating that these closely-related

accessions may have similar genetic control of ovule number (Fig 1D). Interestingly,

several clades were made up of accessions with high, medium, and low ovule numbers.

This suggests that the ovule number trait may be regulated by different loci that have

been selected for in some lineages.

GWAS reveals SNPs linked to natural variation in ovule number

In order to identify genomic regions linked to variation in ovule number, we assessed

whether the average ovule number per flower from 148 accessions was predicted by

Single Nucleotide Polymorphisms (SNP) available in the 1001 Full-sequence dataset;

TAIR 9. Logarithmic transformation was applied to the ovule number data to make the

results more reliable for parametric tests. Associations were tested for each SNP using

a linear regression model (LM) and the results were analyzed using GWAPP(22) (Fig

2A). A significance cutoff value of $-\log_{10}(p$ values) ≥ 6.2 identified at least 9 genomic

regions that are associated with variation in ovule number, while a higher cutoff of $-$

$\log_{10}(p$ values) ≥ 7.5 identifies only four significant genomic regions.

We next determined if known ovule number regulators (9, 15, 17, 23) colocalized with

our GWAS loci. Of the previously described loci, only BIN2 mapped close to a

significant association (Fig S2). This indicates that our GWAS has identified novel

functions for loci in the regulation of ovule number. For further analysis, we focused on

the two loci with the lowest p-values which are both located on the long arm of

chromosome 3 (Fig 2B-C). Genes containing the most significantly associated SNPs as

well as the 10 surrounding genes around the highest peak were considered as

candidates for regulating ovule number. These genes were further prioritized based on

whether they are expressed in developing pistils, by examining publicly available

transcriptome data in ePlant(24) (Fig S3). Based on this prioritization, 35 candidate

genes were selected, and of these 26 insertion mutants were available from the

Arabidopsis Biological Resource Center. All 26 mutants were evaluated for changes in

ovule number compared to the wild-type background, Col-0 (Table S2). Of these, two

had significantly reduced ovule number compared to the Col-0 control (Fig 3A, S4). The

strongest ovule number phenotype was found in insertion mutants in At3g51050, a gene

that was recently identified in a screen for enhancers of exocyst-mediated root

phenotypes and named *NEW ENHANCER OF ROOT DWARFISM 1* (*NERD1*)(25). The

second locus with T-DNA insertions affecting ovule number identified in our screen was

At3g60660, which we call *OVULE NUMBER ASSOCIATED 2* (*ONA2). ONA2* encodes

an unknown protein containing a DUF1395 domain (TAIR) (Fig 2C).

NERD1 is a positive regulator of ovule number

We focused our analysis on *NERD1* since it had the strongest effect on ovule number.

The fruits of homozygous mutants in two T-DNA insertion alleles, *nerd1-2* (described in

(25) and *nerd1-4,* had fewer ovules than wild-type plants (Fig 3A). To confirm that this

resulted from the disruption of NERD1, we generated transgenic plants expressing a

translational fusion of the *NERD1* protein with the red-fluorescent protein tdTomato

under the control of the native *NERD1* promoter. In *nerd1-2* and *nerd1-4* mutants, this

transgene fully rescued the reduced ovule phenotype and resulted in plants with ovule

numbers indistinguishable from Col-0 (Fig 3A). This demonstrates that *NERD1* is a

positive regulator of ovule number.

In addition to reduced ovule number per flower, homozygous *nerd1* mutants had fertility

defects. Homozygous *nerd1-2* mutants had 100% unfertilized ovules and homozygotes

of the less severe *nerd1-4* allele had 75% unfertilized ovules (Fig 3B-C and S5). *nerd1-*

2 mutants displayed both male and female defects. No viable pollen grains were

produced (Fig 3D-E) and embryo sacs were often aborted, leading to 40% unfertilized

ovules in *nerd1-2/nerd1-2* plants pollinated with wild-type pollen (Fig 3F-G and S5). We

performed reciprocal crosses between heterozygous *nerd1-2* mutants and Col-0 to

determine if the fertility defects were gametophytic or sporophytic. When heterozygous

nerd1-2/NERD1 was used as the female, there was no transmission defect,

demonstrating that the reduced female fertility in *nerd1-2* mutants was not

gametophytic. When *nerd1-2/NERD1* was used as the pollen donor, the transmission

efficiency of the mutant allele was reduced to 45%, indicating a male gametophytic

transmission defect (Table 1).

Table 1. Transmission efficiency of the *nerd1-2* allele determined by reciprocal crosses with wild-type Col-0.

Female	Male	NERD1/NERD1	nerd1-2/NERD1	Female TE%	Male TE %
nerd1-2/NERD1	Col-0	43	42	97.67	NA
Col-0	nerd1-2/NERD1	92	41	NA	44.57

NERD1 encodes an integral membrane protein

Phylogenetic analyses indicate that *NERD1* is a member of a low-copy number, highly

conserved gene family that is found throughout the plant kingdom and in cyanobacteria

(Fig 4A-B and (25)). The NERD1 protein is predicted to be an integral membrane

protein with a signal peptide and one transmembrane domain (Fig 5A). The majority of

the protein is predicted to be extracellular, with the transmembrane domain located near

the C-terminus and a 17 amino acid cytoplasmic extension. Transient expression of a

NERD1-GFP fusion in *Nicotiana benthamiana* with subcellular markers confirmed that

NERD1 co-localizes with the Golgi marker and partially colocalizes with a plasma

membrane marker (Fig 5B and D). NERD1 does not co-localize with ER, peroxisome,

and plastid markers (Fig 5C and S6).

NERD1 is expressed throughout Arabidopsis development

The *nerd1* ovule number and fertility phenotypes suggest that *NERD1* should be

expressed in developing flowers. We used a *NERD1$_{pro}$::gNERD1-GUS* fusion to

examine *NERD1* expression throughout Arabidopsis development. In

NERD1$_{pro}$::gNERD1-GUS inflorescences, GUS activity was detected throughout flower

development, including inflorescences, developing and mature anthers, and in the

stigma, ovules, and carpel walls of mature pistils (Fig 6). NERD1-GUS activity was

present in the carpel margin meristem (CMM) in stage 9 flowers, where ovule initiation

occurs (Fig 6C and E). *NERD1* reporter expression in the CMM during pistil

development is consistent with a role for *NERD1* in ovule initiation. During seedling

development, the *NERD1-GUS* reporter was detected in shoot and root apical

meristems (SAM and RAM) and in the vasculature (Fig 6I).

Overexpression of NERD1 increases plant productivity

We examined *NERD1* transcript levels in the low ovule number accession Altai-5

compared to Col-0. *NERD1* transcript accumulation was reduced in Altai-5 buds as

compared to Col-0 but similar in Altai-5 and Col-0 leaves (Fig 7). We hypothesized that

the low ovule number in Altai-5 may be linked to reduced *NERD1* expression in

developing flowers. In order to determine whether increasing *NERD1* expression is

sufficient to increase ovule number, we transformed Col-0 and the low ovule number

accession Altai-5, with a *NERD1-GFP* fusion construct driven by the constitutively

expressed Cauliflower Mosaic Virus 35S promoter (*35S::NERD1-GFP*). Overexpression

of *NERD1* had no effect on ovule number in the Col-0 background, but significantly

increased ovule number in the Altai-5 background (Fig 8A-C), indicating that the *NERD1*

effect on ovule number is background-dependent. The *35S::NERD1-GFP* plants

displayed an even more striking phenotype when overall plant architecture was

examined (Fig 8A). In both the Altai-5 and Col-0 backgrounds, *NERD1* overexpression

led to increased branching (Fig 8D) and shortened internode lengths between flowers,

leading to an overall increase in flower number in the overexpression plants compared

to untransformed controls (Fig 8E-H). Thus, *NERD1* overexpression leads to increased

biomass and reproductive capacity, with up to a 2.5-fold increase in total flower number

over the lifespan of the plant. While all independent transformants displayed increased

branching and flower number, some of the *35S::NERD1* plants were male sterile (Fig

S7). This male sterility correlated with *NERD1* expression levels and plants with higher

NERD1 transcript levels had more severe male sterility (Fig S7). The sterility effect was

more severe in Col-0 than in Altai-5 (Fig S7). The lower endogenous *NERD1*

expression in Altai-5 inflorescences might explain the lower sensitivity of Altai-5 to

NERD1 overexpression with respect to male fertility, demonstrating background-

dependent sensitivity to *NERD1* levels for both ovule number and male sterility.

NERD1 was recently identified in enhancer screen performed on exocyst mutants with

weakly dwarfed roots. *nerd1* mutants have reduced root growth as a result of impaired

cell expansion, indicating that NERD1 may be a positive regulator of exocyst-dependent

root growth(25). Consistently, the *35S::NERD1-GFP* plants had longer roots compared

to the Col-0 control, indicating that root development is also sensitive to *NERD1*

expression levels (Fig S8).

Three SNPs in NERD1 showed significant correlation with ovule number in our GWAS

(Fig 9A-B). Two of them are synonymous SNPs that are not predicted to change the

amino acid sequence, but the third is a non-synonymous SNP (C to A change in

comparison to Col-0 reference) causing a Serine to Tyrosine change at amino acid 230

of NERD1 (Fig 9A-B). This non-synonymous SNP was present in 10 out of the 16

lowest ovule number accessions and not present in the 16 highest ovule number

accessions (Fig 9A). Across all of the accessions in our GWAS panel, the "A" allele at

this position was significantly associated with lower ovule numbers (Fig 9C). However,

some accessions with the "C" allele of *NERD1* have low ovule numbers, including the

Altai-5 accession (Fig 9A), underscoring that multiple mechanisms influence ovule

number in Arabidopsis.

304   Discussion

Our experiments revealed that Arabidopsis displays a wide variation in the number of

ovules per flower, a key component of plant fitness and crop yield. Association genetics

linked this natural variation to the plant-specific gene, *NERD1*, that regulates the

number of ovules produced in Arabidopsis flowers and gametophyte development (Fig

3) and was previously found to alter root growth(25). Overexpression of *NERD1* led to

dramatic effects on plant architecture, indicating that *NERD1* may be involved in

regulating meristem activity during Arabidopsis development.

313   GWAS reveals new ovule number-associated loci in Arabidopsis

A quantitative trait locus (QTL) mapping study utilizing variation between L*er* and Cvi

identified QTLs residing on chromosomes 1, 2 (near the *ERECTA* gene), and two QTL

on chromosome 5(23). No follow-up study has identified the genes underlying these

QTLs. Our population displayed a much larger range of ovule numbers per flower (39-

84) than that seen in Cvi vs L*er* (66 and 56, respectively). We identified 9 significant

associations in our GWAS, but only one potentially overlaps with a known ovule number

QTL (see chromosome 5 in Fig S2). Of the genes with an identified effect on ovule

number, only BIN2 overlapped with a GWAS peak, indicating that our study has

revealed at least seven novel regulators of ovule number in Arabidopsis. We

molecularly identified two new genes that control ovule number that were linked to two

GWAS peaks on chromosome 3. This expands that number of known ovule number

determinants and confirms the value of GWAS as a forward genetics tool.

NERD1 may transmit signals across membranes

NERD1 is a plant-specific, low copy number gene that is found throughout the plant

kingdom. NERD1 was recently identified as an enhancer of root development

phenotypes in weak mutants of the exocyst subunits SEC8 and EXO70A1(25) While a

direct interaction with the exocyst could not be identified, the authors identified Golgi

localization of NERD1 and hypothesized that NERD1 could be involved in synthesizing

or modifying pectins or other polysaccharide components of the cell wall that are

synthesized in the Golgi and transported to the cell wall (25). Changes in cell wall

elasticity mediated by pectin modifications have been correlated with lateral organ

initiation at the shoot apical meristem (26, 27). The decreased ovule number in *nerd1*

mutants and increased lateral branching and flower initiation in *35S::NERD1*

transformants could be consistent with alterations in Golgi-synthesized cell wall

components affecting cell wall mechanics and organ initiation.

Our GFP localization experiments confirmed Golgi localization of the majority of

NERD1-GFP, but also identified fluorescence in the plasma membrane. In Arabidopsis,

the NERD1 protein is predicted to encode a 698 amino acid protein with a signal

peptide, one transmembrane domain, and a short (17 amino acid) cytoplasmic tail at the

C-terminus. This topology is consistent with NERD1 being involved in transducing

signals from the outside of the cell to the inside, perhaps as part of a complex with other

membrane proteins. In animals, α-integrins and β-integrins share a similar topology with

NERD1 and interact with each other to transduce information from the extracellular

matrix to the inside of the cell(28). The existence of integrin-like complexes in plants

remains controversial, since plants lack proteins with clear sequence homology to

animal integrins. However, antibodies to animal β-integrins interact with plant proteins,

indicating structural conservation(29). NERD1 has a predicted beta-propeller with a

calcium-binding pocket similar to α-integrins in its extracellular domain(25), but

determination of whether NERD1 functions in an integrin-like fashion awaits further

investigation.

NERD1 could also interact with other types of proteins to transduce signals. In plants,

receptor-like proteins (RLPs) with short cytoplasmic domains have been shown to

heterodimerize with receptor-like kinases (RLKs) to form a receptor complex that

transduces extracellular signals to intracellular signaling networks. The Arabidopsis

genome contains 57 RLPs with Leucine Rich Repeats (LRRs) in their extracellular

domains(30). These LRR-RLPs have a transmembrane domain near the C-terminus

and a short cytoplasmic domain and are predicted to form complexes with LRR-RLKs.

LRR-RLP/LRR-RLK complexes have been implicated in plant growth and development

and plant immunity (reviewed in(31)). For example, the CLAVATA2 (CLV2) RLP forms a

complex with the RLKs CORYNE (CRN) and/or CLAVATA1 (CLV1) RLK to perceive the

CLAVATA3 (CLV3) ligand and trigger intracellular signaling related to shoot apical

meristem maintenance(32-34). Likewise, the TOO MANY MOUTHS (TMM) RLP

interacts with the LRR-RLKs ERECTA (ER) and ERECTA-LIKE1 (ERL) to regulate

stomatal patterning(35, 36). In contrast to the LRR-RLPs, NERD1 does not share

sequence homology with other proteins in the Arabidopsis genome. Distinguishing

whether NERD1 functions similarly to RLPs, integrins, or some other signaling network

will require identification of NERD1-interacting proteins in future work.

***NERD1* and fertility**

Homozygous *nerd1* mutants are completely male sterile are partially female sterile. This

sterility is due to a lack of pollen production and problems in female gametophyte

development leading to aborted embryo sacs. Transmission efficiency tests using

heterozygous loss-of-function mutants revealed that *nerd1* could be transmitted through

the egg at near 100% efficiency, indicating a sporophytic effect on female gametophyte

development. Ovule development mutants that have defective integument development

such as *short integuments 1* (*sin1*), *bell 1* (*bel1*) and *ant* fail to produce functional

female gametophytes, indicating that female gametophyte development is dependent

on properly differentiated sporophytic cells in the ovule(37-41). Sporophytic

development in ovules seem to be normal in *nerd1* mutants, yet embryo sacs abort. As

a membrane protein, NERD1 could transmit some unknown signal between sporophytic

and gametophytic cells during embryo sac development.

The male fertility defect in *nerd1* plants is more severe than the female defect.

Homozygous *nerd1* anthers are empty, indicating that pollen development is not

initiated or aborted very early. Transmission efficiency tests using pollen from

heterozygous *nerd1* plants crossed to wild-type females revealed that *nerd1* also has

gametophytic effects on pollen function. The sporophytic effects could be related to

early stages of anther development. In particular, specification of the tapetum is critical

for pollen development (reviewed in(42)). In our *35S::NERD1* experiment, transformants

that accumulated the most *NERD1* transcripts were male sterile. Together these results

indicate that pollen development is sensitive to *NERD1* levels, i.e. either too much or

too little *NERD1* is detrimental to pollen development. Future experiments should focus

on determining the stage of anther and/or pollen development that is affected in *nerd1*

mutants and the specific cell types that express *NERD1* in developing anthers.

Even though *NERD1* is expressed broadly throughout the plant, above ground

vegetative development appears to be normal in *nerd1* loss-of-function mutants.

However, *nerd1* roots are shorter than normal and have root hair defects that include

bulging and rupture(25). *NERD1* could have distinct or related developmental functions

in roots and flowers, as is seen for many of the genes involved in hormonal regulation of

development(43).

NERD1 regulates lateral organ formation

NERD1 overexpression under control of the constitutive 35S promoter dramatically

changed plant architecture in both the Col-0 and Altai-5 backgrounds. *NERD1*

overexpression plants produced significantly more branches than wild-type controls and

produced significantly more flowers that were produced at closer intervals along the

stem. Like branches and flowers, ovules are lateral organs produced from meristematic

cells, indicating that *NERD1* may be a positive regulator of meristem activity. Lateral

meristem activation and subsequent branching have been shown to be affected by

several different hormones (reviewed in(44)). Classic experiments in *Vicia faba* showed

that auxin produced in the shoot apex inhibits axillary meristems(45), and more recently

strigolactones were identified as graft-transmissible suppressors of branching (reviewed

in (46)). Gibberellins also repress shoot branching in Arabidopsis, maize, and rice, with

GA-deficient mutants displaying increased branching compared to wild-type controls(47-

50). In contrast, cytokinins and brassinosteroids are both positive regulators of

branching(49-53). The mechanism through which these plant hormones work together

to regulate branching is unknown, but auxin has been proposed to control cytokinin and

strigolactone biosynthesis(54, 55), while the brassinosteroid signaling regulator *BES1*

may inhibit strigolactone signaling to promote branching(56). The promotion of

branching in *35S::NERD1* plants could be related to regulation of one or more of these

hormonal pathways. Like *NERD1*, upregulation of both cytokinin and brassinosteroid

signaling pathways have been shown to positively regulate branching and ovule

number(15, 53, 56, 57), suggesting that *NERD1* may be intimately connected to these

pathways. Future research is needed to explore the intersection between *NERD1* and

hormonal pathways.

NERD1-induced increases in ovule number are background-dependent

Overexpression of *NERD1* in the Col-0 and Altai-5 backgrounds led to increased

branching and flower number, but ovule number was only increased in the Altai-5

background, suggesting a background-dependence on the ovule number trait. In

Arabidopsis, natural accessions were shown to respond differently in their ability to

buffer GA perturbations caused by overexpressing GA20 oxidase 1, which encodes a

rate-limiting enzyme for GA biosynthesis(58). Genetic background dependence has

been shown to be a wide-spread phenomenon in *C. elegans*, with approximately 20% of

RNAi-induced mutations (out of 1400 genes tested) displaying different phenotypic

severity in two different genetic backgrounds(59). Similar to our results with *NERD1*,

where Altai-5 has lower endogenous levels of the *NERD1* transcript in developing

flowers compared to Col-0, the severity of the *C. elegans* phenotypes could be linked to

gene expression levels of either the target gene itself or other genes in the same

pathway(59).

Our GWAS revealed multiple novel loci that control ovule number in Arabidopsis. It

remains to be seen if exocyst function is linked to NERD1's role in ovule number

determination. Identification of NERD1 interacting proteins will provide candidates for

other players in the NERD1 pathway. Some of the other novel loci identified in our

GWAS may well participate in the same signaling pathway as *NERD1*. While *NERD1* is

a promising candidate for engineering plants with increased ovule number, the

differential responses to *NERD1* overexpression seen in Altai-5 and Col-0 suggest that

specific alleles at other loci may be necessary for achieving maximum effect of *NERD1*

overexpression.

Materials and Methods

Plant Material and Growth Conditions

Arabidopsis thaliana accessions and insertion mutants were ordered from the

Arabidopsis Biological Resource Center at Ohio State University (ABRC). ABRC stock

numbers for the accessions and insertion mutants used in our study are listed in

supplementary tables 1 and 2. Seeds were sterilized and plated on ½ Murashige and

Skoog (MS) plates. All plates were sealed and stratified at 4 °C for two days, and then

transferred to the growth chamber (long day conditions, 16h of light and 8 h of dark at

22°C) for germination and growth. After one-week, seedlings were transplanted to soil.

Many of the Arabidopsis accessions require vernalization for flowering(60), we therefore

chose to vernalize all of the accessions in our study for 4 weeks at 4 °C. After

vernalization, the plants were returned to the growth chamber and grown under the long

day conditions described above. Seeds from transformed lines were sterilized and

plated on ½ MS plate with 20mg/L hygromycin for selection of transgenic seedlings,

which were then transplanted to soil and grown in long days.

Ovule Number Phenotyping

For determination of ovule number per flower, we counted four to five siliques per plant

and five plants per accession. Carpel walls were removed with a dissecting needle and

total ovule number (including unfertilized and aborted ovules) was counted with the aid

of a Leica dissecting microscope. To minimize age-related variation in ovule number,

we counted siliques from flowers 6-10 on the primary shoot for all accessions.

Genome-Wide Association Study

GWAS was performed using GWAPP, which is a GWAS web application for Genome-

Wide Association Mapping in *Arabidopsis* (http://gwas.gmi.oeaw.ac.at) (22). In our

study, 148 accessions had single nucleotide polymorphisms (SNPs) data available on

the 1001 Full-sequence dataset; TAIR 9. Logarithmic transformation was applied to

make the results more reliable for parametric tests. A simple linear regression (LM) was

used to generate the Manhattan plot by using GWAPP(22). SNPs with P values ≤ 1 ×

10^{-6} were further considered as candidate loci linked to alleles that regulate ovule

number (a horizontal dashed line in Fig. 2 shows the 5% FDR threshold -$\log_{10}p$

value=6.2, which was computed by the Benjamini-Hochberg-Yekutieli method). SNPs

with <15 minor allele count (MAC) were not considered to help control false positive

rates. 10 genes flanking the highest SNP for each locus were tabulated as candidate

genes for each significant association.

Cloning and Generation of Transgenic Lines

For complementation and overexpression experiments, Gateway Technology was used

to make all the constructs. Genomic DNA fragments corresponding to the coding

regions of candidate genes were amplified from either beginning of the promoter

(defined by the end of the upstream gene) or the start codon to the end of the CDS

(without stop codon) by PCR with primers that had attB1 and attB2 sites from Col-0

genomic DNA (see supplementary table 3 for primer sequences). For amplifying

At3g60660 and At3g51050, PHUSION High-Fidelity Polymerase (NEB, M0535S) was

used. All the PCR products were put into entry vector pDONR207 by BP reactions and

then were recombined into destination vector pMDC83 (GFP) by LR reaction(61).

Native promoter constructs were amplified from ~2KB promoter region to the end

(without stop codon) by PCR with primers that had attB1 and attB2 sites from Col-0

genomic DNA. The PCR products were put into entry vector pDONR207 by BP

reactions and then were recombined into destination vector pMDC32 (tdTomato) and

pMDC163 (GUS) by LR reaction. All constructs were transformed into *Agrobacterium*

tumefaciens strain GV3101 and then used for plant transformation by the floral-dip

method(62).

GUS Staining

GUS staining was performed as previously described(63). Samples were imaged with

Differential Interference Contrast (DIC) on a Nikon Eclipse Ti2-E microscope. 12

independent T1 NERD-GUS transformants were analyzed and showed similar GUS

patterns.

Transient Expression in *N. benthamiana*

Leaves from 3-4 week old *N. benthamiana* were co-infiltrated with 35S::NERD1-GFP

and Golgi-mCherry, ER-mCherry, PM-mCherry, Plastid- mCherry and Peroxisome-

mCherry markers from(64) as previously described(65). Leaves were imaged 2-3 days

after infiltration with a Nikon A1Rsi inverted confocal microscope under 20x dry and 40x

water objectives with GFP excited by a 488nm laser and mCherry excited by a 561nm

laser in normal mode.

Alexander Staining of Pollen

Mature anthers from Col-0 and *nerd1-2/nerd1-2* were dissected under the Leica

dissecting microscope and placed into a drop (20 µl) of Alexander staining solution on a

microscope slide(66). After several minutes of staining, samples were imaged with a

Nikon Eclipse Ti2-E microscope.

Quantitative Real-time RT-PCR

For qRT-PCR, leaves and young flowers were collected from mature plants of Col-0 and

Altai-5. Samples were immediately frozen in liquid nitrogen, ground, and total RNA was

extracted using the E.Z.N.A Plant RNA kit (OMEGA, USA). Oligo-dT primers and

Superscript II reverse transcriptase (Invitrogen) was used for cDNA synthesis. qRT-

PCR reactions were prepared using SYBR Green PCR Master Mix and PCR was

conducted with a StepOnePlus RT-PCR system. Relative quantifications were

performed for all genes with the Actin11 used as an internal reference. The primers

used for qRT-PCR shown in supplementary table 3.

Phylogenetic Analysis

The cladogram tree was generated in MEGA7, which nucleotide distance and neighbor-

join tree file were calculated by PHYlogeny Inference Package (PHYLIP, version 3.696).

The phylogenetic tree of NERD1 was inferred using neighbor-joining method in

MEGA7(67). The associated taxa clustered together with the bootstrap test (1000

replicates)(68). All the branch lengths are in the same units as those of the evolutionary

distances used to generate the phylogenetic tree.

Supplemental Material

Supplementary Figure 1. Statistical analysis of the association between geographic

distribution and ovule number per flower in the Arabidopsis accessions used in the

GWAS.

Supplementary Figure 2. Previously identified ovule number related genes' position in

relation to the Manhattan plot generated from the ovule number GWAS.

Supplementary Figure 3. Heat map of candidate gene expression patterns in

reproductive tissues.

Supplementary Figure 4. Average ovule number per flower of candidate gene T-DNA

mutants on chromosome 3.

Supplementary Figure 5. Homozygous *nerd1* mutants have high levels of infertility can

be partially rescued by pollinating with Col-0 wild-type pollen.

Supplementary Figure 6. Co-localization of 35S::NERD1-GFP with peroxisome and

plastid markers.

Supplementary Figure 7. Male sterility in *35S::NERD1* plants is linked to high *NERD1*

expression.

Supplementary Figure 8. *NERD1* is involved in in root growth.

Supplementary Figure 9. Amino acid changes in and around the *NERD1* gene in low

and high ovule number accessions.

Supplementary table 1. List of Arabidopsis accessions used in the study and their

ovule number data.

Supplementary Figure 2. Chromosome 3 candidate genes and insertion mutants.

Supplementary Figure 3. List of primers used for genotyping and cloning.

References

1.   Palmer RG, Albertsen MC, Heer H. Pollen production in soybeans with respect to
genotype, environment, and stamen position. Euphytica. 1978;27(2):427-33.
2.   Uribelarrea M, Cárcova J, Otegui ME, Westgate ME. Pollen Production,
Pollination Dynamics, and Kernel Set in Maize. Crop Science. 2002;42:1910-8.
3.   Yan J, Chia JC, Sheng H, Jung HI, Zavodna TO, Zhang L, et al. Arabidopsis
Pollen Fertility Requires the Transcription Factors CITF1 and SPL7 That Regulate
Copper Delivery to Anthers and Jasmonic Acid Synthesis. Plant Cell. 2017;29(12):3012-
29.
4.   Coen ES, Meyerowitz EM. The war of the whorls: genetic interactions controlling
flower development. Nature. 1991;353(6339):31-7.
5.   Smyth DR, Bowman JL, Meyerowitz EM. Early Flower Development in
Arabidopsis. Plant Cell. 1990;2(8):755-67.
6.   Reyes-Olalde JI, Zuniga-Mayo VM, Chavez Montes RA, Marsch-Martinez N, de
Folter S. Inside the gynoecium: at the carpel margin. Trends Plant Sci.
2013;18(11):644-55.
7.   Robinson-Beers K, Pruitt RE, Gasser CS. Ovule Development in Wild-Type
Arabidopsis and Two Female-Sterile Mutants. Plant Cell. 1992;4(10):1237-49.
8.   Drews GN, Koltunow AM. The female gametophyte. Arabidopsis Book.
2011;9:e0155.
9.   Cucinotta M, Colombo L, Roig-Villanova I. Ovule development, a new model for
lateral organ formation. Front Plant Sci. 2014;5:117.
10.   Klucher KM, Chow H, Reiser L, Fischer RL. The AINTEGUMENTA gene of
arabidopsis required for ovule and female gametophyte development is related to the
floral homeotic gene APETALA2. Plant Cell. 1996;8(2):137-53.
11.   Azhakanandam S, Nole-Wilson S, Bao F, Franks RG. SEUSS and
AINTEGUMENTA mediate patterning and ovule initiation during gynoecium medial
domain development. Plant Physiol. 2008;146(3):1165-81.
12.   Galbiati F, Sinha Roy D, Simonini S, Cucinotta M, Ceccato L, Cuesta C, et al. An
integrative model of the control of ovule primordia formation. Plant J. 2013;76(3):446-
55.
13.   Nemhauser JL, Feldman LJ, Zambryski PC. Auxin and ETTIN in Arabidopsis
gynoecium morphogenesis. Development. 2000;127(18):3877-88.
14.   Bencivenga S, Simonini S, Benkova E, Colombo L. The transcription factors
BEL1 and SPL are required for cytokinin and auxin signaling during ovule development
in Arabidopsis. Plant Cell. 2012;24(7):2886-97.
15.   Bartrina I, Otto E, Strnad M, Werner T, Schmulling T. Cytokinin regulates the
activity of reproductive meristems, flower organ size, ovule formation, and thus seed
yield in Arabidopsis thaliana. Plant Cell. 2011;23(1):69-80.

16.   Huang HY, Jiang WB, Hu YW, Wu P, Zhu JY, Liang WQ, et al. BR Signal
Influences Arabidopsis Ovule and Seed Number through Regulating Related Genes
Expression by BZR1. Molecular Plant. 2013;6(2):456-69.
17.   Gomez MD, Barro-Trastoy D, Escoms E, Saura-Sanchez M, Sanchez I, Briones-
Moreno A, et al. Gibberellins negatively modulate ovule number in plants. Development.
2018;145(13).
18.   Weigel D. Natural variation in Arabidopsis: from molecular genetics to ecological
genomics. Plant Physiol. 2012;158(1):2-22.
19.   Atwell S, Huang YS, Vilhjalmsson BJ, Willems G, Horton M, Li Y, et al. Genome-
wide association study of 107 phenotypes in Arabidopsis thaliana inbred lines. Nature.
2010;465(7298):627-31.
20.   Wetzsteini HY, Yi WG, Porter JA, Ravid N. Flower Position and Size Impact
Ovule Number per Flower, Fruitset, and Fruit Size in Pomegranate. J Am Soc Hortic
Sci. 2013;138(3):159-66.
21.   Stinchcombe JR, Weinig C, Ungerer M, Olsen KM, Mays C, Halldorsdottir SS, et
al. A latitudinal cline in flowering time in Arabidopsis thaliana modulated by the flowering
time gene FRIGIDA. Proc Natl Acad Sci U S A. 2004;101(13):4712-7.
22.   Seren U, Vilhjalmsson BJ, Horton MW, Meng D, Forai P, Huang YS, et al.
GWAPP: a web application for genome-wide association mapping in Arabidopsis. Plant
Cell. 2012;24(12):4793-805.
23.   Alonso-Blanco C, Blankestijn-de Vries H, Hanhart CJ, Koornneef M. Natural
allelic variation at seed size loci in relation to other life history traits of Arabidopsis
thaliana. Proc Natl Acad Sci U S A. 1999;96(8):4710-7.
24.   Waese J, Fan J, Pasha A, Yu H, Fucile G, Shi R, et al. ePlant: Visualizing and
Exploring Multiple Levels of Data for Hypothesis Generation in Plant Biology. Plant Cell.
2017;29(8):1806-21.
25.   Cole R, Peremyslov V, Van Why S, Moussaoui I, Ketter A, Cool R, et al. A
broadly conserved NERD genetically interacts with the exocyst to affect root growth and
cell expansion. J Exp Bot. 2018;69(15):3625-37.
26.   Peaucelle A, Braybrook SA, Le Guillou L, Bron E, Kuhlemeier C, Hofte H. Pectin-
induced changes in cell wall mechanics underlie organ initiation in Arabidopsis. Curr
Biol. 2011;21(20):1720-6.
27.   Peaucelle A, Louvet R, Johansen JN, Hofte H, Laufs P, Pelloux J, et al.
Arabidopsis phyllotaxis is controlled by the methyl-esterification status of cell-wall
pectins. Curr Biol. 2008;18(24):1943-8.
28.   Barczyk M, Carracedo S, Gullberg D. Integrins. Cell Tissue Res.
2010;339(1):269-80.
29.   Nagpal P, Quatrano RS. Isolation and characterization of a cDNA clone from
Arabidopsis thaliana with partial sequence similarity to integrins. Gene. 1999;230(1):33-
40.
30.   Wang G, Ellendorff U, Kemp B, Mansfield JW, Forsyth A, Mitchell K, et al. A
genome-wide functional investigation into the roles of receptor-like proteins in
Arabidopsis. Plant Physiol. 2008;147(2):503-17.
31.   He Y, Zhou J, Shan L, Meng X. Plant cell surface receptor-mediated signaling - a
common theme amid diversity. J Cell Sci. 2018;131(2).

32.   Bleckmann A, Weidtkamp-Peters S, Seidel CA, Simon R. Stem cell signaling in
Arabidopsis requires CRN to localize CLV2 to the plasma membrane. Plant Physiol.
2010;152(1):166-76.
33.   Brand U, Fletcher JC, Hobe M, Meyerowitz EM, Simon R. Dependence of Stem Cell Fate
in Arabidopsis on a Feedback Loop Regulated by CLV3 Activity. Science.
2000;289(5479):617-9.
34.   Zhu Y, Wang Y, Li R, Song X, Wang Q, Huang S, et al. Analysis of interactions among the
CLAVATA3 receptors reveals a direct interaction between CLAVATA2 and CORYNE in
Arabidopsis. Plant J. 2010;61(2):223-33.
35.   Abrash EB, Davies KA, Bergmann DC. Generation of signaling specificity in Arabidopsis by
spatially restricted buffering of ligand-receptor interactions. Plant Cell. 2011;23(8):2864-79.
36.   Shpak ED, McAbee JM, Pillitteri LJ, Torii KU. Stomatal patterning and differentiation by
synergistic interactions of receptor kinases. Science. 2005;309(5732):290-3.
37.   Elliott RC, Betzner AS, Huttner E, Oakes MP, Tucker WQ, Gerentes D, et al.
AINTEGUMENTA, an APETALA2-like gene of Arabidopsis with pleiotropic roles in ovule
development and floral organ growth. The Plant Cell. 1996;8(2):155-68.
38.   Klucher KM, Chow H, Reiser L, Fischer RL. The AINTEGUMENTA gene of Arabidopsis
required for ovule and female gametophyte development is related to the floral homeotic gene
APETALA2. The Plant Cell. 1996;8(2):137-53.
39.   Modrusan Z, Reiser L, Feldmann KA, Fischer RL, Haughn GW. Homeotic Transformation
of Ovules into Carpel-like Structures in Arabidopsis. The Plant Cell. 1994;6(3):333-49.
40.   Robinson-Beers K, Pruitt RE, Gasser CS. Ovule Development in Wild-Type Arabidopsis
and Two Female-Sterile Mutants. The Plant Cell. 1992;4(10):1237-49.
41.   Schneitz K, Hulskamp M, Kopczak SD, Pruitt RE. Dissection of sexual organ ontogenesis:
a genetic analysis of ovule development in Arabidopsis thaliana. Development.
1997;124(7):1367-76.
42.   Wilson ZA, Zhang DB. From Arabidopsis to rice: pathways in pollen development. J Exp
Bot. 2009;60(5):1479-92.
43.   Schaller GE, Bishopp A, Kieber JJ. The Yin-Yang of Hormones: Cytokinin and Auxin
Interactions in Plant Development. The Plant Cell. 2015;27(1):44-63.
44.   Rameau C, Bertheloot J, Leduc N, Andrieu B, Foucher F, Sakr S. Multiple pathways
regulate shoot branching. Front Plant Sci. 2014;5:741.
45.   Thimann KV, Skoog F. Studies on the Growth Hormone of Plants: III. The Inhibiting
Action of the Growth Substance on Bud Development. Proc Natl Acad Sci U S A. 1933;19(7):714-
6.
46.   Domagalska MA, Leyser O. Signal integration in the control of shoot branching. Nature
Reviews Molecular Cell Biology. 2011;12:211.
47.   Lo SF, Yang SY, Chen KT, Hsing YI, Zeevaart JA, Chen LJ, et al. A novel class of gibberellin
2-oxidases control semidwarfism, tillering, and root development in rice. Plant Cell.
2008;20(10):2603-18.
48.   Silverstone AL, Mak PY, Martinez EC, Sun TP. The new RGA locus encodes a negative
regulator of gibberellin response in Arabidopsis thaliana. Genetics. 1997;146(3):1087-99.
49.   Best NB, Hartwig T, Budka J, Fujioka S, Johal G, Schulz B, et al. nana plant2 Encodes a
Maize Ortholog of the Arabidopsis Brassinosteroid Biosynthesis Gene DWARF1, Identifying

Developmental Interactions between Brassinosteroids and Gibberellins. Plant Physiol.
2016;171(4):2633-47.

50. Best NB, Johal G, Dilkes BP. Phytohormone inhibitor treatments phenocopy
brassinosteroid–gibberellin dwarf mutant interactions in maize. Plant Direct. 2017;1(2).

51. Wickson M, Thimann KV. The Antagonism of Auxin and Kinetin in Apical Dominance.
Physiologia Plantarum. 1958;11(1):62-74.

52. Sachs T, Thimann KV. THE ROLE OF AUXINS AND CYTOKININS IN THE RELEASE OF BUDS
FROM DOMINANCE. American Journal of Botany. 1967;54(1):136-44.

53. Yin Y, Wang Z-Y, Mora-Garcia S, Li J, Yoshida S, Asami T, et al. BES1 Accumulates in the
Nucleus in Response to Brassinosteroids to Regulate Gene Expression and Promote Stem
Elongation. Cell. 2002;109(2):181-91.

54. Foo E, Bullier E, Goussot M, Foucher F, Rameau C, Beveridge CA. The branching gene
RAMOSUS1 mediates interactions among two novel signals and auxin in pea. Plant Cell.
2005;17(2):464-74.

55. Shimizu-Sato S, Tanaka M, Mori H. Auxin-cytokinin interactions in the control of shoot
branching. Plant molecular biology. 2009;69(4):429-35.

56. Wang Y, Sun S, Zhu W, Jia K, Yang H, Wang X. Strigolactone/MAX2-induced degradation
of brassinosteroid transcriptional effector BES1 regulates shoot branching. Developmental cell.
2013;27(6):681-8.

57. Huang HY, Jiang WB, Hu YW, Wu P, Zhu JY, Liang WQ, et al. BR signal influences
Arabidopsis ovule and seed number through regulating related genes expression by BZR1. Mol
Plant. 2013;6(2):456-69.

58. Nam YJ, Herman D, Blomme J, Chae E, Kojima M, Coppens F, et al. Natural Variation of
Molecular and Morphological Gibberellin Responses. Plant Physiol. 2017;173(1):703-14.

59. Vu V, Verster AJ, Schertzberg M, Chuluunbaatar T, Spensley M, Pajkic D, et al. Natural
Variation in Gene Expression Modulates the Severity of Mutant Phenotypes. Cell.
2015;162(2):391-402.

60. Lempe J, Balasubramanian S, Sureshkumar S, Singh A, Schmid M, Weigel D. Diversity of
flowering responses in wild Arabidopsis thaliana strains. PLoS Genet. 2005;1(1):109-18.

61. Curtis MD, Grossniklaus U. A gateway cloning vector set for high-throughput functional
analysis of genes in planta. Plant Physiol. 2003;133(2):462-9.

62. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. 1998;16(6):735-43.

63. Davis TC, Jones DS, Dino AJ, Cejda NI, Yuan J, Willoughby AC, et al. Arabidopsis thaliana
MLO genes are expressed in discrete domains during reproductive development. Plant Reprod.
2017;30(4):185-95.

64. Nelson BK, Cai X, Nebenfuhr A. A multicolored set of in vivo organelle markers for co-
localization studies in Arabidopsis and other plants. Plant J. 2007;51(6):1126-36.

65. Jones DS, Yuan J, Smith BE, Willoughby AC, Kumimoto EL, Kessler SA. MILDEW
RESISTANCE LOCUS O Function in Pollen Tube Reception Is Linked to Its Oligomerization and
Subcellular Distribution. Plant Physiol. 2017;175(1):172-85.

66. Alexander MP. Differential staining of aborted and nonaborted pollen. Stain Technol.
1969;44(3):117-22.

67.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. Mol Biol Evol. 2016;33(7):1870-4.
68.   Felsenstein J. Confidence-Limits on Phylogenies - an Approach Using the Bootstrap.
Evolution. 1985;39(4):783-91.
751

Acknowledgements

We thank Dr. Brian Dilkes, Dr. John Fowler, Dr. Daniel S. Jones, Thomas Davis, Yan

Ju, and Rachel Flynn for helpful discussions and comments on the manuscript. We

thank Dr. Xutong Wang for help with the statistical analysis. Purdue University Start-up

funds and a grant from the Oklahoma Center for the Advancement of Science and

Technology #PS14-008 to S.A.K. supported this work.

Author contributions

Conceptualization: JY SAK.

Data curation: JY.

Formal analysis: JY SAK.

Funding acquisition: SAK.

Investigation: JY.

Methodology: JY.

Project administration: SAK.

Resources: JY SAK.

Supervision: SAK.

Validation: JY.

Visualization: JY.

Writing – original draft: JY.

Writing – review & editing: JY SAK.

Competing Interests

The authors declare no competing interests.

Fig 1. Arabidopsis accessions display natural variation in ovule number per flower. (A) Boxplot of ovule number in 189 Arabidopsis accessions. The detailed names of these accessions are in Supplementary Table 1. (B) Dissected siliques of low, medium, and high ovule number accessions (Bar=2.0mm). (C) Geographic distribution of accessions. Groups 1 to 6 correspond to ovule number ranking (from low to high), with the colors corresponding to panel A. The pie chart indicates the percentage of accessions in each country in each ovule number group, while the size of the pie chart corresponds to the total number of accessions per country. (D) Cladogram of accessions used in this study (generated in MEGA7). The 15 accessions labeled in blue had the lowest ovule numbers, and the 15 accessions labeled with red had the highest ovule numbers. The arrow points to a clade with clustered low ovule number accessions.

Fig 2. GWAS identifies candidate loci associated with ovule number per flower. (A) Manhattan plot for the SNPs associated with ovule number per flower. Chromosomes are depicted in different colors. The horizontal blue dashed line corresponds to a $-\log_{10}(p$ values) ≥ 6.2 and the red dashed line corresponds to $-\log_{10}(p$ values) ≥ 7.5 after Benjamini Hochberg (False Discovery Rate) correction. (B) The genomic region surrounding the two most significant GWA peaks on Chromosome 3. (C) Genes in the genomic region surrounding the two significant GWA peaks in (D). The red boxes are AT3G51050 (*NERD1*) and AT3T60660 (*ONA2*).

Fig 3. *nerd1* mutants have reduced fertility. (A) Bar graph of ovule number per flower

in Col-0, *nerd1-2*, *nerd1-4* and their complementation lines. "***" indicates statistical

significance (*p* value<0.001 determined by Student's t-test). (B-C) *nerd1-2/nerd1-2*

displays lower ovule number, infertile ovules, and short siliques compared to Col-0 and

complemented lines. (D) Alexander stained Col-0 anthers have viable pollen grains (red

indicates viable pollen while the green is non-viable). (E) Alexander-stained *nerd1-*

2/nerd1-2 anthers have no viable pollen. (F) Col-0 mature ovule with a normal embryo

sac. (G) *nerd1-2/nerd1-2* mature ovule with an aborted embryo sac. Bars=2.5mm (B-C),

20μm (D-G).

Fig 4. NERD1 is conserved across plant lineages. (A) Phylogenetic tree from amino

acid alignment of NERD1 generated using the Neighbor-joining method in MEGA7. (B)

NERD1 alignment showing sequence conservation among species. The green boxes

correspond to conserved regions of NERD1 and the lines represent gaps in the

alignment. The red box indicates the FG-GAP domain annotated in NERD1 by

Langhans, et al., 2017.

Fig 5. NERD1 co-localizes with a Golgi marker in *N. benthamiana* epidermal cells.

(A) NERD1 protein domains determined by TMHMM 2.0. (B) NERD1-GFP (green

signal) co-localizes with Golgi-mCherry (magenta signal) in *N. benthamiana* epidermal

cells. (C) NERD1-GFP (green signal) does not co-localize with ER-mCherry (magenta

signal). (D) NERD1-GFP (green signal) partially co-localizes with PM-mCherry

(magenta signal). Bars=15 μm.

Fig 6. NERD1 expression in plant development. The NERD1$_{pro}$::NERD1-GUS

reporter (blue signal) is detected in inflorescence (A-B), the flower in stage 9 (C), the

pistil in stage 8 (D), the pistil in stage 9 (E), mature pistil (F), mature flower (G), mature

anther (H), and SAM and RAM of seedlings (I). Bars=25 µm.

Fig 7. NERD1 expression is reduced in developing flowers of a low ovule number

accession.

(A) qRT-PCR of NERD1 in Col-0 and Altai-5 inflorescences. (B) qRT-PCR of NERD1 in

Col-0 and Altai-5 leaves. "***" indicates statistical significance at p value<0.001

determined by Student's t-test.

Fig 8. Overexpression of NERD1 affects plant architecture and fertility. (A-B)

35S::NERD1-GFP expressed in Col-0 and Altai-5 backgrounds. Bars=5cm. Bar graphs

of total ovule number per flower (C), total branch number (D), total flower number per

plant (E), number of flowers within 15 cm of the main shoot (F), and distance between

adjacent flowers (G-H) in 35S::NERD1-GFP transformants in the Col-0 and Altai-5

backgrounds compared to untransformed controls.

Fig 9. Sequence variation in the *NERD1* locus correlates with ovule number. (A)

SNPs in and around the *NERD1* gene. The blue line indicates the 16 lowest ovule

number accessions and the red line indicates the 16 highest ovule number accessions.

SNPs are identified based on comparison to the Col-0 reference genome. The red

boxes highlight low-ovule number associated SNPs (the turquoise bar indicates two

SNPs in adjacent nucleotides, a non-synonymous C-A SNP and a synonymous C-G

SNP). Only genomes that were available on the SALK 1,001 genomes browser

(http://signal.salk.edu/atg1001/3.0/gebrowser.php) were considered. (B) One of the

ovule number-associated SNPs in NERD1 causes a serine-tyrosine substitution in low

ovule number accessions. (C) Average ovule number in *NERD1* "A" and "C" allele-

containing accessions. The "A" allele is significantly associated with lower ovule

numbers. "***" indicates statistical significance at p value<0.001 determined by

Student's t-test.

Fig 1. Arabidopsis accessions display natural variation in ovule number per flower. (A) Boxplot of ovule number in 189 Arabidopsis accessions. The detailed names of these accessions are in Supplementary Table 1. (B) Dissected siliques of low, medium, and high ovule number accessions (Bar=2.0mm). (C) Geographic distribution of accessions. Groups 1 to 6 correspond to ovule number ranking (from low to high), with the colors corresponding to panel A. The pie chart indicates the percentage of accessions in each country in each ovule number group, while the size of the pie chart corresponds to the total number of accessions per country. (D) Cladogram of accessions used in this study (generated in MEGA7). The 15 accessions labeled in blue had the lowest ovule numbers, and the 15 accessions labeled with red had the highest ovule numbers. The arrow points to a clade with clustered low ovule number accessions.

Figure 2

Fig 2. GWAS identifies candidate loci associated with ovule number per flower.
(A) Manhattan plot for the SNPs associated with ovule number per flower.
Chromosomes are depicted in different colors. The horizontal blue dashed line
corresponds to a $-\log_{10}(p$ values) ≥ 6.2 and the red dashed line corresponds to $-\log_{10}(p$
values) ≥ 7.5 after Benjamini Hochberg (False Discovery Rate) correction. (B) The
genomic region surrounding the two most significant GWA peaks on Chromosome 3.
(C) Genes in the genomic region surrounding the two significant GWA peaks in (D). The
red boxes are AT3G51050 (*NERD1*) and AT3T60660 (*ONA2*).

Fig 3. *nerd1* mutants have reduced fertility. (A) Bar graph of ovule number per flower in Col-0, *nerd1-2*, *nerd1-4* and their complementation lines. "***" indicates statistical significance (*p* value<0.001 determined by Student's t-test). (B-C) *nerd1-2/nerd1-2* displays lower ovule number, infertile ovules, and short siliques compared to Col-0 and complemented lines. (D) Alexander stained Col-0 anthers have viable pollen grains (red indicates viable pollen while the green is non-viable). (E) Alexander-stained *nerd1-2/nerd1-2* anthers have no viable pollen. (F) Col-0 mature ovule with a normal embryo sac. (G) *nerd1-2/nerd1-2* mature ovule with an aborted embryo sac. Bars=2.5mm (B-C), 20μm (D-G).

Figure 4

Fig 4. NERD1 is conserved across plant lineages. (A) Phylogenetic tree from amino acid alignment of NERD1 generated using the Neighbor-joining method in MEGA7. (B) NERD1 alignment showing sequence conservation among species. The green boxes correspond to conserved regions of NERD1 and the lines represent gaps in the alignment. The red box indicates the FG-GAP domain annotated in NERD1 by Langhans, et al., 2017.

Figure 5

Fig 5. **NERD1 co-localizes with a Golgi marker in *N. benthamiana* epidermal cells.**
(A) NERD1 protein domains determined by TMHMM 2.0. (B) NERD1-GFP (green signal) co-localizes with Golgi-mCherry (magenta signal) in *N. benthamiana* epidermal cells. (C) NERD1-GFP (green signal) does not co-localize with ER-mCherry (magenta signal). (D) NERD1-GFP (green signal) partially co-localizes with PM-mCherry (magenta signal). Bars=15 μm.

Fig 6. NERD1 expression in plant development. The NERD1$_{pro}$::NERD1-GUS reporter (blue signal) is detected in inflorescence (A-B), the flower in stage 9 (C), the pistil in stage 8 (D), the pistil in stage 9 (E), mature pistil (F), mature flower (G), mature anther (H), and SAM and RAM of seedlings (I). Bars=25 μm.

Figure 7

Fig 7. NERD1 expression is reduced in developing flowers of a low ovule number accession. (A) qRT-PCR of NERD1 in Col-0 and Altai-5 inflorescences. (B) qRT-PCR of NERD1 in Col-0 and Altai-5 leaves. "***" indicates statistical significance at *p* value<0.001 determined by Student's t-test.

Figure 8

Fig 8. Overexpression of NERD1 affects plant architecture and fertility. (A-B)
35S::NERD1-GFP expressed in Col-0 and Altai-5 backgrounds. Bars=5cm. Bar graphs
of total ovule number per flower (C), total branch number (D), total flower number per
plant (E), number of flowers within 15 cm of the main shoot (F), and distance between
adjacent flowers (G-H) in 35S::NERD1-GFP transformants in the Col-0 and Altai-5
backgrounds compared to untransformed controls.

Figure 9

Position	*p* value	Alleles	Amino	LOCN
18954034	9.55E-07	C-T	R4	Exon
18955603	2.45E-08	C-A	S230Y	Exon
18955604	2.45E-08	T-C	S230	Exon

Fig 9. Sequence variation in the *NERD1* locus correlates with ovule number. (A) SNPs in and around the *NERD1* gene. The blue line indicates the 16 lowest ovule number accessions and the red line indicates the 16 highest ovule number accessions. SNPs are identified based on comparison to the Col-0 reference genome. The red boxes highlight low-ovule number associated SNPs (the turquoise bar indicates two SNPs in adjacent nucleotides, a non-synonymous C-A SNP and a synonymous C-G SNP). Only genomes that were available on the SALK 1,001 genomes browser (http://signal.salk.edu/atg1001/3.0/gebrowser.php) were considered. (B) One of the ovule number-associated SNPs in NERD1 causes a serine-tyrosine substitution in low ovule number accessions. (C) Average ovule number in *NERD1* "A" and "C" allele-containing accessions. The "A" allele is significantly associated with lower ovule numbers. "***" indicates statistical significance at *p* value<0.001 determined by Student's t-test.