1 Simulation-based approaches to characterize metagenome coverage as a function of sequencing effort

2 and microbial community structure

4   Taylor Royalty[a], Andrew D. Steen[a*]

5   [a]Department of Earth and Planetary Sciences, University of Tennessee – Knoxville

6   * Corresponding: asteen1@utk.edu

Abstract

We applied simulation-based approaches to characterize how microbial community structure influences the amount of sequencing effort to reconstruct metagenomes that are assembled from short read sequences. An initial analysis evaluated the quantity, completion, and contamination of complete-metagenome-assembled genome (complete-MAG) equivalents, a bioinformatic-pipeline normalized metric for MAG quantity, as a function of sequencing effort, on four preexisting sequence read datasets taken from a maize soil, an estuarine sediment, the surface ocean, and the human gut. These datasets were subsampled to varying degrees of completeness in order to simulate the effect of sequencing effort on MAG retrieval. Modeling suggested that sequencing efforts beyond what is typical in published experiments (1 to 10 Gbp) would generate diminishing returns in terms of MAG binning. A second analysis explored the theoretical relationship between sequencing effort and the proportion of available metagenomic DNA sequenced during a sequencing experiment as a function of community richness, evenness, and genome size. Simulations from this analysis demonstrated that while community richness and evenness influenced the amount of sequencing required to sequence a community metagenome to exhaustion, the effort necessary to sequence an individual genome to a target fraction of exhaustion was only dependent on the relative abundance of the corresponding organism and its genome size. A software tool, GRASE, was created to assist investigators further explore this relationship. Re-evaluation of the relationship between sequencing effort and binning success in the context of the relative abundance of genomes, as opposed to base pairs, provides a framework to design sequencing experiments based on the relative abundance of microbes in an environment rather than arbitrary levels of sequencing effort.

Background

Background

The reconstruction of high-accuracy short read sequences into metagenome-assembled genomes

(MAGs) is a recent approach to characterize microbial metabolisms within complex communities [1].

The recent creation of ~8,000 MAGs from largely uncultured organisms across the tree of life [2], the

spatial characterization of microbial metabolisms and ecology across Earth's oceans [3], and the

characterization of the potential impact that fermentation-based microbial metabolisms have on

biogeochemical cycling in subsurface sediment environments [4] provide a few examples of how

MAGs have helped to elucidate the relationships between microbial ecology, microbial metabolisms,

and biogeochemistry.

Sampling environmental microbial DNA involves selecting a target environment, sequencing

effort, bioinformatic pipeline software and parameters, metabolism characterization software (i.e., for

gene identification and similarity searches) and databases (Fig. 1). At present, there is little information

to guide how much sequencing is appropriate to achieve scientific goals in such experiments [5]. This

gap in knowledge is partly attributed to the unknown structure of target microbial communities. Further

complications arise in that the accuracy and efficiency of bioinformatic pipelines are often difficult to

characterize, and thus obscure the relationship between sequencing effort and MAG retrieval. Recent

estimates compiled by Quince et al. [5] suggest that typical metagenomic shotgun sequencing

experiments usually sequence between 1 Gbp and 10 Gbp. Researchers require more precise guidance

to select an appropriate shotgun sequencing effort in order to maximize information and minimize cost

for their specific experimental question.

Illumina sequencing technology is currently the most popular platform to generate

metagenomic shotgun sequences [5]. Previous investigators established theoretical relationships

between contig formation rate [6] and single genome coverage [7] as a function of short read

sequencing effort. On the community level, heuristic approaches have been proposed for evaluating community-level coverage to increases in sequencing effort. For example, it has been proposed to utilize short read redundancies as a function of sequencing effort to estimate community level coverage [8]. Without *a priori* knowledge of the microbial community structure, practical application of these methods to estimate MAG retrieval as a function of sequencing effort is hindered. This is particularly true as heuristic methods will only be sensitive towards highly abundant genomes.

Here we present two distinct analyses which constrain the relationship between the quantity of Illumina metagenomic shotgun sequences and the community-level sequence coverage. First, we performed *in silico* experiments to simulate the effect of sequencing effort on retrieved MAG properties for Illumina sequence datasets. For these experiments, we propose a new metric, complete-MAG equivalents, which measures community coverage in response to sequencing effort and is insensitive to splitting versus joining bins in our bioinformatic pipeline. Second, we applied a theoretical model and numerical simulations to estimate the minimum sequencing effort needed to sequence a metagenome to a target fraction of exhaustion. Our theoretical model is unique compared to previous models [6,7] in that we characterize sequencing effort in the context of all physically-distinct DNA fragments in the metagenome. This is important, as complete genome coverage does not necessarily translate into accurate read assembly due to interspecies homologous DNA and intraspecies microdiversity [9]. Coupling results from the two analyses provides a framework for investigators to define sequencing experiments in the context of selecting a rarity and fraction of exhaustion for a desired target genome when sequencing a community. The patterns presented here can be used to guide sequencing effort decisions in future sequencing projects when MAG reconstruction is a primary goal.

Results

The number of 'complete-MAG equivalents' (equation 1) as a function of high-quality bases

empirically fit the Gompertz equation (equation 2; Fig. 2b; parameters in Table 1; Additional file 1).

Complete-MAG equivalents represent the sum of MAGs retrieved, weighted by each MAG's checkM

completeness score [10], and divided by 100% (i.e., a complete MAG). This metric is designed to

highlight the total number of marker genes, used to determine MAG coverage, in an environment while

maintaining the biological relevance of MAGs. Complete-MAG equivalents reduces sensitivity to

whether bioinformatic pipelines tend to lump contigs into fewer, more-complete MAGs versus split

them into fewer, less-complete MAGs.

For each environment, the number of complete-MAG equivalents as a function of simulated

sequencing effort fit the Gompertz equation (equation 2). Here, A, μ, and λ correspond to the maximum

number of complete-MAG equivalents assembled with the pipeline, the maximum rate which

complete-MAG equivalents form with more sequencing, and the "lag bases," or the bases which must

be sequenced prior to a sufficient number of sequence reads existing to generate overlap and form

contigs [6]. The predicted maximum complete-MAG equivalents varied from ~17 in the gut

community to ~97 in the surface ocean. The predicted maximum rate that complete-MAG equivalents

increased varied from ~1.4 to ~5.8. Lastly, the minimum threshold of sequencing necessary prior to

seeing complete-MAG equivalents varied from ~0.6 to ~6.7. The *Tara Oceans* dataset, where

complete-MAG equivalents decreased at sequencing effort >20 Gbp, was an exception. For the estuary,

maize, and human gut datasets, the complete-MAG equivalents yield began to asymptote with

increasing sequencing efforts. The *Tara Oceans* dataset followed a similar pattern at <25 Gbp.

However, when the number of sequenced bases was >25Gbp, the number of complete-MAG

equivalents decreased and became insensitive to sequencing effort. The total number of MAGs

retrieved was considerably higher than the number of complete-MAG equivalents.

Mean MAG completeness also increased towards an asymptote with increasing sequencing

effort (Fig. 2c). Completeness was highest for the human gut dataset, with a maximum of 23.9%, and

increased continuously as sequencing effort increased. The mean MAG completeness reached an

asymptote of ~10-15% for the other three datasets with sequenced bases >10 Gbp. Note that when >10

Gbp were sequenced, the number of complete-MAG equivalents created still increased as new

sequences were added. For all datasets, mean MAG contamination was <2% (Fig. 2d) and did not

depend strongly on sequencing effort.

SIMULATION EXPERIMENTS

Using equation 9, we calculated the number of k-length sequence reads required to sequence all

physically-distinct DNA sequences of length, k (k-mers), in four hypothetical metagenomes, a perfectly

even, moderately uneven, highly uneven, and a lognormally-distributed structure (Fig. 3a-d). Here, we

define physically-distinct DNA as all possible k-mer combinations comprising a genome. Note that k-

mers within homologous regions are mathematically treated independent. The log-transformed

expectation value of sequencing effort (number of sequences) required to fully sequence genomes of

the hypothetical communities was linear with respect to log-transformed size of the metagenome (i.e.,

physically-distinct k-mers in the population, or unique base pairs in a metagenome); this suggests a

power-law relationship between metagenome size and expectation value of sequence reads required to

sequence the metagenome to exhaustion (Fig. 3e). For all community structures, the slope of the

relationship between sequencing effort (log-transformed sequenced reads) and metagenomic richness

(log-transformed physically-distinct number of sequenced reads) was within 1% of 1.06. The structure

of the population strongly influenced the number of reads required such that more-even community

structures required far fewer reads than less-even structures.

As equation 9 only estimates the number of reads to sequence a metagenome to exhaustion, we

used a numerical simulation to estimate the number of k-sized reads to sequence a metagenome to a

target fraction of exhaustion. The numerical simulation and equation agreed on the expected sequences

reads to sequence 100% of a given metagenome (Fig. 4); this supported the use of this simulation. The

log-transformation of both total physically-distinct k-sized reads ($|K_{MG}|$ in equation 8) and sequenced

reads showed a linear response for all target fractions and all community structures. The amount of

sequences required to achieve a given target of physically-distinct k-sized reads was variable for the

different communities shown in Fig. 3a. For instance, the lognormally-distributed community, when

compared to other communities, required the most sequencing effort to sequence a metagenome to

exhaustion but required similar sequencing effort to sequence a metagenome to a target fraction of

50%.

We applied the simulation to demonstrate the effect of community evenness on the number of

reads required to sequence a community to a target fraction of completion. These communities ranged

from perfectly even ($a=0$, equation 11) to more uneven ($a=0.02$, Fig. 5a). Evenness was quantified

using the Pielou evenness index, which expresses Shannon diversity relative to the diversity of a

perfectly even community [11]. The sequencing effort required to characterize genomes depended

strongly on both the evenness and the target fraction of completion (Fig. 5b). Again, less-even

communities required more sequence reads than more-even communities. The strength of this

relationship also depended on the target fraction of completion. A community with Pielou evenness of

0.97 required 3 orders of magnitude more sequence reads to sequence a metagenome to a target

fraction of exhaustion than a perfectly even community while the same community only required about

42% more reads to sequence 50% of the metagenome.

The minimum sequencing effort required to sequence a microbe genome given a combination

of target fraction, genome size, and fraction of the metagenome community was modeled with a

generalized additive model (GAM; Fig. 6). The GAM shows expected sequencing effort required for

microbial genome sizes 0.5, 2, 5, 10, and 20 Mbp, target genome completeness fractions from 0.5 to

1.0, and genome relative abundances from 1 to 0.0001. The smooth dimensions for target fraction,

genome size, and fraction of the metagenome community were 50, 6, and 29, respectively. To

normalize for different sequence read length, sequence reads were converted to bases and ranged from

1×10^{7} to 1×10^{15} total bases. More bases were required to sequence microorganisms 1) when the

genome was relatively rarer in the community, 2) to achieve better coverage of the genome, and 3)

when average genome sizes were larger.

Using the relationship shown in Fig. 6, we can convert sequencing effort to an abundance

cutoff. Fig. 7 shows complete-MAG equivalents retrieved for a genome of a given level of abundance

that is sequenced to a target depth for the human gut, maize soil, estuarine sediment, and the surface

ocean microbiomes analyzed earlier. Correlation coefficients for the regressions used to relate log-

transformed sequencing effort (base pairs) to genome relative abundance were $R=1$ for all three

genome sizes evaluated (1 Mbp, 5 Mbp, and 20 Mbp). Evaluation of 1 Mbp and 20 Mbp define the

range of uncertainty in predicting complete-MAG equivalents as the true size of genomes is unknown.

Unlike the asymptotic response of complete-MAG equivalents to sequencing effort (Fig. 2b), complete-

MAG equivalents increase exponentially as the abundance cutoff decreases (note that the abundance

cutoff is on a log-scale). The target genome completeness fraction was held at a constant of 0.5 for all

regressions and the sensitivity of complete-MAG equivalents will change with respect to the genome

abundance cutoff with different values of target genome completeness fraction.

Discussion

We sought to establish evidence-based guidelines for selecting a sequencing effort during

shotgun metagenomic sequencing experiments. Random subsamples of existing short read datasets

(individually sampled, assembled, and binned) simulated the effect of creating MAGs from datasets of

different sizes and environments. The datasets analyzed here are representative of both the sequencing

effort (1 to 10 Gbp) [5] and the types of target environments that microbial ecologists often investigate

[12]. A variety of software is available for all steps of MAG creation pipelines, and the quantity/quality

of MAGs will depend on software selection, software configuration, and sequenced environment (Fig.

1) [5]. Furthermore, it is best-practice to manually curate algorithmically-created MAG bins [13]. We

do not argue that the pipeline used here is objectively optimal for generating high-quality (high

completeness and low contamination) MAGs. Rather, our pipeline was configured to minimize

contamination (Fig. 2D) associated with retrieved MAGs at the expense of reduced completeness.

Thus, MAG quantity was not directly reported but expressed as complete-MAG equivalents (equation

1). Reporting complete-MAG equivalents reduces bias associated with binning software.

As sequencing effort increased, there was an initial lag in complete-MAG equivalents followed

by a rapid increase in complete-MAG equivalents, and then diminishing returns at higher sequencing

efforts. Previous investigators modeled the response of 16S RNA gene [14–16], Hill's number diversity

[17], taxon-resolved abundance [18], and gene abundance [18] as a function of sequencing effort with

rarefaction curves, or collector's curves. The complete-MAG equivalents as a function of sequencing

effort did not match a traditional collector's curve, which lacks an initial lag. Rather, the data appear

sigmoidal. We modeled the data using the Gompertz function (equation 9), because its parameters can

be interpreted in terms of quantities that are familiar from microbial growth curves (lag time, growth

rate, and maximum density) [19]. The three Gompertz parameters, A, μ, and λ are interpreted as the

maximum possible number of complete-MAG equivalents at infinite sequencing effort, maximum rate

that complete-MAG equivalents increased with increasing sequencing effort, and a minimum

sequencing threshold prior to rapid complete-MAG equivalents retrieval (Fig. 2a), respectively.

Consequently, the Gompertz equation fit parameters may provide means for inter-comparisons between

different metagenomic datasets. We want to emphasize that the datasets analyzed here do not

necessarily reflect the generalized characteristics of their parent communities. More so, these

communities were analyzed as their community properties (richness, abundances, and phylogeny) are

generally different [4,20–22].

The fits to the Gompertz function illustrate that there is an optimal sequencing effort for MAG

creation efforts corresponding to the upper shoulder of the Gompertz curve ("late log phase"). When

sequencing effort is too close to λ, MAGs bin poorly; when sequencing effort is too great, the number

and quality of MAGs per unit sequencing effort (and therefore cost) is low. We speculate that our

choice of pipeline, and specifically the fact that we discarded contigs <3kb, caused poor performance at

higher sequencing effort for the *Tara Oceans* dataset. Species-level microdiversity and interspecies

homologous DNA can cause "bubbles" which impair assembly in larger data sets [9,23]. Improved

assembly would likely have yielded more complete-MAG equivalents for our assembly of the largest

subsets of the *Tara Oceans* data.

As mean MAG completeness converged to an asymptote well below 100% (Fig. 2b), complete-

MAG equivalent yields, or the ratio of the maximum observed complete-MAG equivalents to the

theoretical maximum complete-MAG equivalents (*A*) (Table 1) were close to 100%. The asymptote in

mean MAG completeness (Fig. 2c) and negligible values of mean MAG contamination (Fig. 2d)

demonstrates that the retrieved complete-MAG equivalents at higher sequencing efforts is not

attributed to artifacts of assembly. If the signal in complete-MAG equivalents was attributed to

assembly of false positive marker genes, then either 1) the mean MAG completeness would decrease

with the addition of false positive marker genes which fail to cluster (i.e., more MAGs with low

completeness) or 2) false positive signal would cluster with preexisting MAGs leading to an increase in

mean MAG contamination. Neither of these scenarios occurred. The fact that mean MAG completeness

was at an asymptote <<100% before complete-MAG equivalents began converging suggests more

sequenceable DNA exists in these communities. Thus, we asked how much sequencing was necessary

to sequence a community to exhaustion.

The theoretical probability model (equation 9) demonstrated that the sequencing effort to

sequence a metagenome to exhaustion was predictable, regardless of community structure (Fig. 3).

Furthermore, a simplistic characterization community complexity demonstrates an inherent truth, less

complex communities require less sequencing to sequence all available DNA. This corroborates our

observations where less diverse communities (i.e., human gut) relative to the environmental

communities had greater complete-MAG response to sequencing effort (Fig. 2b). A limitation to

equation 9 is predicting sequencing effort to sequence a metagenome to a fraction of exhaustion. This

is particularly important as 100% genome coverage is not necessary to obtain robust assemblages and

MAGs. As such, a numerical simulation was used for predicting the sequencing effort necessary to

sequence an individual metagenome to a fraction of exhaustion. The numerical simulation agreed with

the theoretical model when predicting the sequencing effort to sequence a metagenome to exhaustion

(Fig. 4). A last analysis explored the semi-quantitative relationship between community evenness and

richness and the necessary sequencing effort to achieve a target fraction of exhaustion for a

metagenome (Fig. 5).

In practice, information about a target community structure may not be available for estimating

sequencing effort. The GAM built here predicts the minimum number of sequences necessary to

sequence a given fraction of a target genome as a function of average genome size and the relative

abundance of the target genome in the community (Fig. 6). Even without knowledge of a target

community's structure, the GAM provides a useful constraint for designing whole-genome shotgun

metagenomic sequencing experiments. Note that the genome sizes selected for this analysis correspond

with typical sizes reported for prokaryotic genomes [12]. Virus and eukaryotic genomes were not

considered, as the primary focus of this study is prokaryotic systems. Contamination from eukaryotic

and virus genomes is usually accounted for using size-fractionation. Nonetheless, the underlying theory

applied here should be applicable to these other genomic systems.

Metagenomic shotgun sequencing experimental designs should be rationally designed such that

sequencing effort is selected to capture a desired fraction of a target microbial genome. Investigators

should be cognizant of the rarest microbial genome desired to be characterized as well as the degree of

characterization of that microbial metagenome when designing a sequencing experiment. To that end,

we have built a tool, `Genome Relative Abundance to Sequencing Effort (GRASE),`

to report estimated sequencing effort required to capture a defined fraction of a genome as a function of

the relative abundance of the corresponding microorganism in the community and average genome

size. This R-based GUI app can be accessed online at http://adsteen.shinyapps.io/grase and is archived

at http://github.com/adsteen/GRASE, from which it can be downloaded and run locally.

When the sequence read datasets analyzed here (Table 2; Fig. 2) are re-evaluated in the context

of the relative abundance of a microbial metagenome (g_{MG}) sequenced to a target fraction (0.5),

complete-MAG equivalents increase appreciably in response to minor increases in deeper

characterization of the community metagenome (Fig. 7). This observation contrasts the complete-MAG

equivalents response to sequencing effort (in base pairs), where substantial increases in sequencing

effort (by contemporary standards) leads to diminishing returns in complete-MAG equivalents. It is

important to note that abundance cutoff and sequencing effort are interchangeable; however, complete-

MAG equivalents response to changes in the respective predictor (i.e., base pairs versus abundance

cutoff) alters the optics of the collector's curve. Modest increases in sequencing effort contribute minor

amounts to extending abundance cutoff. A substantial amount of genomic and metabolic data can be

gained from targeting rarer microbes (metagenomic abundances <0.005), with the caveat that whole-

genome shotgun sequencing technology (as well as computational power) requires significant increases

in either the number or length of reads generated per run.

Materials and Methods

SEQUENCE DATA SOURCES

All sequence data were downloaded from NCBI's Sequence Read Archive (SRA) using the

SRA Toolkit (fastq-dump –split-files) [24]. Exact duplicate reads for both forward and reverse reads

were removed using PRINSEQ (-derep 1; v0.20.4) [25]. All sequencing datasets were limited to

Illumina shotgun metagenomic paired-end reads. Four datasets were analyzed for this analysis. The

first dataset was from oceanic surface water collected at 5m depth in the Caribbean Sea as a part of the

Tara Oceans expedition [20]. The second dataset was from sediment from a depth of 8-10 cm below

the surface (sulfate-rich zone) and collected at the White Oak River Estuary, Station H, North Carolina,

USA [4]. The third dataset was collected from maize soil [21]. The last dataset was collected from

human fecal samples and represented a human gut microbiome [22]. All datasets analyzed in this study

are summarized in Table 1.

MAG ASSEMBLY PIPELINE

The pipeline developed here followed similar pipelines described by other authors [3,26]. All

sequence datasets were analyzed as follows. Trimmomatic (v0.36) [27] removed adapters and trimmed

low-quality bases from individual reads. Read leading and trailing quality scores were required to be

>3. The sliding window was set to 4 base pairs and filtered base pair windows with a mean score <15.

Quality controlled reads were assembled into contigs using MEGAHIT (v1.1.2; --presets meta-large)

[28]. Due to computational limitations, assembled contigs <3000 bp long were excluded from the

analysis. Redundant contigs were removed using CD-HIT (v4.6.8; cd-hi-est -c 0.99 -n 10) [29].

Remaining contigs with high intra-contig similarity based on intra-contig sequence alignments

(Minimus2 -D OVERLAP=100 MINID=95) were removed. The quality-controlled reads (i.e., after

using Trimmomatic) were mapped to the remaining contigs using Bowtie 2 (v2.3.3) [30] to generate a

coverage score for individual contigs.

Resultant contigs were iteratively clustered into MAGs using the unsupervised, clustering

algorithm Binsanity (v0.2.6) [26]. Similar to Tully et al. [3], six initial clustering iterations were

performed with the parameter, *preference* (-p), set to -10 (iteration 1), -5 (iteration 2), -3 (iteration 3-6).

Between iterations, a refinement step (Binsanity-refine) was performed on the putative MAGs with

constant *preference* (-p) of -25. The refined putative MAGs were evaluated for contamination and

completeness using the software CheckM (v1.0.6) [10], which uses HMMER (v3.1) and Prodigal

(v2.6.3) [31]. For this work, we used the recommended lineage-specific marker sets. The lineage-

specific workflow defines marker genes as those appearing in >97% of a single lineage. Contigs

associated with putative MAGs meeting one of the following criteria: 1) had a completeness > 90% and

contamination < 10%, 2) had a completeness > 80% and contamination < 5%, or 3) had a completeness

> 50% and contamination < 5% were treated as high-quality. All other MAGs were considered low-

quality MAGs. MAGs defined as high-quality were not modified any further. Contigs associated with

the high-quality MAGs were not used in the subsequent reclustering and refinement steps. The contigs

associated with low-quality MAGs were pooled together and reclustered during the next iteration of

Binsanity clustering. After the sixth iteration, the remaining MAGs which did not fall into one of the

three categories underwent additional refinement using Binsanity-refine. During this step, MAGs were

iteratively refined with *preference* set to -10 (iteration 1), -3 (iteration 2), and -1 (iteration 3). Between

each refinement step, metrics of contamination and completeness were evaluated using CheckM.

Again, MAGs meeting the criteria of a high-quality category described above were not further

modified. The respective contigs associated with putative MAGs were not used in proceeding

refinement steps. After the last iteration of refinement, all MAGs were reevaluated for completeness,

contamination, and a taxonomic rank using CheckM. Completeness values for MAGs with a minimum

resolved taxonomic rank of phylum were summed together. Complete-MAG equivalents, g_{eq}, were

defined as:

$g_{eq} = \sum_{i=1}^{n} f_i$ (1

where n is the total number of retrieved MAGs and f_i is the corresponding CheckM completeness (%).

Common practice when reporting MAGs involves discarding MAGs not meeting a criterion of

completeness and contamination. In the context of characterizing MAG response to sequencing effort a

MAG pipelines often fails at generating high-quality MAGs. One must discern whether this is a failure

of the bioinformatic pipeline versus insufficient sequencing effort (Fig. 1). We argue that MAGs with

low completeness still provide information regarding the community level coverage when generating

collector's curves. More so, a community has finite number of unique marker genes, and an infinite

sequencing effort would result in sequencing all unique marker genes, regardless of whether or not

these genes cluster to one another.

SUBSAMPLING SEQUENCE READ DATASETS

The effect of decreased sequencing effort was simulated by subsampling the initial sequence read

datasets described above. Downloaded sequence read datasets were randomly sampled at set fractions

of 1%, 10%, 20%, 40%, 60%, 80%, 90%, 95%, and 100%. Each fraction was resampled, assembled,

and binned in triplicate. Each triplicate assembly was binned independently using the MAG assembly

pipeline described above.

MODELING MAG RESPONSE TO SEQUENCING EFFORT

Complete-MAG equivalents (equation 1) as a function of sequencing effort was modeled for

environmental sequence datasets using the Gompertz equation, as reformulated by Zweitering et al.

[19] for use with microbial growth curves:

$$g_{eq}(A, \mu, \lambda, b) = A \times e^{-e^{\frac{\mu \times e}{A}(\lambda - b) + 1}} \tag{2}$$

where A, μ, and λ are fit coefficients and b is high-quality bases (Gbp). From equation 2, MAG yield

could be defined as:

$$MAG\ Yield = \frac{g_{eq,max}}{A} \tag{3}$$

where $g_{eq,max}$ is the largest complete-MAG equivalent derived from the subsampling experiment and

A is from equation 2.

DEFINING THE MICROBIAL METAGENOME AND SEQUENCING PROBABILITY

Here we draw on set theory to provide a theoretical grounding for our *in silico* simulations described

below. The expected number of sequences to sequence a fraction of an individual microbe's genome

can be modeled with probability theory by defining a community metagenome with set theory. Fig. 8a-

e provides a cartoon illustrating the application of this set theory on a hypothetical microbial

population, G. G contains unique genomes (g) with finite abundances (n). The definition of microbial

species is somewhat contentious [32]. Here we define g as the average genome (length and

composition) for all individuals genomes associated with the rank, species, based on an arbitrary

definition. As such, the species richness (s or unique g) of G depends on the definition of g. In the

example community (Fig. 8a-e), s=6 and n=13. Thus, G is represented as (Fig. 8a):

$$G = \{n_1 g_1, n_2 g_2 \dots n_s g_s | n \in \mathbb{N}\} \tag{4}$$

where s is the species richness. When characterizing G via shotgun metagenomics, the i^{th} genome, g_i,

can be sequenced at K unique sections given a characteristic read length, k, and average genome size, l,

in number of base pairs (Fig. 8b). Thus, the number of physically-distinct k-sized reads, K, associated

with the i^{th} genome, g_i, within G is equal to:

$$K_{g_i} = l(g_i) - k + 1 \tag{5}$$

Note that equation 5 considers homologous DNA as unique fragments. From equation 5, the

metagenome, g_{MG}, for g_i is defined as the set of all physically-distinct k-sized reads (Fig. 8c) or:

$$g_{MG,i} = \{g_{i,1,1+k}, g_{i,2,2+k} \dots, g_{i,Kg_i,Kg_i+k}\} \tag{6}$$

where the subscripts for g_i represent a given k-sized read spanning from an arbitrary starting base pair

to the arbitrary starting base pair plus k. By substituting $g_{MG,i}$ into all g for equation 4 (Fig. 8d), the

metagenome for a microbial community, G_{MG}, is derived to be:

$$G_{MG} = \{n_1 g_{MG,1}, n_2 g_{MG,2} \dots n_s g_{MG,s} | n \in \mathbb{N}\} \tag{7}$$

while the population of physically-distinct k-sized reads in the metagenome, G_{MG} (Fig. 8e), is

represented as:

$$K_{MG} = \{g_{MG,1}, g_{MG,2} \dots g_{MG,s}\} \tag{8}$$

From equation 8, one can determine the cardinality, or the total number, of physically-distinct

k-sized reads associated with G_{MG} (expressed as $|K_{MG}|$). To an effect, $|K_{MG}|$ is analogous with

"metagenomic richness" of an environment. When attempting to fully sequence G_{MG} using shotgun

metagenomics, we assume that sampling events (sequence reads) are independent and are sampled with

replacement. In fact, Illumina sequencing technology sequences reads in parallel via the individual

DNA fragments binding to individual clusters. Furthermore, the fragmented DNA cannot be sequenced

twice as the sequencing process is destructive [33]. Nonetheless, the mass of DNA extracted from a

target environment represents a negligible fraction of the total DNA existing in that environment. As

the relative abundance of the k-sized reads in K_{MG} does not change when DNA is extracted from an

environment, sampling events can be treated as independent and thus, DNA sampling reduces to

sampling with replacement. If the proportion DNA mass extracted had a significant impact on the

remaining mass of DNA in the environment, then one would be more suited to sequence all the DNA

versus a smaller proportion of the DNA. The sequencer should have no impact on sampling assuming

no sequencing errors due to misreading or spatial sampling issues (i.e., clonal density issues). These

issues do exist, but for the sake of a first order approximation, these biases can be ignored.

The probability of sequencing all elements in G_{MG} reduces to a coupon collectors problem [34]

by making the above assumptions. Using the general functional form for calculating expected samples

for sampling all unique elements in a set (equation 13b in [8]), one can predict the number of sequences

necessary to sequence all elements in K_{MG}, such that the expected number of sequences, $\mathrm{E}(G_{MG})$, is:

$$\mathrm{E}(G_{MG}) = \int_0^\infty \left(1 - \prod_{j \in K_{MG}}(1 - e^{-p_j t})\right) dt \qquad (9$$

where j is a given element within K_{MG}, t is the number of sampling events, and p_j is equal to the

proportion of the j^{th} k-sized read within a given population of k-sized reads. p_j can be expressed as

follows:

$$p_j = \frac{n_i \times j \in K_{MG}}{|G_{MG}|} \qquad (10$$

where n_i is the respective abundance for the species whose MAG contains the j^{th} k-sized read within

K_{MG}, and $|G_{MG}|$ is the cardinality of G_{MG}, or the total number of k-sized reads in the metagenome, G_{MG}.

MODELING EXPECTED SEQUENCES

Equation 9 provides an estimate for the total number of sequences to sequence all K_{MG}. The

influence of increasing species richness (i.e., s in equation 4) on the expected number of sequences was

tested for four hypothetical communities. The first community had an even structure such that all

physically-distinct DNA fragments were equally distributed across all K_{MG}. In the second community,

90% of the physically-distinct DNA fragments were equally distributed in 50% of K_{MG}, and the remaining

10% of the physically-distinct DNA fragments were distributed equally across the remaining 50% of

K_{MG}. This community represented a community with relatively moderate species evenness. In the third

community, 90% of the physically-distinct DNA fragments were equally distributed across 10% of K_{MG},

and the remaining 10% of the physically-distinct DNA fragments were distributed equally across the

remaining 90% of K_{MG}. This community represented a community with relatively low species evenness.

The last community had 10 equally-sized groups. The abundance of the physically-distinct DNA

fragments in each group was based on the function form of a lognormal community [35] which has been

observed in microbial populations (e.g., [17,36]), such that:

$$S(R) = S_0 e^{-a^2 R^2}$$ (11

where S_0 was treated as the maximum relative of abundance ($S_0 = 1$), a was the inverse width of the

distribution, R was treated as the positive octave range spanning 0 to 9, and $S(R)$ represented the

abundance for a given group. For the lognormal abundance distribution in Fig. 3d, a was set to a value

of 0.2. Each hypothetical community started with $|K_{MG}| = 1 \times 10^6$. $|K_{MG}|$ incrementally increased at 10

equally-spaced, linear steps to a maximum of $|K_{MG}| = 1 \times 10^8$. As $|K_{MG}|$ increased, all community

structures remained constant. Graphical representation of rank abundance in Fig. 3a-d was normalized

by a given $|K_{MG}|$ to reflect that populations retained the same structure even as population size varied.

We defined a normalized rank abundance r_n such that

$$r_n = \frac{r}{s}$$ (12

where r and s are untransformed rank abundance and richness, respectively. For each community, at

each step, the expected number of sequences was calculated using equation 9. The expected number of

sequences as a function of $|K_{MG}|$ were modeled with linear regressions.

Equation 9 gives the expected number of sequences required to sequence any sized community

to exhaustion. Numerical sequencing simulations were performed to determine the number of

sequences necessary to sequence a subset of all physically-distinct DNA (K_{MG}). These numerical

sequencing simulations were applied to four hypothetical community structures described above.

Numerical simulations were performed such that $|K_{MG}| = 3 \times 10^7, \ 4 \times 10^7, 5 \times 10^7, 7 \times 10^7, 9 \times$

10^7, and 1×10^8. During each of these simulations, the parameters read length (k) and average

genome size (l) were set to 100 and 1×10^6, respectively, for all g. Random elements from K_{MG} were

selected with replacement to simulate sequencing events. Numerical simulations were performed until

the percentage of $|K_{MG}|$ sequenced was 50%, 70%, 90%, 95%, 99%, or 100%. A weight distribution

was applied to elements in a given K_{MG}. The weight distribution biased sequencing to reflect the

relative abundances of the four hypothetical communities described above. The percentage of $|K_{MG}|$

sequenced was evaluated every 1×10^7 sequences. Numerical simulations were performed in triplicate

for all $|K_{MG}|$ and all target fractions of $|K_{MG}|$.

428   We explored the influence of community evenness on required sequencing effort by performing

sequencing simulations on 6 different lognormally-distributed communities. The numerical sequencing

simulations followed the simulations described above. The 6 lognormal communities were modeled

such that each community had $S_0 =1$, $R=10$, and $|K_{MG}| = 1 \times 10^7$. Values of a for the 6 lognormal

distributions were $a=0$, $a=0.005$, $a=0.008$, $a=0.01$, $a=0.015$, and $a=0.02$. Evenness was represented

using the Pielou evenness index [11], or the ratio of the Shannon diversity index [37] for an observed

community to an even community of equal richness. Shannon diversity was calculated in the context of

a metagenomes such that:

$H_{MG} = \sum_{j \in K_{MG}} -p_j log(p_j)$ (13

where p_j is the proportion that the j^{th} k-sized read represents among all physically-distinct DNA sequences

in the metagenome. Thus, the Pielou evenness index [11] was calculated such that:

$J = \dfrac{H_{MG}'}{H_{MG,max}}$ (14

where J was the Pielou evenness index, H_{MG}' was the metagenome Shannon diversity index, and

$H_{MG,max}$ represented the metagenome Shannon diversity index when all p_j were equal ($a=0$).

Lastly, numerical simulations were performed to determine the sequencing effort necessary to

achieve a target fraction for an individual metagenome (g_{MG}). Target fractions increased from 0.5 to 1

at 100 linearly-spaced intervals. The fraction of the metagenome community (G_{MG}) that g_{MG}

represented varied from 1% to 100% in 30 lognormally-spaced intervals. The target genome sizes

varied such that $l=0.5\times10^6$, $l=1\times10^6$, $l=2\times10^6$, $l=3\times10^6$, $l=5\times10^6$, $l=10\times10^6$, $l=15\times10^6$, and $l=20\times10^6$.

The sequencing effort for a given combination of target fraction, genome size, and fraction of the

metagenome community was modeled using the GAM function (mgcv R package; [38]). For modeling

purposes, target fraction was raised to the 6[th] power and both genome size and sequences were log-

transformed. The number of smooth dimensions for fraction of community, genome size, and target

fraction were heuristically varied till the resulting fit demonstrated residuals with a normal distribution.

Note that the objective here was not build a predictive model but simply a first order approximation for

simulations performed here. The GAM was extrapolated for genome relative abundances spanning 0.01

to 0.0001 at all sizes and target fractions. The extrapolation was necessary due to the computational

limitation of analyzing genome relative abundances <0.01.

RELATING MAG RESPONSE TO THE THEORETICAL SEQUENCING MODEL

Sequencing effort (in base pairs) used for predicting complete-MAG equivalents for the sequence read

datasets maize soil, an estuarine sediment, the surface ocean, and the human gut, were related to a

genome relative abundance utilizing the GAM presented in the previous section. This was

accomplished by setting a constant genome size and target fraction (0.5) and performing a linear

regression between genome relative abundance (a genome's fraction of a community) and log-

transformed base pairs. The linear regression was performed with genome sizes of 1 Mbp, 5 Mbp, and

20 Mbp. Complete-MAG equivalents were predicted as a function of genome relative abundance

sequenced to a target fraction of 0.5.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing Interests

The authors declare that we have no competing interests.

Funding

This research was supported by National Science Foundation grant OCE-1431598 and a C-DEBI subaward. This is C-DEBI contribution number (contribution number to be determined).

Authors' Contributions

TMR and ADS both conceived the study, performed data analyzed, interpreted results and wrote the manuscript.

Data Availability

All simulations and codes used for modeling sequencing effort are freely available on Github at: https://github.com/taylorroyalty/sequence_simulation_code. All data generated during the subsampling experiment is available in Additional file 1. The GRASE GUI application is available at: http://adsteen.shinyapps.io/grase.

Acknowledgements

We also would like to thank Zachery Stooksbury for last minute assistance with editing figures.

485   **References**

[1]   Knight R, Vrbanac A, Taylor BC, Aksenov A, Callewaert C, Debelius J, et al. Best practices for analysing microbiomes. Nat Rev Microbiol 2018. doi:10.1038/s41579-018-0029-9.

[2]   Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, et al. Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. Nat Microbiol 2017;903:1–10. doi:10.1038/s41564-017-0012-7.

[3]   Tully BJ, Graham ED, Heidelberg JF. The reconstruction of 2,631 draft metagenome-assembled genomes from the global oceans. Sci Data 2018;5:1–8. doi:10.1038/sdata.2017.203.

[4]   Baker BJ, Lazar CS, Teske AP, Dick GJ. Genomic resolution of linkages in carbon, nitrogen, and sulfur cycling among widespread estuary sediment bacteria. Microbiome 2015;3:14. doi:10.1186/s40168-015-0077-6.

[5]   Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. Shotgun metagenomics, from sampling to analysis. Nat Biotechnol 2017;35:833–44. doi:10.1038/nbt.3935.

[6]   Lander ES, Waterman MS. Genomic mapping by end-characterized random clones: a mathematical analysis. Genomics 1988:231–9. doi:10.1016/0888-7543(95)80086-2.

[7]   Wendl MC, Kota K, Weinstock GM, Mitreva M. Coverage theories for metagenomic DNA sequencing based on a generalization of Stevens' theorem. J Math Biol 2013;67:1141–61. doi:10.1007/s00285-012-0586-x.

[8]   Rodriguez-R LM, Konstantinidis KT. Estimating coverage in metagenomic data sets and why it matters. ISME J 2014;8:2349–51. doi:10.1038/ismej.2014.76.

[9]   Ghurye JS, Cepeda-Espinoza V, Pop M. Metagenomic assembly: Overview, challenges and applications. Yale J Biol Med 2016;89:353–62. doi:10.1016/j.joca.2013.10.019.

[10]   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res 2015;25:1043–55. doi:10.1101/gr.186072.114.

[11]   Pielou EC. The measurement of diversity in different types of biological collections. J Theor Biol 1966;13:131–44. doi:10.1016/0022-5193(66)90013-0.

[12]   Land M, Hauser L, Jun S-R, Nookaew I, Leuze MR, Ahn T-H, et al. Insights from 20 years of bacterial genome sequencing. Funct Integr Genomics 2015;15:141–61. doi:10.1007/s10142-015-0433-4.

[13]   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al. Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. Nat Biotechnol 2017;35:725–31. doi:10.1038/nbt.3893.

[14]   Roesch LFW, Fulthorpe RR, Riva A, Casella G, Hadwin AKM, Kent AD, et al. Pyrosequencing enumerates and contrasts soil microbial diversity. ISME J 2007;1:283–90. doi:10.1038/ismej.2007.53.

[15]   Feng BW, Li XR, Wang JH, Hu ZY, Meng H, Xiang LY, et al. Bacterial diversity of water and sediment in the Changjiang estuary and coastal area of the East China Sea. FEMS Microbiol Ecol 2009;70:236–48. doi:10.1111/j.1574-6941.2009.00772.x.

[16]   Rintala A, Pietilä S, Munukka E, Eerola E, Pursiheimo JP, Laiho A, et al. Gut microbiota analysis results are highly dependent on the 16s rRNA gene target region, whereas the impact of DNA extraction is minor. J Biomol Tech 2017;28:19–30. doi:10.7171/jbt.17-2801-003.

[17]   Kang S, Rodrigues JLM, Ng JP, Gentry TJ. Hill number as a bacterial diversity measure framework with high-throughput sequence data. Sci Rep 2016;6:1–4. doi:10.1038/srep38263.

[18]   Zaheer R, Noyes N, Ortega Polo R, Cook SR, Marinier E, Van Domselaar G, et al. Impact of

sequencing depth on the characterization of the microbiome and resistome. Sci Rep 2018;8:5890. doi:10.1038/s41598-018-24280-8.

[19]   Zwietering MH, Jongenburger I, Rombouts FM, Van't Riet K. Modeling of the bacterial growth curve. Appl Environ Microbiol 1990;56:1875–81. doi:10.1111/j.1472-765X.2008.02537.x.

[20]   Karsenti E, Acinas SG, Bork P, Bowler C, de Vargas C, Raes J, et al. A holistic approach to marine Eco-systems biology. PLoS Biol 2011;9:7–11. doi:10.1371/journal.pbio.1001177.

[21]   Howe AC, Jansson JK, Malfatti SA, Tringe SG, Tiedje JM, Brown CT. Tackling soil diversity with the assembly of large, complex metagenomes. Proc Natl Acad Sci 2014;111:4904–9. doi:doi: 10.1073/pnas.1402564111.

[22]   Schirmer M, Smeekens SP, Vlamakis H, Jaeger M, Oosting M, Franzosa EA, et al. Linking the Human Gut Microbiome to Inflammatory Cytokine Production Capacity. Cell 2016;167:1125–1136.e8. doi:10.1016/j.cell.2016.10.020.

[23]   Lonardi S, Mirebrahim H, Wanamaker S, Alpert M, Ciardo G, Duma D, et al. When less is more: "Slicing" sequencing data improves read decoding accuracy and de novo assembly quality. Bioinformatics 2015;31:2972–80. doi:10.1093/bioinformatics/btv311.

[24]   Leinonen R, Sugawara H, Shumway M. The Sequence Read Archive. Nucleic Acids Res 2010;39:2010–2. doi:10.1093/nar/gkq1019.

[25]   Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. Bioinformatics 2011;27:863–4. doi:10.1093/bioinformatics/btr026.

[26]   Graham ED, Heidelberg JF, Tully BJ. BinSanity: unsupervised clustering of environmental microbial assemblies using coverage and affinity propagation. PeerJ 2017;5:e3035. doi:10.7717/peerj.3035.

[27]   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics 2014;30:2114–20. doi:10.1093/bioinformatics/btu170.

[28] Li D, Luo R, Liu C-M, Ting H-F, Sadakane K, Yamashita H, et al. MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies and community practices. Methods 2016;102:3–11. doi:10.1016/J.YMETH.2016.02.020.

[29] Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics 2012;28:3150–2. doi:10.1093/bioinformatics/bts565.

[30] Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods 2012;9:357–9. doi:10.1038/nmeth.1923.

[31] Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics 2010;11:119. doi:10.1186/1471-2105-11-119.

[32] Rosselló-Móra R, Amann R. Past and future species definitions for Bacteria and Archaea. Syst Appl Microbiol 2015;38:209–16. doi:10.1016/j.syapm.2015.02.001.

[33] Bentley DR, Balasubramanian S, Swerdlow HP, Smith GP, Milton J, Brown CG, et al. Accurate whole human genome sequencing using reversible terminator chemistry. Nature 2008;456:53–9. doi:10.1038/nature07517.

[34] Flajolet P, Gardy D, Thimonier L. Birthday paradox, coupon collectors, caching algorithms and self-organizing search. Discret Appl Math 1992;39:207–29. doi:10.1016/0166-218X(92)90177-C.

[35] Magurran AE. Ecological Diversity and Its Measurement. 1st ed. Croom Helm Ltd; 1988. doi:10.1007/978-94-015-7358-0.

[36] Galand PE, Casamayor EO, Kirchman DL, Lovejoy C. Ecology of the rare microbial biosphere of the Arctic Ocean. Proc Natl Acad Sci 2009;106:22427–32. doi:10.1073/pnas.0908284106.

[37]   Shannon CE. A mathematical theory of communication. Bell Syst Tech J 1948;27:379–423. doi:10.1145/584091.584093.

[38]   Wood S. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML Smoothness Estimation 2017.

Tables

Table 1. Estimates of fit coefficients for the Gompertz equation (equation 3) for the complete-MAG

equivalents as a function of sequencing depth in published datasets from ocean surface water, estuarine

sediment, maize soil, and the human gut. *p* values for all coefficients were <<0.05.

Environment	A (±SE)	μ (±SE)	λ (±SE)	Yield*
Ocean Surface Water	97.67 (4.15)	5.84 (0.31)	1.16 (0.33)	0.88
Estuary Sediment	26.25 (2.11)	1.63 (0.06)	3.70 (0.24)	0.86
Maize Soil	43.65 (1.98)	1.43 (0.07)	6.13 (0.60)	0.70
Human Gut	17.49 (1.02)	5.01 (0.46)	0.67 (0.14)	0.90

*Equation 3

Table 2. Summary of sequence datasets analyzed with the MAG pipeline.

Environment	NCBI SRA Accession	Sequencing Platform	Total Reads*	High Quality Bases*	General Notes	Citation
Ocean Surface Water	ERR599029	Illumina HiSeq 2000	3.372×10^8	3.340×10^{10}	Caribbean Sea (5 mbsl)	[20]
Estuary Sediment	SRR5248164	Illumina HiSeq 2000	1.137×10^8	1.589×10^{10}	Sulfate Zone (8-10 cmbsf)	[4]
Maize Soil	SRR351473	Illumina HiSeq 2000	4.727×10^8	3.824×10^{10}	Surface Soil	[21]
Human Gut	SRR5127631	Illumina HiSeq 2000	5.095×10^7	4.847×10^9	--	[22]

* Combined forward and reverse paired-end reads.

Figure Lengeds

Fig. 1. A flow diagram illustrating the workflow for sequencing experiments.

Fig. 2. The influence that the parameters A, μ, and λ had on the Gompertz equation (a). The property of the Gompertz equation that each parameter influences is colored red. Complete-MAG equivalents (in arbitrary units) (b), mean MAG completeness (c), mean MAG contamination (d) as a function of simulated sequencing effort (Gbp) for sequence datasets of the human gut, maize soil, estuarian sediment, and surface ocean microbiomes, using the pipeline described in the methods section. Translucent lines in (b) correspond to nonlinear least squares fits of the Gompertz equation to the respective environmental dataset.

Fig. 3. Sequencing effort, with units of log of sequence reads, required to fully sequence four different community structures, one with relatively high community evenness (a), relatively moderate community evenness (b), relatively low community evenness (c), and one with a lognormal community structure (d), were predicted using linear regressions (e) and the log of $|K_{MG}|$ from equation 9 as a predictor.

Fig. 4. Sequencing effort, with units of log sequence reads, necessary to reach variable target sequencing depths (colors) for four different community structures, one with relatively high community evenness (a), relatively moderate community evenness (b), relatively low community evenness (c), and one with a lognormal community structure (d). Red translucent lines correspond with linear regression curves for the respective community in Fig. 3e.

Fig. 5. Numerical sequencing simulations applied to 6 hypothetical communities with different lognormal distributions that were defined by the parameter, *a*, from equation 10 (a). The sequencing effort, with units of log of sequence reads, necessary to sequence a target fraction of $|K_{MG}|$ (dashed contours) as a function of the Pielou evenness index, *J*, for a given lognormal community structure (b).

Fig. 6. Numerical sequencing simulations show the number of bases (color bar) required to sequence a target fraction of a genome as a function of that genome's relative abundance in the community metagenome. Genomes evaluated were 0.5×10^6 (a), 2×10^6 (b), 5×10^6 (c), 10×10^6 (d), and 20×10^6 (e) base pairs long.

Fig. 7. Complete-MAG equivalents as a function of a target genome's relative abundance when sequenced to a target fraction (0.5) for human gut, maize soil, estuarian sediment, and surface ocean sequence datasets. Sequencing effort was converted to genome relative abundance sequenced to target fraction using the GAM presented in Fig. 6. The translucent shaded areas correspond to uncertainty from the target genome size (1 Mbp, or the lower-bound, to 20 Mbp, or the upper-bound) while the solid lines correspond to genome sizes 5 Mbp.

Fig. 8. A cartoon illustrating an example microbial community (G), the metagenomes for genomes ($g_{MG,i}$) (as defined in equation 5) within G, and the overall metagenome for the given microbial community (G_{MG}). In this example, there are 6 genomes ($s=6$) and a total of 13 individual microbes. (a) Black circles represent individual microbes whose genomes are averaged together, g. The average genome, g, are indicated by different color inner-circles. (b) Individual average genomes can be sequenced at K unique positions depending on the characteristic read length, k, of a sequencer. (c) All unique positions that can be sequenced for a given genome, g, defines the metagenome, g_{MG}, for the i^{th}

genome, g_i. (d) Replacing all individual genomes in (a) with metagenomes, g_{MG}, gives the metagenome of the microbial community, G_{MG}.

Sample Microbial Community

Sequencing Effort

Bioinformatic Pipeline

Metabolic Characterization

<u>Knowns</u>

1) Target Environment
2) Sequencing Depth
3) Software/Assembly Parameters
4) Reference Databases

Community Structure

<u>Unknows</u>

1) Depth and Coverage
2) Pipeline Efficiency
3) Pipeline Accuracy

A

μ

λ

D

High Evenness: 1.06x+1.84
Medium Evenness: 1.06x+3.37
Low Evenness: 1.06x+4.03
Lognormal: 1.07x+4.72

Equation 2

Equation 3

Equation 4

Equation 5

Equation 6