1

2 Chen *et al.* – Genomic investigation of *Euplotes vannus*

5 Genome analysis of the unicellular eukaryote *Euplotes vannus* reveals molecular

6 basis for sex determination and tolerance to environmental stresses

Xiao Chen,[†,1,2] Yaohan Jiang,[†,1] Feng Gao,[*,1,3] Weibo Zheng,[1] Timothy J. Krock,[4] Naomi A.

Stover,[5] Chao Lu,[2] Laura A. Katz,[6] and Weibo Song[1,7]

[1]Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao 266003,

China

[2]Department of Genetics and Development, Columbia University Medical Center, New

York, NY 10032, USA

[3]Key Laboratory of Mariculture (Ministry of Education), Ocean University of China,

Qingdao 266003, China

[4]Department of Computer Science and Information Systems, Bradley University, Peoria, IL

61625, USA

[5]Department of Biology, Bradley University, Peoria, IL 61625, USA

[6]Department of Biological Sciences, Smith College, Northampton, MA 01063, USA.

[7]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine

Science and Technology, Qingdao 266003, China

[†]Equal contributors

*Corresponding author: E-mail: gaof@ouc.edu.cn.

Abstract

As a model organism in studies of cell and environmental biology, the free-living and cosmopolitan ciliated protist *Euplotes vannus* has more than ten mating types (sexes) and shows strong resistance to environmental stresses. However, the molecular basis of its sex determination mechanism and how the cell responds to stress remain largely unknown. Here we report a combined analysis of *de novo* assembled high-quality macronucleus (MAC; i.e. somatic) genome and partial micronucleus (MIC; i.e. germline) genome of *Euplotes vannus*. Furthermore, MAC genomic and transcriptomic data from several mating types of *E. vannus* were investigated and gene expression levels were profiled under different environmental stresses, including nutrient scarcity, extreme temperature, salinity and the presence of free ammonia. We found that *E. vannus*, which possesses "gene-sized" nanochromosomes in its MAC, shares a similar pattern on frameshifting and stop codon usage as *Euplotes octocarinatus* and may be undergoing incipient sympatric speciation with *Euplotes crassus*. Somatic pheromone loci of *E. vannus* are generated from programmed DNA rearrangements of multiple germline macronuclear destined sequences (MDS) and the mating types of *E. vannus* are distinguished by the different combinations of pheromone loci instead of possessing mating type-specific genes. Lastly, we linked the resilience to environmental temperature change to the evolved loss of temperature stress-sensitive regulatory regions of HSP70 gene in *E. vannus*. Together, the genome resources generated in this study, which are available online at *Euplotes vannus* DB (http://evan.ciliate.org), provide new evidence for sex determination mechanism in eukaryotes and common pheromone-mediated cell-cell signaling and cross-mating.

Introduction

Single-celled, ciliated protists are abundant in diverse habitats across the globe, where they are among the most important components of food webs in aquatic ecosystems (Lynn 2009). Ciliate diversity, physiology and abundance has been linked to studies of environmental change (Jiang and Morin 2004; Gong et al. 2005; Xu et al. 2014), pollution monitoring (Yeomans et al. 1997; Gutiérrez et al. 2003; Jiang et al. 2011; Stoeck et al. 2018), biogeography (Borror 1980; Foissner 2006; Petz et al. 2007; Foissner et al. 2008; Liu et al. 2017), adaptive evolution (La Terza et al. 2001; La Terza et al. 2004; La Terza et al. 2007; Clark and Peck 2009; Luo et al. 2017; Wang et al. 2017a; Yan et al. 2017) and epigenetics (Xiong et al. 2016; Wang et al. 2017b; Wang et al. 2017c). *Euplotes* is a genus of free-living marine ciliates that play important roles as both predators of microalgae and preys of multicellular eukaryotes like flatworms (Dhanker et al. 2013). For decades, euplotids, including *Euplotes vannus*, have been widely used as model organisms in studies of predator/prey relationships (Kusch 1993a; Kusch 1993b; Kuhlmann 1994; Kuhlmann and Heckmann 1994; Kusch and Kuhlmann 1994; Kusch 1995; Wiąckowski and Szkarłat 1996), cell signaling (Jerka-Dziadosz et al. 1987; Hadjivasiliou et al. 2015), toxicology of marine pollutants (Persoone and Uyttersprot 1975; Trielli et al. 2007) and experimental ecology (Walton et al. 1995; Xu et al. 2004; Day et al. 2017). For example, a previous molecular study revealed that *Euplotes* species have a large number of genes requiring +1 frameshifts (i.e. addition of base pairs post-transcriptional) for expression at the post-transcriptional level, which is much higher than viruses, prokaryotes and other eukaryotes (Wang et al. 2016).

Ciliates, including the models *Paramecium* and *Tetrahymena*, have been shown to present a wide variety of mating type numbers and modes of inheritance (Orias et al. 2017). *Paramecium* has two mating types and its sex determination (SD) is controlled by scnRNA-dependent excision of the SD gene promoter (Singh et al. 2014). *Tetrahymena* has seven mating types, and different mating types specificities are encoded in the single pair of mating type genes in the MAC from all the six pairs in the MIC (Cervantes et al. 2013). More than ten mating types have been identified in *Euplotes* (Kimball 1942; Nanney et al. 1955; Siegel and Larison 1960; Akada 1985; Dini and Luporini 1985; Heckmann and Kuhlmann 1986), yet the molecular basis and SD mechanism are unknown. Previous

studies identified two subfamilies of mating-type specific pheromones in euplotids, a "shared" pheromone (designated Ec-alpha and named as Type-I pheromone in this work) and a "mating type-specific compositional" subfamily (designated Ec-1, Ec-2 and Ec-3, and named as Type-II pheromone) (Alimenti et al. 2011; Vallesi et al. 2014). They are considered as the key mediator during the cell-cell signaling that regulates cross-mating processes by controlling self/nonself recognition (Di Giuseppe et al. 2011; Vallesi et al. 2016). Previous studies reported that the euplotid model implicates relationships of hierarchical (or serial) dominance among the pheromone alleles (Heckmann 1964; Nobili et al. 1978; Dini and Nyberg 1993). Yet the results from a recent work indicated that these pheromone genes were expressed without relationships of hierarchical dominance (i.e. heterozygous genotypes behaving like homozygous cells) in the *Euplotes* MAC genome (Vallesi et al. 2014).

Euplotids also feature a strong tolerance to environmental stresses. *Euplotes* spp. were reported to have a conserved molecular defense mechanism to heavy metal contamination for homeostasis by modulating mRNA expression (Kim et al. 2018). In contrast, *E. crassus* and *E. focardii* had a barely detecTable inducible HSP70 response to salinity and temperature stresses (La Terza et al. 2001; Kim et al. 2017). These findings add weight to the argument that the lack of the classical heat shock response might be an adaptation strategy of euplotids to extreme environmental stresses. Additional studies reported *E. vannus*, as a microzooplanktonic grazer, had considerably strong tolerance to ammonia, which may enable it to survive in intensive aquaculture ecosystems with high levels of ammonium, potentially causing great damage in microalgal industry (Xu et al. 2004; Day et al. 2017). Thus, it is important to elucidate the molecular mechanism of *Euplotes* cell response to external stresses under the background of global warming.

In this work, we analyze genomic and transcriptomic data of different mating types of *E. vannus* to study its genomic features, which lead us to reveal the molecular basis of sex determination in euplotids. Furthermore, the gene expression profiling of *E. vannus* cells under different environmental stresses allows us to evaluate how this species tolerates the varying harsh conditions that it encounters.

Results

General description of genome sequencing and assembly of Euplotes vannus

In the current work, we acquired the MAC genomic data of four different mating types (EVJ, EVK, EVL and EVM), which were experimentally confirmed (Table S1). The MAC genome assemblies of these mating type had an average size of 164.2 Mb with a mean coverage of 61X (Table 1 and Supplemental Fig. S1 and Table S2).

After the contigs identified as noise (coverage < 5X) or contamination of bacteria and mitochondria were removed, the final genome assembly of *E. vannus* was generated by merging the genome assemblies of these four mating types (assembled genome size is 85.1 Mb and N50 = 2,685 bp, Table 1). We compared these data with those of two other euplotids, *E. crassus* and *E. octocarinatus*. Although *E. vannus* and *E. crassus* shared a similar %GC (36.95% and 38.65%, respectively), the genome size, number of 2-telomere contigs and N50 value of *E. vannus* were more comparable with those of *E. octocarinatus* (Table 1).

The contig N50 values of these three *Euplotes* species were all smaller than 3 kb, because chromosomes of *Euplotes* species are "nanochromosomes", similar to that of *Oxytricha trifallax* (Wang et al. 2016). The 2-telomere contig percentage of the merged genome of *E. vannus* is 66.7%, consistent with *E. octocarinatus* in which 70.3% contigs contained telomeres on both ends. Among the four mating types with genomic sequencing data, the genome assembly of EVJ was of highest quality with a 2-telomere contig percentage of 81.6% and N50 of 2,954 bp (Supplemental Table S2).

To evaluate the completeness of the genome assembly of *E. vannus*, gene content from single-copy orthologs of protists was identified by BUSCO, and the result indicated that the current assembly had a comparable percentage of complete ortholog sequences with other species (Fig. 1 and Supplemental Fig. S2). Furthermore, the majority of genomic DNA sequencing reads of four mating types (EVJ, EVK, EVL and EVM) and RNA-seq reads of six mating types (EVJ, EVK, EVL, EVM, EVP and EVX) in both starvation and vegetative stages can successfully be mapped back to the merged reference genome assembly with a mean mapping ratio of 80.1% (Supplemental Table S3). Furthermore, 109 tRNAs which consist of 48 codon types for 20 amino acids, were detected in the final genome assembly (Supplemental Table S4). These results indicated that our genome assembly of *E. vannus* was largely complete. The information of repeat regions and

functional annotation of genes are summarized in Supplemental Fig. S3, Table S5 and
Table S6.

The size distribution of complete chromosomes of *E. vannus* (i.e. those bearing
telomeric repeats "C4A4" and "T4G4" on both ends) is quite close to that of another two
euplotids, *E. octocarinatus* and *E. crassus*, with the peak values around 1.5 kb (Fig. 1B). A
similar result was found in the size distribution of telomeres, in which most telomeres of all
three euplotids had a length of 28 bp, with an increment of 8 bp (Fig. 1C). Most identified *E.*
vannus introns were around 25 bp in length, with a canonical sequence motif 5'-GTR
(N)nYAG-3' at either end (Fig. 1D, E and Supplemental Fig. S4).

Evolution and synteny/comparative genomic analyses among euplotids

Most chromosomes (37501/38245, 98.1%) in *E. vannus* are nanochromosomes, or so-
called gene-sized chromosomes, bearing a single gene on each. A similar result was
observed in *E. octocarinatus* (40396/41980, 96.2%). There was a small proportion of
chromosomes that contains more than one gene (Fig. 2A). These "joint"
nanochromosomes were then divided into two groups by the consistency and
inconsistency of the transcription directions of the genes on them (cis and trans,
respectively). The trans-joint nanochromosomes had close numbers, considering the total
chromosome numbers were similar in these two euplotids (Table 1). However, *E.*
octocarinatus possessed 2-fold more cis-joint nanochromosomes than *E. vannus*.

E. vannus and *E. crassus* shared 25026 closely related contigs (E-value cutoff = 1e-
5), which was equivalent to 65.4% and 44.2% of total contigs for each species,
respectively (Fig. 2B). However, only 469 contigs were shared between these two species
and *E. octocarinatus*. Furthermore, *E. vannus* and *E. crassus* shared not only more
homologous contigs, but also more sequence identity (Fig. 2C). For example, the
sequence identity between *E. vannus* and *E. crassus* of the corresponding region on the
chromosome that contains the coding gene of dynein heavy chain protein was 99.6%,
while the sequence identity between *E. vannus* and *E. octocarinatus* was 71.2%.

Frameshifting events in *E. vannus* and *E. octocarinatus* were detected by identifying
the adjacent region between two BLASTX hits that are in different frames, targeting to a

same protein sequence (illustrated by Supplemental Fig. S5). The E-value cutoff (1e-5)
ensured the accuracy of the prediction process and a small inner distance cutoff (10 nt)
was applied to get rid of the interference from introns, because all introns of euplotids were
larger than 20 nt as described above (Fig. 2D). The result indicated that the high frequency
of +1 programmed ribosomal frameshifting (PRF) was a conserved feature in euplotids.
However, intriguingly, more +2 and -1 PRF events were found in *E. vannus* (16.6%) than
in *E. octocarinatus* (4.4%). With more cases of +2 and -1 PRF events being spotted, a
novel motif rather than the canonical motif 5'-AAA-TAR-3' (R = A or G) was revealed as 5'-
WWW-TAR-3' (W = A or T) (Fig. 2E).

Based on the genomic and transcriptomic sequencing data, the stop codon usage
was analyzed in *E. vannus* and *E. octocarinatus* and compared between the regular
transcripts and the slippery sites of PRF transcripts (Fig. 2F). In these two euplotids, UAA
was preferentially used in the regular termination signal (73.7% and 76.0%, respectively)
and in the slippery signal (91.3% and 91.0%, respectively). Moreover, the frequency of
UAA codon usage in slippery signal is significantly higher than that in the regular
termination signal (p = 0.005024 < 0.01, Analysis of variance), which suggested that UAA
may be favorable for frameshifting in both *E. vannus* and *E. octocarinatus*.

Profiling and development of Euplotes pheromone genes

Pheromone alleles were successfully identified in MAC genomes of four *E. vannus* mating
types (Supplemental Table S7). Other than the pheromones Ev-1, Ev-2, Ev-3 and Ev-
alpha that were orthologous to Ec-1, Ec-2, Ec-3 and Ec-alpha in *E. crassus*, respectively, a
novel Type-II pheromone Ev-4 and a novel Type-I pheromone Ev-beta were found in *E.*
vannus (Fig. 3A and Supplemental figs. S6, S7 and S8). Although Ev-4 used "TAG" as
stop codon rather than using "TAA" in other three Type-II pheromones (Supplemental Fig.
S6), together with Ev-beta, they showed a significant sequence similarity with the other
three known pheromones, especially in the pre- and pro-regions of the cytoplasmic
precursor (marked by red arrows in Fig. 3A) and retained highly conserved cysteine
residues in the secreted region (marked by red dots), which was an important feature of
pheromone allele sequences.

The phylogenetic analysis of *Euplotes* pheromones, including the homologs in each mating type of *E. vannus* and the corresponding consensus sequences, was performed (Fig. 3A). Although these MAC loci were generated by alternative processing of MIC regions in each species, the result indicated that the pheromones of *E. vannus* clustered together with those of *E. crassus*, distinct from those of other Euplotids, *E. octocarinatus*, *E. nobilii* and *E. raikovi*.

Chromatin profiling of pheromone genes indicated that the combination of Type-II pheromone genes, Ev-1, Ev-2, Ev-3 and Ev-4, exhibited a mating type-specific feature on genic level (Fig. 3B and Supplemental Fig. S9). Different mating types retained 1-3 Type-II pheromone genes in the MAC genome and no duplicate events existed. Gene expression profiling of pheromone genes consistently showed that both two Type-I pheromones Ev-alpha and Ev-beta were highly expressed in all six mating types in *E. vannus* and confirmed the mating type-specific chromatin profiling of Type-II pheromone genes in different mating types at the transcriptional level. As another independent verification, PCR amplification of pheromone loci was carried out to verify the presence of the pheromone-related contigs in the MAC genome of each mating type (Fig. 3C). The results of PCR amplification of pheromone loci indicated that Type-II pheromone gene Ev-1 was absent from the mating types EVL, EVM and EVP, Ev-2 was absent from EVK, EVM, EVP and EVX, Ev-3 was absent from EVJ and Ev-4 was absent from EVK and EVP, which were mostly consistent with the results from chromatin and gene expression profiling (Fig. 3B). In brief, each of the six mating types we identified contained a unique combination of four Type-II pheromone genes.

To further study the development process of the pheromone genes during the programmed DNA rearrangement from germline MIC to somatic MAC in *E. vannus*, MIC genomic DNA was acquired by single-cell sequencing and its draft genome was assembled (Table 2). Then the germline genome (MIC) origins of Ev-1 (MAC Contig16568), Ev-2 (Contig28896), Ev-3 (Contig29423) and Ev-4 (Contig34058) were mapped to MIC contigs (Fig. 3C and Supplemental Table S8, E-value cutoff = 1e-5). The coding region of pheromone genes Ev-1, Ev-2 and Ev-3 consisted of three MDS regions from the MIC genome while Ev-4 consisted of at least two MDS regions. However, the germline source of a part of the pro-region and the secreted region of Ev-4 was not found.

Molecular basis of strong tolerance to extreme environmental stresses

Starvation stress (i.e. nutrient scarcity) had limited impact on the expression pattern of different mating types by global transcription profile (Fig. 4A and Supplemental Fig. S10). However, differential gene expression analysis revealed some mating type-specific transcripts under nutrient scarcity (Fig. 4B). Transcripts induced by starvation tended to be associated with mating type-specific genes (42.8%; see Supplemental Fig. S11). Gene functional annotation reveals that these starvation-induced and mating type-specific transcripts are related to protein transport and phosphorylation process in cells (Fig. 4C and Supplemental Fig. S12). The expression of these genes may facilitate the cell response to pheromone-mediated cell-cell signaling and cross-mating behavior under the stress from nutrient scarcity.

PCA analysis based on the differential gene expression of *E. vannus* EVJ cells under different extreme environmental stresses was performed (Supplemental Fig. S13A). The result revealed changes in the gene expression profile of cells under high temperature (35 $^{\circ}$C), low temperature (4 $^{\circ}$C), high or low salinity (60 and 10 psu, respectively). Also, the presence of free ammonia had substantial impact on transcription patterns (Supplemental Fig. S13). Surprisingly, cells under high salinity and low salinity shared a similar gene expression profile (Supplemental Fig. S13A).

To further dissect the relationships between co-expression of genes associated with the regulation of cellular processes and pathways under extreme environmental changes, a weighted gene co-expression eigengene network was constructed (Fig. 5A). The network clustered different eigengenes into six modules based on their co-expression profile (Supplemental Fig. S13B). A strongly co-expressed eigengene module was up-regulated in cells under both high and low salinity stresses (colored in steel blue in Fig. 5A and Supplemental Fig. S13B). This module was involved with an extensive activation of many pathways, mainly related to tRNA aminoacylation, tRNA and rRNA processing, nucleosome assembly and pseudouridine synthesis (p.adjust < 0.05). In addition, two small eigengene modules were up-regulated in cells under high salinity stress (purple) and low salinity stress (dark green), respectively, and low salinity stress activated an extra pathway related to the glutamine metabolic process.

Intriguingly, low temperature stress induced a large module cluster of eigengenes
related to small GTPase mediated signal transduction, while very few eigengenes co-
expressed under the high temperature (blue and purple, respectively, in Fig. 5 and
Supplemental Fig. S13B). The homolog of the highly conserved heat-shock protein 70
(Hsp70), which many organisms upregulate under environmental stress, was identified in
E. vannus and compared with its counterparts in *E. nobilii* and *E. focardii* (Fig. S14). The
result revealed that only the *E. focardii* Hsp70 sequence had numerous amino acid
substitutions within its two major functional domains, i.e. the ATP-binding and substrate-
binding domains (Fig. 6A). However, the transcription of the HSP70 gene in *E. vannus* did
not respond to temperature stresses while being responsive to other stresses like salinity
and chemical stresses (Fig. 6B).

To gain a better understanding of the molecular basis of the lack of change in
response of the HSP70 gene to temperature stress in *E. vannus*, we analyzed the
structure of non-coding regions flanking the gene in *E. vannus* and compared the structure
of this gene between *E. vannus* and *E. focardii* (Fig. 6C, D). The result indicated that no
substantial difference was detected in the 5' promoter region between the HSP70 genes of
these two species, both bearing canonical regulatory *cis*-acting elements that bind
transcriptional trans-activating factors, including heat-shock elements (HSE) and stress-
response elements (StRE) (Fig. 6C). However, the sequence of HSEs in these two species
is poorly conserved. Furthermore, neither *E. vannus* nor *E. focardii* retained an mRNA
destabilization adenine-rich element (ARE) in their 3' promoter region (Fig. 6D).

In contrast, the gene expression of cells under the presence of free ammonia were
very similar to those under high temperature stress, and there was a small cluster of
eigengenes that responded related to the lipid metabolic process (Fig. 5 and Supplemental
Fig. S13B). However, chemical stress activated the expression of HSP70 gene in *E.
vannus* significantly (Fig. 6B).

Discussion

Genomes of ciliates are divergent to each other and largely unexplored

In the current work, we present a high-quality macronuclear and a partial micronuclear genome assemblies of a unicellular eukaryote, *Euplotes vannus*, which possesses "gene-sized" MAC nanochromosomes. Comparative genome analysis reveals that *E. vannus* shares similar patterns on frameshifting and stop codon usage with *E. octocarinatus* and is undergoing incipient sympatric speciation with *E. crassus* (figs. 2 and 3). Besides euplotids, we also evaluated the assembly completeness of genomes or transcriptomes of other ciliates and found a large divergence among them (Fig. 1A and Supplemental Fig. S2). It reflects two facts: 1) ciliates have great genetic distances, even those closely related on phylogeny; 2) ciliate genomic data sequenced so far has not been collected by assembly completeness evaluation tools like BUSCO (the only ciliate covered is *Tetrahymena thermophila*). Nevertheless, among these ciliates, the completeness of transcriptome assembly of three species, *Anophryoides haemophila*, *Uronema* sp. and *Condylostoma magnum*, were not evaluated. One of the most likely reasons is that they possess small genome volume and thus share few genes with other ciliates (Chen et al. 2018). For instance, *Anophryoides* and *Uronema* are well-known parasitic species that cause disease in fish and lobsters in aquaculture facilities and have very small genome sizes (Cheung et al. 1980; Dragesco et al. 1995; Munday et al. 1997; Iglesias et al. 2001; Acorn et al. 2011). Another reason is that stop codon rearrangement occurs in some ciliates, such as *Condylostoma magnum*, and all standard stop codons are reassigned to amino acids in a context-dependent manner (Heaphy et al. 2016; Swart et al. 2016). These unusual features could dramatically increase the difficulty to precisely evaluate the assembly completeness of ciliate genomes/transcriptomes. Overall, the genomic investigation in ciliates is still waiting to be explored and the evaluation of ciliate genome assemblies calls for further improvement.

Genomic data and pheromone gene assay supports the hypothesis that E. vannus and E. crassus may be undergoing incipient sympatric speciation

An early study reported that *E. vannus* could interbreed with *E. crassus* under laboratory conditions (Valbonesi et al. 1988). Other studies reported that *E. vannus* and *E. crassus* might be undergoing sympatric speciation (Dobzhansky 1940; Zhao et al. 2018). The result in the current work increased the weight of this argument in three aspects: 1) *E. vannus*

and *E. crassus* are closely clustered with each other among euplotid species (Fig. 1A); 2)

these two species shared both a large number of homologous sequences and at high

levels of sequence identity, in contrast to comparisons with *E. octocarinatus* (Fig. 2B, C); 3)

the homologs of pheromones of *E. vannus* and *E. crassus* are closely related and distinct

from those from other *Euplotes* species (Fig. 3A, B). Orthologs of each pheromone allele

from different mating types of *E. vannus* share identical sequences on both gene and

protein levels in most cases except pheromone gene Ev-2 (Fig. 3A and Supplemental figs.

S6, S7 and S8). Our findings might describe a pattern that mating type loci evolve rapidly

after a recent speciation between *E. crassus* and *E. vannus.*

Molecular basis of the sex determination in E. vannus: the combination of Type-II

pheromone loci is mating type-specific

Type-I pheromone Ec-alpha has large sequence differences with the Type-II pheromones

and has been considered as an "adaptor" that interacts with the other pheromones, as it

has a strong propensity to oligomerize and retains a hydrophilic domain for putative

interaction (Vallesi et al. 2014; Vallesi et al. 2016). This argument is supported by the

results of expression profiling for pheromone genes in the current work (Fig. 3B).

Studies on the pheromones from other euplotids, including *E. raikovi* (Brown et al.

1993; Luginbühl et al. 1994; Mronga et al. 1994; Ottiger et al. 1994; Weiss et al. 1995; Liua

et al. 2001; Zahn et al. 2001), *E. nobilii* (Pedrini et al. 2007; Placzek et al. 2007; Di

Giuseppe et al. 2011; Vallesi et al. 2012) and *E. octocarinatus* (Schulze Dieckhoff et al.

1987; Kuhlmann et al. 1997; Brünen-Nieweier et al. 1998; Möllenbeck and Heckmann

1999), revealed that highly enriched and conserved cysteine residues in the secreted

region is the most outstanding sequence motif of *Euplotes* pheromones (Vallesi et al.

2016). The novel pheromone Ev-4 identified in this study from *E. vannus* retains 10

cysteines, as same as other Type-II pheromones in *E. vannus* and *E. crassus,* and thus

matches this characteristic perfectly (Fig. 3A).

As the Ec-1 and the other two Type-II pheromones, Ec-2 and Ec-3, were identified

in different mating types of *E. crassus* by pheromone purification and molecular mass

determination after chromatographic separation, confirmed by PCR amplification and

sequencing, Type-II pheromones have been considered as mating type-specific (Alimenti et al. 2011; Vallesi et al. 2014). However, our study demonstrated that six *E. vannus* mating types retain different combinations of Type-II pheromone loci in their MAC genomes (Fig. 3C), and thus they exhibited highly different pheromone gene expression profiling instead of possessing exclusive, mating type-specific genes (Fig. 3B). Furthermore, mating types EVL and EVP have the same set of pheromone genes but with different abundance (Fig. 3C). Therefore, it suggests that the mating type-specific combination of the Type-II pheromone loci might not be an all-or-none phenomenon, but a manner related to the composition or copy number of Type-II pheromone genes. Although further studies are expected, the observations of mating type-specific combination in the current study support the allelic codominance or non-hierarchical dominance relationship among signaling pheromone genes in euplotids (Vallesi et al. 2014; Vallesi et al. 2016).

Taken together, the current work revealed that euplotids have a novel SD manner by which mating types are determined through mating type-specific combination of four Type-II pheromone genes (Fig. 7C). Unlike *Paramecium tetraurelia*, there is no excision event on promoter regions of *E. vannus* pheromone genes (figs. 3C, 7A and Supplemental Fig. S6). On the other hand, the MAC of *E. vannus* does not possess exclusive mating type-specific SD loci as *Tetrahymena thermophila* (Fig. 3 and 7B). This mating type-specific feature of Type-II pheromones comes from the programmed DNA rearrangement between germline and somatic genomes. Intriguingly, none of the *E. vannus* mating types we have identified possesses all four Type-II pheromone genes in MAC (Fig. 3B and Supplemental Fig. S9). Thus, the results of current study described a third SD type in ciliates (Fig. 7C).

Molecular basis of the HSP70's lack of response to temperature stress

A previous study indicated that response of the HSP70 gene expression to temperature change, no matter gradually or abruptly, was divergent between *E. nobilii* and *E. focardii* (La Terza et al. 2001). When transferred from 4 to 20 °C, a strong transcriptional activity of HSP70 genes was induced in *E. nobilii* cells, while no measurable change was found in cells of *E. focardii*. In contrast, HSP70 expression increased with oxidative and chemical stresses such as tributyltin and sodium arsenite (La Terza et al. 2004). Furthermore,

together with the results from the previous studies (Dobzhansky 1940), the current work
strongly suggests the HSP70 gene of *E. vannus*, which is largely divergent from that of *E.*
focardii, does not carry unique amino acid substitutions of potential significance for cold
adaptation (Fig. 6A).

The HSP70 gene of the cosmopolitan species *E. vannus* was only activated when
facing with chemical stresses but not thermal changes (Fig. 6B). The study in Antarctic
psychrophilic euplotid *E. focardii* also reported similar Hsp70 transcriptional activity which
exclusively responded to non-thermal stresses (La Terza et al. 2004). *E. focardii* harbors
the *cis*-acting elements like heat-shock elements (HSE) and stress-response elements
(StRE) in 5' promoter region, which are known to be targets of trans-acting transcriptional
activators characterized in a variety of organisms in association with their stress-inducible
genes (Kobayashi and McENTEE 1993; Fernandes 1994; Ruis and Schüller 1995). It is
also argued that the HSE-modulated HSP70 gene transcription is more specific for a
response to temperature stress while the StRE-modulated HSP70 gene transcription is
more specific for a response to a broader range of non-temperature stresses (La Terza et
al. 2007). However, the corresponding HSE element of *E. vannus* is poorly conserved,
which may impair its sensitivity to temperature stress (Fig. 6C). Furthermore, the absence
of an adenine-rich element (ARE) in the HSP70 3' regulatory region which would exclude a
rapid mRNA degradation was also observed in *E. vannus* (Fig. 6D), as reported in the
previous study of *E. focardii* (La Terza et al. 2007). Combining these data with
observations in the current work, we argue that structural divergence of transcriptional
trans-activating factors underlies that lack of change of HSP70 gene expression in
response to temperature stress in *E. vannus* and *E. focardii*.

Conclusions

In summary, the results of the current study indicate that *Euplotes vannus* has a set of
orthologous pheromones to the reported ones in *E. crassus*, as well as a novel type of
pheromone named as Ev-4 which also shares close homology with *E. crassus*, and thus
explains the hybridization between these two species on the molecular level. Chromatin
and expression profiling of pheromone genes indicated that the combination of these
genes is mating type-specific on the genic level. Our work provides new evidence for sex

determination mechanism and supports the allelic codominance or non-hierarchical
dominance relationship among pheromone loci during *Euplotes* pheromone-mediated cell-
cell signaling and cross-mating. Furthermore, the homologous search between MAC and
MIC genomes reveals that pheromone genes in *E. vannus* develop by programmed DNA
rearrangement. According to the analyses of transcriptomes under different environmental
stresses, although the HSP70 gene of *E. vannus* does not carry unique amino acid
substitutions of potential significance for cold adaptation, it has evolved to be insensitive to
temperature change by losing the temperature stress-sensitive HSE element and mRNA
destabilization ARE elements in the regulatory region of HSP70.

Methods

Methods

Cell culture

Six mating types of *Euplotes vannus (*EVJ, EVK, EVL, EVM, EVP and EVX) were collected
from seawater along the coast of Yellow Sea at Qingdao (36°06′ N, 120°32′ E), China.
Cells of each mating type were cultured separately in filtered marine water at 20°C for 10
days, with a monoclonal population of *Escherichia coli* as the food source, until reaching
10^6 cells.

Experimental treatment simulating environmental stresses

To simulate the stress from nutrient scarcity, 10^6 cells of each mating type (EVJ, EVK, EVL,
EVM, EVP and EVX) of *E. vannus* were starved for 48 hours before harvest. For stresses
from low and high temperature, 10^6 cells of *E. vannus* mating type EVJ were cultured
under the temperature of 4 °C and 35 °C, respectively, for 6 hours before harvest. For
stress from low and high salinity, 10^6 cells of mating type EVJ were cultured under the
salinity of 10 psu and 60 psu, respectively, for 6 hours before harvest. For stress from the
presence of free ammonia, 10^6 cells of mating type EVJ were cultured in filtered marine
water with 100 mg/L NH_4Cl (pH 8.3, 20 °C and 35 psu), as described in the previous study
(Xu et al. 2004). Cells in two negative control groups were cultured under pH 7.8 and pH
8.2, respectively, in filtered marine water under 20 °C and 35 psu. Each group had two
biological replicates.

High-throughput sequencing and data processing

For regular genomic and transcriptomic sequencing to acquire macronucleus (MAC) genome information, cells were harvested by centrifugation at 300 g for 3 min. The genomic DNA was extracted using the DNeasy kit (QIAGEN, #69504, Germany). The total RNA was extracted using the RNeasy kit (QIAGEN, #74104, Germany) and digested with DNase. The rRNA fraction was depleted using GeneRead rRNA Depletion Kit (QIAGEN, #180211, Germany).

For single-cell whole-genome amplification to acquire micronucleus (MIC) genome information, a single vegetative cell of the mating type EVJ of *E. vannus* was picked and washed in PBS buffer (without Mg^{2+} or Ca^{2+}) and its MIC genomic DNA was enriched and amplified by using REPLI-g Single Cell Kit (QIAGEN, #150343, Germany), which was based on the whole-genome amplification (WGA) technology and tended to amplify longer DNA fragments.

Illumina libraries were prepared from MAC genomic DNA, mRNA and amplified single-cell MIC genomic DNA of *E. vannus* according to manufacturer's instructions and paired-end sequencing (150 bp read length) was performed using an Illumina HiSeq4000 sequencer. The sequencing adapter was trimmed and low-quality reads (reads containing more than 10% Ns or 50% bases with Q value <= 5) were filtered out.

Genome assembly and annotation

Genomes of four mating types (EVJ, EVK, EVL and EVM) were assembled using SPAdes v3.7.1 (-k 21,33,55,77), respectively (Bankevich et al. 2012; Nurk et al. 2013). Mitochondrial genomic peptides of ciliates and genome sequences of bacteria were downloaded from GenBank as BLAST databases to remove contamination caused by mitochondria or bacteria (BLAST E-value cutoff = 1e-5). CD-HIT v4.6.1 (CD-HIT-EST, -c 0.98 -n 10 -r 1) was employed to eliminate the redundancy of contigs (with sequence identity threshold = 98%) (Fu et al. 2012). Poorly supported contigs (coverage < 5 and length < 300 bp) were discarded by a custom Perl script.

A final genome assembly of *E. vannus* was merged from the genome assemblies of
four mating types (EVJ, EVK, EVL and EVM) by CAP3 v12/21/07 (Huang and Madan
1999). Completeness of genome assembly was evaluated based on expectations of gene
content by BUSCO v3 (dataset "Alveolata") and the percentage of both genomic and
transcriptomic reads mapping to the final assembly by HISAT2 v2.0.4 (Kim et al. 2015;
Simao et al. 2015). Reads mapping results were visualized on GBrowse v2.0 (Stein 2013).
Genome assemblies of *E. crassus* (accession numbers: GCA_001880385.1) and *E.*
octocarinatus (accession numbers: PRJNA294366) and their annotation information were
acquired from NCBI database and the previous studies (Wang et al. 2016; Lobanov et al.
2017; Wang et al. 2018).

Telomeres were detected by using a custom Perl script which recognized the
telomere repeat 8-mer 5'- (C4A4)n-3' at the ends of contigs, as described in a previous
study (Swart et al. 2013). The repeats in the merged genome assembly were annotated by
combining *de novo* prediction and homology searches using RepeatMasker (-engine
wublast -species '*Euplotes vannus*' -s -no_is) (Tarailo-Graovac and Chen 2009). *De novo*
genome-wide gene predictions were performed using AUGUSTUS v3.2.2 (--species =
euplotes, modified from the model "tetrahymena", rearranging TAA/TAG as stop codon,
TGA as Cys) (Stanke et al. 2006). ncRNA genes were detected by tRNAscan-SE v1.3.1
and Rfam v11.0 (Lowe and Eddy 1997; Burge et al. 2013).

Gene modeling and functional annotation

After mapping RNA-seq data of each mating type of *E. vannus* back to the merged
reference genome assembly, the transcriptome of six mating types as well as a merged
transcriptome were acquired by using StringTie v1.3.3b (Pertea et al. 2015). Annotation of
predicted protein products were matched to domains in Pfam-A database by InterProScan
v5.23 and ciliate gene database from NCBI GenBank by BLAST+ v2.3.0 (E-value cutoff =
1e-5) (Camacho et al. 2009; Jones et al. 2014).

Comparative genomic analysis

BLAST+ v2.3.0 was employed to search ciliate gene database from NCBI GenBank to
identify corresponding homologous sequences in euplotids (E-value cutoff = 1e-1 and
match length cutoff = 100 nt) (Camacho et al. 2009). "Joint" chromosomes were detected
by using a custom Perl script which recognized the chromosomes containing multiple
genes (cutoff of distance between two genes = 100 nt). Frameshifting events were
detected by using a custom Perl script which recognized the frame change between two
BLASTX hits (E-value cutoff = 1e-5 and inner distance <= 10 nt), modified from the
protocol in a previous study (Wang et al. 2016), with the addition of a strict criterion (the
distance between two adjacent hits with different frames <= 10 bp) to make sure no intron
was involved. 30 bp sequences from the upstream and downstream of each type of
frameshifting site (+1, +2 or -1) were extracted to identify the motif. Local motifs of nearby
frameshifting sites were illustrated by WebLogo 3 (Crooks et al. 2004). The frequency of
stop codon usage was estimated by a custom Perl script which recognized the stop codon
TAA or TAG in transcripts of euplotids.

Differential gene expression analysis

Transcript abundances were estimated and differential gene expression was analyzed by
using featureCounts (Liao et al. 2013) and R packages "Ballgown" and "DESeq2" (p.adjust
< 0.01) (Love et al. 2014; Frazee et al. 2015). Starvation induced genes were defined as
the average value of RPKM of gene expression from starved samples > 1 and the average
value of RPKM of gene expression from vegetative samples < 0.1. Mating type-specific
transcripts were defined as the average value of RPKM of gene expression from starved
samples > 5 and the average value of RPKM of gene expression from vegetative samples
< 0.1. Weighted gene co-expression eigengene network analysis was performed by
WGCNA (Langfelder and Horvath 2008). Gene Ontology (GO) term enrichment analysis
was performed by using BiNGO v3.0.3 (p.adjust < 0.05), which was integrated in
Cytoscape v3.4.0, and the plot was generated by the R package, ggplot2 (Maere et al.
2005; Kohl et al. 2011; Wickham 2016).

Homolog detection of pheromone genes and environmental stress-related genes

Homologous pheromone gene sequences in *E. vannus* were acquired by using BLAST+

v2.3.0 (E-value cutoff = 1e-5), according to the pheromone sequences of *E. crassus*

(Alimenti et al. 2011; Vallesi et al. 2014). Genomic DNA samples were harvested from

vegetative cells of six mating types of *E. vannus*. Type-II pheromone loci in MAC were

amplified using Q5 High-Fidelity 2X Master Mix (NEB, #M0492S, US) with 10 cells of each

mating type and genotyping primers (PCR annealing temperature was 64.5 ℃, sequences

of genotyping primers see Supplemental Table S9).

Homologous HSP70 gene sequences in *E. vannus* was acquired by using BLAST+

v2.3.0 (E-value cutoff = 1e-5), according to the Hsp70 protein sequences of *E. focardii* and

E. nobilii from the previous studies (GenBank accession number: AAP51165 and

ABI23727, respectively) (La Terza et al. 2001; La Terza et al. 2007). The complete

sequences of the *E. focardii* and *E. nobilii* HSP70 genes are available at NCBI with the

accession numbers AY295877 and DQ866998, according to the previous studies (La

Terza et al. 2004; La Terza et al. 2007). The essential amino acid positions of Hsp70 were

reported in previous studies (Sriram et al. 1997; Morshauser et al. 1999). The consensus

amino acids sequence of Hsp70 was according to the previous reports (La Terza et al.

2004; La Terza et al. 2007).

Phylogenetic analysis

The DNA and amino acid sequences of *Euplotes* pheromones homologous genes were

acquired from NCBI, according to the previous work (Vallesi et al. 2014), and aligned by

MUSCLE v3.8.31 and ClustalW v2.1, respectively (Chenna et al. 2003; Edgar 2004).

Maximum Likelihood tree based on amino acid sequences was reconstructed by MEGA

v7.0.20, using the LG model of amino acid substitution, 500 bootstrap replicates (Le and

Gascuel 2008; Kumar et al. 2016).

For phylogenomic analysis by supertree approach, predicted protein sequences of

Euplotes vannus by us and 31 other ciliates from previous works or transcriptome

sequencing by the Marine Microbial Eukaryote Transcriptome Sequencing Project (data

available on iMicrobe: http://imicrobe.us/, accession number and gene ID see

Supplemental Table S10) (Aeschlimann et al. 2014; Keeling et al. 2014; Gentekaki et al.

2017; Slabodnick et al. 2017; Wang et al. 2018) were used to generate the concatenated

dataset. Maximum Likelihood tree based on the concatenated dataset covering 157 genes

was reconstructed by using GPSit v1.0 (relaxed masking, E-value cutoff = 1e-10,

sequence identity cutoff = 50%) (Chen et al. 2018) and RAXML-HPC2 v8.2.9 (on CIPRES

Science Gateway, LG model of amino acid substitution + Γ distribution + F, four rate

categories, 500 bootstrap replicates) (Stamatakis 2014). Trees were visualized by MEGA

version 7.0.20 (Kumar et al. 2016).

Data access

578   *Euplotes vannus* MAC genome assembly and gene annotation data including coding

579   regions and predicted protein sequences are available at *Euplotes vannus* DB (EVDB,

580   http://evan.ciliate.org).

Acknowledgements

The authors would like to thank the following people for assistance with this study:

Tengteng Zhang and Ruitao Gong (Ocean University of China, China), for the assistance

of experimental verification; Dr. Fengbiao Mao (University of Michigan, USA), for the

advice on data visualization; Dr. Pierangelo Luporini (University of Camerino, Italy) and Dr.

Estienne Swart (Max Planck Institute, Germany) for the advice on the preparation of the

manuscript. This work was supported by the Aoshan Science and Technology Innovation

Program of the Qingdao National Laboratory for Marine Science and Technology, Natural

Science Foundation of China (project No. 31772428), Young Elite Scientists Sponsorship

Program by CAST (2017QNRC001) and the Fundamental Research Funds for the Central

Universities (201841013 and 201762017). Research reported in this publication was also

supported by the grants of the National Institutes of Health (award No. P40OD010964) and

the National Science Foundation (grant No. 1158346) to N.A.S. and two grants to L.A.K.

(NSF DEB-1541511 and NIH 1R15GM113177-01). The content is solely the responsibility

of the authors and does not necessarily represent the official views of the National

Institutes of Health. Any opinions, findings, and conclusions or recommendations

expressed in this material are those of the author(s) and do not necessarily reflect the

views of the National Science Foundation.

Supplemental materials

Figure S1. K-mer analysis of *Euplotes vannus* mating types to estimated genome size.

Figure S2. Genome assembly completeness evaluation of ciliates by BUSCO.

Figure S3. Venn diagram shows the genes annotated by BLASTX and Interproscan.

Figure S4. Schematic representation of the exon/intron boundaries with WebLogo in all

78661 introns in *E. vannus* mating type EVJ. The GTR and YAG motifs are well conserved.

Figure S5. A schema illustrates the criteria for detecting +1 frameshifting events. Blue

boxes indicate the different BLASTX hits of a CDS region to a same target protein

sequence (E-value cutoff = 1e-5). Grey boxes indicate the adjacent region between two

BLASTX hits of a CDS region (inner distance cutoff = 10 nt). The brackets above denote

the 0-frame codons and the brackets underneath denote the +1-frame codons. Yellow dots

denote the nucleotides while the red ones denote the slippery site where frameshifting

events occur.

Figure S6. Sequence alignment of the reverse complements of the MAC contigs

containing Type-II pheromone coding genes in *E. vannus* and *E. crassus*. Blue and red

boxes denote the start and stop codon of the coding region of pheromone genes,

respectively.

Figure S7. Sequence alignment of the Type-I pheromone protein sequences in *E. vannus*

and *E. crassus*. Identical residues are shadowed in black and similar residues are shaded

in grey. Asterisks mark the positions of stop codons. Filled and light arrowheads indicate

the extension positions of the pre- and pro-regions, respectively. Red dots denote the

conserved cysteine residues in the secreted region. Numbers indicate the progressive

amino acid positions in the sequences.

Figure S8. Sequence alignment of the reverse complements of the MAC contigs

containing Type-I pheromone coding genes in *E. vannus* and *E. crassus*. Blue and red

boxes denote the start and stop codon of the coding region of pheromone genes, respectively.

Figure S9. GBrowse snapshots of genomic and transcriptomic reads mapping on pheromone gene-related chromosomes in different mating types.

Figure S10. Overall gene expression level in starved or vegetative cells of different mating types.

Figure S11. Venn diagram shows a large part of mating type-specific transcripts is also starvation induced.

Figure S12. Species relationship and functional annotation of the mating type-specific and starvation induced transcripts.

Figure S13. Differential gene expression analysis under temperature, salinity and free ammonia stresses (relative to Figure 5). (A) PCA analysis on gene expression of mating type EVJ under different stresses. (B) Different environmental stresses activated or deactivated different gene groups.

Figure S14. Sequence alignment of Hsp70 protein sequences.

Table S1. Mating pattern observed when cultures of two mating types are mixed and genomic and transcriptomic (mRNA) data accessibility of six mating types of *E. vannus*.

Table S2. Genome assembly information of four mating types of *Euplotes vannus*.

Table S3. Genomic and transcriptomic reads mapping information.

Table S4. List of identified ncRNAs.

Table S5. Annotation information of repeats in the merged genome assembly of *E. vannus*.

Table S6. Expression and annotation information of *Euplotes vannus* genes.

Table S7. Homologs of mating-type loci in each mating type.

Table S8. Homologous search results by BLASTN reveal the relationship between coding regions of four pheromone genes in MAC genome and the corresponding MDS regions in MIC genome of *E. vannus*.

Table S9. PCR primers for genotyping of Type-II pheromone genes in *E. vannus* and
determine mating types.

Table S10. Information of accession of genome/transcriptome assemblies of 32 ciliates.

References

Acorn AR, Clark KF, Jones S, Després BM, Munro S, Cawthorn RJ, Greenwood SJ. 2011. Analysis
of expressed sequence tags (ESTs) and gene expression changes under different growth
conditions for the ciliate *Anophryoides haemophila*, the causative agent of bumper car
disease in the American lobster (*Homarus americanus*). *J Invertebr Pathol* **107**: 146-154.
Aeschlimann SH, Jonsson F, Postberg J, Stover NA, Petera RL, Lipps HJ, Nowacki M, Swart EC.
2014. The draft assembly of the radically organized *Stylonychia lemnae* macronuclear
genome. *Genome Biol Evol* **6**: 1707-1723.
Akada R. 1985. Mating types and mating-inducing factors (gamones) in the ciliate *Euplotes patella*
syngen 2. *Genet Res* **46**: 125-132.
Alimenti C, Vallesi A, Federici S, Di Giuseppe G, Fernando D, Carratore V, Luporini P. 2011.
Isolation and structural characterization of two water-borne pheromones from *Euplotes*
crassus, a ciliate commonly known to carry membrane-bound pheromones. *J Eukaryot*
Microbiol **58**: 234-241.
Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD. 2012. SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing. *J Comput Biol* **19**: 455-477.
Borror AC. 1980. Spatial distribution of marine ciliates: Micro-Ecologic and Biogeographic aspects
of Protozoan Ecology. *J Eukaryot Microbiol* **27**: 10-13.
Brown L, Mronga S, Bradshaw R, Ortenzi C, Luporini P, Wüthrich K. 1993. Nuclear magnetic
resonance solution structure of the pheromone Er-10 from the ciliated protozoan *Euplotes*
raikovi. *J Mol Biol* **231**: 800-816.
Brünen-Nieweier C, Weiligmann JC, Hansen B, Kuhlmann H-W, Möllenbeck M, Heckmann K. 1998.
The pheromones and pheromone genes of new stocks of the *Euplotes octocarinatus*
species complex. *Eur J Protistol* **34**: 124-132.
Burge SW, Daub J, Eberhardt R, Tate J, Barquist L, Nawrocki EP, Eddy SR, Gardner PP, Bateman
A. 2013. Rfam 11.0: 10 years of RNA families. *Nucleic Acids Res* **41**: 3.
Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 1471-2105.
Cervantes MD, Hamilton EP, Xiong J, Lawson MJ, Yuan D, Hadjithomas M, Miao W, Orias E. 2013.
Selecting one of several mating types through gene segment joining and deletion in
Tetrahymena thermophila. *PLoS biology* **11**: e1001518.
Chen X, Wang Y, Sheng Y, Warren A, Gao S. 2018. GPSit: An automated method for evolutionary
analysis of nonculturable ciliated microeukaryotes. *Mol Ecol Resour* **4**: 1-14.
Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. 2003. Multiple
sequence alignment with the Clustal series of programs. *Nucleic Acids Res* **31**: 3497-3500.
Cheung P, Nigrelli R, Ruggieri G. 1980. Studies on the morphology of *Uronema marinum* Dujardin
(Ciliatea: Uronematidae) with a description of the histopathology of the infection in marine
fishes. *J Fish Dis* **3**: 295-303.
Clark MS, Peck LS. 2009. HSP70 heat shock proteins and environmental stress in Antarctic marine
organisms: a mini-review. *Marine genomics* **2**: 11-18.
Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo generator.
Genome Res **14**: 1188-1190.

Day JG, Gong Y, Hu Q. 2017. Microzooplanktonic grazers - A potentially devastating threat to the
commercial success of microalgal mass culture. *Algal Research* **27**: 356-365.
Dhanker R, Kumar R, Tseng L-C, Hwang J-S. 2013. Ciliate (*Euplotes* sp.) predation by
Pseudodiaptomus annandalei (Copepoda: Calanoida) and the effects of mono-algal and
pluri-algal diets. *Zool Stud* **52**: 34.
Di Giuseppe G, Erra F, Dini F, Alimenti C, Vallesi A, Pedrini B, Wüthrich K, Luporini P. 2011.
Antarctic and Arctic populations of the ciliate *Euplotes nobilii* show common pheromone-
mediated cell-cell signaling and cross-mating. *Proc Natl Acad Sci* **108**: 3181-3186.
Dini F, Luporini P. 1985. Mating-type polymorphic variation in *Euplotes minuta* (Ciliophora:
Hypotrichida). *The Journal of protozoology* **32**: 111-117.
Dini F, Nyberg D. 1993. Sex in ciliates. In *Adv Microb Ecol*, (ed. KE Nelson), pp. 85-153. Springer.
Dobzhansky T. 1940. Speciation as a stage in evolutionary divergence. *The American Naturalist*
74: 312-321.
Dragesco A, Dragesco J, Coste F, Gasc C, Romestand B, Raymond J-C, Bouix G. 1995.
Philasterides dicentrarchi, n. sp.,(Ciliophora, Scuticociliatida), a histophagous opportunistic
parasite of *Dicentrarchus labrax* (Linnaeus, 1758), a reared marine fish. *Eur J Protistol* **31**:
327-340.
Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **32**: 1792-1797.
Fernandes M. 1994. Structure and regulation of heat shock gene promoters. *The biology of heat*
shock proteins and molecular chaperones: 375-393.
Foissner W. 2006. Biogeography and dispersal of micro-organisms: a review emphasizing protists.
Acta Protozool **45**: 111-136.
Foissner W, Chao A, Katz LA. 2008. Diversity and geographic distribution of ciliates (Protista:
Ciliophora). *Biodivers Conserv* **17**: 345-363.
Frazee AC, Pertea G, Jaffe AE, Langmead B, Salzberg SL, Leek JT. 2015. Ballgown bridges the
gap between transcriptome assembly and expression analysis. *Nat Biotechnol* **33**: 243-246.
Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. *Bioinformatics* **28**: 3150-3152.
Gentekaki E, Kolisko M, Gong Y, Lynn D. 2017. Phylogenomics solves a long-standing
evolutionary puzzle in the ciliate world: The subclass Peritrichia is monophyletic. *Mol*
Phylogenet Evol **106**: 1-5.
Gong J, Song W, Warren A. 2005. Periphytic ciliate colonization: annual cycle and responses to
environmental conditions. *Aquat Microb Ecol* **39**: 159-170.
Gutiérrez JC, Martín-González A, Díaz S, Ortega R. 2003. Ciliates as a potential source of cellular
and molecular biomarkers/biosensors for heavy metal pollution. *Eur J Protistol* **39**: 461-467.
Hadjivasiliou Z, Iwasa Y, Pomiankowski A. 2015. Cell-cell signalling in sexual chemotaxis: a basis
for gametic differentiation, mating types and sexes. *J R Soc Interface* **12**: 20150342.
Heaphy SM, Mariotti M, Gladyshev VN, Atkins JF, Baranov PV. 2016. Novel ciliate genetic code
variants including the reassignment of all three stop codons to sense codons in
Condylostoma magnum. *Mol Biol Evol* **33**: 2885-2889.
Heckmann K. 1964. Experimentelle Untersuchungen an *Euplotes crassus. Zeitschrift für*
Vererbungslehre **95**: 114-124.
Heckmann K, Kuhlmann HW. 1986. Mating types and mating inducing substances in *Euplotes*
octocarinatus. J Exp Zool **237**: 87-96.
Huang X, Madan A. 1999. CAP3: A DNA sequence assembly program. *Genome Res* **9**: 868-877.
Iglesias R, Paramá A, Alvarez M, Leiro J, Fernández J, Sanmartín M. 2001. *Philasterides*
dicentrarchi (Ciliophora, Scuticociliatida) as the causative agent of scuticociliatosis in
farmed turbot *Scophthalmus maximus* in Galicia (NW Spain). *Dis Aquat Organ* **46**: 47-55.

Jerka-Dziadosz M, Dosche C, Kuhlmann HW, Heckmann K. 1987. Signal-induced reorganization
of the microtubular cytoskeleton in the ciliated protozoon *Euplotes octocarinatus*. *J Cell Sci*
87: 555-564.
Jiang L, Morin PJ. 2004. Temperature-dependent interactions explain unexpected responses to
environmental warming in communities of competitors. *J Anim Ecol* **73**: 569-576.
Jiang Y, Xu H, Hu X, Zhu M, Al-Rasheid KA, Warren A. 2011. An approach to analyzing spatial
patterns of planktonic ciliate communities for monitoring water quality in Jiaozhou Bay,
northern China. *Mar Pollut Bull* **62**: 227-235.
Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A,
Nuka G et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**: 1236-1240.
Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, Armbrust EV, Archibald JM,
Bharti AK, Bell CJ et al. 2014. The Marine Microbial Eukaryote Transcriptome Sequencing
Project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans
through transcriptome sequencing. *PLoS Biol* **12**.
Kim B-M, Rhee J-S, Choi I-Y, Lee Y-M. 2018. Transcriptional profiling of antioxidant defense
system and heat shock protein (Hsp) families in the cadmium- and copper-exposed marine
ciliate *Euplotes crassus*. *Genes & Genomics* **40**: 85-98.
Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods* **12**: 357-360.
Kim S-J, Kim J-H, Ju S-J. 2017. Adaptation responses of individuals to environmental changes in
the ciliate *Euplotes crassus*. *Ocean Science Journal* **52**: 127-138.
Kimball R. 1942. The nature and inheritance of mating types in *Euplotes patella*. *Genetics* **27**: 269.
Kobayashi N, McENTEE K. 1993. Identification of cis and trans components of a novel heat shock
stress regulatory pathway in *Saccharomyces cerevisiae*. *Mol Cell Biol* **13**: 248-256.
Kohl M, Wiese S, Warscheid B. 2011. Cytoscape: software for visualization and analysis of
biological networks. *Methods Mol Biol* **696**: 291-303.
Kuhlmann H-W. 1994. Escape response of *Euplotes octocarinatus* to turbellarian predators. *Archiv*
für Protistenkunde **144**: 163-171.
Kuhlmann H-W, Brünen-Nieweler C, Heckmann K. 1997. Pheromones of the ciliate *Euplotes*
octocarinatus not only induce conjugation but also function as chemoattractants. *J Exp Zool*
277: 38-48.
Kuhlmann H-W, Heckmann K. 1994. Predation risk of typical ovoid and 'winged' morphs of
Euplotes (Protozoa, Ciliophora). *Hydrobiologia* **284**: 219-227.
Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis version
7.0 for bigger datasets. *Mol Biol Evol* **33**: 1870-1874.
Kusch J. 1993a. Behavioural and morphological changes in ciliates induced by the predator
Amoeba proteus. *Oecologia* **96**: 354-359.
Kusch J. 1993b. Induction of defensive morphological changes in ciliates. *Oecologia* **94**: 571-575.
Kusch J. 1995. Adaptation of inducible defense in *Euplotes daidaleos* (Ciliophora) to predation
risks by various predators. *Microb Ecol* **30**: 79-88.
Kusch J, Kuhlmann HW. 1994. Cost of Stenostomum-induced morphological defence in the ciliate
Euplotes octocarinatus. *Arch Hydrobiol* **130**: 257-267.
La Terza A, Miceli C, Luporini P. 2004. The gene for the heat-shock protein 70 of *Euplotes focardii*,
an Antarctic psychrophilic ciliate. *Antarct Sci* **16**: 23-28.
La Terza A, Papa G, Miceli C, Luporini P. 2001. Divergence between two Antarctic species of the
ciliate *Euplotes*, *E. focardii* and *E. nobilii*, in the expression of heat-shock protein 70 genes.
Mol Ecol **10**: 1061-1067.
La Terza A, Passini V, Barchetta S, Luporini P. 2007. Adaptive evolution of the heat-shock
response in the Antarctic psychrophilic ciliate, *Euplotes focardii*: hints from a comparative
determination of the hsp70 gene structure. *Antarct Sci* **19**: 239-244.

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis.
BMC Bioinformatics **9**: 559.
Le SQ, Gascuel O. 2008. An improved general amino acid replacement matrix. *Mol Biol Evol* **25**:
1307-1320.
Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**: 923-930.
Liu W, Jiang J, Xu Y, Pan X, Qu Z, Luo X, El-Serehy HA, Warren A, Ma H, Pan H. 2017. Diversity
of free-living marine ciliates (Alveolata, Ciliophora): faunal studies in coastal waters of
China during the years 2011-2016. *Eur J Protistol* **61**: 424-438.
Liua A, Luginbühla P, Zerbea O, Ortenzib C, Luporinib P, Wüthricha K. 2001. NMR structure of the
pheromone Er-22 from *Euplotes raikovi*. *J Biomol NMR* **19**: 75-78.
Lobanov AV, Heaphy SM, Turanov AA, Gerashchenko MV, Pucciarelli S, Devaraj RR, Xie F,
Petyuk VA, Smith RD, Klobutcher LA et al. 2017. Position-dependent termination and
widespread obligatory frameshifting in *Euplotes* translation. *Nat Struct Mol Biol* **24**: 61-68.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**: 014-0550.
Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes
in genomic sequence. *Nucleic Acids Res* **25**: 955-964.
Luginbühl P, Ottiger M, Mronga S, WÜthrich K. 1994. Structure comparison of the pheromones Er-
1, Er-10, and Er-2 from *Euplotes raikovi*. *Protein Sci* **3**: 1537-1546.
Luo X, Gao F, Yi Z, Pan Y, Al-Farraj SA, Warren A. 2017. Taxonomy and molecular phylogeny of
two new brackish hypotrichous ciliates, with the establishment of a new genus (Ciliophora,
Spirotrichea). *Zool J Linn Soc-lond* **179**: 475-491.
Lynn D. 2009. Ciliates. In *Encyclopedia of Microbiology (Third Edition)*,
doi:https://doi.org/10.1016/B978-012373944-5.00248-0 (ed. M Schaechter), pp. 578-592.
Academic Press, Oxford.
Maere S, Heymans K, Kuiper M. 2005. BiNGO: a Cytoscape plugin to assess overrepresentation of
gene ontology categories in biological networks. *Bioinformatics* **21**: 3448-3449.
Möllenbeck M, Heckmann K. 1999. Characterization of two genes encoding a fifth so far unknown
pheromone of *Euplotes octocarinatus*. *Eur J Protistol* **35**: 225-230.
Morshauser RC, Hu W, Wang H, Pang Y, Flynn GC, Zuiderweg ER. 1999. High-resolution solution
structure of the 18 kDa substrate-binding domain of the mammalian chaperone protein
Hsc701. *J Mol Biol* **289**: 1387-1403.
Mronga S, Luginbühl P, Brown L, Ortenzi C, Luporini P, Bradshaw R, Wüthrich K. 1994. The NMR
solution structure of the pheromone Er-1 from the ciliated protozoan *Euplotes raikovi*.
Protein Sci **3**: 1527-1536.
Munday B, O'Donoghue P, Watts M, Rough K, Hawkesford T. 1997. Fatal encephalitis due to the
scuticociliate *Uronema nigricans* in sea-caged, southern bluefin tuna *Thunnus maccoyii*. *Dis*
Aquat Organ **30**: 17-25.
Nanney D, Caughey PA, Tefankjian A. 1955. The genetic control of mating type potentialities in
Tetrahymena pyriformis. *Genetics* **40**: 668.
Nobili R, Luporini P, Dini F. 1978. Breeding systems, species relationships and evolutionary trends
in some marine species of Euplotidae (Hypotrichida Ciliata). *Marine organisms: genetics,*
ecology, and evolution Plenum, New York: 591-616.
Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin
A, Sirotkin A, Sirotkin Y. 2013. Assembling single-cell genomes and mini-metagenomes
from chimeric MDA products. *J Comput Biol* **20**: 714-737.
Orias E, Singh DP, Meyer E. 2017. Genetics and Epigenetics of mating type determination in
Paramecium and *Tetrahymena*. *Annu Rev Microbiol* **71**: 133-156.
Ottiger M, Szyperski T, Luginbühl P, Ortenzi C, Luporini P, Bradshaw R, Wüthrich K. 1994. The
NMR solution structure of the pheromone Er-2 from the ciliated protozoan *Euplotes raikovi*.
Protein Sci **3**: 1515-1526.

Pedrini B, Placzek WJ, Koculi E, Alimenti C, LaTerza A, Luporini P, Wüthrich K. 2007. Cold-
adaptation in sea-water-borne signal proteins: sequence and NMR structure of the
pheromone En-6 from the Antarctic ciliate *Euplotes nobilii*. *J Mol Biol* **372**: 277-286.
Persoone G, Uyttersprot G. 1975. The influence of inorganic and organic pollutants on the rate of
reproduction of a marine hypotrichous ciliate: *Euplotes vannus* Muller. *Revue Internationale*
d'Oceanographie Medicale **37**: 125-152.
Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* **33**:
290-295.
Petz W, Valbonesi A, Schiftner U, Quesada A, Cynan Ellis-Evans J. 2007. Ciliate biogeography in
Antarctic and Arctic freshwater ecosystems: endemism or global distribution of species?
Fems Microbiol Ecol **59**: 396-408.
Placzek WJ, Etezady-Esfarjani T, Herrmann T, Pedrini B, Peti W, Alimenti C, Luporini P, Wüthrich
K. 2007. Cold-adapted signal proteins: NMR structures of pheromones from the antarctic
ciliate *Euplotes nobilii*. *Iubmb Life* **59**: 578-585.
Ruis H, Schüller C. 1995. Stress signaling in yeast. *Bioessays* **17**: 959-965.
Schulze Dieckhoff H, Freiburg M, Heckmann K. 1987. The isolation of gamones 3 and 4 of
Euplotes octocarinatus. *The FEBS Journal* **168**: 89-94.
Siegel R, Larison L. 1960. The genic control of mating types in *Paramecium bursaria*. *Proc Natl*
Acad Sci **46**: 344-349.
Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*
31: 3210-3212.
Singh DP, Saudemont B, Guglielmi G, Arnaiz O, Goût J-F, Prajer M, Potekhin A, Przybòs E,
Aubusson-Fleury A, Bhullar S. 2014. Genome-defence small RNAs exapted for epigenetic
mating-type inheritance. *Nature* **509**: 447.
Slabodnick MM, Ruby JG, Reiff SB, Swart EC, Gosai S, Prabakaran S, Witkowska E, Larue GE,
Fisher S, Freeman RM, Jr. et al. 2017. The macronuclear genome of *Stentor coeruleus*
reveals tiny introns in a giant cell. *Curr Biol* **27**: 569-575.
Sriram M, Osipiuk J, Freeman B, Morimoto R, Joachimiak A. 1997. Human Hsp70 molecular
chaperone binds two calcium ions within the ATPase domain. *Structure* **5**: 403-414.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**: 1312-1313.
Stanke M, Tzvetkova A, Morgenstern B. 2006. AUGUSTUS at EGASP: using EST, protein and
genomic alignments for improved gene prediction in the human genome. *Genome Biology* **7**:
1.
Stein LD. 2013. Using GBrowse 2.0 to visualize and share next-generation sequence data. *Brief*
Bioinform **14**: 162-171.
Stoeck T, Kochems R, Forster D, Lejzerowicz F, Pawlowski J. 2018. Metabarcoding of benthic
ciliate communities shows high potential for environmental monitoring in salmon
aquaculture. *Ecological Indicators* **85**: 153-164.
Swart EC, Bracht JR, Magrini V, Minx P, Chen X, Zhou Y, Khurana JS, Goldman AD, Nowacki M,
Schotanus K et al. 2013. The *Oxytricha trifallax* macronuclear genome: a complex
eukaryotic genome with 16,000 tiny chromosomes. *PLoS Biol* **11**: 29.
Swart EC, Serra V, Petroni G, Nowacki M. 2016. Genetic codes with no dedicated stop codon:
context-dependent translation termination. *Cell* **166**: 691-702.
Tarailo-Graovac M, Chen N. 2009. Using RepeatMasker to identify repetitive elements in genomic
sequences. *Curr Protoc Bioinformatics* **4**: 4-10.
Trielli F, Amaroli A, Sifredi F, Marchi B, Falugi C, Corrado MU. 2007. Effects of xenobiotic
compounds on the cell activities of *Euplotes crassus*, a single-cell eukaryotic test organism
for the study of the pollution of marine sediments. *Aquat Toxicol* **83**: 272-283.

Valbonesi A, Ortenzi C, Luporini P. 1988. An integrated study of the species problem in the *Euplotes crassus-minuta-vannus* group. *J Eukaryot Microbiol* **35**: 38-45.

Vallesi A, Alimenti C, Federici S, Di Giuseppe G, Dini F, Guella G, Luporini P. 2014. Evidence for gene duplication and allelic codominance (not hierarchical dominance) at the mating-type locus of the ciliate, *Euplotes crassus*. *J Eukaryot Microbiol* **61**: 620-629.

Vallesi A, Alimenti C, Luporini P. 2016. Ciliate pheromones: primordial self-/nonself-recognition signals. In *Lessons in Immunity*, (ed. L Ballarin, M Cammarata), pp. 1-16. Elsevier.

Vallesi A, Alimenti C, Pedrini B, Di Giuseppe G, Dini F, Wüthrich K, Luporini P. 2012. Coding genes and molecular structures of the diffusible signalling proteins (pheromones) of the polar ciliate, *Euplotes nobilii*. *Marine genomics* **8**: 9-13.

Walton BM, Gates MA, Kloos A, Fisher J. 1995. Intraspecific variability in the thermal dependence of locomotion, population growth, and mating in the ciliated protist *Euplotes vannus*. *Physiol Zool* **68**: 98-113.

Wang C, Zhang T, Wang Y, Katz LA, Gao F, Song W. 2017a. Disentangling sources of variation in SSU rDNA sequences from single cell analyses of ciliates: impact of copy number variation and experimental error. *Proc R Soc B* **284**: 20170425.

Wang R, Miao W, Wang W, Xiong J, Liang A. 2018. EOGD: the *Euplotes octocarinatus* genome database. *BMC Genomics* **19**: 63.

Wang R, Xiong J, Wang W, Miao W, Liang A. 2016. High frequency of +1 programmed ribosomal frameshifting in *Euplotes octocarinatus*. *Sci Rep* **6**: 21139.

Wang Y, Chen X, Sheng Y, Liu Y, Gao S. 2017b. N6-adenine DNA methylation is associated with the linker DNA of H2A. Z-containing well-positioned nucleosomes in Pol II-transcribed genes in *Tetrahymena*. *Nucleic Acids Res* **45**: 11594-11606.

Wang Y, Wang Y, Sheng Y, Huang J, Chen X, AL-Rasheid KA, Gao S. 2017c. A comparative study of genome organization and epigenetic mechanisms in model ciliates, with an emphasis on *Tetrahymena*, *Paramecium* and *Oxytricha*. *Eur J Protistol* **61**: 376-387.

Weiss MS, Anderson DH, Raffioni S, Bradshaw RA, Ortenzi C, Luporini P, Eisenberg D. 1995. A cooperative model for receptor recognition and cell adhesion: evidence from the molecular packing in the 1.6-A crystal structure of the pheromone Er-1 from the ciliated protozoan *Euplotes raikovi*. *Proc Natl Acad Sci* **92**: 10172-10176.

Wiąckowski K, Szkarłat M. 1996. Effects of food availability on predator-induced morphological defence in the ciliate *Euplotes octocarinatus* (Protista). *Hydrobiologia* **321**: 47-52.

Wickham H. 2016. *ggplot2: elegant graphics for data analysis*. Springer.

Xiong J, Gao S, Dui W, Yang W, Chen X, Taverna SD, Pearlman RE, Ashlock W, Miao W, Liu Y. 2016. Dissecting relative contributions of cis- and trans-determinants to nucleosome distribution by comparing *Tetrahymena* macronuclear and micronuclear chromatin. *Nucleic Acids Res* **44**: 10091-10105.

Xu H, Song W, Warren A. 2004. An investigation of the tolerance to ammonia of the marine ciliate *Euplotes vannus* (Protozoa, Ciliophora). *Hydrobiologia* **519**: 189-195.

Xu H, Zhang W, Jiang Y, Yang EJ. 2014. Use of biofilm-dwelling ciliate communities to determine environmental quality status of coastal waters. *Sci Total Environ* **470**: 511-518.

Yan Y, Rogers AJ, Gao F, Katz LA. 2017. Unusual features of non-dividing somatic macronuclei in the ciliate class Karyorelictea. *Eur J Protistol* **61**: 399-408.

Yeomans W, Chubb J, Sweeting R. 1997. Use of protozoan communities for pollution monitoring. *Parassitologia* **39**: 201-212.

Zahn R, Damberger F, Ortenzi C, Luporini P, Wüthrich K. 2001. NMR structure of the *Euplotes raikovi* pheromone Er-23 and identification of its five disulfide bonds. *J Mol Biol* **313**: 923-931.

Zhao Y, Yi Z, Warren A, Song WB. 2018. Species delimitation for the molecular taxonomy and ecology of the widely distributed microbial eukaryote genus *Euplotes* (Alveolata, Ciliophora). *Proc R Soc B* **285**: 20172159.

955 **Tables**

956 **Table 1.** MAC genome assembly and transcriptome-improved gene annotation of *Euplotes*
957 *vannus* in comparison with that of other euplotids.

	E. vannus	*E. crassus*	*E. octocarinatus*
Genome size (Mb)	85.1	58.6	88.9
%GC	37.0	38.7	28.2
Contig #	38245	56587	41980
Contig N50 (bp)	2685	1581	2947
2-telomere contig #	25519	13783	29532
1-telomere contig #	7835	20646	4842
0-telomere contig #	4890	22158	7606
2-telomere contig percentage (%)	66.7	24.4	70.3
Genome size (with telomere) (Mb)	75.9	44.9	83.1
Scaffold (with telomere) #	33354	34429	34374
%Scaffold (with telomere)	87.2	60.8	81.9
Scaffold N50 (bp)	2714	1830	2999
Gene #	32755	-	29076
Exon #	175735	-	96843
Transcript #	43040	-	29076

960 **Table 2.** MIC genome assembly information of *Euplotes vannus* and recognition of MDS-
961 containing contigs and those that contain multiple MDSs.

MIC genome	Total	With MDS	Multi-MDS
Genome size (Mb)	120.0	49.8	31.8
%GC	36.0	35.7	35.9
Contig #	104988	13140	5166
Contig N50 (bp)	1953	5597	7718

Figures

Figure 1. Genome assembly of *E. vannus*. (*A*) Maximum likelihood phylogenetic tree by
supermatrix approach and assembly completeness evaluation of ciliate
genomes/transcriptomes by BUSCO. Dark blue blocks represent the percentage
of complete and single-copy genes among protists, and the steel-blue blocks
represent that of complete and duplicated genes in each species. Genomic data
of *Euplotes vannus* sequenced in the current work is marked by the red triangle.
S: class Spirotrichea. L: class Litostomatea. O: class Oligohymenophorea. C:
class Colpodea. P1: class Protocruzia. M: class Mesodiniea. P2: class
Prostomatea. H: class Heterotrichea. The scale bar corresponds to 10
substitutions per 100 nucleotide positions. (*B*) Size distribution of 2-telomere
scaffolds of *E. vannus*, *E. crassus* and *E. octocarinatus*. (*C*) Size distribution of
telomeres of *E. vannus*, *E. crassus* and *E. octocarinatus*. (*D*) Size distribution of
introns in *E. vannus* mating type EVJ. (*E*) Sequence motif of 8792 tiny introns
with the size of 25 nt. Weblogo was generated and normalized to neutral base
frequencies in intergenic regions.

Figure 2. Evolution and synteny/comparative genomic analyses. (*A*) "Joint"
nanochromosome detection in *E. vannus* and *E. octocarinatus*. Red bars denote
the frequency of joint nanochromosomes containing genes in a same
transcription direction (cis) and purple bars denote that of joint
nanochromosomes containing genes in opposite transcription directions (trans).
(*B*) Closely related contigs among three euplotids. (*C*) Homologous comparison
of the contigs containing dynein heavy chain coding gene among *E. vannus* (as
reference), *E. crassus* (blue) and *E. octocarinatus* (red). (*D*) Frameshifting
detection & comparison with *E. octocarinatus*. (*E*) Conserved sequence motif
associated with frameshift sites. Sizes of letters denote information content, or
sequence conservation, at each position. The analysis is based on the alignment
of 30 bp upstream and downstream the frameshifting motif from 1236 predicted
+1 frameshifting events that involves stop codon TAA or TAG. Note the canonical
motif 5'-AAA-TAR-3' (R = A or G) in +1 PRF and noncanonical motif 5'-WWW-
TAR-3' (W = A or T) in +2 and -1 PRF. (*F*) Stop codon usage in slippery and non-
slippery transcripts of *E. vannus* (Ev) and *E. octocarinatus* (Eo). Asterisks denote
the significant difference (*p* < 0.01).

1000

1001

Figure 3. Genomic investigation revealed new pheromone loci Ev-4 and Ev-beta and a mating type-specific combination of these loci in *E. vannus*. (*A*) Phylogenetic analysis of *Euplotes* pheromones and sequence alignment of Type-II pheromones of *E. vannus* and *E. crassus*. Identical residues are shadowed in black and similar residues are shaded in grey. Asterisks mark the positions of stop codons. Filled and light arrowheads indicate the extension positions of the pre- and pro-regions, respectively. Red dotes denote the 10 conserved cysteine residues in secreted region. Numbers indicate the progressive amino acid positions in the sequences. (*B*) Chromatin and gene expression profiling based on the RPKM of pheromone loci in each mating type of *E. vannus* by genome and transcriptome sequencing. The Tables are colored by RPKM values. (*C*) The primer design (left) and PCR amplification results (right) of *E. vannus* Type-II pheromone genes. Boxes in orange, blue and grey denote the telomeres of the MAC contigs containing the pheromone genes, the MDS regions and regions those loci in MIC genome still unknown (N/A), respectively. Arrow heads in green denote the positions of the primers. Arrows in blue denote the coding regions of pheromone genes. PCR amplifications of each pheromone gene in each mating type are conducted with three biological replicates.

1020

1021

Figure 4. Gene function analysis of mating type-specific transcripts of *E. vannus* indicates
they are starvation-induced and related to protein transport and phosphorylation
process in cells. (*A*) PCA analysis on transcript enrichment of different mating
types. (*B*) Cross plot of differential expression of mating type-specific and
starvation-regulated transcripts. (*C*) GO enrichment analysis of mating type-
specific and starvation induced transcripts.

1028

1029

1030 **Figure 5.** Differential gene expression analysis reveals several large cohorts of co-
1031 expressed genes under temperature, salinity and free ammonia stresses. (*A*)
1032 Heatmap of weighted gene co-expression network, in accordance with different
1033 stress-response gene groups. (*B*) GO term enrichment analysis on different
1034 stress-response gene groups.

1035

1036

1037   **Figure 6.** Gene expression and sequence analysis of HSP70 in *E. vannus* shows it has
1038   evolved to be insensitive to temperature change. (*A*) Amino acid substitutions
1039   that occur in *E. focardii* at the level of its HSP70 ATP- and substrate-binding
1040   domains and are unique with respect to *E. nobilii* and other organisms. Asterisks
1041   denote identities. Numbers indicate essential amino acid positions of Hsp70. (*B*)
1042   Up-regulation fold change in cells under different environmental stresses (4 °C,
1043   35 °C, 10 psu, 60 psu, pH 8.2 or with the presence of free ammonia) with respect
1044   to the control (25°C, 35 psu and pH 7.8). (*C*) Nucleotide sequence alignment of
1045   the 5' regulatory regions of the *E. vannus* (gene "MSTRG.11315" on contig
1046   "Contig21532") and *E. focardii* HSP70 genes. The identities are shaded; the
1047   telomeric C4A4 repeats and transcription initiation ATG codons are in yellow;
1048   putative sites for the transcription initiation are boxed; sequence motifs bearing
1049   agreement with HSE and StRE elements are indicated by red and blue bars,
1050   respectively. (*D*) Nucleotide sequence alignment of the 3' regulatory regions of
1051   the *E. vannus* and *E. focardii* HSP70 genes and neither of them carries ARE
1052   elements in the 3' regulatory region. The identities are shaded; the telomeric
1053   G4T4 repeats and stop TAA codons are in yellow; putative polyadenylation motifs
1054   are boxed.

1055

1056

Figure 7. Current models (simplified) for sex determination in *Paramecium tetraurelia*, *Tetrahymena thermophila* and *Euplotes vannus*. (*A*) One of the two *P. tetraurelia* mating types, mating type E depends on expression of the mtA gene during sexual reactivity. The other mating type O is determined during macronuclear development by excision of the mtA promoter (box in orange) as an internal eliminated sequence (IES), preventing expression of the gene. Adapted from Extended Data Figure 10 of Reference (Singh et al. 2014). (*B*) Mating type gene pairs in *Tetrahymena thermophila* macronuclear are assembled by joining mating type-specific macronucleus destined sequence (MDS) from micronuclear to reproduce six mating types. Segments filled with grey represent conserved transmembrane regions. Adapted from Figure 3 of Reference (Cervantes et al. 2013). (*C*) *E. vannus* shows a mating type-specific feature on the combination of different pheromone genes in MAC. Red dots denote chromosome breakage site (CBS). Segments filled with solid orange, light colors and grey denote telomeres, MDSs and conserved transmembrane regions, respectively. Dashed lines denote putative IESs.

1073