Occurrence of diamondback moths *Plutella xylostella* and their parasitoid wasps

***Cotesia vestalis* in mizuna greenhouses and their surrounding areas**

Junichiro Abe[1†], Masayoshi Uefune[2†], Kinuyo Yoneya[3], Kaori Shiojiri[4] and Junji

Takabayashi[5]

[1] National Agricultural Research Center for Western Region, Ayabe, Kyoto, 623-0035,

Japan

[2] Department Agrobiological Resources, Faculty of Agriculture, Meijo University,

Nagoya, Aichi 468-8502, Japan

[3] Entomological Laboratory, Faculty of Agriculture, Kindai University, 3327-204,

Nakamachi, Nara 631-8505, Japan

[4] Department of Agriculture, Ryukoku University, 1-5 Ooe, Otsu, Shiga 520-2194,

Japan

[5] Center for Ecological Research, Kyoto University, Otsu, Shiga, 520-2113, Japan

[†] Both are equally contributed to this paper

Correspondence: Junichiro Abe, National Agricultural Research Center for Western

Region, Ayabe, Kyoto, 623-0035, Japan; Tel: +81-84-923-4100 Fax: +81-84-924-7893

E-mail: abejun@affrc.go.jp

Author Contribution

25   JA, MU and JT conceived research. A, MU, YK and KS conducted

experiments. JA and MU analysed field data and conducted statistical analyses. JT

wrote the manuscript. JT secured funding. All authors read and approved the

manuscript.

Acknowledgements

We thank the owner of the greenhouses for his kind acceptance of our research. This

study was supported by the Bio-oriented Technology Research Advancement Institution

and by a Grant-in-Aid for Scientific Research (S) (No. 19101009), (B) (No. 26292030),

(A) (No. 18H03952) and the Naito Foundation

.

Short title: Seasonal variations in pest insects and parasitoid wasps

Abstract

Pest insects invade commercial greenhouses from the surrounding areas. We aimed to

clarify whether (1) the seasonal population dynamics of local pest insects affects their

greenhouse invasions and (2) carnivorous natural enemies of the pests mirror this

invasion. We compared the occurrence of diamondback moth [DBM: *Plutella xylostella*

(Lepidoptera: Plutellidae)] larvae and their native parasitoid wasps Cotesia vestalis

(Hymenoptera: Braconidae) on mizuna plants [*Brassica rapa* var. *laciniifolia*

(Brassicales: Brassicaceae)] in commercial greenhouses, with their population dynamics

on a wild cruciferous weed *Rorippa indica* (Brassicales: Brassicaceae) in satoyama in

Kyoto, Japan. *C. vestalis* followed the occurrence of DBM larvae on both mizuna and *R.*

indica; no *C. vestalis* were recorded in greenhouses free from DBM larvae. *C. vestalis*

females were more attracted to volatiles emitted from DBM-infested than from

uninfested mizuna. However, the presence of DBM in greenhouses could not always be

explained by its seasonal population dynamics in the surroundings.

Keywords

Plutella xylostella (Lepidoptera: Plutellidae), *Cotesia vestalis* (Hymenoptera:

Braconidae), invasion, plant volatiles, attractants, seasonal variation

Introduction

The rural "satoyama" forest and village landscape in Japan consists of areas of

small-scale wet rice paddy fields, crop fields, and greenhouses (Kobori et al. 2003).

Similar agricultural landscapes are also found in other countries (Takeuchi et al. 2003).

One of the ecological characteristics of satoyama environments is that populations of

several pest insects are harbored in the surrounding natural areas (Katoh et al. 2009),

and invasions of these pest insects in greenhouses from the surroundings are common.

Here, an intriguing question is whether the seasonal population dynamics of pest insects

in the surrounding natural environment affects the incidence of pest invasions in

greenhouses.

69   In satoyama areas, carnivorous natural enemies of pest insects also live in the

natural environment (Kagawa and Maeto 2009). Several studies have shown that, in

response to damage caused by herbivorous arthropods, plants start emitting so-called

"herbivory-induced plant volatiles (HIPVs)" that attract the carnivorous natural enemies

of the currently infesting herbivores (Arimura et al. 2009; Dicke et al. 1990; Hare 2011;

McCormick et al. 2012; Takabayashi and Dicke 1996). The attraction capability of

some of these HIPVs has been confirmed under field conditions (James 2003; James

and Grasswitz 2005; James and Price 2004; Rodriguez-Saona et al. 2011; Uefune et al.

2012). Most of the crops infested by pest insects in greenhouses start emitting

carnivore-attractive HIPVs. However, the question of whether such HIPVs attract

natural enemies from the surroundings into the greenhouses remained unanswered.

80   To clarify the above two questions, we conducted field experiments in

greenhouses in the "Miyama" satoyama area in the Kyoto Prefecture of Japan (35.3°N,

135.5°E), where a cruciferous crop, "mizuna" [*Brassica rapa* var. *laciniifolia*

(Brassicales: Brassicaceae)] was produced; pesticides were not routinely applied in

these greenhouses, but only to control observed outbreaks of pest insects as part of an

integrated pest management strategy. Populations of diamondback moth (DBM)

[*Plutella xylostella* (Lepidoptera: Plutellidae)] live in the surrounding area, and DBM

larvae are one of the most important pests of mizuna plants in greenhouses in Miyama

(J. Abe, personal observation). Populations of *Cotesia vestalis* (Hymenoptera:

Braconidae), a native parasitoid wasp of DBM larvae (Furlong et al. 2013; Talekar and

Shelton 1993), are also harbored in the Miyama area (J. Abe, personal observation). We

have previously reported that HIPVs emitted from komatsuna plants (*B. rapa*

var. *perviridis*, i.e., the same species as mizuna but a different cultivar) infested by

DBM larvae attract *C. vestalis* under both laboratory and greenhouse conditions (Ohara

et al. 2017; 2018; Uefune et al. 2012; Yoneya et al. 2018).

We observed the occurrence of DBM and *C. vestalis* on mizuna plants in four

commercial greenhouses and on *Rorippa indica* (Brassicales: Brassicaceae) plants in the

surrounding area in Miyama. Further, we tested whether mizuna plants infested by

DBM larvae attracted *C. vestalis*. Based on these data, together with our previously

reported results on the olfactory response of *C. vestalis* to DBM larvae-infested crucifer

plants, we discuss the relationship between the occurrence of DBM larvae and *C.*

vestalis in greenhouses and the surrounding areas.

Materials and Methods

Field observation

We used four greenhouses owned by one farmer, set in a "dice four" arrangement with

2–3 m distance between each. They were surrounded by open agricultural fields and a

thicket with a river, and DBM and *C. vestalis* were believed to inhabit these areas

(Supplemental Figure 1). Mizuna plants were grown in these greenhouses, but since the

growth stages in the four greenhouses differed, we treated the data from each

greenhouse independently.

The occurrence of DBM larvae and *C. vestalis* cocoons on mizuna plants in

the greenhouses and their surrounding areas were observed. Observations were made

approximately every 7–14 days during the observation period. In greenhouses, when

plants had fewer than 10 leaves, 11 to 30 leaves, or more than 30 leaves, we assessed

100, 50, and 20 plants, respectively. We also counted the numbers of DBM larvae and *C.*

vestalis cocoons on a wild cruciferous species, *Rorippa indica*, which was growing in

the surrounding area, up to around 3 m away from the tested greenhouses. DBM larvae

found on *R. indica* plants and mizuna plants were reared in a climate-controlled room in

the laboratory (25 ± 2 °C, 50–60% RH, 16L:8D), to check the incidence of parasitism.

Laboratory experiments

Insects and plants

Plutella xylostella larvae were collected from fields in Ayabe, Kyoto, Japan (35°N,

135°E) in 2001, and were reared with potted komatsuna plants [*Brassica rapa*

var. *perviridis* (Brassicales: Brassicaceae)] in a climate-controlled room (25 ± 3°C, 60 ±

10% RH, 16L:8D). The laboratory colony of DBM was reared on potted komatsuna

plants in a climate-controlled room (25 ± 3 °C, 60 ± 10% RH, 16L:8D) to obtain eggs.

Newly emerged adults of DBM were maintained in acrylic cages (35 cm × 25 cm × 30

cm high) in a climate-controlled room (25 ± 3 °C, 60 ± 10% RH, 16L:8D) and provided

with a 50% (v/v) honey solution as food and potted komatsuna plants, to ensure mating.

Komatsuna plants with eggs were collected daily, and hatched larvae were reared on cut

komatsuna plants in small cages (25 cm × 15 cm × 10 cm high).

C. vestalis were obtained from parasitized DBM larvae collected in the same

field. Adults of the parasitoid species were maintained separately in plastic cages (30

cm × 20 cm × 13 cm high) with a 50% (v/v) honey solution as food in a

climate-controlled room (18 ± 3 °C, 60 ± 10% RH, 16L:8D) for 3 days to ensure mating.

The second stadium DBM larvae that were newly parasitized by *C. vestalis* were put in

a polypropylene box (25 cm × 15 cm × 10 cm high) with detached komatsuna leaves for

food; the leaves were replaced by fresh ones every other day until the egression of *C.*

vestalis larvae from DBM larvae. After egression, *C. vestalis* formed cocoons in the

polypropylene box. Cocoons were collected and kept in closed end glass tubes until

emergence. To ensure mating, emerged females were kept together with males in a

plastic cage for 3 days. Thereafter, they were maintained in glass tubes at 18 °C to

prolong their lifespan, and in continuous darkness to suppress flight. They were a

maximum of 10 days old since emergence from the host and were acclimatized for 1–2

h in the climate room before the experiments were started.

Mizuna (*Brassica rapa* var. *nipposinica* 'Jounan-Sensuji') and komatsuna (*B.*

rapa var. *perviridis* L. 'Rakuten') plants were cultivated in a greenhouse (25 ± 3 °C, 60

± 10% relative humidity, 16L:8D). Four plants were cultivated in a plastic pot

(diameter: 72 mm, depth: 65 mm) for 4–5 weeks; these potted plants were used as the

odor sources in the laboratory experiments.

Response of C. vestalis to DBM-infested plants

Prior to each experiment, the potted mizuna or komatsuna plants had either remained

uninfested or had received damage from one second stadium DBM larva per plant

which had been allowed to feed for 24 hours to produce plants with at least one infested

edge/leaf. Prior to the tests, the larvae, their silk, and their feces were removed from the

infested plants with the aid of a fine brush.

C. vestalis females were tested for their flight responses towards a pot of

DBM-infested mizuna plants versus a pot of uninfested mizuna plants in an acrylic cage

(25 × 30 × 35 cm; 3 nylon gauze-covered windows and one door) under fluorescent

light (20 W, 3000 lux) in a climate-controlled room (25 ± 3 °C, 60 ± 10% RH, 16L:8D).

In the cage, there was no wind. The results of Shiojiri et al. (2000, 2010) show that

visual cues are not involved in the flight responses of *C. vestalis* in the choice chamber.

Females were released individually from a glass tube (25 mm inner diameter,

120 mm length) positioned halfway between two plant pots. Upon their first visit to one

of the tested plants (defined as landing), they were removed with an insect aspirator.

Ten wasps were tested using the same set of two potted seedlings. Each wasp was only

tested once; the experiments were repeated on three or four experimental days with new

sets of parasitoids and plants. We also compared the flight response of the parasitoid

wasps toward a pot of DBM-infested mizuna plants versus a pot of DBM-infested

komatsuna plants in the same manner. Two-choice data under laboratory conditions

were analyzed using the replicated *G*-test (Sokal and Rohlf 1995); parasitoids that made

no choice for either plant were discarded from this analysis.

Results

Field observations

In the surrounding areas, *R. indica* plants were detected throughout the observation

period (Figure 1a: surroundings), with the highest number on August 17. We observed

seasonal variation in the occurrence of DBM larvae in the surroundings, with two major

peaks (May 9: 246 plants, 0.13 DBM/plant; June 6: 243 plants, 0.20 DBM/plant).

Seasonal variation in the occurrence of *C. vestalis* was also observed, with two peaks

that were synchronous with those of the DBM (Figure 1: surroundings).

In the four greenhouses (Figure 1b-e), most of the plants were mizuna, with a

few *R. indica* plants found solely around the edges of the greenhouses. The cyclical

changes in the numbers of mizuna plants in the greenhouses represented the planting

and harvesting of the mizuna plants. The control threshold (level of infestation at which

pesticides were applied to the crop) of DBM in a mizuna greenhouse was set at 0.05

DBM/plant (after Abe et al. 2007); hereafter, we define the occurrence of more than

0.05 DBM/plant as a peak.

In greenhouse 1 (Figure 1b), the occurrence of DBM larvae had five peaks.

The dominant occurrence was at the second peak (June 21: 9451 plants; 0.51

DBM/plant) and was detected after the dominant peak for DBM had been recorded in

the surroundings. At each peak, the occurrence of *C. vestalis* was also detected.

In greenhouse 2 (Figure 1c), the occurrence of DBM larvae had three peaks.

The dominant occurrence was again at the second peak (June 26: 12331 plants; 0.64

DBM/plant). *C. vestalis* was detected at each peak.

In greenhouse 3 (Figure 1d), the occurrence of DBM larvae had four peaks.

This time, the dominant occurrence was the third peak (July 4: 10962 plants; 0.36

DBM/plant); however, the second most dominant peak (June 13: 3053 plants; 0.18

DBM/plant) was detected after the dominant peak for DBM had been recorded in the

surroundings. *C. vestalis* was detected at each peak.

In greenhouse 4 (Figure 1e), there were five peaks in the emergence of DBM

larvae, and mizuna production was stopped between May 16 to June 6 and July 19 to

October 14, due to outbreaks of DBM. As the occurrence of September 20 consisted of

one DBM on an *R. indica* weed when no mizuna plants were being grown in the

greenhouse, this was not classified as a peak. The third peak was observed when *C.*

vestalis was recorded in the surroundings (July 12: 2859 plants; 3.2 DBM/plant). *C.*

vestalis was detected in the greenhouse at each peak.

Olfactory responses of *C. vestalis* to DBM larvae-infested plants

We offered DBM-infested mizuna plants versus uninfested mizuna plants to *C.*

vestalis females in a choice chamber. *C. vestalis* females preferred infested mizuna

plants over uninfested mizuna plants ($G_P = 7.1034$, df = 1, $P = 0.0077$; $G_H = 0.5977$, df

= 2, $P = 0.7417$; $G_T = 7.7011$, df = 3, $P = 0.0526$, replicated *G*-test) (Figure 2: upper

bar). We then offered DBM-infested mizuna plants versus DBM-infested komatsuna

plants to *C. vestalis* females in a choice chamber. *C. vestalis* females showed an equal

distribution between the two odor sources ($G_P = 1.6108$, df = 1, $P = 0.2044$; $G_H =$

0.8402, df = 2, $P = 0.8398$; $G_T = 2.4511$, df = 3, $P = 0.6534$, replicated *G*-test) (Figure

2: lower bar).

Discussion

Throughout the observation period, DBM larvae were followed by *C. vestalis* on both

mizuna plants and *R. indica* plants. This synchronized occurrence was observed in all

four greenhouses when DBM occurred at a density of more than 0.05 DBM/plant.

Though the seasonal occurrences of DBM larvae and *C. vestalis* in the greenhouses

were similar to their seasonal population dynamics in the surrounding areas, some could

not be explained by their population dynamics. Thus, while the incidence of the

invasion was affected by the surroundings, some took place irrespective of their density

in the surrounding areas. Further, there were some differences in the occurrence patterns

of DBM even within the four adjacent greenhouses with a dice four arrangement. This

could be explained by the differences in the growth stages of the mizuna plants in the

four greenhouses. Studies on the time series observations of the invasions of pest insects

and their natural enemies in greenhouses, coupled with their seasonal population

dynamics in the surrounding satoyama areas, as shown in this study, are not

accumulating to date. However, further studies are needed to evaluate the seasonal

invasions of pests and the efficacy of native natural enemies that are harbored in the

surroundings. Shimomoto (2002) reported that the invasion of five native parasitoid

species was observed in eggplant greenhouses in which leafminers (*Liriomyza trifolii*)

occurred on eggplants.

Notably, native *C. vestalis* were recorded only in the presence of DBM larvae

in greenhouses. This specific shadowing by *C. vestalis* would in part be explained by

their response to plant volatiles. *C. vestalis* are attracted to volatiles emitted from

various crucifer plants infested by DBM larvae under both laboratory (Reddy et al.

2002; Shiojiri et al. 2000, 2010; Yoneya et al. 2018) and field conditions (Uefune et al.

2012). In the present study, we showed that mizuna plants started attracting *C. vestalis*

after being damaged by DBM larvae, and that the attractiveness was equal to that of

DBM-infested komatsuna plants (the same species as mizuna). Under experimental field

conditions, Uefune et al. (2012) reported that *C. vestalis* females were attracted to

HIPVs emitted from DBM-infested cabbage plants. Taken together, volatiles from

DBM-infested mizuna plants that are attractive to *C. vestalis* would be one of the factors

causing the co-occurrence of DBM larvae and *C. vestalis* in greenhouses.

253   It was concluded that the synchronized invasion of *C. vestalis* did not

effectively suppress the DBM larvae in the greenhouses in our study because the DBM

density in the greenhouses was more than 0.1 larva/plant at each occurrence. However,

Abe et al. (2007) reported that the single release of five *C. vestalis* in experimental

greenhouses with 0.05 DBM per komatsuna plant successfully suppressed two

successive generations of DBM (i.e., for ca 40 days); in their experiments, there were

outbreaks of DBM in the control greenhouses. Based on their study, we hypothesized

that, in the present study, the number of *C. vestalis* invading from the surroundings was

not high enough to meet the conditions for successful suppression, and that the artificial

recruitment of additional *C. vestalis* from the surroundings into greenhouses would be

one way to control the DBM. Four volatile compounds [(*Z*)-3-hexenyl acetate, α-pinene,

sabinene, and *n*-heptanal] emitted from DBM-infested cabbage plants have been shown

to attract *C. vestalis* under both experimental greenhouse and field conditions (Ohara et

al. 2017; 2018; Uefune et al. 2012). Whether the use of the volatiles that recruit native *C.*

vestalis into greenhouses will successfully suppress DBM is the subject of a subsequent

study.

Conflicts of Interest

The authors declare no conflicts of interest.

References

Abe J, Urano S, Nagasaka K, Takabayashi J. (2007) Release ratio of *Cotesia vestalis* to

diamondback moth larvae (*Plutella xylostella*) that suppresses the population

growth of diamondback moth feeding *Brassica* leaf vegetables in a greenhouse.

Bullet Nat Agr Res Centr West Region. 6:125-132. (In Japanese.)

Arimura G, Matsui K, Takabayashi J. (2009) Chemical and molecular ecology of

herbivore-induced plant volatiles: Proximate factors and their ultimate functions.

Plant Cell Physiol 50:911-923, doi: 10.1093/pcp/pcp030

Dicke M, Sabelis MW, Takabayashi J, Bruin J, Posthumus MA. (1990) Plant strategies

of manipulating predator prey interactions through allelochemicals: Prospects for

application in pest control. J Chem Ecol 16(11):3091-118. doi:

10.1007/BF00979614

Furlong MJ, Wright DJ, Dosdall LM, (2013) Diamondback moth ecology and

management: problems, progress, and prospects. Annu Rev Entomol 58: 517-541,

doi: 10.1146/annurev-ento-120811-153605

Hare, JD (2011) Ecological role of volatiles produced by plants in response to damage

by herbivorous insects. Annu Rev Entomol 56: 161-80. doi:

10.1146/annurev-ento-120709-144753

James DG (2003) Field evaluation of herbivore-induced plant volatiles as attractants for

beneficial insects: Methyl salicylate and the green lacewing, *Chrysopa nigricornis*.

J Chem Ecol 29: 1601-1609, https://doi.org/10.1023/A:1024270713493

James, DG and Grasswitz, T R (2005). Field attraction of parasitic wasps, *Metaphycus*

sp. and *Anagrus* spp. using synthetic herbivore-induced plant volatiles. BioControl

50: 871-880, DOI 10.1007/s10526-005-3313-3

James DG, Price TS (2004) Field-testing of methyl salicylate for recruitment and

retention of beneficial insects in grapes and hops. J Chem Ecol 30: 1613-1628

doi:10.1023/B:JOEC.0000042072.18151.6f

Katoh K, Sakai S, Takahashi T (2009) Factors maintaining species diversity in

satoyama, a traditional agricultural landscape of Japan. Biol Conserv 142:

1930-1936, doi.org/10.1016/j.biocon.2009.02.030

Kagawa Y, Maeto K (2009) Spatial population structure of the predatory ground beetle

Carabus yaconinus (Coleoptera: Carabidae) in the mixed farmland-woodland

satoyama landscape of Japan. Eur. J. Entomol 106: 385–391,

doi: 10.14411/eje.2009.049

Kobori H, Primack RB (2003) Participatory conservation approaches for satoyama, the

traditional forest and agricultural landscape of Japan. Ambio. Jun;32: 307-11

McCormick AC, Unsicker SB and Gershenzon J (2012) The specificity of

herbivore-induced plant volatiles in attracting herbivore enemies. Trends Plant Sci

17: 303-310

Ohara, Y., Uchida, T., Kakibuchi, K., Uefune, M. and Takabayashi, J. (2017) Effects of

an artificial blend of host-infested plant volatiles on plant attractiveness to specialist

parasitic wasps. J Appl Entomol 141: 231–234, doi: 10.1111/jen.12328

Ozawa R., Ohara, Y., Shiojiri K., Uchida, T., Kakibuchi, K., Kugimiya S., Uefune, M.

and Takabayashi, J. (2018) Uninfested plants and honey enhance the attractiveness

of a volatile blend to a parasitoid *Cotesia vestalis*. J Appl Entomol (in press)

Reddy, GVP, Holopainen JK, Guerrero A (2002) Olfactory responses of *Plutella*

xylostella natural enemies to host pheromone, larval frass, and green leaf cabbage

volatiles. J Chem Ecol, 28: 131–143

Rodriguez-Saona C, Kaplan I, Braasch J, Chinnasamy D, Williams L (2011) Field

responses of predaceous arthropods to methyl salicylate: a meta-analysis and case

study in cranberries. Biol Control 59:294-303

doi.org/10.1016/j.biocontrol.2011.06.017

Shimomoto, M. (2002) Control of *Liriomyza trifolli* (Burgess) by indigeous and

imported parasitoids on eggplant in forcing culture. Bull. Kochi Agric, Res. Cent.

11: 37-44

Shiojiri, K, Takabayashi, J, Yano, S and Takafuji A (2000) Flight response of parasitoid

toward plant-herbivore complexes: a comparative study of two parasitoid-herbivore

systems on cabbage plants. Appl Entomol Zool 35: 87-92,

doi.org/10.1303/aez.2000.87

Shiojiri K, Ozawa R, Kugimiya S, Uefune M, van Wijk M, Sabelis MW and

Takabayashi J (2010) Herbivore-specific, density-dependent induction of plant

volatiles: Honest or "cry wolf" signals? PLoS One e12161

doi.org/10.1371/journal.pone.0012161

Sokal RR and Rohlf FJ (1995) Biometry: the principles and practice of statistics in

biological research, 3rd ed. W.H. Freeman and Company; New York.

Takabayashi J, Dicke M (1996) Plant-carnivore mutualism through herbivore-induced

carnivore attractants. Trends Plant Sci 1:109-113,

doi.org/10.1016/S1360-1385(96)90004-7

Talekar NS, Shelton AM, 1993. Biology, ecology, and management of the diamondback

moth. Ann Rev Entomol 38, 275-301.

Takeuchi, K., Brown, R. D., Washitani, I., Tsunekawa, A. and Yokohari, M. (2003)

Satoyama: The traditional rural landscape of Japan. Springer-Verlag

Uefune M, Choh Y, Abe J, Shiojiri K, Sano K and Takabayashi J (2012) Application of

synthetic herbivore-induced plant volatiles causes increased parasitism of

herbivores in the field. J Appl Entomol 136: 561-567, doi:

10.1111/j.1439-0418.2011.01687.x

Yoneya K, Uefune M, Takabayashi J (2018) Parasitoid wasps' exposure to

host infested plant volatiles affects their olfactory cognition of host infested

plants. Anim Cogn 21:79–86, doi: 10.1007/s10071-017-1141-3

Figure Captions

Fig. 1 Seasonal changes in the occurrence of diamondback moth (larva and pupal

stages) and its larval parasitoid wasp *Cotesia vestalis* in greenhouses and in their

surroundings. The numbers of host plants (mizuna and *R. indica*) in both areas are

also shown. There were no plants on October 24 (Greenhouse 1), July 19 and

September 14-20 (Greenhouse 3), and May 30 (Greenhouse 4). In Greenhouse 4,

mizuna production was stopped from May 16 to June 6, and July 19 to October 14,

due to outbreaks of DBM

Fig. 2 Olfactory responses of *C. vestalis* females to komatsuna crop plants with

different treatments. The experiment was repeated on 3 or 4 experimental days

(upper and lower bars, respectively); the data were pooled and subjected to a G-test.

Numbers next to bars indicate the numbers of *C. vestalis* that responded to the

volatiles. ** $0.01 > P > 0.001$; ns: not significantly different

Supplemental Fig 1 Map of the four greenhouses. This map was revised from an

original one by Google Maps (Google, Menlo Park, California, USA)

(a) **Surroundings**

(b) **Greenhouse 1**

(c) **Greenhouse 2**

(d) **Greenhouse 3**

(e) **Greenhouse 4**

Figure 1

No choice

6   **   19   5

16   ns   24   0

80   40   0   40   80

% Response of female *C. vestalis* to crucifer crop plants

☐ Uninfested mizuna plants   ▨ DBM-infested mizuna plants   ▨ DBM-infested komatsuna plants

383   Figure 2