1

Universal metabolic constraints on the thermal tolerance of marine phytoplankton

Samuel Barton[1*], James Jenkins[1], Angus Buckling[1], C.-Elisa Schaum[1], Nicholas Smirnoff[2] &
Gabriel Yvon-Durocher[1*]

[1] Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, Cornwall
TR10 9EZ, UK,

[2] Biosciences, College of Life and Environmental Sciences, Geoffrey Pope Building University of
Exeter, Exeter, EX4 4QD, UK

*for correspondence, email: G.Yvon-Durocher@exeter.ac.uk , sb384@exeter.ac.uk

Keywords: Phytoplankton physiology, climate change, evolutionary ecology, metabolic
temperature dependence, thermal performance curves

Abstract

Marine phytoplankton are responsible for over 45% of annual global net primary
production. Ocean warming is expected to drive massive reorganisation of
phytoplankton communities, resulting in pole-ward range shifts and sharp declines in
species diversity, particularly in the tropics. The impacts of warming on phytoplankton
species depend critically on their physiological sensitivity to temperature change,
characterised by thermal tolerance curves. Local extinctions arise when temperatures
exceed species' thermal tolerance limits. The mechanisms that determine the
characteristics of thermal tolerance curves (e.g. optimal and maximal temperatures)
and their variability among the broad physiological diversity of marine phytoplankton
are however poorly understood. Here we show that differences in the temperature
responses of photosynthesis and respiration establish physiological trade-offs that
constrain the thermal tolerance of 18 species of marine phytoplankton, spanning
cyanobacteria as well as the red and green super-families. Across all species we found
that rates of respiration were more sensitive to increasing temperature and typically
had higher optimal temperatures than photosynthesis. Consequently, the fraction of
photosynthetic energy available for allocation to growth (carbon-use efficiency) declined
exponentially with rising temperatures with a sensitivity that was invariant among the
18 species. Furthermore, the optimal temperature of growth was generally lower than
that of photosynthesis and as a result, supra-optimal declines in growth rate were
associated with temperature ranges where the carbon-use efficiency exhibited
accelerated declines. These highly conserved patterns demonstrate that the limits of
thermal tolerance in marine phytoplankton are underpinned by common metabolic
constraints linked to the differential temperature responses of photosynthesis and
respiration.

Significance Statement

The impacts of warming on marine phytoplankton depend on their sensitivity to rising
temperatures, yet there is currently limited understanding of the mechanisms that limit
thermal tolerance among the diversity of marine phytoplankton. Using a comparative study
on the dominant, ecologically important lineages of marine phytoplankton –
Bacillariophyceae, Dinophyceae, Cyanophyceae, Prasinophyceae, Prymnesiophyceae – we

show that rates of respiration are consistently more sensitive to increasing temperature than

photosynthesis. Consequently, the fraction of photosynthetic energy available for growth

declines with rising temperatures with a sensitivity that is invariant among species. Our

results suggest that declines in phytoplankton performance at high temperatures are driven by

universal metabolic constrains linked to rising respiratory costs eventually exceeding the

supply of reduced carbon from photosynthesis.

Introduction

The planet's oceans are changing at an unprecedented rate (1); over the past half-

century average sea surface temperatures have been increasing by 0.1 ºC per decade (2) and

are projected to rise by a further 3ºC or more by the end of the century (3). Ocean warming is

thought to be a key driver of recent declines in phytoplankton productivity (4–6), and models

of marine biogeochemistry predict further reductions in productivity over the 21st century as

temperatures exceed limits of thermal tolerance and nutrient limitation increases in warmer,

more stratified oceans (7). Thermal tolerance curves of marine phytoplankton (like all

ectotherms) exhibit characteristic unimodality and left-skew, meaning that fitness declines

more sharply above the optimum temperature than below (8). Marine phytoplankton species

exhibit substantial variability in their thermal tolerance. Optimal temperatures for growth

range between approximately 2 to 38ºC and are positively correlated with the average

temperature of the environment, indicating a global pattern of thermal adaptation (8, 9).

Ocean warming is expected to result in major reorganisation of marine phytoplankton

communities as temperatures exceed the thermal optima of some species but not others. In

particular, tropical and sub-tropical regions are projected to experience pronounced declines

in species diversity and productivity (8, 9) because many of the taxa in these areas already

exist close to their limits of thermal tolerance. Despite its importance for predicting the

impacts of global warming on marine phytoplankton communities, we currently understand

very little about the physiological processes that determine the limits of thermal tolerance in

marine phytoplankton.

To address this fundamental knowledge gap we carried out a large-scale experiment

to investigate the physiological mechanisms that set the limits of thermal tolerance in marine

phytoplankton. Our experiments span a representative sample of the broad physiological and

phylogenetic diversity of the marine phytoplankton; including 18 species belonging to

ecologically important functional groups – Cyanobacteria, Diatoms, Dinoflagellates,
Coccolithophores, Rhodophytes, Chlorophytes and Prasinophytes (Table 1, SI). These
species were chosen to encompass the putative primary and secondary endosymbionts of both
the red and green super-families, and thus reflect the complex evolutionary histories of
marine phytoplankton (10, 11). This allowed for us to investigate whether, in spite of such
physiological diversity in plastid evolution, similar physiological constraints underpin the
limits of thermal tolerance.

<u>Results & Discussion</u>

We first characterised variability in thermal tolerance curves among taxa by
measuring growth rates for each species across a temperature gradient spanning 15 to 37°C
and fitting the Sharpe-Schoolfield equation for high temperature inactivation to the data using
non-linear mixed effects modelling (12, 13) (Fig.1). The upper limits of thermal tolerance
varied across the taxa, with $T_{max}{}^{\mu}$ (maximum temperature of observed growth), ranging from
27°C to 37°C. The optimal temperature of growth, $T_{opt}{}^{\mu}$, ranged from 23.8°C to 34.0°C and
the activation energy, $E_a{}^{\mu}$ – which characterises the increase in rate up to $T_{opt}{}^{\mu}$ – ranged from
0.40 eV to 1.46 eV, with an average $E_a{}^{\mu}$ of 0.77eV (95% CI: 0.58 to 0.97) (Fig.1, and Table
2, SI). These $E_a{}^{\mu}$ values highlight that the temperature dependence of growth at the species
level is significantly higher than previously reported temperature dependence parameters,
such as the canonical Eppley coefficient (equivalent to $E_a{}^{\mu} \approx 0.3$ eV), that are derived by
comparing maximum growth rates across many species and are the standard way in which the
impacts of warming on phytoplankton productivity are represented in models of marine
biogeochemistry (14–16). These findings suggest that the Eppley coefficient (and other
values from similar analyses (16)), which capture the broad-scale, macroecological impacts
of temperature along geographic gradients, might significantly under estimate the impacts of
temperature fluctuations on phytoplankton growth at local to regional scales (see
Supplementary Information : additional text, S1, for further discussion).

To determine the physiological processes that shape the thermal tolerance curves, in
particular those that determine the optimum temperature and supra-optimal declines in
growth rate, it is essential to understand how the key metabolic pathways that drive biomass
synthesis respond to warming. Despite having diverse evolutionary histories, all unicellular

phytoplankton share common, key metabolic pathways (17) and their ability to sequester

carbon, and therefore grow, is ultimately determined by photosynthesis and respiration (18,

19). The light-dependent reactions of photosynthesis account both for the processes that

convert inorganic carbon to organic carbon stores and those that facilitate the production of

ATP and reductant used to fuel biomass synthesis (20). The dark reactions in respiration can

be conceptually divided into 'growth' and 'maintenance' components (18–21). 'Growth-

respiration' provides the ATP, reductant and carbon skeletons required for producing new

biomass and is expected to be proportional to the rate of growth. By contrast, 'maintenance-

respiration' provides the ATP for macromolecular turnover and the maintenance of solute

gradients, and is proportional to cell biomass (20). Whilst dark respiration clearly plays an

important role in photolithotrophic growth in microalgae, the majority of the energy used to

fuel biosynthesis (between $60 - 90\%$) is thought to derive from photosynthesis (20, 21). To

understand the physiological constraints that shape the variability in phytoplankton thermal

tolerance, we quantified temperature-dependent variation in rates of photosynthesis and dark

respiration in the 18 species of marine phytoplankton.

For each species, we measured the acute responses of gross photosynthesis and dark

respiration across a temperature gradient spanning 7°C to 49°C, and quantified the resultant

thermal response curves by fitting the Sharpe-Schoolfield equation for high temperature

inactivation to the data using non-linear mixed effects modelling (see Methods). We found

consistent differences in the parameters characterising the thermal responses of

photosynthesis and respiration across all the species in this study despite their diverse

evolutionary histories (Fig.2, Fig.3 and 4, SI). The activation energy for respiration was

greater than that of photosynthesis (i.e. $E_a^R > E_a^P$; Fig. 3B, Fig. 3 and 4, SI) in all 18

species. Pooling the parameters across species yielded an average activation energy for

photosynthesis of $E_a^P = 0.74$ eV (95% CI: 0.69 to 0.79), whilst the average for respiration was

$E_a^R = 1.07$ eV (95% CI: 0.98 to 1.15). Critically, the average activation energy for

photosynthesis was statistically indistinguishable from that of growth rate ($E_a^\mu = 0.77$eV, 95%

CI: 0.58 to 0.97). These results demonstrate that respiratory costs become an increasingly

large proportion of photosynthetic carbon fixation and biomass synthesis as temperatures rise

toward the peak of the thermal response curves. We also found that for most species, the

optimum temperature for respiration was higher than that of photosynthesis (i.e. $T_{opt}^R >$

T_{opt}^P), with the average thermal optimum for photosynthesis, $T_{opt}^P = 31.18$°C ± 0.83 (s.e.m.)

and respiration, $T_{opt}^R = 32.91$°C ± 0.48 (s.e.m.) (Fig. 3C, Fig. 3 and 4, SI). Furthermore, in

all species, the deactivation energy, which characterises the speed that rates decline past the

optimum, was lower for respiration relative to photosynthesis (i.e. $E_h{}^P > E_h{}^R$), with the

average across species for photosynthesis $E_h{}^P$ = 6.08 (95% CI: 5.04 to 7.12) and respiration

$E_h{}^R$ = 2.62 (95% CI: 2.31 to 2.93) (Fig. 3D, Fig. 3 and 4, SI). Thus, as temperatures rise

beyond T_{opt}, rates of photosynthesis decline faster than rates of respiration. Overall these

findings show remarkable consistency across diverse taxa (Fig. 3 and 4, SI) in how

differences in the parameters that characterise the thermal responses of photosynthesis and

respiration result in increasing respiratory expenditure of carbon fixed by photosynthesis as

temperatures rise.

The carbon-use efficiency (CUE = 1-*R*/*P*), is an estimate of the fraction of

photosynthetic energy (*P*) that can be allocated to growth after accounting for respiration (*R*).

Recent work on both marine and freshwater phytoplankton species suggests that declines in

CUE at high temperature may be linked to impaired performance at supra-optimal

temperature (22, 23). Furthermore, observations that the evolution of elevated thermal

tolerance are coupled with adaptive shifts in metabolic traits that increase CUE at high

temperature (22–24), imply an important role for CUE in constraining thermal tolerance that

could provide a general explanation for high-temperature impairment of growth across the

diversity of the phytoplankton. To determine whether the differential thermal responses of

photosynthesis and respiration can help explain the physiological processes that constrain the

thermal tolerance curves of diverse phytoplankton, we quantified how the CUE varied as a

function of temperature. Consistent with previous work, we found that the CUE decreased

with increasing temperature in all 18 species. Declines in the CUE with rising temperature

were however highly non-linear, with the fall in CUE dramatically accelerating at high

temperatures. Because $T_{opt}{}^R > T_{opt}{}^P$ and $E_h{}^P > E_h{}^R$ for most species, as temperature rose

beyond $T_{opt}{}^P$ the CUE exhibited an accelerated decline at high temperatures. To quantify this

non-linear response and the location of the inflection point where declines in CUE become

accelerated, we fitted a break-point model to the thermal responses of the CUE. We found a

significant break-point in the thermal response of the CUE for all 18 species that was tightly

coupled with $T_{opt}{}^P$ (Fig.3). As $E_a{}^R > E_a{}^P$ for all species, temperature dependent declines in

CUE up to the break-point were universal across the species (Fig.4) with an average

activation energy, $E_a{}^{CUE}$, of -0.12eV (95% CI: -0.16 to -0.08). Furthermore, in all 18 species

the optimum temperature for growth ($T_{opt}{}^\mu$) either coincided with the CUE break-point (i.e.

the 95% CIs of the CUE break-point included $T_{opt}{}^{\mu}$), or was lower than the CUE break-point

(Fig.3). This finding suggests that temperature-driven declines in the CUE, linked to

fundamental differences in the intrinsic thermal responses of photosynthesis and respiration,

could play an important role in constraining the thermal tolerance of diverse marine

phytoplankton. Because the metabolic costs for repair and maintenance are largely accounted

for by dark respiration (20, 21) the temperature-driven declines in the CUE likely reflect

increases in the costs associated with maintenance and repair of heat-induced cellular damage

that eventually exceed the rate of substrate supply by photosynthesis, causing rates of growth

to decline at supra-optimal temperatures.

It is important to note that our experiments were conducted under nutrient replete

conditions. A recent study has suggested that the temperature sensitivities of photosynthesis

and respiration (25) in some marine phytoplankton may decline under nutrient limitation and

that the differential temperature sensitivities of photosynthesis and respiration may be

negligible under limited conditions. This work however quantified the temperature

sensitivities of photosynthesis and respiration at only 3 or 4 temperatures leading to estimates

of thermal sensitivities with large error margins and a high probability of generating type II

errors (i.e. accepting the null hypothesis of no difference in the thermal sensitivity of

photosynthesis and respiration). Furthermore, measurements were made only under resource

limited conditions precluding a quantitative comparison with nutrient replete conditions via

the same methodology. Whilst we expect that the absolute values of the thermal sensitivities

of photosynthesis and respiration are likely to decline under resource limitation, it is highly

improbable that the intrinsic differences between photosynthesis and respiration documented

in this study under nutrient replete conditions will be erased under nutrient limitation. Indeed,

our analyses demonstrate that light limitation had a negligible impact on the temperature

sensitivity of photosynthesis and in particular, the fundamental differences in the impacts of

temperature on photosynthesis and respiration were preserved under light limited conditions

(see Fig.2 and Table 6, SI). We therefore anticipate that the supra-optimal declines in growth

linked to temperature-driven decoupling between photosynthesis, respiration and biomass

synthesis that we have shown here, apply equally under nutrient replete and limited

conditions. Whilst large areas of the global ocean are under nutrient limited conditions for

long periods (26), understanding the impacts of temperature under nutrient replete conditions

(as we have done here) remains critically important because a large proportion of marine

primary productivity occurs during episodic bloom events driven by short periods of

increased nutrient concentrations (27–29). Clearly, significant further work is required to

understand the interplay between temperature and nutrient availability on phytoplankton

physiology and to assess whether the patterns we have shown here apply to conditions of

nutrient limitation, given that current experimental evidence (25) is not sufficient to draw

meaningful conclusions.

Conclusions

Overall, our findings highlight marked similarities in the temperature dependence of

photosynthesis and respiration across diverse taxonomic groups, spanning the cyanobacteria

and red and green super families and suggest that common physiological trade-offs underpin

the thermal tolerance of marine phytoplankton. We found that rates of respiration were more

sensitive to temperature, had higher thermal optima and declined less abruptly past the

optimum than those of photosynthesis. Consequently, the fraction of photosynthetic energy

available for allocation to growth (the CUE) exhibited an accelerated decline with rising

temperatures in a manner that was highly conserved among the 18 species investigated. We

also found that the optimal temperature for growth coincided with, or was lower than, an

inflection point in the temperature dependence of the CUE, which marked a transition that led

to accelerated declines at high temperatures. These patterns suggest that universal metabolic

constraints driven by the differential temperature sensitivity of photosynthesis and respiration

play a key role in setting the limits of thermal tolerance of diverse marine phytoplankton. Our

results therefore help pave the way for improving representations of phytoplankton

biodiversity in models of ocean biogeochemistry by providing a process-based understanding

of the factors that shape the limits of temperature tolerance for diverse species of marine

phytoplankton, which can be used to aid predictions of immigration and local extinctions

driven by global warming.

Methods

Culturing of marine phytoplankton strains

18 marine phytoplankton strains were obtained from CCAP (The Culture Collection of Algae

and Protozoa) and RCC (Roscoff Culture Collection) between autumn 2015 and spring 2016.

Strains of eukaryotic phytoplankton were selected from phylogenetic groups of both the red

and green superfamilies (10, 11), in addition to two strains of cyanobacteria. We tried to

work with organisms that had been well studied in the literature, were known to be globally

abundant and play crucial roles for marine ecology and global carbon cycling. The strains

were originally isolated from a range of latitudes and some have been in culture for up to 65

years (Table 1, SI). Stocks of each of the strains were cultured on their previous culture

collection medium (Table 1, SI) using artificial sea water. The following media were used:

Guillard's F/2 and F/2 + Si, Keller's K, K + Si and K/2, and PCR-S11 Red Sea medium (with

Red Sea salts). All stock cultures were incubated in Infors HT incubators at 20°C, under a

12:12 hour light-dark cycle with a PAR intensity of 45-50 µmol $m^2 s^{-1}$ and shaken at 65RPM.

Where possible we tried to obtain strains from the culture collections that matched, or were

close to, these conditions. The red alga *Porphyridium purpureum* was an exception, which we

cultured at 20-25 µmol $m^2 s^{-1}$. Cultures were kept under exponential, nutrient replete, growth

conditions for ~ 2 months before any physiological data was collected.

Measuring the thermal tolerance curve

For each species, a minimum of 3 technical replicates were inoculated with the same starting

density into fresh growth medium across a range of temperatures (15°C - 37°C). Cell counts

were made daily using flow cytometry (Accuri C6 flow cytometer, BD Scientific), and

population density was tracked until cultures reached carrying capacity. Per capita growth

rates (μ) were quantified from a modified Baranyi growth model without the lag phase(30),

using non-linear least squares regression via the 'nlsMicrobio' package in R statistical

software (v3.3.1). Models were fitted using the 'nlsLoop' function in the R github package

'nlsLoop'. This draws on the 'nlsLM' function in the 'minpack.lm' R package, which uses a

modified Levenberg-Marquardt optimisation algorithm. Model parameters were determined

by running 2000 random combinations of estimated starting parameters, which were then

selected using the Akaike Information Criterion (AIC) to determine the set of parameters that

best characterised the data. Growth rates derived for each technical replicate at each growth

temperature were then used to determine the thermal tolerance curves (Fig.1A).

Estimates of Cell Carbon and Nitrogen

270   For each species, an exponentially growing culture from the 20°C stock was divided into 3

271   technical replicates and centrifuged at 3500RPM, at 4°C for 30 minutes. The resultant pellets

were rinsed with deionised water and re-spun 3 times to remove any artificial sea water residue. For the calcifying organisms (*Emiliania huxleyi, Gephyrocapsa oceanica, Thoracosphaera heimii* i.e. those with a calcium carbonate coccoliths) it was necessary to dissolve the extra-cellular inorganic carbon (31, 32). This was achieved by treating these pellets with 0.5 mL of 3M HCl for 1 hour before being rinsed with deionised water and re-pelleted. All pellets were freeze-dried using a CoolSafe (95-15 PRO, ScanVac) over 24 hours and then weighed to obtain dry weight. Samples were placed in tin cups and sent to Elemtex (Elemtex Ltd, Cornwall, UK, PL17 8QS) for elemental analysis of %C and %N using a SerCon Isotope Ratio Mass Spectrometer (CF-IRMS) system (continuous flow mode). For each technical replicate we then calculated the C:N ratio as well as $\mu g\ C\ cell^{-1}$ (Table 3, SI).

Measuring the metabolic thermal response curves

Measurements of photosynthesis and dark respiration were collected across a range of assay temperatures (7°C to 49°C) for a minimum of 3 biological replicates per species. We used a clark-type oxygen electrode as part of a Chlorolab 2 system (Hansatech Ltd, King's Lynn, UK) to measure net rates of oxygen evolution in the light (net primary production, NP) and oxygen consumption in the dark (dark respiration); both in units of $\mu mol\ O_2\ mL^{-1}\ s^{-1}$. All biological replicates were sampled from the stock cultures, which had all been growing at 20°C and were taken at the mid-logarithmic growth phase to ensure that the samples were not substrate limited. To improve the signal to noise ratio when measuring rates, all biological replicate samples were concentrated by centrifugation at 1500rpm, 20°C, for 15 minutes and re-suspended into an adequate volume of fresh growth medium. Prior to running a sample at each assay temperature, all samples were given ~ 15 minutes to pre-acclimate to the assay temperature in the dark before any data was collected. This also gave the electrode system sufficient time to stabilise before metabolic rates were measured. This was necessary for two reasons, i) as the sample adjusts to the assay temperature this will naturally cause changes in the dissolved oxygen concentration, ii) the electrode system results in oxygen signal drift, and this too is temperature dependent. We measured rates of oxygen depletion from 21 sterilised artificial seawater samples across a range of temperatures 4°C - 44°C and found that the impact of drift was minimised after ~15 minutes of stabilisation time. Nevertheless, signal drift was linearly temperature dependent after this time. To account for drift in our dataset we corrected all our raw data using the following empirically derived relationship:

$drift = (-0.392 \times T) - 6.51$ [1]

Where T is assay temperature (°C), and *drift* is the non-biological depletion in oxygen

concentration measured in units $\mu mol O_2$ mL^{-1} s^{-1} after approximately 15 minutes of

stabilisation. The raw O_2 flux data was then corrected by subtracting the estimated drift.

Rates of net photosynthesis, measured as O_2 evolution, were collected across a range of light

intensities from 0 to 1800 μmol m^2 s^{-1} with increments of 50 μmol m^2 s^{-1} between 0 to 200

μmol m^2 $s^{-1,}$ 100 μmol m^2 s^{-1} between 200 and 1000 μmol m^2 s^{-1}, followed by 1200 μmol m^2 s^{-}

^{1}, 1500 μmol m^2 s^{-1} and finally 1800 μmol m^2 s^{-1}. This enabled us to model a photosynthesis-

irradiance (PI) curve for each assay temperature, and therefore obtain an estimate of light

saturated net photosynthesis, NP_{max}, see Eq. 2. Respiration (R) was measured as oxygen

consumption in the dark, over a 3-minute period directly following the light response outlined

above. The photosynthesis-irradiance curve was then quantified by fitting Eiler's

photoinhibiton model to the data using non-linear least squares regression (as described

above) (33, 34):

$NP(I) = \dfrac{NP_{max}I}{\frac{NP_{max}}{\alpha I_{opt}^2}I^2 + \left(1 - 2\frac{NP_{max}}{\alpha I_{opt}}\right)I + \frac{NP_{max}}{\alpha}}$ [2]

Where $NP(I)$ is the rate of net primary production at light intensity, I, NP_{max} is the

maximum rate of NP at the optimal light intensity, I_{opt}, and α is the rate in which NP

increases up to NP_{max}.

Light saturated gross primary production (P) was then calculated for each assay temperature

as:

$P = NP_{max} + R$ [3]

To investigate the effect of light limitation on the temperature dependence of photosynthesis

we used Eq. 2 to determine the predicted NP at half the light saturated irradiance ($0.5 \times$

I_{opt}). Thus replacing in NP_{max} in Eq. 3 with this prediction we derived $P_{0.5}$, a light limited

value of gross primary production at half the saturating irradiance for each assay temperature

response.

Metabolic rates were then converted from units μmol O_2 mL^{-1} s^{-1} to μg C μg C^{-1} $hour^{-1}$. We

achieved this using the following equation:

$$b(\mu g\ C\ \mu g\ C^{-1}\ h^{-1}) = \frac{b(\mu mol\ O_2\ cell^{-1}\ h^{-1})\times 32\ \times\ M\ \times\ (\frac{12}{44})}{\mu gC\ cell^{-1}} \qquad [4]$$

Where b is the metabolic rate (either P or R), 32 is the molecular weight of O_2, M is a species

specific assimilation quotient for CO_2:O_2 (35) which is used to describe consumption or

fixation of C in the cell per unit of O_2 , and 12/44 is the ratio of molecular weight of C to CO_2,

thus $32\ \times\ M\ \times\ \frac{12}{44}$ converts from $\mu mol\ O_2$ to μgC. Samples from each strain were analysed

to determine species-specific $\mu g\ C\ cell^{-1}$ values and the number of cells mL^{-1} was

measured for each biological replicate using flow cytometry. The calculation of M is based

on the assumption that NO_3^- is the main nitrogen source in the growth medium and that there

is a balanced growth equation, where:

$$nCO_2 + (n+1)H_2O + HNO_3\ \rightarrow (CH_2O)_n\ NH_3 + (n+2)O_2 \qquad [5]$$

If the C:N ratio (n) of the phytoplankton is calculated in moles then the ratio of CO_2:O_2, or M,

will be equal to $n/n+2$ (35). Our calculated values of M ranged from ~0.71 to ~0.89 (Table 3,

SI).

Quantifying the thermal response curves

The thermal response curves for rates of growth, photosynthesis (at both saturated and half

saturated irradiance) and respiration were quantified using a modified version of the Sharpe-

Schoolfield equation (12, 13):

$$\ln(b(T)) = E_a\left(\frac{1}{kT_c} - \frac{1}{kT}\right) + \ln(b(T_c)) - \ln\left(1 + e^{E_h\left(\frac{1}{kT_h} - \frac{1}{kT}\right)}\right) \qquad [6]$$

where b is either the rate of growth (d[-1]), photosynthesis or respiration ($\mu g\ C\ \mu g\ C^{-1}\ h^{-1}$), k is

Boltzmann's constant (8.62×10^{-5} eV K[-1]), E_a is the activation energy (eV), indicative of the

steepness of the slope leading up to the thermal optima, T is temperature in Kelvin (K), E_h is

the deactivation energy which characterizes temperature-induced decrease in rates above T_h

where half the enzymes have become non-functional and $b(T_c)$ is rate normalized to an

arbitrary reference temperature, here $T_c = 20$ºC (+ 273.15), where no low or high temperature

inactivation is experienced. Eq. 6 can be used to derive an optimum temperature where the

maximum rate is predicted:

$$T_{opt} = \frac{E_h T_h}{E_h + kT_h\ln\left(\frac{E_h}{E_a}-1\right)} \qquad [7]$$

The parameters $b(T_c)$, E_a, E_h, T_h, and T_{opt}, can be considered as traits that characterise the

unimodal response of biological rates to temperature change. We expect these traits to differ

across the diverse taxa analysed in this study, owing to their diverse evolutionary histories

and ancestral temperature regimes (given that they have been isolated from different

latitudes/oceans). To test this assumption, we fitted the data for growth, photosynthesis and

respiration across all species to Eq. 6 using non-linear mixed effects modelling with the

'nlme' package in R. We used separate analyses to assess the thermal responses of growth,

photosynthesis and respiration. All models included each of the parameters in Eq. 6 as fixed

effects, which quantify the average value of the parameter across all species and replicates.

For the analysis of the thermal response of growth rate, we included 'species' as a random

effect on each parameter, which quantifies species-specific deviations from the average

across all species (i.e. the fixed effect) that are assumed to be normally distributed with a

mean of zero. For the analyses of photosynthesis and respiration, we included 'replicate'

nested within 'species' to account for the fact that we measured a minimum 3 replicate

thermal response curves for each species. Here the random effect quantifies species-specific

deviations from the fixed effects as well as those attributable to variance among the replicates

of each species.

Because the Sharpe-Schoolfield equation can only take non-zero and positive rate values, in

instances where either no observed growth rate, or a negative growth rate were measured

(typically the highest and lowest temperature) we set the rate to the minimum value measured

for the species in order to fit the model.

Quantifying the carbon-use efficiency and modelling the break-point temperature

The carbon-use efficiency (CUE) was calculated as:

$\text{CUE} = 1 - R/P$ [8]

Due to the non-linear temperature response of the CUE, with accelerated declines at high-

temperatures, we fitted a segmented linear regression model to estimate the break-point in the

temperature response after which the CUE exhibited an accelerated decline. We fitted the

segmented linear regression model to CUE values derived from the fitted Sharpe-Schoolfield

curves for each species enabling us to derive an estimate of CUE at every 1°C increment

across the range of assay temperatures where metabolic rates were measured for each species

390 (Fig 2. Main text). We fitted the break-point model to the CUE values using the 'segmented'

391 package in R, where the breakpoint estimate is defined in the segmented model as the

392 intersection where there is significant difference in slopes (36), determined by the Davies test

393 for performing hypothesis (37). It is for this reason that it was necessary to use the predicted

394 values of respiration and photosynthesis to derive the break-point, as the measured data in

395 most cases only provided one or two data points beyond the inflection point, and this would

396 not have been sufficient to accurately model the second slope beyond this point (Fig.3, Main

397 text). The model returned an estimate of the CUE break-point temperature and the 95%

398 confidence intervals surrounding this value for each species (Table 7, SI).

Determining the temperature dependence of the CUE

400 We characterized the temperature dependence of the CUE up to the CUE breakpoint

401 temperature for each species using the Arrhenius equation,

402 $\ln \text{CUE}(T) = E_a{}^{CUE}\left(\frac{1}{kT_c} - \frac{1}{kT}\right) + \ln \text{CUE}(T_c)$   [9]

403 where $\ln \text{CUE}(T)$ is the natural logarithm of the CUE at temperature T (in Kelvin), $E_a{}^{CUE}$ is

404 the apparent activation energy characterising the temperature dependence of CUE. We

405 centred the temperature data using an arbitrary reference temperature T_c = 283 K = 20ºC, so

406 that $\ln \text{CUE}(T_c)$ is the CUE at T_c. We fitted Eq. 9 to all the measurements of CUE, up to the

407 CUE break-point temperature identified for each species (Fig.3 Main text, Table 7, SI) using

408 a linear mixed effects model. This allowed us to derive an average value for $E_a{}^{CUE}$ and

409 $\ln \text{CUE}(T_c)$ across the 18 species. We also included random effects of 'replicate' nested

410 within 'species' in the model to account for the fact we measured a minimum of 3 replicate

411 responses of respiration and photosynthesis for each species. This allowed us to capture the

412 species-specific and replicate specific estimates $E_a{}^{CUE}$ and $\ln \text{CUE}(T_c)$.

Acknowledgements

414 This study was supported by a grant from the Leverhulme Trust (RPG-2013-335) awarded to

415 G.Y.-D., A.B. and N.S.

Author contributions

417 S.B. and G.Y.-D conceived the study. S.B., G.Y.-D, A.B. and N.S. designed the experimental

418 work. S.B., J.J. and C.-E.S. conducted the experimental work. S.B. and G.Y.-D analysed the

419 data. S.B. and G.Y.-D wrote the manuscript and all authors contributed to revisions.

References

421   1.   Henson S A, et al. (2017) Rapid emergence of climate change in environmental drivers of
422   marine ecosystems. *Nat Commun* 8(5020):1–9.

423   2.   Rhein, M. *et al.*(2013) Observations: Ocean Pages. *Climate Change 2013: The Phyical.*
424   *Science Basis. Contribution to Working Group I to the Fifth Assessment Report of the*
425   *Intergovernmental Panel on Climate Change,* Cambridge University Press, Cambridge,
426   UK and New York, pp 255–316.

427   3.   Collins, M. *et al.*(2013) Long-term Climate Change: Projections, Commitments and
428   Irreversibility. *Climate Change 2013: The Phyical. Science Basis. Contribution to*
429   *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on*
430   *Climate Change,* Cambridge University Press, Cambridge, UK and New York, pp
431   1029–1136.

432   4.   Boyce DG, Lewis MR, Worm B (2010) Global phytoplankton decline over the past century.
433   *Nature* 466(7306):591–596.

434   5.   Behrenfeld MJ, et al. (2006) Climate-driven trends in contemporary ocean productivity.
435   *Nature* 444(December):752–755.

436   6.   Capuzzo E, et al. (2018) A decline in primary production in the North Sea over 25 years,
437   associated with reductions in zooplankton abundance and fish stock recruitment. *Glob Chang*
438   *Biol* 24:e352–e364.

439   7.   Laufkotter C, et al. (2015) Drivers and uncertainties of future global marine primary
440   production in marine ecosystem models. *Biogeosciences* 12:6955–6984.

441   8.   Thomas MK, Kremer CT, Klausmeier C a., Litchman E (2012) A Global Pattern of Thermal
442   Adaptation in Marine Phytoplankton. *Science* 338:1085–1088.

443   9.   Thomas MK, Kremer CT, Litchman E (2016) Environment and evolutionary history determine
444   the global biogeography of phytoplankton temperature traits. *Glob Ecol Biogeogr* 25:75–86.

445   10.   Falkowski PG, et al. (2004) The Evolution of Modern Eukaryotic. *Science* 305:354–360.

446   11.   Keeling PJ (2004) Diversity and evolutionary history of plastids and their hosts. *Am J Bot*
447   91(10):1481–1493.

448   12.   Sharpe, PJH, DeMichele, DW (1977) Reaction Kinetics of Poikilotherm Development.
449   *J. Theor. Biol.* 64, 649–670 .

450   13.   Schoolfield RM, Sharpe PJH, Magnuson CE (1981) Non-linear regression of biological
451   temperature-dependent rate models based on absolute reaction-rate theory. *J Theor Biol*
452   88:719–731.

453   14.   Eppley RW (1972) Temperature and phytoplankton growth in the sea. *Fish Bull* 70(4):1063–
454   1085.

15. Chen B, Laws E a. (2017) Is there a difference of temperature sensitivity between marine
phytoplankton and heterotrophs? *Limnol Oceanogr* 62:806–817.

16. Kremer CT, Thomas MK, Litchman E (2017) Temperature- and size-scaling of phytoplankton
population growth rates: Reconciling the Eppley curve and the metabolic theory of ecology.
Limnol Oceanogr 62:1658–1670.

17. Quigg A, et al. (2003) The evolutionary inheritance of elemental stoichiometry in marine
phytoplankton. *Nature* 425(6955):291–4.

18. Shuter B (1979) A model of physiological adaption in unicellular algae. *J Theor Biol* 78:519–
552.

19. Raven J A., Geider RJ (1988) Temperature and algal growth. *New Phytol* 110:441–461.

20. Raven J A (1976) The Quantitative Role of ' Dark ' Respiratory Processes in Heterotrophic
and Photolithotrophic Plant Growth. *Ann. Bot.* 40:587–602.

21. Geider RJ, Osborne B A (1989) Respiration and microalgal growth : a review of the
quantitative relationship. *New Phytol* 112:327–341.

22. Padfield D, Yvon-Durocher G, Buckling A, Jennings S, Yvon-Durocher G (2016) Rapid
evolution of metabolic traits explains thermal adaptation in phytoplankton. *Ecol Lett* 19:133–
142.

23. Schaum C-E, Buckling A, Smirnoff N, Studholme DJ, Yvon-Durocher G (2018)
Environmental fluctuations accelerate molecular evolution of thermal tolerance in a marine
diatom. *Nat Commun*:1–14.

24. Schaum C-E, et al. (2017) Adaptation of phytoplankton to a decade of experimental warming
linked to increased photosynthesis. *Nat Ecol Evol* 1:0094.

25. Marañón E, Lorenzo MP, Cermeño P, Mouriño-Carballido B (2018) Nutrient limitation
suppresses the temperature dependence of phytoplankton metabolic rates. *ISME J*.
doi:10.1038/s41396-018-0105-1.

26. Moore CM, et al. (2013) Processes and patterns of oceanic nutrient limitation. *Nat Geosci*
6(9):701–710.

27. Bruland KW, Rue EL, Smith GJ, DiTullio GR (2005) Iron, macronutrients and diatom blooms
in the Peru upwelling regime: Brown and blue waters of Peru. *Mar Chem* 93:81–103.

28. Townesend DW, Cammen LM, Holligan PM, Campbell DE, Pettigrew NR (1994) Causes and
consequences of variability in the timing of phytoplankton blooms. *Deep Res Part I*
41(5):747–765.

29. Pingree RD, Pugh PR, Holligan PM, Forster (1975) Summer phytoplankton blooms and red
tides along tidal fronts in the approaches to the English Channel. *Nature* 258:672–677.

30. Baranyi J, Roberts T a. (1994) A dynamic approach to predicting bacterial growth in food. *Int
J Food Microbiol* 23:277–294.

31. Raven J A (1976) The Quantitative Role of ' Dark ' Respiratory Processes in Heterotrophic
and Photolithotrophic Plant Growth. *Ann. Bot.* 40:587–602.

32. Ho T, et al. (2003) the Elemental Composition of Some Marine Phytoplankton. *J Phycol*
39:1145–1159.

33. Eilers PHC, Peeters JCH (1988) A model for the relationship between light intensity and the
rate of photosynthesis in phytoplankton. *Ecol Modell* 42:199–215.

34. Edwards KF, Thomas MK, Klausmeier C a., Litchman E (2016) Phytoplankton growth and the
interaction of light and temperature: A synthesis at the species and community level. *Limnol*
Oceanogr 61:1232–1244.

35. Falkowski PG, Gan R, Wyman K (1985) Growth-irradiance relationships in phytoplankton .
Limnol. Oceanogr. 30(2):311–321.

36. Muggeo VMR (2003) Estimating regression models with unknown break-points. *Stat Med*
22:3055–3071.

37. Davies RB (1977) Hypothesis Testing when a Nuisance Parameter is Unidentified Under the
Alternative. *Biometrika* 64:247–254.

Fig. 1. Thermal tolerance curves for 18 species of marine phytoplankton. (A) Thermal reaction norms for all 18 species used in this study. The data points presented are the natural logarithm of per capita growth rate, μ, for each replicate (n= minimum of 3 technical replicates per assay temperature for each species). The fitted lines are from the predicted random effect of species derived from non-linear mixed effects modelling with a Sharpe-Schoolfield model. The vertical dashed lines correspond with the optimal temperatures of growth. **(B)** Boxplot distributions of optimal growth temperatures (T_{opt}) and maximum temperatures of growth (T_{max}) across all 18 species. **(C)** Boxplot distribution of growth activation energy, or temperature dependence (E_a^μ), across all 18 species (Table 2, SI). The bold horizontal line corresponds to the median value, the top and bottom of the box correspond to the 75[th] and 25[th] percentiles and the whiskers extend to the largest and smallest values no greater or less than 1.5 × the interquartile range, beyond which the points are plotted as outliers.

Fig. 2. Thermal performance curves for respiration and gross photosynthesis in 18 species of marine phytoplankton. (A) Metabolic thermal performance curves for all 18 species used in this study. Green colouring denotes gross photosynthesis, red colouring denotes respiration. The data points presented are the natural logarithm of mean metabolic, with error bars denoting ± s.e.m (n = minimum of 3 biological replicates per response for each species). The fitted lines for each species are from the random effects of a non-linear mixed effects model fitted to the rate data using the Sharpe-Schoolfield equation (see Methods). The vertical dashed lines correspond with the optimal temperatures for each metabolic flux, with the black dashed line added to show optimal growth temperature. **(B, C and D)** Boxplots showing the distribution of the estimated values for activation energy (E_a), optimal temperature (T_{opt}) and deactivation energy (E_h) for photosynthesis and respiration across the 18 species (Tables 4 and 5, SI). The bold horizontal line corresponds to the median value, the top and bottom of the box correspond to the 75th and 25th percentiles and the whiskers extend to the largest and smallest values no greater or less than 1.5 × the interquartile range, beyond which the points are plotted as outliers.

Figure. 3. Carbon-use efficiency breakpoints constrain the optimal temperature of growth. (A)
Segmented linear regression models fitted to the predicted carbon use efficiency (CUE), derived from
the thermal performance parameters of respiration and photosynthesis for each species (Fig 2). The
modelled response is presented here alongside the calculated mean CUE at each assay temperature,
with with error bars denoting ± s.e.m (n = minimum of 3 biological replicates per response for each
species). The dashed vertical dashed blue line represents the predicted break-point in the model,
where there was a significant change in the slope of the CUE thermal response. The dashed vertical
black line represents the estimate optimal temperature of growth (Fig.1). In most cases this either
coincides with the break-point, falling within the 95% CIs of the break-point, or was lower than the
break-point. **(B)** Boxplots showing the distribution of the estimated values for the CUE break-point
temperature, optimal temperature of gross photosynthesis (T_{opt}^{P}) and optimal temperature of growth
(T_{opt}^{μ}) across the 18 species (Tables 2, 4 and 5, SI). The bold horizontal line corresponds to the
median value, the top and bottom of the box correspond to the 75[th] and 25[th] percentiles and the
whiskers extend to the largest and smallest values no greater or less than 1.5 × the interquartile
range, beyond which the points are plotted as outliers. **(C)** The significant coupling between the CUE
and T_{opt}^{P}, illustrating that the sharp declines in CUE are determined by the universal metabolic
constrains identified in Fig.2.

Figure. 4. The temperature dependence of the carbon-use efficiency. (A) A scatterplot showing
the relationship between the natural logarithm of the carbon-use efficiency (CUE) and standardised
Boltzmann temperature up to the CUE break-point (Fig.3) for the pooled dataset of 18 species, where
$T_c = 20°C$ and k is the Boltzmann constant (8.62×10^{-5} eV). The fitted line represents the fixed
effect of a linear mixed effects model fitted to the data using the Boltzmann-Arrhenius equation (see
Methods). Values of ln(CUE) have been standardised by dividing by the species-specific intercept
derived from the random effects of the mixed effects model. This standardisation was for visualisation
of the data only. The plot demonstrates that the CUE decreases up to the CUE break-point
temperature with a consistent temperature dependence, equating to an average activation energy (
E_a^{CUE}) of -0.12eV. (B) Boxplot of the species-specific E_a^{CUE} values derived from the linear mixed
effects model. The bold horizontal line corresponds to the median value, the top and bottom of the
box correspond to the 75[th] and 25[th] percentiles and the whiskers extend to the largest and smallest
values no greater or less than 1.5 $\times$ the interquartile range, beyond which the points are plotted as
outliers.

570