Programmed knock-out of the *omega-1* gene of the blood fluke *Schistosoma mansoni*

Wannaporn Ittiprasert[1], Victoria H. Mann[1], Shannon E. Karinshak[1], Apisit Chaidee[1,2], Toshihiko Tanno[3], Chutima Kumkhaek[4], Pannathee Prangtaworn[1,5], Margaret Mentink-Kane[6], Geetha Sankaranarayan[7], Avril Coghlan[7], Christina J. Cochran[1], Gabriel Rinaldi[7], Patrick Driguez[7], Nancy Holroyd[7] Alan Tracey[7], Rutchanee Rodpai[1,2], Bart Everts[8], Cornelis H. Hokke[8], Karl F. Hoffmann[9], Matthew Berriman[7], Paul J. Brindley[1,*]

1. Department of Microbiology, Immunology & Tropical Medicine, & Research Center for Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington University, Washington, D.C. 20037, USA

2. Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

3. Department of Surgery and the Institute of Human Virology, University of Maryland, Baltimore, MD 21201, USA

4. Cellular and Molecular Therapeutics Laboratory, National Heart, Lungs and Blood Institute, National Institutes of Health, Bethesda, MD 20814 USA

5. Department of Parasitology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand 10700

6. Schistosomiasis Resource Center, Biomedical Research Institute, Rockville, MD 20850, USA

7. Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, CB10 1SA, UK

8. Department of Parasitology, Leiden University Medical Center, Leiden, Netherlands

9. Institute of Biological, Environmental & Rural Sciences (IBERS), Aberystwyth University, Aberystwyth, SY23 3DA, UK

Running title   Genome editing of schistosome *omega-1*

Keywords: *Schistosoma mansoni*; genome editing; Cas9; omega1; ribonuclease; flatworms; Platyhelminthes; functional genomics; lentivirus; homology directed repair; non-homologous end joining; double stranded break; Th2 phenotype; monocytic macrophage cell; granuloma;

*Correspondence, Paul Brindley, pbrindley@gwu.edu

Abstract

Soluble egg antigen (SEA) and excretory-secretory (ES) products of the egg of *Schistosoma*
mansoni contain a glycosylated T2 family ribonuclease termed omega-1 (ω1). Following release
from the egg, ω1 instructs antigen presenting cells to induce naïve CD4$^+$ T cells to mature into T
helper 2 effectors that, in turn, establish the Th2 immunological phenotype characteristic of
schistosomiasis. Here, schistosome eggs were either transiently exposed to recombinant Cas9
complexed with a synthetic guide RNA (sgRNA) of 20 nt complementary to exon 1 of ω1 by
electroporation or transduced with pseudotyped lentivirus encoding Cas9 and this sgRNA.
Subsequently, some eggs also were transduced with a single stranded oligodeoxynucleotide
donor transgene that coded for a stop codon in all six frames, flanked by 50 nt-long 5'- and 3'-
homology arms matching the predicted Cas9-catalyzed double stranded break (DSB) within *ω1*.
CRISPResso analysis of amplicons spanning the predicted DSB revealed ~4.5% of the reads
(total reads >2×10^6 per treated sample) were mutated by insertions, deletions and/or
substitutions, with an efficiency for homology directed repair of 0.19% insertion of the donor
transgene. Messenger RNAs encoding *ω1* were reduced >80%, and soluble egg antigen (SEA)
from ω1-mutated eggs exhibited markedly reduced ribonuclease activity, indicative that
programmed Cas9 cleavage mutated the *ω1* gene. Following injection of schistosome eggs into
the tail vein of BALB/c mice, the volume of pulmonary granulomas surrounding wild type eggs
was 18-fold greater than ω1-mutated eggs, $6.21×10^{-2} ± 1.61×10^{-3}$ versus $0.34×10^{-2} ± 0.12×10^{-4}$
mm^3, respectively. In parallel, whereas wild-type SEA polarized Th2 cytokine responses
including IL-4 and IL-5 in human monocyte/T cell co-cultures, significantly reduced levels of
these cytokines followed the exposure to ω1-mutated SEA. In overview, programmed genome
editing was active in schistosomes, Cas9-catalyzed chromosomal breakage was repaired by
homology directed repair and/or non-homologous end-joining, and mutation of ω1 impeded the
capacity of schistosome eggs to drive Th2 polarization and to provoke formation of pulmonary
circumoval granulomas. These findings demonstrated for the first time the utility of
CRISPR/Cas9-based genome editing for functional genomics for schistosomes.

Introduction

Schistosomiasis is considered the most problematic of the human helminth diseases in terms of morbidity and mortality [1-4]. The past decade has seen major advances in knowledge and understanding of the pathophysiology, developmental biology, evolutionary relationships and genome annotation of the human schistosomes [5-16]. As with other diseases, adapting CRISPR/Cas9 genome editing for functional genomics of schistosomes will drive innovation to enable deeper knowledge of schistosome pathogenesis, biology and evolution [17], and provide stable site-specific gene mutation and phenotype in contrast to retroviral transgene based RNA interference [16]. The schistosome egg plays a central role both in the transmission of the disease and in pathogenesis [1]. The appearance of *S. mansoni* eggs by six to eight weeks following infection coincides with profound polarization to a granulomatous, T helper 2 (Th2) phenotype [18-22]. Several egg proteins have been characterized, with >1,000 identified in soluble extracts (SEA, soluble egg antigen) [23-26]. In the viable egg, about 30 of these soluble proteins are located outside the developing miracidium and encompass the complement of secreted antigens (egg-secreted proteins, ESP) that interact with host tissues to facilitate passage of the egg from the mesenteric veins to the lumen of the intestines [27].

The T2 ribonuclease ω1 (omega-1) is the principal Th2-inducing component of ESP with Th2-polarizing activity requiring both RNase activity and ω1 glycosylation [19, 20, 28]. This RNase is hepatotoxic [29], secreted by the egg into the granuloma, and regulates the pattern recognition receptor signaling pathways in dendritic cells that, in turn, prime Th2 responses from $CD4^+$ cells [30]. Secreted ω1 provokes granulomatous inflammation around eggs traversing the wall of the intestines, and trapped in hepatic sinusoids and elsewhere, driving fibrosis that eventually results as hepatointestinal schistosomiasis [1, 31]. Here, we investigated the feasibility of programmed genome editing using CRISPR/Cas9 [32, 33] in schistosomes, aiming to edit the *ω1* locus for both gene knockout and knock-in. The investigation revealed that programmable genome editing catalyzed by the bacterial endonuclease Cas9 was active in schistosomes, with chromosomal double stranded breaks (DSB) repaired by homologous recombination directed repair (HDR) using a donor, single stranded oligonucleotide template bearing short homology arms and/or by non-homologous end joining (NHEJ). The programmed mutagenesis decreased levels of ω1 mRNA and induced distinct *in vitro* and *in vivo* phenotypes, including a substantial loss of capacity of SEA from ω1-mutated eggs to polarize Th2 cytokine responses (IL-4 and IL-5) in co-cultured dendritic and T cells and loss of capacity to provoke formation of pulmonary granulomas in BALB/c mice. The findings highlighted the utility of programmed genome editing for functional genomics for schistosomes.

Results
Omega-1, a multicopy locus on chromosome 1 of *S. mansoni*

Five genomic copies of *ω1* (*omega-1*) on chromosome 1 were identified in the genome of *S. mansoni*, versions 5 and 7 (Fig. S1) [12]. A single copy of *ω1* selected for genome editing includes four exons separated by three introns and spans 2,569 nt (Smp_193860, Uniprot Q2Y2H0_SCHMA) (Fig. 1A). Other copies have discrete exon/intron structure (that may partly reflect imprecise annotation) (Fig. S1) although the predicted amino acid sequences of copies are highly conserved. The predicted open reading frame (ORF) of Smp_193860 encodes a T2 ribonuclease of ~15 kDa, which lacks 5′ exons coding for a conserved amino acids region 1 (CASI) catalytic region, although it does include the CASII, and is smaller than the 31 kDa reported for ω1 [34]. The sgRNA targeted residues 164 to 183 of the predicted exon 1 of Smp_193860 adjacent to the protospacer adjacent motif (PAM), AGG, with the predicted Cas9 cleavage site located at three residues upstream of this PAM. The AGG and the nucleotide sequence complementary to this sgRNA were present on two of the other copies of *ω1*, Smp_179960 and Smp_184360, which shared >99% identity to Smp_193860. The coding sequences of the gene copies of *ω1* differed by several substitutions (Fig. S2, panels A-D).

NHEJ pathway genes in schistosomes
Artemis and DNA-PKcs are essential NHEJ factors in vertebrates [35, 36]. By searching for *S. mansoni* genes with the same Pfam domain content known for human NHEJ pathway genes, candidates for homologues six of seven human NHEJ pathway genes in *S. mansoni* were identified (Table S1) although a putative homologue of Cernunnos/XLF was not apparent in the genome of *S. mansoni* [36]. The human Cernunnos/XLF gene includes the Pfam XLF domain (PF09302), but this domain appears to be absent from genes in any flatworm species using the dataset of the International Helminth Genomes Consortium [37].

Site-specific integration of exogenous DNA confirmed CRISPR-Cas9 activity in schistosomes
The activity and efficiency of CRISPR/Cas9 to edit the schistosome genome, by targeting the ω1 locus, was explored using two approaches. First, a ribonucleoprotein complex (RNP) comprised of sgRNA mixed with recombinant Cas9 endonuclease was delivered into schistosome eggs isolated from livers of infected mice (egg preparation termed 'LE', liver eggs) by square wave electroporation. In addition, Homology Directed Repair (HDR) of CRISPR/Cas9-induced DSBs at the *ω1* locus in the presence of a donor DNA template was investigated [38-40]. Therefore, a single stranded oligodeoxynucleotide (ssODN) of 124 nt in length was delivered to some LE as a template for HDR of chromosomal double stranded breaks (DSBs) (Fig. 1B). The ssODN included a short transgene coding for six stop codons flanked by 5'- and 3'-homology arms, each arm 50 nt in length, complementary to the genome sequence of exon 1 on both the 5′ and 3′ sides of the programmed Cas9 cleavage site (Fig. 1A, B). After 72 hours in culture, total RNA was recovered from LE to quantify the ω1expression. In a second approach, LE were maintained for 24 hours in culture medium containing pseudotyped pLV-ω1X1 virions [41] and transfected with the ssODN repair template (above) by electroporation. Specifically, the pLV-ω1 plasmid encoding Cas9 driven by the mammalian translational elongation factor 1 (tEF1) promoter, and the sgRNA (20 nt) targeting exon 1 driven by the human U6 promoter [42], was assembled in the lentiviral vector pLV-U6g-EPCG (Sigma) and termed pLV-ω1X1 (Fig. 1C).

146
Given that the donor ssODN included a short transgene that would facilitate genotyping, PCR
was performed using template genomic DNAs from the CRISPR/Cas9-treated LE [38] to reveal
the extent by which knock-in (KI) had occurred. A forward primer termed Smω1X1-6stp-cds-F
specific for the ssODN transgene was paired with three discrete reverse primers, termed Smω1-
R1, Smω1-R2 and Smω1-R3, at increasing distance from the predicted HDR insertion site in *ω1*
(Table S2). Amplicons of the expected sizes of 184, 285 and 321nt were observed in genome-
edited eggs but not from control eggs (Fig. 2A, B; Fig. S3), a diagnostic pattern indicative of the
insertion of the ssODN transgene into ω1 and, in turn, indicating that the resolution of the DSB
at the ω1 locus from CRISPR/Cas9 had been mediated by HDR. Amplification using a control
primer pair, termed Smω1-control-F/R, which spanned the predicted DSB also was carried out,
yielding amplicons of the expected size of 991 nt. Similar findings were observed with genome
editing delivered by RNPs and by LV (Fig. S3). Sanger sequence analysis of the knocked-in
amplicons (KI-R1, KI-R2 and KI-R3) confirmed the presence of the transgene inserted into the
ω1 locus at the site predicted for programmed Cas9 cleavage (Fig. 2C).

Programmed mutations detected in exon 1 of *ω1*
The activity of CRISPR/Cas9 was first evaluated by a quantitative PCR (qPCR) approach that
relies on the inefficient binding of a primer overlapping the gRNA target, i.e. where mutations
are expected to have occurred, compared to the binding efficiency of flanking primers, i.e.
outside the mutated region [43, 44]. The overlapping (OVR) primer pair shared the reverse
primer with OUT primer (Fig. S4). Genomic DNA template was used for qPCR to quantify the
efficiency of CRISPR-mediated mutagenesis at the target locus; the ratio between the OVR
products and OUT products provides an estimate of the frequency of CRISPR-induced indels in
the gRNA sequence. We observed that there were 2.2, 6.3, 3.7 and 11.4% estimated indels in
genomic DNAs of gRNA/Cas9 RNP complex treated, gRNA/Cas9 RNA complex with ssODN
treated, pLV-ω1X1 treated and pLV-ω1X1 with ssODN treated samples. Differences were not
apparent among the control groups, untreated LE. Opti-MEM only, heated- killed pLV-ω1X1
with and without donor ssODN, and ssODN only.
174
In addition, to identify and quantify the scope and nature of the mutation(s) that arose in the
genome of *ω1* gene edited-eggs, we constructed amplicon NGS Illumina barcoded libraries from
pooled genomic DNAs of six independent exposures of LE to pLV- ω1X1 virions following by
transfection with the donor ssODN. The amplicon libraries were subjected to deep sequencing on
the MiSeq Illumina platform. Interrogation of the nature and impact of CRISPR/Cas9 at *ω1* was
undertaken with the CRISPResso pipeline [45]. At the outset, ~2 million reads from the
GENEWIZ reads of the pLV-ω1X1+ ssODN group were compared to the reference sequence,
which revealed numerous deletions proximal to the programmed mutation site in the
Smp_193860 locus (Fig. 2D). Subsequently, CRISPResso compared >56 million sequenced
reads to the reference sequence of the Smp_193860 locus (Table S3). The analysis revealed that
71% were WT (unmodified DNA) whereas 29 % reads exhibited apparent mutations (Fig. 2E)
across the 202 bp amplicon, with 0.13% insertions, 0.58% deletions and 28.2% substitutions
(sample 9 in Table S3). In contrast, in a control eggs-only sample, 76% were WT, and 24% of
reads exhibited apparent mutations, with 0.14% insertions, 0.33% deletions, and 24.0%
substitutions (Fig. 2E; sample 2 in Table S3). Thus, subtracting the rate of apparent mutations in
the control, we estimate that approximately 0.25% and 4.2% of reads in the experimental sample
carried programmed CRISPR-induced deletions and substitutions, respectively.

INDELs of 1-2 bp (or multiples thereof) in coding DNA cause frame-shifts, and consistent with
its higher rate of indels, the CRISPR-treated sample displayed a higher rate of frame-shifts
compared to a control eggs-only sample (2.0% for sample 9 versus 1.4% for sample 2; Table S3;
Fig. 2E). A caveat was that some apparent substitutions may reflect variation among the copies
of $\omega1$ rather than programmed mutations. The Illumina sequence reads revealed several common
variants, such as adjacent 'TA' substitutions instead of 'CC' at positions 152-153 of the
amplicon, which causes a change from Q to K at the amino acid level. Gene Smp_193860 has
'AT' at this position in the V5 assembly, as does mRNA XP_018647487.1 from the NCBI
database, whereas Smp_193860, Smp_184360 and Smp_179960 all have 'CC' at this position in
the V7 assembly (Fig. S4 panels A, B), whereas 'CC' was also found in KI fragments using
Sanger direct sequencing (Fig. 2C). Similarly, a second common dinucleotide substitution from
'AC' to 'TT' at positions 60-61 encodes a change from T to F. Both dinucleotide substitutions
were found together in 8% of reads in the control sample (sample 2, Table S3) and 4% of reads
in the treated sample (sample 9). Both these nonsynonymous substitutions are proximal to the
CAS-2 catalytic site, and hence may have functional significance (Fig. S2, panels C, D).
207
Along with the predicted NHEJ-catalyzed mutations, CRISPResso determined the rate of HDR-
mediated ssODN knock-in (Table S3; Fig. 2F). Here, insertion of the 24 bp transgene was
confirmed in 0.19% of the reads at the sgRNA programmed CRISPR/Cas9 target site (Fig. 2F;
sample 9 in Table S3). Some of the MiSeq reads containing the knock-in sequence also had the
'CC' to 'TA' substitutions at positions 152-153 and 'AC' to 'TT' substitutions at position 60-61
(Fig. S2; Table S3). This suggests that the indels catalyzed by NHEJ and/or KI by HDR occurred
in at more than one copy of $\omega1$, probably Smp_193860, Smp_184360 and/or Smp_179960, and
perhaps all three and also even in other homologues. The qPCR approach estimated the indel
frequency at 11.4% in the pLV-$\omega1$X1 with ssODN treatment group (Fig. S4), whereas
CRISPResso analysis of the pooled NGS reads indicated a frequency of indel mutation of <1%
(Table S3).

Programmed gene editing reduced the expression of ω1
LE transfected with RNP complexes, with or without ssODN, displayed a down regulation of the
$\omega1$-specific transcript of ~45% and 81%, respectively, compared to controls ($P \leq 0.05$; $n = 11$;
control groups of LE were treated with Opti-MEM medium alone (mock), Cas9 only, sgRNA
only, ssODN only and WT-LE. Additionally, LE transduced with pLV-$\omega1$X1 virions with or
without ssODN showed a reduction of the $\omega1$-specific transcript of 67% and by 83%
respectively, when compared to controls (Fig. 3A). LE exposed to heat-inactivated virions with
ssODN, LE exposed to ssODN only, LE exposed to Opti-MEM only, and WT-LE all were
included as controls. Similar outcomes were seen in all biological replicates undertaken ($n =11$).
This outcome indicated that resolution of chromosomal DSB by NHEJ plus HDR provided
enhanced programmed gene knockout compared to NHEJ-mediated chromosomal repair alone.
Nevertheless, both RNPs and pLV virions efficiently delivered programmed gene editing to
schistosomes but pLV virions delivered enhanced performance with a stronger gene silencing, in
particular when the donor repair template was provided (Fig. 3A). When examined at later time
points (days 5 and 7 following manipulation of the LE), further reduction in $\omega1$ abundance was
not apparent (Fig. 3B). We decided to investigate whether silencing of $\omega1$ was associated with a
reduction in the copy number of this multi-copy locus due to large DNA deletions as has been
described for other helminth species [39] A relative copy number estimation by qPCR performed

as described [46] did not show significant differences in the number of gene copies of ω1 among
the genome-edited eggs and controls (Fig. S5).
239
Diminished ribonuclease activity in CRISPR/Cas9 mutated Soluble Egg Antigen
preparation (SEA)
The ribonuclease activity of the ω1 glycoprotein in SEA is associated with the Th2-polarizing
immune response that characterizes the appearance of schistosome egg granulomata [19, 22].
Ribonuclease activity of SEA from control and experimental groups on substrate yeast RNA was
investigated following CRISPR/Cas9 programmed mutation of *ω1* mediated by the RNP and by
the pLV approaches with or without ssODN (genome editing condition that led to the highest
level of mRNA reduction) [20, 47]. Intact yeast RNA was evident in the DNase-RNase free-
condition (negative control), indicative of absence of RNase activity in the reagents (200 ng
yeast RNA at the outset). There was near complete hydrolysis of yeast RNA following exposure
to RNase A (positive control); ~1.4 ng RNA remained intact. Wild type SEA exhibited marked
RNase activity against the yeast RNA; ~70 ng RNA remained after one hour, corresponding
to >60% digestion. Incubation of the RNA with Δω1-SEA from the experimental groups, RNP,
RNP+ssODN, pLV-ω1X1, and pLV-ω1X1+ssODN, resulted ~30% substrate digestion, with
124, 140, 135 and 153 ng RNA remaining, respectively. All conditions for programmed genome
editing resulted in less digestion of the yeast RNA than for WT SEA ($P \leq 0.0001$) (Fig. 3C, D).
Moreover, the Δω1-SEA with programmed knock-in exhibited less RNase activity than Δω1-
SEA prepared without the donor ssODN repair template ($P \leq 0.01$).
258
Depleting SEA of ω1 down-regulated inflammatory activities of macrophages to induce
Th2 response
The ω1 ribonuclease alone is capable of conditioning human monocyte-derived dendritic cells to
drive Th2 polarization [22] and enhanced CD11b[+] macrophage [30] modulation of intracellular
toll like receptor (TLR) signaling. ω1 ribonuclease inhibited TLR-induced cytokine production
of IL-1β and shifted the TLR signaling outcome towards an anti-inflammatory posture via
mannose receptor (MR) and dectin [22, 48, 49]. A co-culture of a human monocytic cell line,
THP-1, and a human CD4[+] T cell line, THP-1, was employed to study the interaction of antigen
presenting cells and T cells [50, 51]. First, the THP-1 cells were activated with PMA for 48
hours, then pulsed with SEA or Δω1–SEA for 48 hours, after which the Jurkat CD4[+] T cells were
added to the wells. Thereafter, the co-culture continued for 72 hours. Th2 cytokines and other T
helper cells cytokines including IL-4, IL-5, IL-13, IL-2, IL-6, IL-10, TNF-α and IFN-γ were
quantified in culture supernatants of the co-cultured cells (Fig. 4). SEA from ω1-mutated eggs
stimulated reduced levels of Th2 cytokines; IL-4 and IL-5 in comparison to wild-type SEA ($P \leq$
0.01). There also was less IL-13 but this was not statistically significant (Fig. 4). Production of
IL-6 and TNF-α were reduced ($P \leq 0.01$, whereas significant differences in IL-10. IL-2 and IFN-
γ were not apparent between the WT- and mutant-SEA co-cultures.
276
Granulomatous inflammation markedly reduced in lungs of mice injected with Δω1 eggs
Following the entrapment of eggs in the intestines, liver and eventually lungs, the glycosylated
ω1 ribonuclease represents the principal stimulus that provokes the circumoval granuloma,
necessary for extravasation of the eggs [52]. A long established model of the schistosome egg
granuloma employs tail vein injection of eggs into mice, which leads to formation of circumoval
granuloma in the mouse lung [53-55]. The latter approach has been extensively employed for

immunopathogenesis-related studies of ω1 [16]. Accordingly, to elicit circumoval granulomas,
~3,000 WT or Δω1 LE were injected into the lateral vein of the tail of BALB/c mice, and the
mice euthanized 10 days later. The entire left lung was removed, fixed, sectioned, and stained for
histological analysis (Fig. 5). Representative digital microscopic images of whole mouse lungs
acquired through the 2D ultra-high-resolution digital scanner from Leica Aperio Slide Scanner
are shown in Fig. 5, panels A-F. At low magnification (2×), severe and widespread pulmonary
inflammation was visible in the mice exposed to WT eggs compared to Δω1eggs. In addition,
more intense and dense focal circumoval inflammation was induced by WT compared to Δω1
eggs (Fig. 5A). Granulomas were not seen in control naïve mice not exposed to schistosome eggs
(Fig. 5C). At 20× magnification, considerable disparity in volume of the circumoval granulomas
was observed for WT versus Δω1 LE (Fig. 5A1-A2, 5D, 5E vs Figs. 5B1-B2, 5F, G). The
volume of granulomas surrounding single schistosome eggs was quantified; those surrounding
WT eggs in lungs of the mice was 18-fold greater than for ω1-mutated eggs, means $6.21 \times 10^{-2} \pm$
1.61×10^{-3} mm^3 and $0.34 \times 10^{-2} \pm 0.12 \times 10^{-4}$ mm^3 (17-26 granulomas per mouse), respectively ($P <$
0.0001) (Fig. 5H). The experiment was repeated, with 3-4 mice per group, with similar findings.
These findings documented a marked deficiency in the induction of pulmonary granulomas by
the Δω1 compared to WT eggs of *S. mansoni*.
300
GenBank/EMBL/DDBJ

Database accession Bioproject; PRJNA415471,
https://www.ncbi.nlm.nih.gov/bioproject/415471. Biosample SAMN07823308; *Schistosoma*
mansoni omega1 gene editing by CRISPR/Cas9
SAMN07823308 • SRS2756667 • Sequencing data for the mutation profiles are available at the
Sequence Read Archive (SRA) accessions SRR6374209, SRR6374210.
307

Discussion

In the current study, programmed CRISPR/Cas9 genome editing in *Schistosoma mansoni* was undertaken to investigate its feasibility in a representative platyhelminth species. Genome editing was successfully achieved as evidenced by site-specific mutations in the ω1 locus in chromosome 1. The chromosomal lesion induced by CRISPR/Cas9 was repaired by the non-homologous end joining (NHEJ) pathway [56] in the absence of a donor oligonucleotide, and was repaired by homology directed repair (HDR) when a single stranded donor oligonucleotide was provided as the repair template and/or by NHEJ [57-59]. To our knowledge, this report provides the first description and characterization of CRISPR/Cas programmed genome engineering in schistosomes and in any species of the phylum Platyhelminthes.

At least five intact copies of ω1 are localized on chromosome 1 of *S. mansoni*, and although the open reading frame and amino acid sequences of these copies are conserved, the copies of *ω1* display variation in exon/intron structure (although some of these differences may be due to imprecise annotation). In order to edit the *ω1* locus, we decided first to deliver the CRISPR/Cas9 components to the mature egg stage (liver eggs, LE) of the schistosome since the ω1 T2 ribonuclease is tightly expressed only at this stage of schistosome development [20, 27, 29]. The LE preparation includes a spectrum of egg development ranging from newly laid, immature, fully mature and dead eggs, although the proportion of LE that is mature and ready to hatch when placed in water is > 70% [11]. When released from the adult female schistosome into the mesenteric veins, the schistosome ovum contains a single celled zygote surrounded by 30 to 40 vitelline cells [27]. By six or so days later, the miracidium within the eggshell is a mobile, ciliated larva comprised of organs, tissues, muscles and nerves [25, 60-62].

Somatic genome engineering was undertaken by delivering the CRISPR/Cas9 machinery by two methods to schistosome eggs recovered from livers of experimentally infected mice. First, CRISPR/Cas9 was delivered as a ribonucleoprotein particle (RNP) complex consisting of recombinant Cas9 from *Streptococcus pyogenes* complexed with a single guide RNA (sgRNA) specific for exon 1 of the ω1 locus. The second approach was based on lentiviral virion delivery where both the Cas9 and sgRNA were included as cargo in the construct. Delivery of CRISPR/Cas9 by RNP, by lentiviral virions, and other methods have been thoroughly employed in cells lines, tissues and entire organisms [63-66]. Immediate editing leading to DSB is an advantage with delivery of RNPs in cells whereas the short half-life of the RNP components may be disadvantageous. By contrast, delivery of the CRISPR components on plasmids and by viral mediated infection can provide more sustained activity, transgene integration in non-dividing cells, and other attributes [33, 67]. With mature schistosome eggs, LV-mediated delivery was expected to increase the performance and efficiency of the somatic genome editing given that the provirus integrates into the genome after infection, the transgene would be transmitted by mitosis, and extended expression would be feasible. RNPs were delivered to LE by square wave electroporation, an established delivery method for this developmental stage [68]. For LV, we used VSVG-pseudotyped HIV virions, since the lentiviral provirus will integrate into schistosome chromosomes [41]. Moreover, delivery by LV provides a hands-free approach that enables scaling of gene editing and potentially to reach less accessible and/or differentiated cells.

Genomic DNAs recovered from ~ 10,000 CRISPR/Cas9-treated LE included many thousands of
copies of the schistosome genome, and in addition, several copies of the gene itself are found in
each genome. Analysis assisted by the CRISPResso pipeline of the deep sequence reads of
amplicons spanning the predicted DSB site revealed that ~6% of the reads were mutated by
insertions, deletions, and substitutions, with an efficiency for HDR of 0.18% insertion of the
donor transgene. The target locus was mutated by knock-in (KI) of a ssODN repair template
bearing short homology arms to mediate homology directed repair following DSB at $\omega 1$.
360
Concerning the mutation profile of CRISPR/Cas9-induced repair, the numerous substitutions
compared to deletions and insertions was noteworthy. On the other hand, this is a multi-copy
gene and some substitutions may reflect the multi-family nature of ω1. The sequenced reads
from the amplicon NGS libraries revealed micro- and macro-variability in the sequences at the
following repair of the DSB. Given that this genome editing investigation involved multicellular
targets, it is an example of somatic (rather than germ line) genome editing. [67]. Given also the
multiple copies of $\omega 1$, the numerous cells in each of the ~10,000 LE, the spectrum of
development of these eggs, the diverse mutation profile of NHEJ, and the other contributions to
variability (below), mosaicism of mutations was expected, i.e. some displayed HDR, some NHEJ
but not HDR, other cells and NHEJ and/or HDR, and many retained the wild type (non-mutated).
As well as the expected relative ease of access by the RNPs, virions and ssODN to the cells of
syncytial inner envelope compared to the cells within the miracidium, other factors may have
contributed to variability of CRISPR efficiency among the eggs and among the cells of a single
egg. Cells exposed to CRISPR/Cas9 components may not have been undergoing cell dividing;
HDR can proceed only at cell division, with the cell at late S phase or G2 following completion
of DNA replication and access to sister chromatids as the repair templates [69]. Otherwise,
NHEJ proceeds to repair the DSB. Whereas incubating the eggs in virions would be expected to
ensure that virions contacted every egg, the coverage of infection was not assessed, both for the
coverage by RNPs and the subsequent transduction by ssODN.
380
In addition to the mutations identified in the DNA we investigated their impact at the mRNA
level of ω1. The levels of ω1-encoding mRNA were reduced by up to 83% (relative to the
control), indicating that Cas9 catalyzed mutation of $\omega 1$, and that programmed Cas9-induced
double stranded chromosomal breaks had been resolved by non-homologous end joining (NHEJ)
and/or homology direct repair (HDR) events. The knock-in of a ssODN repair template in
cultured eggs induced 81-83 % reduction in $\omega 1$-specific mRNA levels, whereas a down
regulation of 67 to 83 % was observed following exposure to RNP/LV without ssODN. A
tentative explanation of the paradox of the >80% reduction in transcription of ω1 side-by-side
with <1 % efficiency in gene editing (based on NGS reads) may be tight tissue specific
expression of ω1, which only occurs within the syncytial 'inner envelope' of squamous,
epidermal-like cells surrounding the miracidium [60]. Expression of $\omega 1$ appears to be tightly
developmentally regulated; based on comparison of expression in the egg to expression in the
miracidium, immunostaining and transmission electron microscopy of the mature egg, all or
most expression of ω1 takes place in the mature egg and not in immature eggs or other larval or
adult stages [20, 25, 27, 29, 70]. Within the mature egg, the fully developed miracidium is
surrounded by a squamous, syncytial epithelium termed variously as the envelope, the inner
envelope, or von Lichtenberg's envelope [25, 27, 60, 61]. This syncytial layer is metabolically
active [27], and likely the site of synthesis of T2 ribonuclease ω1 that is released from the egg

[25, 27, 29], along with other secreted/ excreted proteins that help facilitate egress of the egg
from the blood vessels and through the wall of the intestine [27]. The virions may not have
infected many of the cells within mature eggs, because after entry into the egg the virus would be
expected to contact the syncytial envelope, rather than traveling further to directly contact cells
of the miracidium. That is, the efficiency of gene editing may not have been high across all cells
of an egg but may (due to ease of access to certain cells) have had highest efficiency in those
cells in which ω1 is expressed. This may explain the large reduction in *ω1* transcripts in the
CRISPR-treated eggs relative to the control eggs.

The inconsistency between the substantially reduced mRNA levels and the lower frequency of
mutations observed in MiSeq sequence data might also reflect CRISPR/Cas9 induced large size
deletions (>500 bp) that would include the primer sites and hence not be detected by the
sequencing approach (i.e. PCR-amplicon next generation sequencing) as reported in
Strongyloides stercoralis [39]. Large deletions would explain the transcript-knock down due to a
reduction in the copy number of the target gene. However, based on the qPCR findings
differences in *ω1* copy number were not evident among CRISPR/Cas9-treated LE and controls.
Notwithstanding, indels arising following Cas9-catalyzed DSBs frequently result in frameshift
mutations that, in turn, lead to nonsense-mediated mRNA decay (NMD) of the mutant transcript
[71], which effectively knocks out the gene [72].

Whereas wild-type SEA polarized Th2 cytokine responses including IL-4 and IL-5 in human
monocyte/T cell co-cultures, significantly reduced levels of these cytokines were observed when
the cells were exposed to ω1-mutated SEA. Moreover, following introduction of eggs into the
tail vein of mice, pulmonary circumoval granulomas were substantially reduced in size around
Δω1 eggs compared to granulomas and the inflammation provoked by wild type eggs. Whereas
this outcome extends earlier findings using lentiviral transduction of eggs of *S. mansoni* to
deliver microRNA-adapted short hairpin RNAs aiming to silence expression of ω1 [16], the
reduction in granuloma volume was far more prominent in the present study. Residual
granulomas around mutant eggs could be due to prostaglandin E_2 levels [73], in addition to
incomplete disruption of all copies of *ω1*. Given that the T2 ribonuclease ω1 is the principal type
2-polarizing protein among egg-secreted proteins [19, 20], our findings of a phenotype
characterized by absent or diminutive granulomas provide functional genomics support to this
earlier advance [22].
430
The findings confirmed that somatic genome editing of the schistosome egg, lead to knockout of
the ω1 T2 ribonuclease, and mosaicism of mutant and wild type cells within the schistosome
egg. The Cas9 nuclease catalyzes programmed cleavage of schistosome chromosomes, and the
nuclease and sgRNA can be provided from schistosome cells following infection with lentiviral
virions. The Cas9 gene edited schistosome eggs remained viable but exhibited loss of function of
ω1. Programmed mutation of *ω1* using CRISPR/Cas9 not only achieved our aim to establish
applicability and tractability of genome editing for functional genomics of schistosomes but also
demonstrated manipulation of a gene expressed in the schistosome egg, the developmental stage
of the pathogen central to the pathophysiology and immunobiology of schistosomiasis. This
report provides a blueprint that can be applied to other schistosome genes, and those of
trematodes and platyhelminths at large. In addition, future studies that combine pseudotyped
virions to deliver transgene-encoded programmed genome editing to the schistosome germ line,
to derive lines of mutant parasites [11] would enable more comprehensive understanding of the

pathogenesis and pathophysiology of this neglected tropical diseases pathogen that will lead to
the discovery of novel strategies for control.
446
Materials and Methods

Ethics statement

Mice experimentally infected with *S. mansoni*, obtained from the Biomedical Research Institute,
Rockville (BRI), MD were housed at the Animal Research Facility of the George Washington
University Medical School, which is accredited by the American Association for Accreditation
of Laboratory Animal Care (AAALAC no. 000347) and has an Animal Welfare Assurance on
file with the National Institutes of Health, Office of Laboratory Animal Welfare, OLAW
assurance number A3205-01. All procedures employed were consistent with the Guide for the
Care and Use of Laboratory Animals. The Institutional Animal Care and Use Committee of the
George Washington University approved the protocol used for maintenance of mice and
recovery of schistosomes. Studies with BALB/c mice involving tail vein injection of schistosome
eggs and subsequent euthanasia using overdose of sodium pentobarbitone was approved by the
IACUC of the Biomedical Research Institute (BRI), protocol 18-04, AAALAC no. 000779 and
OLAW no. A3080-01.

Schistosome eggs

Mice were euthanized seven weeks after infection with *S. mansoni*, livers were removed at
necropsy, and schistosome eggs recovered from the livers, as described [74]. The eggs termed
'LE' were maintained in RPMI medium supplemented with 10% heat inactivated fetal bovine
serum (FBS), 2% streptomycin/penicillin and protease inhibitor cocktail (Sigma) at 37°C under
5% CO_2 in air [75, 76]. Polymyxin B to 10 μg/ml was added to the cultures twice daily to
neutralize lipopolysaccharide (LPS) [77]. Soluble egg antigen (SEA) was prepared from these
eggs, as described [23, 53]. In brief, the homogenate of eggs in 1×PBS containing protease
inhibitor cocktail (Sigma) was frozen and thawed twice, clarified by centrifugation at 13,000
rpm, 15 min, 4°C, the supernatant passed through a 0.22-μm pore size membrane. Protein
concentration of the supernatant (SEA) were determined by the Bradford Protein Assay [78] and
aliquots of the SEA stored at -80°C.

Guide RNAs, Cas9, and single stranded DNA repair template

Single guide RNA (sgRNA) was designed using the web-based tools at
http://bioinfogp.cnb.csic.es/tools/breakingcas/ [79] to predict cleavage sites for the *Streptococcus*
pyogenes Cas9 nuclease within the genome of *S. mansoni*. The sgRNA targeted exon 1 of the ω1
gene, Gene ID Smp_193860, www.genedb.org, residues 164-183, adjacent to the protospacer
adjacent motif, AGG (Fig. 1A). This is a multi-copy gene with five copies of ω1 located in
tandem on chromosome 1. The Smp_193860 copy of ω1 includes four exons interspersed with
three introns (2,569 nt) (Fig. 1A). Synthetic gRNA (sgRNA), ω1-sgRNA was purchased from
Thermo Fisher Scientific (Waltham, MA). A double stranded DNA sequence complementary to
the sgRNA was inserted into lentiviral gene editing vector, pLV-U6g-EPCG (Sigma), which
encodes Cas9 from *S. pyogenes* driven by the eukaryotic (human) translation elongation factor 1

alpha 1 (tEF1) promoter and the sgRNA driven by the human U6 promoter (Fig 1C). (The pLV-
U6g-EPCG vector is tri-cistronic and encodes the reporter genes encoding puro and GFP, in
addition to Cas9, using the 2A peptide approach [80].) This gene-editing construct, targeting
exon 1 of $\omega1$ Smp_193860, was termed pLV-$\omega1$X1. A single stranded oligodeoxynucleotide
(ssODN), which included homology arms of 50 nt each in length at the 3' (position 131-180 nt)
and 5'(181-230 nt) flanks and a small transgene (5'-TAAGTGACTAGGTAACTGAGTAG-3',
encoding stop codons (six) in all reading frames) (Fig. 1B), was synthesized by Eurofin
Genomics (KY, USA). An oligonucelotide primer of the same 24 nt sequence was also employed
as the forward primer to investigate the presence of CRISPR/Cas9-programmed insertion of the
transgene [81] (Table S2).
496

Transfection of schistosome eggs with a Cas9/guide RNA complex

For the riboncleoprotein (RNP) complex of the $\omega1$-sgRNA and recombinant Cas 9, three µg $\omega1$-
sgRNA and 3 µg Cas9-NLS nuclease (Dharmacon, Lafayette, CO) were mixed in 100 µl Opti-
MEM (Sigma) to provide 1:1 ratio w/w RNP. The mixture was incubated at room temperature
for 10 min, pipetted into a 4 mm pre-chilled electroporation cuvette containing ~10,000 LE in
~150 µl Opti-MEM, subjected to square wave electroporation (one pulse of 125 volts, 20
milliseconds) (BTX ElectroSquarePorator, ECM830, San Diego, CA), thereafter incubated for 5
min at room temperature, and subsequently maintained at 37°C, 5% CO_2 in air for 3, 5 and 7
days. To investigate whether homology-directed repair (HDR) could catalyze the insertion of a
donor repair template, 3 µg ssODN was mixed with RNP and the LE before electroporation. In
the second approach (above), the ssODN was delivered to LE by electroporation at ~24 hours
after lentiviral transduction of the LE. The eggs were collected 3, 5 and 7 days later and genomic
materials recovered from LE. The negative controls included LE subjected to electroporation in
the presence of only Opti-MEM (mock), only Cas 9, only sgRNA, and only ssODN.
511

Transduction of schistosome eggs with lentiviral particles

Escherichia coli Zymo 5α (Zymo Research) cells were transformed with lentiviral plasmid pLV-
$\omega1$X1 and cultured in LB broth in 100 µg/ml ampicillin at 37°C, agitated at 225 rpm for ~18
hours, after which plasmid DNA was recovered (GenElute Plasmid purification kit, Invitrogen).
A lentiviral (LV) packaging kit (MISSION, Sigma-Aldrich) was used to prepare LV particles in
producer cells (human 293T cell line). In brief, 2.3×10^5 of 293T cells/well were seeded 6-well
tissue culture plate in DMEM supplemented with 10% heat- inactivated fetal bovine serum
(FBS), 2 mM L-glutamine, 2% penicillin/streptomycin and cultured at 37°C, 5% CO_2 for 18
hours. The producer cells were transfected using FUGENE 6 (Promega) with pLV-$\omega1$X1 and LV
packaging mix containing two additional plasmids, one to express HIV structural and packaging
genes and the other to express the pseudotyping envelope protein, VSVG (vesicular stomatitis
virus glycoprotein). Subsequently, the transfection mixture (37.5 µl; 500 ng plasmid DNA, 4.6 µl
packaging mix, 2.7 µl of FUGENE 6 in Opti-MEM) was dispensed drop wise into each well on
the plate. Sixteen hours later, the media were removed from the transfected cells, replaced with
pre-warmed complete DMEM, and cells cultured for 24 hours. The supernatant, containing
VSVG-pseudotyped LV particles was filtered through 22 µm pore size membranes [41], and
stored at 4°C. Additional pre-warmed complete DMEM was added to the well, for culture for a
further 24 hours. The supernatant was collected, as above, and added to the first supernatant, and

concentrated (Lenti-X concentrator, Takara Bio, Mountain View, CA). Virion titer was
estimated by two methods: first, by use of Lenti-X-GoStix (Takara Bio) to establish the presence
of functional virions at $>10^5$ infectious units (IFU)/ml, and second, by reverse transcriptase assay
[41, 82] to quantify levels of active virions. Virions with counts of ~4×10^6 count per minute
(cpm)/ml were aliquoted and stored at -80°C.

To transduce LE with LV, ~10,000 eggs were incubated for 24 hours in complete DMEM
containing 500 μl of ~4×10^6 cpm/ml VSVG-LV virions. Thereafter, the LE were washed three
times in 1×PBS and transfected with ssODN by square wave electroporation or further steps. LV
virions heat-killed by incubation at 70°C for two hours [41] with subsequent transfection with
the ssODN, transfection with ssODN in the absence of virions, and Opti-MEM only, all served
as negative controls.

PCR amplification of diagnostic transgene to detect knock-in into exon 1 of ω1

For each DNA sample, four separate PCR assays using four distinct primer pairs (Table S2) were
carried out. The first ω1 primer pair, to amplify locations 107-1,097 nt of Smp_193860, was
served as a positive control for the presence of genomic DNA with the Smp_193860 copy of *ω1*.
The other primer pairs shared one forward primer complementary to the knock-in 24 nt transgene
with three reverse primers, Sm ω1-R1, -R2 and -R3 at positions 346-364, 446-465 and 482-501
nt, respectively, bind complementary to three sites downstream the ω1 target site DSB (Fig. 2A;
Table S2) [81]. The PCR mix included 10 μl Green GoTaq DNA polymerase mix (Promega)
with 200 nM of each primer and 10 ng genomic DNA. Thermal cycling conditions involved
denaturation at 95°C, 3 min followed by 30 cycles of 94°C, 30 sec, 60°C, 30 sec and 72°C, 30
sec and a final extension at 72°C for 5 min. Amplicons were visualized following agarose gel
electrophoresis (1.2% agarose/TAE), and those of the expected sizes were recovered from gels
and cloned into pCR4-TOPO (Thermo Fisher). *E. coli* Zymo 5α competent cells were
transformed with the ligation products, three single colonies of each transformant on LB
agar/ampicillin were amplified, and insert sequences determined to confirm the presence and
knock-in of the transgene (Fig. 1C).

Illumina sequencing

Pooled LE DNA samples from six independent KI experiments of pLV-ω1X1 with ssODN were
used as the template to amplify the on-target DNA fragment using MiSeq primers (Fig. 2A) with
High Fidelity *Taq* DNA polymerase (Thermo Fisher). The expected size of the amplicon
flanking predicted DSB was 205 bp. Amplicons of this size were purified by Agencourt AMPure
XP (Beckman Coulter) and prepared for sequencing using the Illumina MiSeq platform. Paired-
end read libraries were constructed using the QIAseq 1-Step Amplicon Library Kit (Qiagen) and
GeneRead Adaptor I Set A 12-plex (Qiagen) using discrete index adaptors. These libraries were
pooled, and the library pool was quantified using the QIAseq Library Quant System (Qiagen).
Paired end, MiSeq Illumina sequencing of the pooled amplicons was carried out either at
GENEWIZ (South Plainfield, NJ) and/or the Wellcome Genome Center.

Samples (10) (Table S3) were multiplexed and each run on four MiSeq lanes. After sequencing,
the fastq files for each particular sample were merged. Samples 7 and 9 were prepared by a

slightly modified approach (higher PCR anneal temperature) and in a separate laboratory than
the other libraries, and (for as yet unclear reason) included an extra 10 bp at the start of the
MiSeq sequences, 'GTTTTAGGTC', present upstream of the 5' primer in the genomic DNA).
We trimmed this sequence from the reads using cutadapt v1.13 [83]. To detect HDR events,
CRISPResso v1.0.9 [45, 84] was employed using a window size of 500 bp (-w 500) with the
reference amplicon according to gene *Smp_193860* in the *S. mansoni* V7 assembly, and with the
--exclude_bp_from_left 25 and --exclude_bp_from_right 25 options in order to ignore the (24
bp) primer regions on each end of the amplicon when indels are being quantified. A window size
of 500 nt was employed include the entire amplicon. In order to search for HDR events,
CRISPResso checked for HDR events (using –e and –d options) in treatment groups including
controls. To infer frameshifts using CRISPResso the –c option was used, giving CRISPResso the
coding sequence from positions 42-179 of the amplicon. To confirm the insertions of the knock-
in sequences reported by CRISPResso (column L in Table S3), we took all insertions of 20-28 bp
reported by CRISPResso, and calculated their percent identity to the expected knock-in sequence
using ggsearch v36.3.5e in the fasta package [85], and an insertion was considered 'confirmed' if
it had ≥75% identity to the expected knock-in sequence.
588

Copy number estimation for ω1

A quantitative PCR to estimate the relative copy number of *ω1* was performed using Kapa qPCR
mastermix SYBRfast (KK4602) on 1 ng of gDNA templates isolated from control and test
samples, in 20 μl volumes. Primer pair OMGgRNA1F and OMGgRNA1R was used to amplify
the ω1 gRNA target region and SmGAPDH (Smp_056970) as reference single-copy gene
(primers shown in Table S2). The PCR efficiencies for primer pairs were estimated by titration
analysis to be 100% ±5 [86] and qPCRs were performed in triplicate, using 96-well plates, with a
denaturation step at 95°C of 3 min followed by 40 cycles of 30 s at 95°C and 30 s at 55°C, in
thermal cycler fitted with a real time detector (StepOnePlus, Applied Biosystem). The relative
quantification assay $2^{-\Delta\Delta Ct}$ method [87] was used to ascertain the relative copy number of *ω1*.
Relative ω1 copy number in CRISPR/Cas9 treated groups reflects the fold change of *ω1* copy
number normalized to the Sm GAPDH reference gene and relative to the untreated control group
(calibrator sample with relative *ω1* copy number = 1) [46].
602

Analysis of RNase activity of ω1

A stock solution of yeast RNA (Omega Bio-tek, Norcross, GA) was prepared at 1.0 μg/μl, 50
mM Tris-HCl, 50 mM NaCl, pH 7.0. Yeast RNA (200 ng) was incubated with 2 μg SEA from
control and experimental groups individually at 37°C for 60 min. (SEA investigated here, named
Δω1-SEA, was extracted LE transduced with pLV- ω1X1 virions and ssODN, pooled from six
biological replicates.) RNase A, an endoribonuclease from bovine pancreas (Thermo Fisher)
served as a positive control enzyme whereas yeast RNA in reaction buffer only served as the
negative control. The RNase activity of ω1 in wild type SEA or Δω1-SEA was analyzed by
visualizing and quantifying the substrate that remained following enzymolysis by agarose gel
electrophoresis and staining with ethidium bromide. The yeast RNA digestion by control SEAs
or Δω1-SEA were set up in triplicates, with quantity of residual RNA determined by
densitometry [47].
615

Schistosome egg induced primary pulmonary granuloma in mice

For induction of circumoval, egg-induced granulomas in the lungs of mice, 8 week old female [25] BALB/c mice were injected with 3,000 WT egg or $\Delta\omega1$-egg or 1×PBS as negative control by tail vein, as described [55]. The mice (6-9 mice/group) were euthanized 10 days later. For histopathological assessment of granuloma formation, the left lung was removed at necropsy and fixed in 10% formalin in pH 7.4 buffered saline for 24 hours, after which it was dehydrated in 70% ethanol, and clarified in xylene. The fixed lung tissue was embedded in paraffin and sectioned at 4-μm-thickness by microtome [55]. Thin sections of the left lung lobe were mounted on glass slides and fixed at 58-60°C. Subsequently, rehydrated sections were stained with hematoxylin-eosin (H&E) for evaluation of inflammatory infiltrates and cellularity of granulomas. The longest (R) and shortest (r) diameters of each granuloma containing a single egg were measured with an ocular micrometer, and the volume of the granuloma calculated assuming a prolate spherical shape, using $4/3 \, \pi \, Rr^2$ [25]. All granulomas in all slides from the left lung of the mice, 15 slides per treatment group, were measured, in total >100 granulomas from each treatment group. Digital images were obtained using a 2D glass slide digital scanner (Aperio Slide Scanner, Leica Biosystems, Vista, CA), and examined at high magnification using the Aperio ImageScope (Leica) software [54, 88].

Macrophage polarization by WT or $\Delta\omega1$-SEA and T-cell activation *in vitro*

Human monocytic THP-1 cells were maintained in Roswell Park Memorial Institute medium (RPMI) 1640 with L-glutamine, HEPES (Thermo Fisher Scientific) containing 10% (v/v) FBS with 4 mM glutamine, 25 mM HEPES, 2.5 g/L D-glucose at 37°C in 5% CO_2 in air. THP-1 cells were non-adherent cells. In a 6-well plate, THP-1 monocytes (3×10^5 cells in each well) were differentiated into macrophages (MΦ) by incubation in 150 nM phorbol 12-myristate 13-acetate (PMA) (Sigma) for 48 hours [89]. MΦ were treated with SEA (50 ng/ml) or $\Delta\omega1$-SEA (50 ng/ml) (from LE transduced with pLV-ω1X1 virions and ssODN) for 48 hours. The SEA or $\Delta\omega1$-SEA treated MΦ were collected and processed for RNA extraction to examine cytokine transcription. To investigate macrophage and T cell interactions, MΦ cells were pulsed with SEA or $\Delta\omega1$-SEA and thereafter co-cultured with Jurkat (human $CD4^+$) T cells. Nine $\times10^5$ Jurkat cells were added to MΦ and incubated for an additional 72 hours. Cell-free supernatants from the co-cultures were collected to quantify secretion of T helper cell cytokines including IL-4, IL-5, IL-13, IL-10, TNF-α, IL-6, IL-2 and IFN-γ by enzyme linked immunosorbent assay (ELISA) (Qiagen) [90]. The ELISA test was performed using three biological replicates, with positive control for each analytes provided in the kit (Qiagen).

Differential gene expression for ω1 mRNA and cytokine mRNAs

Total RNAs from either schistosome were extracted using the RNAzol RT reagent (Molecular Research Center, Inc), which eliminates contaminating DNA [91]. A NanoDrop ND-1000 spectrophotometer was used for the RNA concentration and purity ($OD_{260/280}$ ~2.0). RNA (500 ng) from was reverse transcribed using iScript Reverser Transcript (Bio-Rad) and the first strand cDNA was employed as template for qPCR using SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) and run in triplicates by iQ5 Real time PCR instrument (Bio-Rad).

To evaluate the levels of ω1 transcripts, RT-qPCR reaction mixtures included 2 μl first stand
cDNA, 5 μl SsoAdvanced universal SYBR Green Supermix, and 300 nM schistosome gene
specific primers. Table S2 provides details of the oligonucleotide primers. Thermal cycling
conditions were denaturation at 95°C for 30 sec, 40 amplification cycles each consisting of
denaturation at 95°C for 15 sec, annealing/extension at 60°C for 30 sec, and a final melting
curve. The output was analyzed employing the iQ5 software. The relative expression of genes
was calculated using ΔΔCt method and normalized to either schistosome GAPDH expression
[87]. Data are presented as transcript levels (three replicates) compared to the WT (100%) LE,
and fold change reported as mean relative expression ± SD.

Statistics

669   Means for experimental groups were compared to controls by one-way ANOVA, and, where
670   appropriate, by a two-tailed Student's *t*-test or Welsh's *t*-test (GraphPad Prism, La Jolla, CA). *P*
671   values of ≤ 0.05 were considered to be statistically significant.

Acknowledgements

We thank Dragana Jankovic, Alan Sher, Thomas Wynn, Michael Bukrinsky, Larisa Dubrovsky,
Arnon Jurberg, Robert Thompson, Meredith Brindley and Luca Pinello for advice and technical
assistance. Schistosome-infected mice and snails were provided by the NIAID Schistosomiasis
Resource Center of the Biomedical Research Institute, Rockville, Maryland through NIH-NIAID
Contract HHSN272201000005I for distribution through BEI Resources. These studies were
supported by Wellcome Trust Strategic Award number 107475/Z/15/Z (Hoffmann, K.F.,
Principal Investigator), award R21AI109532 (GR, PJB) award from NIAID, National Institutes
of Health, and in part by generous support from MaxMind Inc./ TJ Mather.

 Figure legends

Figure 1. Genomic structure of the locus encoding ω1 (omega-1) in the genome of
***Schistosoma mansoni,* and CRISPR/Cas9 encoding construct and guide RNA.** Exon-intron
map of the copy of *ω1* at GeneDB Smp_193860, showing the position of its four exons, three
introns and flanking region spanning 2,569 bp on chromosome 1 (panel **A**). Nucleotide sequence
in exon 1 indicating location and sequence of gRNA target site, predicted double stranded break
(DSB) (arrow), protospacer adjacent motif (PAM) (AGG, blue box), and 124-nucleotide
sequence of the single stranded DNA donor template provided for DSB repair by homologous
recombination. Homology arms of 50 nt span a central 24 nt of six-stop-codon transgene (**B**).
Linear map of pLV-ω1X1 showing position of regulatory and coding regions for CRISPR/Cas9
editing; the positions of human U6 promoter to drive gRNA, translational elongation factor EFα-
1 promoter driving Cas9 from *Streptococcus pyogenes*, and the left and right long terminal
repeats of the lentiviral vector derived from HIV-1 (**C**).

Figure 2. Programmed chromosomal break at *ω1* locus repaired by homologous
recombination from donor template. Schematic diagram to indicate positions of primer
binding sites (blue arrows), with the foreign gene cassette as the forward primer (Smω1X1-6-
stop codons cassette-F) paired with three discrete reverse primers, Smω1-R1, Smω1-R2 and
Smω1-R3 from the *ω1* locus and a primer pair for target amplicon NGS library amplification;
miSeq-F and miSeq-R (panel **A**). The control PCR amplicon was generated using the Sm**ω**1-
control-F and –R primers. The green box shows the location of 5' and 3' homology arms, red
box and arrow indicate the stop codon bearing transgene. **B**, PCR products visualized in ethidium
bromide-stained agarose gel demonstrating Cas9-catalyzed target site-specific insertional
mutagenesis in exon 1 of the *ω1* gene. Evidence for transgene knocked-in into programmed
target site revealed by amplicons of the expected sizes in lanes R1, R2 and R3, of 184, 285 and
321 bp, respectively (arrows at left) spanned the mutated site in the genomic DNAs pooled from
schistosome eggs, including a positive control flanking the insert site (991 bp). The control
DNA result shown in this gel was isolated from heat inactivated-pLV-ω1X1+ssODN treated LE.
Similar findings were obtained when programmed gene editing was executed by lentiviral virion-
delivered Cas9 and *ω1*-gRNA transgenes and by ribonucleoprotein complex (RNP) delivered by
square wave electroporation (supporting information). The non-KI control groups (sgRNA only,
heat inactivated pLV-ω1X1 only, ssODN only) showed similar results, with no amplicons by
stop cassette-KI primers with R1, R2 or R3 (Fig. S4). **C**, Multiple sequence alignments
confirmed the presence of the 24 nt transgene inserted precisely into the exon 1 of ω1 locus from
KI-R1, -R2 and -R3 fragments comparing with *ω1* wild type (WT). The white box on ω1-WT
indicates the absence of the transgene sequence and white boxes on KI-R1, -R2 and –R3
fragments show locations of substitutions relative to the other *ω1* copies (Smp_184360): 2 bp
(AT to CC) mismatches at positions 253-254 nt. All three contained the (knock-in) insertion
sequences (white box), which was the targeted mutation of the *ω1* gene. **D-F.** Illumina deep
sequence analysis of amplicon library from eggs of *S. mansoni* revealed Cas9 induced on-target
repair of programmed gene mutation of the ω1 locus by deletions, insertions, and substitutions
by CRISPResso analysis. **D**. Frequency distribution of position dependent deletions and deletion
sizes, which vary from point mutations to more than 20 bp around the DSB. The dotted line
indicates the predicted site of the programmed double stranded break. **E.** Frequency of frameshift
versus in-frame mutations reported by CRISPResso**.** This shows the fraction of all mutations

(INDELs and substitutions) in coding region (positions 42-179) of the amplicon that cause
frameshifts (i.e. indels of 1-2 bp, or multiples thereof). Here (a) is sample 2 (control), which only
contained eggs, and (b) is sample 9, which was treated with lentivirus and ssODN (CRISP-
treated). Findings for all control and treated samples are provided in Table S3. **F.** Insertions of
the knock-in sequence. The y-axis is the number of amplicon reads containing an insertion of the
knock-in sequence (with ≥75% identity to it), and the x-axis is the position of the insertion in the
reference amplicon. The expected Cas9 cut site is between positions 102 and 103. Samples 3, 7
and 9 are independent amplicon libraries (technical replicates) made from the same sample of six
biological replicates treated with lentivirus and ssODN. The insert shows a sequence logo,
created using WebLogo [92], of the sequences of the 3,826 sequence reads from samples 3, 7
and 9, with insertions of 24 bp at position 102, showing that most match the expected knock-in
sequence, TAAGTGACTAGGTAACTGAGTAGC.
740
Figure 3. Reduction of ω1 transcript and ribonuclease T2 activity after ω1 genome editing
by CRISPR/Cas9. Panel **A.** ω1 mRNA abundance was reduced up to ~ 70% after genome
editing by sgRNA/Cas9 complex and lentivirus systems, and markedly reduced >80% with the
addition of ssODN as the DNA repair donor. The % ω1 transcripts represented mean ±SD, 11
biological replicates, at three days following CRISPR/Cas manipulation; asterisks **** indicate
statistical significance at $P ≤ 0.0001$ (n=11), one-way, multiple comparison ANOVA. **B.** Stable
reduction of *ω1* transcripts at days 5 and 7 after treatment (three biological replicates) in four
experimental groups; RNP (black), RNP + ssODN (red), pLV-ω1X1 (green), pLV-ω1X1 +
ssODN (blue). **C, D.** Reduction of RNase activity, assessed by hydrolysis of yeast RNA [41].
Residual yeast following exposure to SEA, visualized after gel electrophoresis; lane 1, buffer
(negative control); 2, RNase A; 3-8, WT SEA and other control SEAs; 9-12, Δω1-SEA from
RNP, RNP + ssODN, pLV-ω1X1 and pLV-ω1X1 + ssODN, respectively (C). Intact yeast RNA
(nanograms) remaining following incubation with SEA, mean ±SD (n = 6). Significantly more
RNA remained following incubation with Δω1-SEA in all groups, RNP, RNP+ssODN, pLV-
ω1X1 and pLV-ω1X1+ssODN treated SEA) (blue) than in the WT SEA controls ($P ≤ 0.0001$).
Moreover, among the gene edited experimental groups, more RNA remained where donor
template was introduced at the same time as RNP or pLV-ω1X1 virions ($P ≤ 0.01$). Differences
were not apparent among WT SEA control groups (D).
759
Figure 4. Reduced Th2 cytokine production following treatment with Δω1-SEA. Panel **A.**
Reduction in Th2 cytokines IL-4 and IL-5 but not IL-13 [93, 94] following pulsing of MΦ
(PMA induced-THP-1 cells) with Δω1-SEA prior to co-culture with human CD4$^+$ T cells (Jurkat
cell line) compared with WT-SEA pulsed-MΦ (top panels). In addition, levels of IL-6 and TNF-
α were reduced where MΦ were first pulsed with Δω1-SEA but not WT SEA. Differences were
not evident for IL-10. The assay was carried out in triplicate. $P < 0.0001, ≤ 0.0001, 0.0038$ and
0.0252 indicated as ****, ***, ** and *, respectively (one way ANOVA, multiple comparison).
767
Figure 5. Pulmonary circumoval granulomas revealed attenuated granulomatous response
to Δω1 schistosome eggs. Schistosome eggs (~3,000 eggs) that had been transduced with
lentivirus virions encoding *ω1*-specific sgRNA and Cas9 in tandem with ssODN, were
introduced via the tail vein into mice. The mice were euthanized 10 days later; thin sections of
the left lung were stained with H&E, and circumoval granulomas counted and measured.
Representative 2D scanned micrographs of granulomas inoculated with WT eggs (panel **A**) (2×

magnification) and 20× magnification (A1 and A2), and with Δω1 eggs; **B** (2×), B1 and B2
(20×). **C**, Control mouse lung. **D, E** Individual control egg induced-granuloma that were counted
assessed for granuloma volume. **F, G**, Δω1 egg induced-granuloma. All single egg induced-
granuloma from WT and Δω1 eggs were measured and granuloma volume estimated [25]. **H**
presents scatter plots of the volume (mm^3) for individual granuloma, with mean ± SEM (red).
Granulomas induced by Δω1 eggs were far smaller than those surrounding WT eggs (Welch's *t*-
test, $P \leq 0.0001$, *n* >100).
781

References

1.   Gryseels B, Polman K, Clerinx J, Kestens L: **Human schistosomiasis.** *Lancet* 2006,
368:1106-1118.
2.   Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J: **Helminth**
infections: the great neglected tropical diseases. *J Clin Invest* 2008, **118:**1311-1321.
3.   Hotez PJ, Alvarado M, Basanez MG, Bolliger I, Bourne R, Boussinesq M, Brooker SJ,
Brown AS, Buckle G, Budke CM, et al: **The global burden of disease study 2010:**
interpretation and implications for the neglected tropical diseases. *PLoS Negl Trop*
Dis 2014, **8:**e2865.
4.   Colley DG, Bustinduy AL, Secor WE, King CH: **Human schistosomiasis.** *Lancet* 2014,
383:2253-2264.
5.   Berriman M, Haas BJ, LoVerde PT, Wilson RA, Dillon GP, Cerqueira GC, Mashiyama
ST, Al-Lazikani B, Andrade LF, Ashton PD, et al: **The genome of the blood fluke**
Schistosoma mansoni. *Nature* 2009, **460:**352-358.
6.   Schistosoma japonicum Genome S, Functional Analysis C: **The Schistosoma japonicum**
genome reveals features of host-parasite interplay. *Nature* 2009, **460:**345-351.
7.   Young ND, Jex AR, Li B, Liu S, Yang L, Xiong Z, Li Y, Cantacessi C, Hall RS, Xu X, et
al: **Whole-genome sequence of Schistosoma haematobium.** *Nat Genet* 2012, **44:**221-
225.
8.   Lepesant JM, Grunau C, Cosseau C: **Towards an understanding of the epigenetics of**
schistosomes: a comparative epigenomic study. *Mem Inst Oswaldo Cruz* 2011,
106:823-830.
9.   Geyer KK, Rodriguez Lopez CM, Chalmers IW, Munshi SE, Truscott M, Heald J,
Wilkinson MJ, Hoffmann KF: **Cytosine methylation regulates oviposition in the**
pathogenic blood fluke Schistosoma mansoni. *Nat Commun* 2011, **2:**424.
10.   Vanderstraete M, Gouignard N, Cailliau K, Morel M, Hahnel S, Leutner S, Beckmann S,
Grevelding CG, Dissous C: **Venus kinase receptors control reproduction in the**
platyhelminth parasite Schistosoma mansoni. *PLoS Pathog* 2014, **10:**e1004138.
11.   Rinaldi G, Eckert SE, Tsai IJ, Suttiprapa S, Kines KJ, Tort JF, Mann VH, Turner DJ,
Berriman M, Brindley PJ: **Germline transgenesis and insertional mutagenesis in**
Schistosoma mansoni mediated by murine leukemia virus. *PLoS Pathog* 2012,
8:e1002820.
12.   Protasio AV, Tsai IJ, Babbage A, Nichol S, Hunt M, Aslett MA, De Silva N, Velarde GS,
Anderson TJ, Clark RC, et al: **A systematically improved high quality genome and**
transcriptome of the human blood fluke Schistosoma mansoni. *PLoS Negl Trop Dis*
2012, **6:**e1455.
13.   Valentim CL, Cioli D, Chevalier FD, Cao X, Taylor AB, Holloway SP, Pica-Mattoccia L,
Guidi A, Basso A, Tsai IJ, et al: **Genetic and molecular basis of drug resistance and**
species-specific drug action in schistosome parasites. *Science* 2013, **342:**1385-1389.
14.   Wang B, Collins JJ, 3rd, Newmark PA: **Functional genomic characterization of**
neoblast-like stem cells in larval Schistosoma mansoni. *Elife* 2013, **2:**e00768.
15.   Collins JJ, 3rd, Wang B, Lambrus BG, Tharp ME, Iyer H, Newmark PA: **Adult somatic**
stem cells in the human parasite Schistosoma mansoni. *Nature* 2013, **494:**476-479.
16.   Hagen J, Young ND, Every AL, Pagel CN, Schnoeller C, Scheerlinck JP, Gasser RB,
Kalinna BH: **Omega-1 knockdown in Schistosoma mansoni eggs by lentivirus**
transduction reduces granuloma size in vivo. *Nat Commun* 2014, **5:**5375.

17.   Hoffmann KF, Brindley PJ, Berriman M: **Medicine. Halting harmful helminths.**
Science 2014, **346:**168-169.

18.   Pearce EJ, MacDonald AS: **The immunobiology of schistosomiasis.** *Nat Rev Immunol*
2002, **2:**499-511.

19.   Steinfelder S, Andersen JF, Cannons JL, Feng CG, Joshi M, Dwyer D, Caspar P,
Schwartzberg PL, Sher A, Jankovic D: **The major component in schistosome eggs**
responsible for conditioning dendritic cells for Th2 polarization is a T2 ribonuclease
(omega-1). *J Exp Med* 2009, **206:**1681-1690.

20.   Everts B, Perona-Wright G, Smits HH, Hokke CH, van der Ham AJ, Fitzsimmons CM,
Doenhoff MJ, van der Bosch J, Mohrs K, Haas H, et al: **Omega-1, a glycoprotein**
secreted by Schistosoma mansoni eggs, drives Th2 responses. *J Exp Med* 2009,
206:1673-1680.

21.   Pearce EJ, C MK, Sun J, J JT, McKee AS, Cervi L: **Th2 response polarization during**
infection with the helminth parasite Schistosoma mansoni. *Immunol Rev* 2004,
201:117-126.

22.   Everts B, Hussaarts L, Driessen NN, Meevissen MH, Schramm G, van der Ham AJ, van
der Hoeven B, Scholzen T, Burgdorf S, Mohrs M, et al: **Schistosome-derived omega-1**
drives Th2 polarization by suppressing protein synthesis following internalization
by the mannose receptor. *J Exp Med* 2012, **209:**1753-1767, S1751.

23.   Dunne DW, Jones FM, Doenhoff MJ: **The purification, characterization, serological**
activity and hepatotoxic properties of two cationic glycoproteins (alpha 1 and omega
1) from Schistosoma mansoni eggs. *Parasitology* 1991, **103 Pt 2:**225-236.

24.   Curwen RS, Ashton PD, Johnston DA, Wilson RA: **The Schistosoma mansoni soluble**
proteome: a comparison across four life-cycle stages. *Mol Biochem Parasitol* 2004,
138:57-66.

25.   Ashton PD, Harrop R, Shah B, Wilson RA: **The schistosome egg: development and**
secretions. *Parasitology* 2001, **122:**329-338.

26.   Cass CL, Johnson JR, Califf LL, Xu T, Hernandez HJ, Stadecker MJ, Yates JR, 3rd,
Williams DL: **Proteomic analysis of Schistosoma mansoni egg secretions.** *Mol*
Biochem Parasitol 2007, **155:**84-93.

27.   Mathieson W, Wilson RA: **A comparative proteomic study of the undeveloped and**
developed Schistosoma mansoni egg and its contents: the miracidium, hatch fluid
and secretions. *Int J Parasitol* 2010, **40:**617-628.

28.   Wilbers RH, Westerhof LB, van Noort K, Obieglo K, Driessen NN, Everts B, Gringhuis
SI, Schramm G, Goverse A, Smant G, et al: **Production and glyco-engineering of**
immunomodulatory helminth glycoproteins in plants. *Sci Rep* 2017, **7:**45910.

29.   Fitzsimmons CM, Schramm G, Jones FM, Chalmers IW, Hoffmann KF, Grevelding CG,
Wuhrer M, Hokke CH, Haas H, Doenhoff MJ, Dunne DW: **Molecular characterization**
of omega-1: a hepatotoxic ribonuclease from Schistosoma mansoni eggs. *Mol*
Biochem Parasitol 2005, **144:**123-127.

30.   Ferguson BJ, Newland SA, Gibbs SE, Tourlomousis P, Fernandes dos Santos P, Patel
MN, Hall SW, Walczak H, Schramm G, Haas H, et al: **The Schistosoma mansoni T2**
ribonuclease omega-1 modulates inflammasome-dependent IL-1beta secretion in
macrophages. *Int J Parasitol* 2015, **45:**809-813.

31.   Wynn TA, Thompson RW, Cheever AW, Mentink-Kane MM: **Immunopathogenesis of**
schistosomiasis. *Immunol Rev* 2004, **201:**156-167.

32.   Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E: **A programmable**
875   **dual-RNA-guided DNA endonuclease in adaptive bacterial immunity.** *Science* 2012,
876   **337:**816-821.

877   33.   Hsu PD, Lander ES, Zhang F: **Development and applications of CRISPR-Cas9 for**
878   **genome engineering.** *Cell* 2014, **157:**1262-1278.

879   34.   Murare HM, Dunne DW, Bain J, Doenhoff MJ: **Schistosoma mansoni: control of**
880   **hepatotoxicity and egg excretion by immune serum in infected immunosuppressed**
881   **mice is schistosome species-specific, but not S. mansoni strain-specific.** *Exp Parasitol*
882   1992, **75:**329-339.

883   35.   Lee AH, Symington LS, Fidock DA: **DNA repair mechanisms and their biological**
884   **roles in the malaria parasite Plasmodium falciparum.** *Microbiol Mol Biol Rev* 2014,
885   **78:**469-486.

886   36.   Deriano L, Roth DB: **Modernizing the nonhomologous end-joining repertoire:**
887   **alternative and classical NHEJ share the stage.** *Annu Rev Genet* 2013, **47:**433-455.

888   37.   International Helminth Genomes Consortium; Coghlan A, Mireva, M., Berriman, M.:
889   **Comparative genomics of the major parasitic worms.** *bioRxiv* 2017,
890   https://www.biorxiv.org/content/early/2017/12/20/236539

891   38.   Lok JB, Shao H, Massey HC, Li X: **Transgenesis in Strongyloides and related**
892   **parasitic nematodes: historical perspectives, current functional genomic**
893   **applications and progress towards gene disruption and editing.** *Parasitology* 2017,
894   **144:**327-342.

895   39.   Gang SS, Castelletto ML, Bryant AS, Yang E, Mancuso N, Lopez JB, Pellegrini M,
896   Hallem EA: **Targeted mutagenesis in a human-parasitic nematode.** *PLoS Pathog*
897   2017, **13:**e1006675.

898   40.   Chen F, Pruett-Miller SM, Davis GD: **Gene editing using ssODNs with engineered**
899   **endonucleases.** *Methods Mol Biol* 2015, **1239:**251-265.

900   41.   Suttiprapa S, Rinaldi G, Tsai IJ, Mann VH, Dubrovsky L, Yan HB, Holroyd N, Huckvale
901   T, Durrant C, Protasio AV, et al: **HIV-1 Integrates Widely throughout the Genome of**
902   **the Human Blood Fluke Schistosoma mansoni.** *PLoS Pathog* 2016, **12:**e1005931.

903   42.   Duvoisin R, Ayuk MA, Rinaldi G, Suttiprapa S, Mann VH, Lee CM, Harris N, Brindley
904   PJ: **Human U6 promoter drives stronger shRNA activity than its schistosome**
905   **orthologue in Schistosoma mansoni and human fibrosarcoma cells.** *Transgenic Res*
906   2012, **21:**511-521.

907   43.   Shah AN, Davey CF, Whitebirch AC, Miller AC, Moens CB: **Rapid reverse genetic**
908   **screening using CRISPR in zebrafish.** *Nat Methods* 2015, **12:**535-540.

909   44.   Yu C, Zhang Y, Yao S, Wei Y: **A PCR based protocol for detecting indel mutations**
910   **induced by TALENs and CRISPR/Cas9 in zebrafish.** *PLoS One* 2014, **9:**e98282.

911   45.   Pinello L, Canver MC, Hoban MD, Orkin SH, Kohn DB, Bauer DE, Yuan GC:
912   **Analyzing CRISPR genome-editing experiments with CRISPResso.** *Nat Biotechnol*
913   2016, **34:**695-697.

914   46.   Suttiprapa S, Rinaldi G, Brindley PJ: **Prototypic chromatin insulator cHS4 protects**
915   **retroviral transgene from silencing in Schistosoma mansoni.** *Transgenic Res* 2012,
916   **21:**555-566.

917   47.   Ke XD, Shen S, Song LJ, Yu CX, Kikuchi M, Hirayama K, Gao H, Wang J, Yin X, Yao
918   Y, et al: **Characterization of Schistosoma japonicum CP1412 protein as a novel**

member of the ribonuclease T2 molecule family with immune regulatory function.
Parasit Vectors 2017, **10:**89.

48. Zaccone P, Burton OT, Gibbs SE, Miller N, Jones FM, Schramm G, Haas H, Doenhoff
MJ, Dunne DW, Cooke A: **The S. mansoni glycoprotein omega-1 induces Foxp3**
expression in NOD mouse CD4(+) T cells. *Eur J Immunol* 2011, **41:**2709-2718.

49. Ritter M, Gross O, Kays S, Ruland J, Nimmerjahn F, Saijo S, Tschopp J, Layland LE,
Prazeres da Costa C: **Schistosoma mansoni triggers Dectin-2, which activates the**
Nlrp3 inflammasome and alters adaptive immune responses. *Proc Natl Acad Sci U S*
A 2010, **107:**20459-20464.

50. Qin Z: **The use of THP-1 cells as a model for mimicking the function and regulation**
of monocytes and macrophages in the vasculature. *Atherosclerosis* 2012, **221:**2-11.

51. Fuentes L, Hernandez M, Fernandez-Aviles FJ, Crespo MS, Nieto ML: **Cooperation**
between secretory phospholipase A2 and TNF-receptor superfamily signaling:
implications for the inflammatory response in atherogenesis. *Circ Res* 2002, **91:**681-
688.

52. Doenhoff MJ, Hassounah O, Murare H, Bain J, Lucas S: **The schistosome egg**
granuloma: immunopathology in the cause of host protection or parasite survival?
Trans R Soc Trop Med Hyg 1986, **80:**503-514.

53. Boros DL, Warren KS: **Delayed hypersensitivity-type granuloma formation and**
dermal reaction induced and elicited by a soluble factor isolated from Schistosoma
mansoni eggs. *J Exp Med* 1970, **132:**488-507.

54. Eltoum IA, Wynn TA, Poindexter RW, Finkelman FD, Lewis FA, Sher A, Cheever AW:
Suppressive effect of interleukin-4 neutralization differs for granulomas around
Schistosoma mansoni eggs injected into mice compared with those around eggs laid
in infected mice. *Infect Immun* 1995, **63:**2532-2536.

55. Wynn TA, Eltoum I, Cheever AW, Lewis FA, Gause WC, Sher A: **Analysis of cytokine**
mRNA expression during primary granuloma formation induced by eggs of
Schistosoma mansoni. *J Immunol* 1993, **151:**1430-1440.

56. Lieber MR: **The mechanism of double-strand DNA break repair by the**
nonhomologous DNA end-joining pathway. *Annu Rev Biochem* 2010, **79:**181-211.

57. Paquet D, Kwart D, Chen A, Sproul A, Jacob S, Teo S, Olsen KM, Gregg A, Noggle S,
Tessier-Lavigne M: **Efficient introduction of specific homozygous and heterozygous**
mutations using CRISPR/Cas9. *Nature* 2016, **533:**125-129.

58. Zhang WW, Matlashewski G: **CRISPR-Cas9-Mediated Genome Editing in**
Leishmania donovani. *MBio* 2015, **6:**e00861.

59. Yoshimi K, Kunihiro Y, Kaneko T, Nagahora H, Voigt B, Mashimo T: **ssODN-mediated**
knock-in with CRISPR-Cas for large genomic regions in zygotes. *Nature*
communications 2016, **7:**10431.

60. Jurberg AD, Goncalves T, Costa TA, de Mattos AC, Pascarelli BM, de Manso PP,
Ribeiro-Alves M, Pelajo-Machado M, Peralta JM, Coelho PM, Lenzi HL: **The**
embryonic development of Schistosoma mansoni eggs: proposal for a new staging
system. *Dev Genes Evol* 2009, **219:**219-234.

61. Neill PJ, Smith JH, Doughty BL, Kemp M: **The ultrastructure of the Schistosoma**
mansoni egg. *Am J Trop Med Hyg* 1988, **39:**52-65.

62. Mann VH, Suttiprapa S, Rinaldi G, Brindley PJ: **Establishing transgenic schistosomes.**
PLoS Negl Trop Dis 2011, **5:**e1230.

63.   Kosicki M, Rajan SS, Lorenzetti FC, Wandall HH, Narimatsu Y, Metzakopian E, Bennett
EP: **Dynamics of Indel Profiles Induced by Various CRISPR/Cas9 Delivery**
Methods. *Prog Mol Biol Transl Sci* 2017, **152:**49-67.

64.   Holmgaard A, Askou AL, Benckendorff JNE, Thomsen EA, Cai Y, Bek T, Mikkelsen
JG, Corydon TJ: **In Vivo Knockout of the Vegfa Gene by Lentiviral Delivery of**
CRISPR/Cas9 in Mouse Retinal Pigment Epithelium Cells. *Mol Ther Nucleic Acids*
2017, **9:**89-99.

65.   Yu Z, Chen Q, Chen W, Zhang X, Mei F, Zhang P, Zhao M, Wang X, Shi N, Jackson S,
Hong Y: **Multigene editing via CRISPR/Cas9 guided by a single-sgRNA seed in**
Arabidopsis. *J Integr Plant Biol* 2018, **60:**376-381.

66.   Luo JJ, Bian WP, Liu Y, Huang HY, Yin Q, Yang XJ, Pei DS: **CRISPR/Cas9-based**
genome engineering of zebrafish using a seamless integration strategy. *FASEB J*
2018**:**fj201800077RR.

67.   Shalem O, Sanjana NE, Zhang F: **High-throughput functional genomics using**
CRISPR-Cas9. *Nat Rev Genet* 2015, **16:**299-311.

68.   Kines KJ, Rinaldi G, Okatcha TI, Morales ME, Mann VH, Tort JF, Brindley PJ:
Electroporation facilitates introduction of reporter transgenes and virions into
schistosome eggs. *PLoS Negl Trop Dis* 2010, **4:**e593.

69.   Heyer WD, Ehmsen KT, Liu J: **Regulation of homologous recombination in**
eukaryotes. *Annu Rev Genet* 2010, **44:**113-139.

70.   Schramm G, Hamilton JV, Balog CI, Wuhrer M, Gronow A, Beckmann S, Wippersteg V,
Grevelding CG, Goldmann T, Weber E, et al: **Molecular characterisation of kappa-5, a**
major antigenic glycoprotein from Schistosoma mansoni eggs. *Mol Biochem Parasitol*
2009, **166:**4-14.

71.   Brogna S, Wen J: **Nonsense-mediated mRNA decay (NMD) mechanisms.** *Nat Struct*
Mol Biol 2009, **16:**107-113.

72.   Pankowicz FP, Jarrett KE, Lagor WR, Bissig KD: **CRISPR/Cas9: at the cutting edge of**
hepatology. *Gut* 2017, **66:**1329-1340.

73.   Kaisar MMM, Ritter M, Del Fresno C, Jonasdottir HS, van der Ham AJ, Pelgrom LR,
Schramm G, Layland LE, Sancho D, Prazeres da Costa C, et al: **Dectin-1/2-induced**
autocrine PGE2 signaling licenses dendritic cells to prime Th2 responses. *PLoS Biol*
2018, **16:**e2005504.

74.   Dalton JP, Day SR, Drew AC, Brindley PJ: **A method for the isolation of schistosome**
eggs and miracidia free of contaminating host tissues. *Parasitology* 1997, **115 (Pt**
1):29-32.

75.   Mann VH, Morales ME, Rinaldi G, Brindley PJ: **Culture for genetic manipulation of**
developmental stages of Schistosoma mansoni. *Parasitology* 2010, **137:**451-462.

76.   Mann VH, Suttiprapa S, Skinner DE, Brindley PJ, Rinaldi G: **Pseudotyped murine**
leukemia virus for schistosome transgenesis: approaches, methods and perspectives.
Transgenic Res 2014, **23:**539-556.

77.   Cardoso LS, Araujo MI, Goes AM, Pacifico LG, Oliveira RR, Oliveira SC: **Polymyxin B**
as inhibitor of LPS contamination of Schistosoma mansoni recombinant proteins in
human cytokine analysis. *Microb Cell Fact* 2007, **6:**1.

78.   Bradford MM: **A rapid and sensitive method for the quantitation of microgram**
quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*
1976, **72:**248-254.

79. Oliveros JC, Franch M, Tabas-Madrid D, San-Leon D, Montoliu L, Cubas P, Pazos F:
Breaking-Cas-interactive design of guide RNAs for CRISPR-Cas experiments for
ENSEMBL genomes. *Nucleic Acids Res* 2016, **44:**W267-271.
80. Minskaia E, Nicholson J, Ryan MD: **Optimisation of the foot-and-mouth disease virus**
2A co-expression system for biomedical applications. *BMC Biotechnol* 2013, **13:**67.
81. Lok JB, Shao H, Massey HC, Li X: **Transgenesis in Strongyloides and related**
parasitic nematodes: historical perspectives, current functional genomic
applications and progress towards gene disruption and editing. *Parasitology* 2016**:**1-
16.
82. Marozsan AJ, Fraundorf E, Abraha A, Baird H, Moore D, Troyer R, Nankja I, Arts EJ:
Relationships between infectious titer, capsid protein levels, and reverse
transcriptase activities of diverse human immunodeficiency virus type 1 isolates. *J*
Virol 2004, **78:**11130-11141.
83. Martin M: **Cutadapt removes adapter sequences from high-throughput sequencing**
reads. *EMBnetjournal North America* 2011,
http://journal.embnet.org/index.php/embnetjournal/article/view/200/458.
84. Canver MC, Haeussler M, Bauer DE, Orkin SH, Sanjana NE, Shalem O, Yuan GC,
Zhang F, Concordet JP, Pinello L: **Integrated design, execution, and analysis of**
arrayed and pooled CRISPR genome-editing experiments. *Nat Protoc* 2018, **13:**946-
986.
85. Pearson WR, Lipman DJ: **Improved tools for biological sequence comparison.** *Proc*
Natl Acad Sci U S A 1988, **85:**2444-2448.
86. Ginzinger DG: **Gene quantification using real-time quantitative PCR: an emerging**
technology hits the mainstream. *Exp Hematol* 2002, **30:**503-512.
87. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time**
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001, **25:**402-408.
88. Cheever AW, Finkelman FD, Caspar P, Heiny S, Macedonia JG, Sher A: **Treatment**
with anti-IL-2 antibodies reduces hepatic pathology and eosinophilia in Schistosoma
mansoni-infected mice while selectively inhibiting T cell IL-5 production. *J Immunol*
1992, **148:**3244-3248.
89. Genin M, Clement F, Fattaccioli A, Raes M, Michiels C: **M1 and M2 macrophages**
derived from THP-1 cells differentially modulate the response of cancer cells to
etoposide. *BMC Cancer* 2015, **15:**577.
90. Schmid MC, Varner JA: **Myeloid cells in the tumor microenvironment: modulation of**
tumor angiogenesis and tumor inflammation. *J Oncol* 2010, **2010:**201026.
91. Chomczynski P: **A reagent for the single-step simultaneous isolation of RNA, DNA**
and proteins from cell and tissue samples. *Biotechniques* 1993, **15:**532-534, 536-537.
92. Crooks GE, Hon G, Chandonia JM, Brenner SE: **WebLogo: a sequence logo generator.**
Genome Res 2004, **14:**1188-1190.
93. Gieseck RL, 3rd, Wilson MS, Wynn TA: **Type 2 immunity in tissue repair and**
fibrosis. *Nat Rev Immunol* 2018, **18:**62-76.
94. Wynn TA: **Type 2 cytokines: mechanisms and therapeutic strategies.** *Nat Rev*
Immunol 2015, **15:**271-282.
95. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,
Valentin F, Wallace IM, Wilm A, Lopez R, et al: **Clustal W and Clustal X version 2.0.**
Bioinformatics 2007, **23:**2947-2948.

1057

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5