Structure-aware *M. tuberculosis* functional annotation uncloaks resistance, metabolic, and virulence genes

Running title: Structure-aware *M.tb* annotation uncloaks key functions

[a]Samuel J Modlin, [a]Afif Elghraoui, [a]Deepika Gunasekaran, [a]Alyssa M Zlotnicki, [b]Nicholas A Dillon, [a]Nermeeta Dhillon, [a]Norman Kuo, [a]Cassidy Robinhold, [a]Carmela K Chan, [b]Anthony D Baughn, [a]Faramarz Valafar*

[a] Laboratory for Pathogenesis of Clinical Drug Resistance and Persistence, San Diego State University, San Diego, CA 92182

[b] Department of Microbiology and Immunology, University of Minnesota Medical School, Minneapolis, MN, 55455, USA

* Corresponding Author

ABSTRACT Accurate and timely functional genome annotation is essential for translating basic pathogen research into clinically impactful advances. Here, through literature curation and structure-function inference, we systematically update the functional genome annotation of *Mycobacterium tuberculosis* virulent type strain H37Rv. First, we systematically curated annotations for 589 genes from 662 publications, including 282 gene products absent from leading databases. Second, we modeled 1,711 under-annotated proteins and developed a semi-automated pipeline that captured shared function between 400 protein models and structural matches of known function on protein data bank, including drug efflux proteins, metabolic enzymes, and virulence factors. In aggregate, these structure- and literature-derived annotations update 940/1,725 under-annotated H37Rv genes and generate hundreds of functional hypotheses. Retrospectively applying the annotation to a recent whole-genome transposon mutant screen provided missing function for 48% (13/27) of under-annotated genes altering antibiotic efficacy and 33% (23/69) required for persistence during mouse TB infection. Prospective application of the protein models enabled us to functionally interpret novel laboratory generated Pyrazinamide-resistant (PZA) mutants of unknown function, which

implicated the emerging Coenzyme A depletion model of PZA action in the mutants' PZA resistance. Our findings demonstrate the functional insight gained by integrating structural modeling and systematic literature curation, even for widely studied microorganisms. Functional annotations and protein structure models are available at https://tuberculosis.sdsu.edu/H37Rv in human- and machine-readable formats.

IMPORTANT *Mycobacterium tuberculosis,* the primary causative agent of tuberculosis, kills more humans than any other infectious bacteria. Yet 40% of its genome is functionally uncharacterized, leaving much about the genetic basis of its resistance to antibiotics, capacity to withstand host immunity, and basic metabolism yet undiscovered. Irregular literature curation for functional annotation contributes to this gap. We systematically curated functions from literature and structural similarity for over half of poorly characterized genes, expanding the functionally annotated *Mycobacterium tuberculosis* proteome. Applying this updated annotation to recent *in vivo* functional screens added functional information to dozens of clinically pertinent proteins described as having unknown function. Integrating the annotations with a prospective functional screen identified new mutants resistant to a first-line TB drug supporting an emerging hypothesis for its mode of action. These improvements in functional interpretation of clinically informative studies underscores the translational value of this functional knowledge. Structure-derived annotations identify hundreds of high-confidence candidates for mechanisms of antibiotic resistance, virulence factors, and basic metabolism; other functions key in clinical and

basic tuberculosis research. More broadly, it provides a systematic framework for improving prokaryotic reference annotations.

KEYWORDS *Mycobacterium tuberculosis*, annotation, structure, virulence factors, functional genomics, pyrazinamide, resistance

INTRODUCTION

Manual curation remains the gold standard for annotating function from literature[1], yet requires massive effort from highly specialized researchers. UniProt curators alone evaluate over 4,500 papers each year[1]. Literature annotation is typically complemented with functional inference by sequence homology, but this approach fails to identify distant relatives (remote homologs) or convergently evolved proteins of shared function (structural analogs).

These challenges hinder the study of *Mycobacterium tuberculosis,* the etiological agent of tuberculosis. The *M. tuberculosis* virulent type strain is H37Rv, a descendant of strain H37, was isolated from a pulmonary TB patient in 1905 and kept viable through repeated subculturing[2]. Following sequencing of the H37Rv genome, function was assigned to 40% of its open reading frames (ORFs)[3], then expanded to 52% in 2002 following re-annotation[4]. New annotations continued to be added by TubercuList (now part of Mycobrowser, https://mycobrowser.epfl.ch/) until March 2013. To date, one-quarter of the H37Rv genome (1,057 genes) lacks annotation entirely, listed in "conserved hypotheticals" or "unknown" functional categories and hundreds more minimally describe product function (e.g. "possible membrane protein"). Though other databases have emerged in recent years[5–9] Mycobrowser remains the primary resource for

gene annotation for TB researchers(10) yet lacks functional characterizations reported in the

literature.

Moreover, many proteins key to *M. tuberculosis* pathogenesis are challenging to ascribe

function by sequence similarity. For instance, transport proteins—many of which allow *M.*

tuberculosis to tolerate drug exposure by effluxing drug out of the cell(11)—have membrane-

embedded regions under relaxed constraint compared to globular proteins and diverge in

sequence more rapidly as a result(12). This rapid divergence challenges their characterization

through homology. Limitations of sequence-based approaches to detect and annotate *M.*

tuberculosis proteins motivates an alternative approach to annotating *M. tuberculosis* gene

function.

One alternative approach is identifying protein homologs and analogs through shared

structure, which offers considerable advantages. First, it removes bias toward *a priori*

assumptions by not limiting search space to evolutionarily close relatives, facilitating discovery

of functions unexpected by conventional predictions. Second, structure-based functional

annotation can infer analogy between proteins structures. This ability to detect analogy is

especially valuable for inferring function at the host-pathogen interface, which is challenging to

recapitulate in the laboratory. Moreover, analogous relationships between proteins of shared

structure/function cannot be resolved by sequence homology because they often evolve

convergently with low amino acid (AA) similarity rather than from common ancestry (13).

Iterative Threading ASSEmbly Refinement (I-TASSER)(14), builds three-dimensional protein

structure from sequence through multiple threading alignment of Protein Data Bank (PDB)(15)

templates, followed by iterative fragment assembly simulations. I-TASSER accurately predicts structure (16–20), provides metrics for model quality(21) (C-score) and pairwise structural similarity(22) (TM-score), and integrates function and structure prediction tools(23) (COACH and COFACTOR) comprising Gene Ontology (GO) terms(24), Enzyme Commission (EC) numbers(25), and Ligand Binding Sites (LBS)(26).

EC numbers and GO terms partially or completely define gene function and are widely incorporated into mainstream databases. EC numbers describe catalytic function hierarchically through a four-tiered numerical identifier system that funnels from general enzyme class (e.g. ligase, oxidoreductase) down to substrate specificity with atomic precision(25). GO terms add to EC number content: they describe gene products by where they function, the processes they are involved in, and their specific molecular function in species-independent form(27, 28). This cross-species unification is particularly useful for reconciling annotation transfers of analogs and distant homologs into gene product names.

Previous hypothetical gene annotation efforts for *M. tuberculosis* have not included a systematic manual literature curation component and have drawn from inferential techniques such as protein homology and fold similarity(29, 30), aggregating gene orthology server predictions(31), metabolic pathway gap-filling(32), and STRING interactions(33), lacking inclusion criteria based on benchmarked likelihood of correctness. Measured interpretation of annotated gene functions requires the source of the annotation and the reliability of the evidence warranting it to be described explicitly. We strived to provide this resource by reconciling the H37Rv annotation on Mycobrowser with published functional characterization and systematically inferring function

from structural similarity to annotate genes challenging to characterize through experiment and sequence analysis. We include orthogonal validation measures to confidently capture unexpected functions while minimizing "overannotation"(34, 35).

We report our findings in three sections. First, we establish the set of under-annotated genes, describe our systematic manual literature curation protocol, and summarize the novelty of the resulting annotations with respect to popular functional databases. Next, we describe our structural modelling pipeline, orthogonal validation and quality assurance methods, and two illustrative examples of manually curated functional annotations from structural inferences unsupported by an established method of detecting remote functional homology. Finally, we summarize the updated annotation and genes remaining to be characterized, and demonstrate the value added of this annotation through its application to previously published and novel functional screens.

RESULTS

Numerous genes lack annotation in all common *M. tuberculosis* databases. First, we defined a set of 1,725 under-annotated genes (Dataset 1) based on their TubercuList entry. We included:

(1) Genes in "conserved hypothetical" or "unknown" functional categories.

(2) Genes qualified by an adjective connoting low confidence (e.g. "predicted" or "possible").

(3) Genes described by something other than function: (e.g. "Alanine-rich protein" or "isoniazid-inducible protein").

(4) PE/PPE family genes – A largely uncharacterized, polymorphic protein family unique to Mycobacteria with proline-glutamine or proline-proline-glutamine N-terminal domains.

Next, we asked how many of these genes lacked annotations across commonly referenced

databases (Table 1). Although BioCyc and UniProt had more genes with GO terms than

TubercuList, and UniProt, and Mtb Network Portal had fewer hypothetical proteins than

TubercuList, all databases had over one-quarter of coding sequences (CDS) annotated as

hypothetical, demonstrating the need for systematic manual annotation.

Table 1. Comparison among frequented annotation resources. "Functional assignments" refer to annotations
that describe protein function, and excluded hypothetical, unknown/uncharacterized, and PE/PPE family proteins.
Counts reflect database content on May 17, 2017 for RefSeq(36) https://www.ncbi.nlm.nih.gov/refseq/, PATRIC(6)
https://www.patricbrc.org/,and Mtb Network Portal(9) http://networks.systemsbiology.net/mtb/ and June 23, 2017
for   KEGG(120)   https://www.kegg.jp/kegg/genome/pathogen.html   and   UniProt(116)
https://www.uniprot.org/uniprot/ . The number of CDSs in KEGG is reported as 3906 because they include only
protein coding genes. The source of annotations for *M. tuberculosis* protein-coding genes in KEGG is
TubercuList(125).

Metric	TubercuList	PATRIC	RefSeq	Mtb Network Portal	UniProt	KEGG	BioCyc
Coding sequences	4038	4367	3989	4038	3997	3906	4031
Proteins with functional assignments	2815	3007	2341	2853	2906	1750	2571
Hypothetical proteins	1223	1360	1648	1185	1091	2156	1460
Proteins with ≥ 1 GO term	2629	969	0	2460	3305	0	3557
Proteins with EC number(s) assigned	1293	1074	1081	1003	1138	1050	1018

146
147

148   **Frequently consulted annotation sources lack experimentally demonstrated functions.**
149   We devised a manual curation protocol (*Text S1*, Fig. S1) that:

150   1) Assigns qualifying adjectives that connote confidence

151   2) Assigns Enzyme Commission ("EC") numbers

152   3) Requires multiple reviewers per paper to hedge against human error, and an additional

153   quality control curator to check formatting and annotation consistency.

We systematically reviewed over 5,000 publications according to this protocol, furnishing

annotations for one-third of under-annotated genes (575) with product function or functional

notes (Dataset 1). Of these, 282 were annotated with product function absent from TubercuList,

including 122 enzymes and 28 regulatory proteins. These annotations include fourteen oxidative

stress response genes, 22 proteins mediating RNA and DNA functions, and eight transport/efflux

proteins.

Next, we evaluated whether these missing annotations were restricted to TubercuList or more

widespread. We checked our curations against four frequented annotation resources: UniProt

(Dataset 2), Mtb Network Portal(9), PATRIC(6), and RefSeq(36) (Dataset 2). Product function

information was absent from 172/282 (61%) of these genes on UniProt (Dataset 2) and 118 (64

of which are antigens, *Text S1,* Fig. S4A) were more thoroughly annotated than in any of the

examined databases (Dataset 2). This novelty underscores the value of these manual curations

and highlights critical information that these databases lack (Table 2 and Dataset 1). After

excluding antigens, 25.2% of genes with function curated from literature were absent from all

five annotation resources. To identify enzymatic functions unannotated elsewhere, we compared

our manual EC number assignments to commonly referenced databases (*Text S1*, Fig. S4). This

comparison revealed 59/98 (60.2%) of genes assigned EC numbers have EC numbers only in our

annotation. These missing annotations include functions affecting drug-resistance, features of *in*

vivo infection, and other important functions. Examples include a rare instance where a PE/PPE

gene has demonstrated catalytic function(37) (Rv1430), a probable peptidoglycan hydrolase

implicated in isoniazid (INH) and PZA resistance and biofilm formation(38) (Rv0024), a rhomboid

protease with roles in biofilm formation and ciprofloxacin and novobiocin resistance(39)

(Rv1337), and an oxidoreductase important for in-host survival of *M. tuberculosis*(40) (41, 42)

(Rv3005c). Additional findings pertinent to pathogenesis, host-pathogen interaction, and

antibiotic resistance were noted across under-annotated genes (S1 Annotation).

Table 2. Genes annotated through systematic manual curation that expand upon annotations from major
databases. Annotations are separated into completely novel, those with similar annotations but with greater
specificity, and those with an additional, orthogonal annotation compared to evaluated databases (*Text S1*, Fig. S4A).
Pubmed IDs (PMID) from which annotations for each product were derived are included. Members of the PE/PPE
family are indicated by asterisk.

Gene	product	PMID	TubercuList	UniProt	Mtb Net Portal
			Novel		
Rv0309	Adhesin/Putative L,D-transpeptidase	23922800; 23889607; 26201501	Possible conserved exported protein	Possible conserved exported protein	
Rv0394c	Hyaluronidase/ Chondrosulfatase	23465892	Possible secreted protein	Possible secreted protein	Possible membrane protein
Rv0431	Probable vesiculogenesis/immune response regulator	24248369; 21170273; 17436267; 26324094; 27765619	Putative tuberculin related peptide	Putative tuberculin related peptide	Tuberculin related peptide
Rv1430*	Esterase	23383323	PE family protein PE16	PE family protein PE16	PE family protein
Rv1993c	Putative chaperone	21925112	Conserved protein	Uncharacterized protein	
Rv2345	Probable phosphatase	25782739	Possible conserved transmembrane protein	UPF0603 protein	Possible membrane protein
Rv2923c	Probable Osmotically induced bacterial protein C (OsmC, a probable peroxide reductase)	22088319	Conserved protein	Uncharacterized protein	
Rv2954c	Probable Methyltransferase	23536839	Hypothetical protein	Uncharacterized protein	
Rv2969c	Periplasmic disulfide-bond forming (Dsb) Enzyme	24100317; 18539140	Possible conserved membrane or secreted protein	Membrane protein (Possible conserved membrane or secreted protein)	POSSIBLE CONSERVED MEMBRANE OR SECRETED PROTEIN
Rv3528c	Probable serine hydrolase	26853625	Unknown protein	Uncharacterized protein	
			Greater specificity		
Rv0059	Probable toxin DarT/Probable DNA ADP-ribosyl transferase	27939941	Hypothetical protein	Uncharacterized protein	
Rv0060	Probable antitoxin DarG/Probable DNA ADP-ribosyl glycohydrolase	27939941	Conserved hypothetical protein	Uncharacterized protein	ADP-ribose 1-phosphate phophatase related protein
Rv1337	Probable rhomboid protease/integral membrane protein	19165721; 23029216	Probable integral membrane protein	Uncharacterized protein	Rhomboid membrane family protein
Rv1357c	Cyclic diguanylate phosphodiesterase	21151497	Conserved hypothetical protein	Uncharacterized protein	Sensory box/GGDEF family protein
Rv1566c	Probable non-catalytic peptidoglycan binding RipD protein/probable antigen	24107184; 26481294	Possible Inv protein	Possible Inv protein	Invasion protein
Rv2024c	Restriction enzyme/m-6-adenine DNA methyltransferase (Mycobacterial adenine methyltransferase B "MamB")	26704977	Conserved hypothetical protein	Uncharacterized protein	putative helicase

			Conserved hypothetical alanine rich protein	Conserved hypothetical alanine rich protein	
Rv2695	Probable serine hydrolase	26853625	Conserved hypothetical alanine rich protein	Conserved hypothetical alanine rich protein	
Rv2991	Probable flavin/deazaflavin oxidoreductase	26434506	Conserved protein	Conserved protein (F420-dependent protein)	
Rv3036c	Esterase	25224799	Probable conserved secreted protein TB22.2	Probable conserved secreted protein TB22.2	Possible membrane protein
Rv3354	protein kinase	25139900	Conserved hypothetical protein	Lipoprotein	Possible lipoprotein LprJ
Orthogonal annotation					
Rv0256c *	B cell Antigen/probable INOS promoter binding protein	23827809; 28071726	PPE family protein PPE2	Uncharacterized PPE family protein PPE2	Predicted cobalt transporter in Mycobacteria
Rv2204c	probable serine hydrolase	26298037; 26853625; 26536359	Conserved protein	Protein Rv2204c	probable iron binding protein from the HesB_IscA_SufA family
Rv3779	polyprenylphosphomannosyl synthase/galactosaminyltransferase	21030587; 19717608	Probable conserved transmembrane protein alanine and leucine rich	Membrane protein (Probable conserved transmembrane protein alanine and leucine rich)	

Annotating function from structure similarity. Next, we modeled protein structures and developed a procedure to annotate function based on shared structure according to the likelihood that two proteins shared function (i.e. precision; Equation 1, Methods and Materials). To inform our annotation methods we first assessed whether we could

(1) Reliably infer precision according to similarity.

(2) Differentiate between precision thresholds at different levels of functional detail (e.g. EC number tiers).

To make these assessments we benchmarked precision as a function of template-modelling score ("TM-score"), a measure of structural similarity independent of protein length, and sequence similarity (Amino acid identity, "AA%"), using a set of 363 genes with known function (Methods & Materials) through the standalone version of I-TASSER. TM-score and AA% were predictive of precision and mutually correlated ($R=0.784$, Pearson correlation coefficient) among both concordant and discordant EC numbers (*Text S1*, Fig. S2B). We accounted for TM-score and

201 AA% simultaneously by their geometric mean (μ_{geom}) to estimate precision. Precision of EC

202 number prediction increased monotonically as a function of μ_{geom} for all 4 EC tiers, and regression

203 lines for the 4 degrees of EC functional specificity did not intersect (Fig. 1).

204
Fig. 1. Determining similarity thresholds for annotation inclusion criteria. Precision of EC number as regressed
against the geometric mean of TM-score and AA% ("μ_{geom}") for each specificity tier. Horizontal lines define (50%
and 75%) thresholds, the points where precision intersects with regression lines for each EC specificity curve
(labelled). Circles at the bottom and top are individual data points (incorrect = 0 and correct = 1, y-axis: precision,
x-axis: μ_{geom}). Circles are rendered at 10% opacity to visualize observation density. Only templates with AA% < 40%
were included.

From these properties we concluded that we could reliably estimate precision from μ_{geom} with

distinct thresholds for each EC tier. We defined thresholds as the μ_{geom} value where logistic

regression lines intersected with 50% for receiving "putative" and 75% for receiving "probable"

as qualifying adjectives (Fig. 1). Through this procedure we defined distinct thresholds for

ascribing "putative" or "probable" status to enzymatic function at each of the 4 tiers of EC

specificity. We incorporated EC numbers and GO terms from similar structures deposited on

Protein Data Bank (PDB) hierarchically, according to evidence reliability (*Text S1*, Fig. S3). After

the quality control pipeline described below, we recorded annotations in NCBI Table File Format

and according to Genbank Prokaryotic Annotation Guide

(www.ncbi.nlm.nih.gov/genbank/genomesubmit_annotation/) syntax and guidelines (integrated

with manual curations from literature), and collated them into a unified functional annotation in

GFF3 format (Fig. 2).

Fig. 2. Information flow for producing annotations from structural similarity. The flow of information and
procedures for acquiring, processing, filtering, and representing information, running from retrieval of amino acid
sequences to the final updated H37Rv annotation. Some details are omitted for clarity. The 1,725 amino acid
sequences were retrieved from TubercuList and run through a local installation of I-TASSER v5.1. 1,711 of 1,725
Amino Acid sequences had models generated successfully. Comparison metrics for sequence (Amino acid identity)
and Structure (TM-score) were extracted from I-TASSER output. To set criteria for annotation transfer, precision
(Equation 1) of GO Term and EC number concordance between similar matches on PDB and true function of 363
positive controls with GO terms and EC numbers of known function were regressed against extracted similarity
metrics to generate a curve relating the geometric mean of TM-score and Amino acid similarity to precision. These
informed inclusion thresholds for transferring GO and EC annotations from structures on PDB similar to the 1,711
modeled structures. CATH Topology folds were transferred according to a previous precision curve based on TM-
score. This threshold was also used for inclusion of proteins classes that vary in sequence more than structure (e.g.
transporters) and as criteria for transferring annotations from structures that were not annotated with EC numbers
or GO terms. Annotations derived only from structure were passed through orthogonal validation and manual
structure analysis for verification that transferred annotations were reasonable. All annotations were
programmatically collated into an updated H37Rv reference genome annotation.

Although using μ_{geom} to determine inclusion criteria is useful for proteins with PDB entries of

somewhat homologous sequence, it would not capture relationships by structural analogy or

remote homology (because their low AA% would lower their score). To identify potential analogs

and remote homologs, we used "TM_{ADJ}", an adjusted TM-score that accounts for model quality

to conservatively estimate the TM-score between the true structure of a modelled protein and

its putative homolog/analog of solved structure (Materials & Methods). We re-examined hits

with TM_{ADJ} values that indicated matching topology according to previous benchmarks(21) (*Text*

S1), and annotated function with EC numbers, GO terms, and product names (*Text S1* and

Methods and Materials).

Validating structure-based annotations. To validate our structure-based functional inference

approach, we ran proteins with annotations derived only from structural similarity (n=366

through HHpred (43) (Fig. 3), a server that detects remote homology between proteins by

comparing hidden Markov model profiles(43). We compared enzymatic structure-derived

annotations (those with EC numbers, n=335 distinct EC number annotations from 271 proteins)

programmatically, and non-enzymatic annotations manually (n=95, Dataset 3, Materials &

Methods). Evaluating only the annotations to at least the 2nd EC number level (n=325), most

structure-inferred predictions were partially (288/335, 86.0%) or wholly (266/335, 79.4%)

corroborated by HHpred (Fig. 3C), substantiating the validity of our structure-based approach to

functional inference. Partially corroborated annotations (e.g. 3.1.2.4 to the level of 3.1.2.-, but not

the 4th level of EC specificity) were revised to reflect the less specific, HHpred-supported level of

functional detail, and manually reconciled in cases where multiple EC numbers were

corroborated (Materials & Methods).

Fig. 3. Orthogonal validation and quality assurance for structure-to-function inference. (**A**) Quality assurance and validation protocol assignment and (**B**) Decision workflow for retaining functional annotations inferred from structural similarity to proteins of solved structure and known function. (**C**) Heat map depicting fraction of EC number inferences corroborated by HHpred at each level of specificity. Fraction denominator is binned according to the number of E.C digits annotated (x-axis). (**D,E**) Structure quality assurance. Distribution of fraction of residues in "most favorable" region of Ramachandran plot prior to (**D**) and following application of a heuristic threshold (**E**) to discard biophysically improbable structural models.

Next, we assessed protein structure model quality using the fraction of residues in "most

favored" regions of Ramachandran plots (Materials & Methods). Screening for abnormally low

fractions can identify models with sterically untenable residue configurations, signaling low model quality(44). A threshold of 90% is often used for solved proteins(45), but we expected deviation from 90% even in quality models (since they are models rather than solved structures). To determine an acceptable threshold, we compared the distribution of residue fractions in "most favorable" regions among models with functions fully corroborated by HHpred with that of 29 wholly uncorroborated by HHpred. Fractions for HHpred-corroborated proteins distributed unimodally and peaked around 90% of residues falling in the "most favorable" region (median = 89.15%). This observation is consistent with HHpred-corroborated proteins having high-quality structures and informs us of the range of fractions characteristic of high-quality structural models. Models with functions wholly uncorroborated by HHpred, meanwhile, distributed bimodally, with one mode resembling the fully corroborated distribution and the second mode peaking at a lower fraction (Fig. 3D). This bimodal distribution is consistent with a mixture of quality models and truly poor models. To distinguish between poor- and high-quality models in the wholly uncorroborated set, we implemented a heuristic threshold at the intersection of the two distributions (75%, Fig. 3D). After removing models below the threshold, the remaining uncorroborated structures formed a single peak that resembled the HHpred-corroborated proteins (Fig. 3E). We used this threshold (75%) as the minimum acceptable fraction for HHpred-uncorroborated proteins to be considered for structure-based functional annotation.

Seven of the protein models with exclusively wholly unsupported structure-based annotations (n=29) were PE_PGRS protein models that resembled fatty acid synthase (FAS) subunit protein structures (particularly *Saccharomyces cerevisiae* PDB template 2pff). All seven failed

Ramachandran filtering. This underscores the importance of these QC steps and suggests they

excluded models implicating false functional analogies as intended. These annotations were

likely artifactual, owing to glycine-abundant, low complexity regions of PE_PGRS proteins

aligning to the hydrophobic regions of large eukaryotic synthases, inflating their similarity score

and spuriously implying structural similarity.

Since HHpred is designed to detect homology between proteins(43) (but not necessarily

analogy), there may be genuine functions inferred by our structural similarity pipeline that

HHpred did not corroborate. To preserve such annotations while ensuring annotation quality,

we manually inspected HHpred-uncorroborated annotations (Fig. 3B) for protein models that

passed Ramachandran filtering (n=22). To accept annotations, we verified template protein

quality, structural alignment of regions underlying function, and conservation of structural

features and key functional residues. This step salvaged structure-derived functional annotations

for nine proteins (Table 3, Dataset 3), two of which we highlight in detail in Fig. 4.

Table 3. Functional annotation of HHpred-uncorroborated enzymes following manual structural analysis.

Rv number	PDB ID	AA %	Inferred EC	Updated EC	Change	Recommended product name	Product name (Mycobrowser)
Rv0036c	2nsg	14%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved protein
Rv0738	2nsg	17%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved protein
Rv1632c	2p12	59%	3.6.1.-	3.6.1.-	Verified	putative cytidylyl-2-hydroxypropylphosphonate hydrolase	Hypothetical protein
Rv1727	2nsg	12%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein
Rv1734c	3l60	26%	1.2.4.-	2.3.-.-	Changed function	putative acyltransferase	Conserved hypothetical protein
Rv1775	3hwp	29%	3.-.-.-	3.7.1.-	Increased specificity	putative 3-oxo-carboxylic acid hydrolase	Conserved hypothetical protein
Rv2036	2nsg	15%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein
Rv2968c	3kp9	21%	1.1.4.-	1.17.4.-	Nomenclature change	putative Vitamin K epoxide reductase	Probable conserved integral membrane protein
Rv3224B	1dbx	38%	3.1.1.-	None	Reduced specificity	putative trans-editing enzyme	Conserved hypothetical protein

In the first example (Fig. 4A-B), manual structural analysis fully corroborates the HHpred-uncorroborated function inferred from structure and extends annotation specificity. Originally, Rv1775 was ascribed putative hydrolase function by our structure-function inference pipeline. Its structural model is globally similar (TM_{ADJ}=0.817) to 2,4-diacetylphloroglucinol hydrolase PhlG (EC 3.7.1.24) from *Pseudomonas fluorescens*(46) (PfPhlG) despite only modest sequence similarity (27.6%). Comparison of Rv1775 to PfPhlG (Fig. 4A) and potential mycobacterial homolog (TM_{ADJ}=0.753, AA=28.4%) phloretin hydrolase (EC 3.7.1.4) of *Mycobacterium abscessus* (47) (MaPhlG) showed conserved Zn^{2+}-coordinating and catalytic residues in the Rv1775 protein model (Fig. 4 A,B). These conserved features suggest Rv1775 encodes a hydrolase acting on C— C bonds (EC 3.7.-.-), an uncommon class of catalytic activity(46). The only subsubclass within EC 3.7.-.- is 3.7.1.-, suggesting Rv1775 is a 3-oxoacid carboxylase.

Protein	Length (AA)	TM$_{ADJ}$	Seq. identity	Predicted Ligands	Metal-coordinating triad	Speculated function(s)	Evidence
Rv0036c	257	0.63	14%	L-Glutamic acid; Ni^{2+}	His43, His142, Val146	Intraphagosomal stress response; detoxification	Increased abundance in intraphagosomally-grown bacilli
Rv0738	182	0.75	17%	Zn^{2+}	His43, Asp112, His116	Hypoxic stress response; detoxification	More abundant in biofilm than in planktonic bacilli
Rv1727	189	0.69	12%	Fe^{3+}	His43, Ser121, His125	Iron starvation stress response	Co-transcribed transferrin-sequestering protein (Rv1872c) under carbon limitation
Rv2036	213	0.74	15%	Zn^{2+}	His52, Asp117, His121	Hypoxic stress response; detoxification	Highly correlated (R=0.72) expression with co-operonic transcriptional activator of hypoxic response (DosR) and virulence (PhoP) master regulators

Fig. 4. Manual structural analysis refines functional annotations uncorroborated by HHpred. **(A-B)** Conservation of structure- and sequence-features essential for C—C bond hydrolysis supports the inferred hydrolase function of Rv1775. **(A)** Structural alignment of modeled Rv1775 and its closest structural match (PDB 3hpw), a 2,4-diacetyl-phloroglucinolhydrolase of *Pseudomonas fluorescens* (*Pf*PhlG), The structures are superposed (top). Zoomed and re-oriented images of *Pf*PhlG Zinc-coordinating (box on left) and catalytic (pop-out) residues superposed with analogous *Mt*Rv1775 residues. **(B)** Comparison of functional and structural features between *Mt*Rv1775 and a putative PhlG homolog of *M. absecssus* (*Ma*PhlG), phloretin hydrolase, which catalyzes C—C bond hydrolysis of a different substrate. Comparison carried out in a similar scheme as in **A**. Superposition of the putative

homologs, color-annotated with conserved residues essential for phloroglucol moiety recognition and for phloretin substrate specificity in *Ma*PhlG(47). The structural similarity and conserved Zinc-coordinating and catalytic residues affirm Rv1775 as a *bona fide* C—C hydrolase, potentially with a substrate that includes a phloroglucol moiety, but likely not phloretin. Conservation of structure- and sequence-features characteristic of DinB-like metalloenzymes exemplified by structural homology of Rv2036 and a mycothiol-dependent maleylpyruvate isomerase from *Corynebacterium glutamicum* (*C. glu* MDMPI) (**C-G**). (**C**) Superposition of Rv2036 structure model and *C. glu* MDMPI (PDB: 2nsf). Conserved Zn^{2+}-coordinating (pink) and catalytic residues are highlighted (green). (**D**) Highly conserved residues Arg^{222} (C-terminal domain, Arg^{193} in Rv2036) and Asp^{151} (N-terminal domain, in Asp^{124} Rv2036) are in close proximity (4.22Å), suggesting conservation of their proposed role as inter-domain protein stabilizers(51) (**E**) Spatial conservation of Zn^{2+}-coordinating residues of the catalytic triad (Asp and Glu are observed interchangeably) are consistent with conserved catalytic function. (**F**) Surface hydrophobicity of Rv2036 model and 2nsf shows that the hydrophilic core proposed to underlie MDMPI catalysis(51) is relatively conserved. (**G**) Structure-based sequence alignment of Rv2036 and *C. glu* MDMPI with conserved residues were manually annotated according to prior work(51). (**H**) Summary of relevant genomic context potentially informative of function, protein similarity metrics between putative *M. tub* MDMPI homologs and *C. glu* MDPMI, and predicted protein features. All structural images were rendered in PyMol. Structurally homologous sequence alignments are based on TM-align(22) (****<5Å between residues; *<10Å between residues).

Although the precise substrate(s) of Rv1775 are indiscernible from structural comparison alone, examining its structure suggests a potential role in lipid metabolism. It shares the phloroglucinol moiety recognition residues conserved across R-phloroglucinol hydrolases but lacks conserved residues required for phloretin hydrolysis (Fig. 4B). This suggests Rv1775 is not a phloretin hydrolase but may act on substrate(s) containing a phloroglucinol moiety or similar aromatic chemical species. Considering reports of *M. tuberculosis* utilizing cholesterol as a carbon source(48), known C—C hydrolytic enzymes in *M. tuberculosis* cholesterol catabolism(49), and gaps in the current understanding of cholesterol catabolism(50), cholesterol ring species are plausible C—C hydrolysis substrates.

In the second example (Fig. 4C-H), we examine one of four HHpred-uncorroborated proteins structurally resembling mycothiol-dependent maleylpyruvate isomerase (MDMPI, a DinB superfamily protein; PDB: 2nsf) of *Corynebacterium glutamicum* (*C. glu* MDMPI). This example illustrates the case when manual inspection corroborates conserved structural features yet

precise molecular function remains indiscernible. Manual structural analysis of the putative

MDMPI homologs validated that—despite low sequence homology (12-17% similarity)—

structural features characteristic of DinB-like enzymes are conserved (shown for Rv2036, Fig. 4).

All four putative DinB-like enzyme were highly structurally similar to *C. glu* MDMPI (TM_{ADJ} = 0.63-

0.75, Fig. 4C) with a conserved hydrophilic core (Fig. 4F), predicted metal-binding sites (Fig. 4E),

retained catalytic triad residues(51) (Fig. 4G), and conserved residues that form a salt bridge

between the C- and N-domains(51) of MDMPI (Fig. 4D). However, DinB superfamily proteins

comprise several functions (52), making even putative inference of a precise molecular function

challenging. Most functionally characterized bacterial DinB-like enzymes are thiol-

dependent(52) and the putative MDMPI homologs' closest structural match was a mycothiol-

dependent DinB-like enzyme, suggesting thiol-dependence of these four proteins is probable,

likely with mycothiol as the thiol-cofactor (the predominant mycobacterial low molecular weight

thiol). We annotated these genes as "putative thiol-dependent DinB-like metalloenzymes" and

note as "potential (myco)thiol-dependent S-transferase (EC 2.-.-.-)" (53). For such cases, where

structural modelling confidently ascribes protein family and features of structure but not

function, integrating knowledge of the function of structural orthologs', expression data, and

genomic context can inform rational speculation about their function (Fig. 4H, *Text S1*).

Hundreds of annotations inferred by structural similarity. Our structural annotation

pipeline inferred function from structure for 400/1,725 under-annotated genes (23.2%, Dataset

1). Structure-derived annotations (mean C-score = 0.39) came from higher quality models (*P* =

1.83×10^{-163}, student's t-test) than proteins without passing matches (mean C-score = -1.91),

and more specific annotations tended to come from higher-quality models (*Text S1*, Fig. S5).

Structure-based annotation captured putatively shared function for numerous previously

unannotated proteins lacking appreciable sequence similarity (Table 4, Dataset 3).

Table 4. Novel annotations transferred through structural similarity despite low sequence similarity. Selected
proteins with modelled structures highly similar to solved PDB structures of known function. Sequence similarities
range in the "twilight zone" of sequence similarity, below which remote homology is undetectable by sequence
similarity(126). TM_{ADJ} above 0.52 indicates that template and under-annotated gene share structural folds.
Annotations from UniProt, Mtb systems Portal, and TubercuList are shown, along with the highest error-adjusted
structural similarity match, its identifier ("PDB"), and final product annotation. "Affirmatory" corroborate the
annotations in UniProt or Mtb Network Portal. "Novel" are entirely novel annotations to those in UniProt and Mtb
Network Portal, while "More specific" are in accord with annotations in other databases but describe product
function in greater detail.

Rv number	Top I-TASSER hit	AA%	TM_{ADJ}	PDB	Final annotation	MycoBrowser	UniProt	Mtb Net Portal	Type
Rv1139c	Integral membrane Methyltransferase	18	0.86	4a2n	putative integral Membrane Methyltransferase	Conserved hypothetical Membrane protein	Conserved hypothetical membrane protein (Membrane protein)	None	Novel
Rv1766c	Copper-sensing transcriptional Repressor CsoR	29	0.84	4m1p	Putative Transcription factor	conserved protein	Conserved protein	None	Novel
Rv3192c	5,10-methylenetetrahydro Methanopterin Reductase	16	0.83	1z69	Putative Monooxygenase	Conserved hypothetical alanine and Proline-rich protein	Conserved hypothetical Alanine and Proline-rich protein	Oxidoreductase	More specific
Rv2141c	M20 family Metallo peptidase	20	0.82	2pok	putative linear amide hydrolase	conserved protein	conserved protein	FIG016551: Putative peptidase	Affirmatory
Rv1775	2,4-diacetylphloroglucinol hydrolase	29	0.82	3hwp	putative 3-oxo-carboxylic acid hydrolase	Conserved hypothetical protein	Uncharacterized protein	None	Novel
Rv0052c	Isonitrile hydratase	33	0.81	3noo	putative hydro-lyase/putative deglycase	Conserved protein	Conserved protein	ThiJ/PfpI Family protein	Novel
Rv2036	mycothiol-dependent maleylpyruvate isomerase	15	0.73	2nsg	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein	DinB family protein	None	More specific

395
396

397 These remote homologs and structural analogs include an integral membrane methyltransferase,

398 which can modify of mycolic acids embedded in the *M. tuberculosis* cell wall essential for

399 virulence(54) and redox response-related functions (Rv0052 and Rv3192) critical for enduring

400 host immune defenses in macrophage(55).

Putative efflux and transport proteins uncovered through structural similarity. Membrane-spanning regions of transport proteins vary in sequence relative to structure far more than globular proteins(12, 56), making them good subjects for structure-based functional inference. Twenty-four proteins were identified as transport proteins and corroborated by HHpred (Dataset 3), including several matches with drug transport proteins (n=8). Eight HHpred-corroborated proteins were not annotated with any transport function in UniProt (Table 5). Rv1510 and Rv3630 exclusively match drug transporters and are uncharacterized across functional databases. Rv3630 mutations have been reported in POA-resistant mutants but no clear causal link was identified(57). Rv1510 is a *Mycobacterium tuberculosis* complex marker in diagnostic assays(58) and its loss-of-function induces autophagy(59), suggesting Rv1510 is an autophagy antagonist important for human-adapted tuberculosis. Verapimil, a potent efflux pump inhibitor, induces autophagy(60), consistent with the putative function of Rv1510 in drug efflux, which could contribute to drug tolerance(58). These putative transporters might contribute to intrinsic efflux-mediated drug resistance and tolerance in *M. tuberculosis* (11). Other putative novel transport proteins may serve important homeostatic roles in the dynamic host microenvironment(61, 62), and could make attractive drug(63) and vaccine(64) targets.

Table 5: Putative transport proteins. Matches between proteins encoded by under-annotated genes (Locus) and transport proteins structure entries on Protein Data Bank (PDB). Only matches undescribed as transport proteins on Uniprot are included (see Dataset 3E for all such matches). The top two matches are shown, if they exceed adjusted TM-score (TM$_{ADJ}$) > 0.52 (the TM-score corresponding to matching topologies > 50% of the time). AA% refers to the amino acid identity shared between the aligned region of the protein in *M. tuberculosis* its match on PDB.

Rv number	Product annotation	AA (%)	PDB macromolecule name	PDB ID	TM-score	TM$_{ADJ}$
Rv1085c	putative membrane transporter receptor protein	0.073	Chloride pumping rhodopsin	5b2nA	0.71	0.59
		0.058	Sodium pumping rhodopsin	4xtlA	0.71	0.59
Rv1462	putative transporter	0.173	ABC transporter, ATP-binding protein	4dn7A	0.79	0.67
Rv1510	putative Na+/H+ antiporter drug efflux protein	0.104	Putative drug/sodium antiporter	4z3nA	0.89	0.60
		0.088	Multi antimicrobial extrusion protein (Na(+)/drug antiporter) MATE-like MDR efflux pump	3mktA	0.84	0.54
Rv1680	putative phosphonate transporter component	0.151	PhnD, subunit of alkylphosphonate ABC transporter	3p7iA	0.91	0.77
		0.165	Binding protein component of ABC phosphonate transporter	3n5lA	0.89	0.76
Rv2325c	putative transport protein	0.165	Putative cobalt ABC transporter, permease protein	5d3mD	0.87	0.60
		0.215	Energy-coupling factor transporter transmembrane protein EcfT	4huqT	0.81	0.54
Rv2508c	putative MFS membrane transporter	0.104	Solute carrier family 2, facilitated glucose transporter member 3	5c65A	0.79	0.63
		0.119	D-xylose-proton symporter	4gbyA	0.78	0.63
Rv3630	putative Na+/H+ antiporter drug efflux protein	0.087	Multi antimicrobial extrusion protein (Na(+)/drug antiporter) MATE-like MDR efflux pump	3mktA	0.84	0.63
		0.108	Putative drug/sodium antiporter	4z3nA	0.83	0.62

An updated *M. tuberculosis* reference genome functional annotation. Through manual curation (n=282) and structural inference (n=400), we annotated 623 gene products, reducing under-annotated genes by 36.1%. Including annotated CATH (Class, Architecture, Topology and Homologous superfamily) topologies, functional notes, and Ligand-Binding Sites (LBS) totals 940 (54.5%) with original annotation (Fig. 5B). For genes lacking specific product annotations CATH (Dataset 3L) and LBS assignments (Dataset 3D) can refine functional hypotheses and, in some cases, imply function directly(65). Tetracycline repressor folds (n=17, Dataset 3M), for instance, function nearly exclusively as concentration-dependent transcriptional activators and vary in sequence yet are structurally homogeneous(66). CATH annotations were not used to inform product annotations nor to assign EC numbers in this annotation, however.

Fig. 5. An updated H37Rv functional annotation. (A-C) Circos plots illustrating annotation coverage prior to the annotation effort (left) and following it (right), colored according to annotation status. In plots A and C, all 4,031 CDSs are represented as segments of equal width whereas (B) segments the ring into only the 1,725 under-annotated genes. Black genes reflect what was on TubercuList, are considered "annotated", and are mutually exclusive from the 1,725 under-annotated genes (white). (B) shows only the 1,725 under-annotated genes, whereas A and C include all 4,031 original CDS. Inside the circos rings are stacked bar charts with genes in 100kb bins according to gene start position. The terminal-proximal (± 250 kilobases) region is marked with dashed lines and labelled (*ter*). (D) Cumulative number of genes annotated, by annotation type. LBS = Ligand binding site. Percentages refer to under-annotated genes annotated/1,725 initial under-annotated genes. Genes are binned into mutually exclusive categories hierarchically: manually curated product name > structure-derived > literature notes > CATH > LBS. Manually curated and literature notes categories are combined as "Literature-curated" in the visualization. For the purposes of these counts, functional notes from publications implicating many proteins but not clearly establishing function were not counted (e.g. PMIDs 26536359, 21085642, and 22485162).

Our updated annotation provides function for 34.4% (45/131) of genes with hypothetical function identified in a recent systems resource as broadly conserved across *Mycobacteria*(67) (Dataset 3 contains full set). Mycobacterial core genes annotated include functions well-established experimentally, such as essential component of the mycobacterial transcription initiation complex RbpA (https://gitlab.com/LPCDRP/Mtb-H37Rv-annotation/-/blob/master/features/Rv2050.tbl), and others not evident from extant literature but of potential clinical relevance, like the host-directed effector function inferred for Rv3909 (https://gitlab.com/LPCDRP/Mtb-H37Rv-annotation/-/blob/master/features/Rv3909.tbl). These annotations came in similar numbers from published experimental evidence (n=21) and structural inferences (n=24).

Updated annotations distribute across all segments of the chromosome (Fig. 5A) and implicate efflux proteins (Table 5), metabolic functions (Fig. 6), virulence factors, and functions key to survival during infection (Table 6) and under drug pressure (Table 7). Yet many under-annotated genes remain without products or functional notes assigned (n=785). Of these 785 remaining under-annotated genes (Dataset 1), 190 have quality models (C-score > -1.5) but lacked annotations meeting inclusion criteria. Meanwhile, 182 of those remaining have product annotations in UniProt or *Mtb* Network portal. Remaining still, however, are 466 under-annotated genes with neither quality structure nor functional annotation in these databases. These genes frequently cluster consecutively along the genome (99 genes across 23 clusters, Dataset 1G), forming syntenic blocks of unknown function. Genomic context suggests several of these clusters have roles in virulence and drug tolerance (Dataset 1G).

470
Fig. 6. Functional annotations across *M. tuberculosis* metabolism. Annotated EC numbers for manually curated
and structure-inferred products were compared with annotations for each under-annotated gene in popular
databases **(A)** Set analysis of under-annotated genes (UAG) with an EC number assigned in this study compared to
popular databases. **(B)** Novelty of EC numbers for UAG annotated in this study with respect to popular databases.
(**C & D**) Distribution of EC numbers annotated across KEGG (**C**) Subsystems and (**D**) Pathways. Generic KEGG
subsystems are depicted. All pathways with at least three genes have the number of EC numbers displayed. For
subsystems with no pathways with three or more genes, the highest total pathway is displayed.

Table 6. Updated annotations add functional knowledge to genes required for optimal fitness during TB infection. Source annotation is the annotation listed by Bellerose and colleagues(83) and new annotation derived from the current project. Protein data Bank identifiers (PDB ID) of the protein structures matching H37Rv protein models are listed for structure-based annotations. PubMed IDs are listed for the papers from which functional annotations were manually curated.

Rv number	Source annotation	New annotation	PubMed ID	PDB ID
Literature annotations				
Rv1205	hypothetical protein	riboside monophosphate phosphoribohydrolase	25728768	
Rv2018	hypothetical protein	probable antitoxin VapB/Antigen	28066388, 23140854	
Rv2272	transmembrane protein	Probable gamma delta T-cell activator	23389928	
Rv2525c	tat pathway signal sequence	probable peptidoglycan hydrolase	16952959, 25869294, 25260828	
Rv2923c	hypothetical protein	probable Osmotically induced bacterial protein C (OsmC, a probable peroxide reductase)	22088319	
Rv3632	membrane protein	Putative flippase	21030587	
Rv3763	lipoprotein LpqH	adhesin/antigen LpqH	16098710, 12594264	
Rv3788	hypothetical protein	probable secondary channel binding-factor of RNA polymerase	22194445	
Literature & Structural inference				
Rv0191	integral membrane protein	Putative efflux pump	25690361, 12520088, 22132058	5c65, 4gby
Rv1433	exported protein	Probable l,d-transpeptidase LdtMt3	24041897	3tur, 3vae, 4jmn
Rv1769	hypothetical protein	T-cell antigen/putative aldehyde-lyase	26853625, 15102765	4v15
Rv3722c	hypothetical protein	probable serine hydrolase, probable aspartate transaminase		3ppl
Structural inference				
Rv0047c	hypothetical protein	Putative transcriptional regulator		1yg2a, 3l9f
Rv0259c	hypothetical protein	Putative lyase		2jh3, 4ccs
Rv0323c	hypothetical protein	Putative hydro-lyase/Putative linear amide hydrolase		1q7t, 5cgz
Rv0449c	hypothetical protein	Putative oxidoreductase		2ive, 1sez, 3nks, 3i6d, 3lov
Rv0767c	hypothetical protein	putative transcription factor		3mnl
Rv1085c	hemolysin-like protein	putative membrane transporter receptor protein		5b2nA, 4xtlA
Rv2052c	hypothetical protein	putative endodeoxyribonuclease		3igh
Rv2160A	hypothetical protein	putative transcription factor		2hyj
Rv3226c	hypothetical protein	putative peptidase		2icu
Rv3433c	hypothetical protein	putative hydro-lyase/putative isomerase		2ax3
Rv3719	hypothetical protein	putative amide-bond oxidoreductase		3dq0, 2exr, 4o95

Table 7. Updated annotation enriches functional interpretation of under-annotated genes affecting drug sensitivity. Source annotation is the annotation listed by Bellerose and colleagues(83). Enriched conditions are the drugs' exposure under which differential mutant abundance was observed. Sources of updated annotation are listed in the evidence column. INH=Isoniazid; RIF=Rifampicin; EMB=Ethambutol; PZA=Pyrazinamide; HRZE=Combination regimen of INH, RIF, PZA, and EMB. "(+)" indicates enrichment observed at multiple time points.

Rv number	Source annotation	New annotation	Enriched condition(s)	Evidence
Rv0998	hypothetical protein	cAMP-dependent lysine acetyltransferase	EMB, RIF, HRZE	23553634
Rv1205	hypothetical protein	riboside monophosphate phosphoribohydrolase	EMB (+)	25728768
Rv0767c	hypothetical protein	putative transcription factor	INH	3mnl, 3bjb (PDB ID)
Rv3131	hypothetical protein	putative nitroreductase	INH	27094446; 28261197
Rv2140c	hypothetical protein	Phosphatidylethanolamine-binding protein TB18.6	INH	23907008; 27895634; 26238929
Rv2061c	hypothetical protein	probable serine hydrolase	INH	26853625; 26536359
Rv3267	hypothetical protein	probable peptidoglycan-arabinogalactan ligase	INH	27486192
Rv1770	hypothetical protein	probable serine hydrolase	PZA	26853625
Rv3005c	hypothetical protein	Probable membrane oxidoreductase component (MRC) DoxX	RIF	26067605
Rv1184c	exported protein	mycoacyltransferase	RIF	25331437; 25124040
Rv3036c	secreted protein	secreted esterase	RIF, INH	25224799
Rv2190c	hypothetical protein	peptidoglycan peptidase RipC/Antigen	RIF	24843173; 22952680; 28241799
Rv0079	hypothetical protein	Putative Dormancy Associated Translation Inhibitor (DATIN)	RIF (+)	22719925; 23819907; 28261197

Genes remaining without any form of annotation (Fig. 5D) were overrepresented (P = 0.0011, odds ratio=1.35, Fisher's Exact) near the terminus (±250 kilobases from half the genome length) of the chromosome (*ter*-proximal genes, Dataset 1H). An even stronger bias for uncharacterized genes can be seen for genes transcribed opposite the direction of replication (P = 1.14 x 10^{-7}, odds ratio=1.53; Fisher's exact). To ensure that circumstantial factors such as PE/PPE or insertion element density were not accounting for the apparent orientation and spatial trends across the chromosome, we removed all PE/PPE and insertion seq and phage genes and repeated the analysis. The trend strengthened for the *ter*-proximal gene bias (P = 0.0034, odds ratio=1.44; Fisher's exact) and decreased only marginally (P = 2.53 x 10^{-6}, odds ratio=1.49; Fisher's exact) for the orientation bias.

These biases are consistent with three previously noted trends that could influence the

likelihood of gene characterization. First is the general trend of decreased gene expression as a

function of distance from the *oriC* in bacteria(68). On average, highly expressed genes are more

amenable to functional characterization. Second is the strong bias for symmetric inversions

around the terminus(69), particularly in Actinobacteria(70). Hypotheses leading to experimentally

determined functions are often informed by orthology, which can be inferred by conserved

synteny between species(71). Common inversions around the terminus can disrupt this synteny

with increased frequency. Disruption can occur globally—through moving across the

chromosome by inversion—and locally, by inversion boundaries interrupting operons or other

syntenic features. Third, genes transcribed opposite the direction of replication frequently collide

with the replication machinery, making them more mutable than genes with transcription and

replication co-oriented(72). This increases the likelihood of weakened promoters or loss-of-

function mutations evolving *in vitro* for genes non-essential in H37Rv. One potential confound

is that genes encoding virulence/toxin proteins are enriched on the lagging strand(72). As these

genes operate in the context of infection, they are challenging to functionally characterize, which

may contribute to the observed enrichment of uncharacterized genes on the lagging strand.

Turning our attention to metabolism, 381 under-annotated genes were annotated with EC

numbers (Dataset 1I Methods & Materials), over two-thirds of which were absent from other

databases (Fig. 6A,B). Fully specific (4th EC digit) EC numbers (n=92) were ascribed to 85 genes.

These newly annotated reactions span diverse metabolic pathways and subsystems (Fig. 6,

Dataset 1J), many implicated in mediators of *M. tuberculosis'* virulence such as lipid and

polyketide and terpene metabolism(73–75), which are integral to the unique composition of the mycomembrane. Proteins of these pathways have important immunity-subverting functions(76) at the host-pathogen interface(77). For instance, terpenes play an immunomodulatory role early in *M. tuberculosis* infection and phagosomal maturation(78–80), are potential agonists of antibiotics for TB treatment(81), and include cell membrane surface-expressed molecular species unique to *M. tuberculosis*(82). The numerous carbohydrate-metabolizing products (Fig. 6D) may identify alternative metabolic pathways in *M. tuberculosis* and aid in gap-filling efforts in *M. tuberculosis* metabolic reconstructions.

Integration with recently published functional screens. Next, we assessed how much novel functional information our annotation added to ambiguously or hypothetically annotated genes from a recent transposon mutagenesis study that sought to identify specific bacterial functions limiting drug efficacy during a mouse model of infection(83). We assessed two sets of genes identified in the study. In the first set of under-annotated genes—those newly reported to as essential for optimal growth in mouse infection—one-third (23/69) could be updated by our annotations (Table 6). Fifteen were structural inferences, demonstrating the value structure-based inference of putative function where the difficulty of recapitulating complexities of the host environment challenges functional elucidation through experiment. Notably, following its inference based on structure, Rv3722c has since been confirmed to indeed encode an aspartate transaminase(84) and Rv1085c has been found likely not to encode hemolysin(85), substantiating the structure-derived functional annotations in Table 6.

Our annotation functionally described 13/27 under-annotated genes affecting drug sensitivity (Table 7). Notably, some genes affecting drug sensitivity have published functions consistent with the mechanism of action of the drug of interest but listed without annotation. For instance, the authors noted cell wall permeability as a central theme among genes dictating sensitivity to rifampicin (RIF), and disruption of Rv2190c—a peptidoglycan hydroase—rendered mutants hypersusceptible to RIF, consistent with an effect on cell wall permeability. Others (e.g. Rv1184c) were unannotated in their primary data, but their functional ties was discussed in the text, suggesting the function was curated from literature. Our updated annotation centralizes such functional knowledge.

Structural models enable functional interpretation of novel PZA-resistant mutants. Next, we applied our annotations prospectively to a new resistance screen, querying the molecular basis of pyrazinamide (PZA) resistance in *M. tuberculosis*. PZA is a cornerstone of modern tuberculosis therapy, yet the mechanism by which it exerts its anti-tubercular activity remains elusive. PZA is a pro-drug that must be converted to its active form pyrazinoic acid (POA) by a mycobacterial amidase[86]. While multiple explanations for POA action have been proposed[87–89], many of these models have not held up to scrutiny[90–92]. Recently, several groups have shown that POA either directly or indirectly disrupts mycobacterial coenzyme A (CoA) biosynthesis[93–95]. Identification of novel resistance mechanisms could shed additional light on the elusive action of this drug. Thus, a library of 10^5 transposon-mutagenized *M. tuberculosis* H37Rv was used to select for POA resistant isolates. While the frequency of spontaneous resistance to POA is approximately 10^{-6}, the frequency of resistance from our transposon

mutagenized library was 10^{-3}. Four mutant strains chosen for further characterization of drug

resistance profile and transposon insertion site (Fig. 7) showed insertions in genes of unknown

function. Each of these strains showed ≥2-fold resistance to PZA and POA (Fig. 7A-D) and no

change in INH susceptibility (Fig. 7A-D) when compared to wild-type H37Rv.

Fig. 7. Annotation of genes involved in pyrazinamide resistance. A library of *M. tuberculosis* H37Rv transposon
insertion mutants was used to select for strains that were resistant to POA. The transposon insertion sites were
mapped and the strains were characterized for their susceptibility to PZA, POA, and INH in comparison with wild-
type H37Rv (MIC$_{90}$: 50 µg/mL PZA, 200 µg/mL POA, 0.0625 µg/mL INH). (A) H37Rv *Rv2705c::Tn.* (B) H37Rv
Rv2706c::Tn. (C) H37Rv *Rv3256c::Tn.* (D) H37Rv *Rv3916c::Tn.* Error bar depicts standard deviation across triplicates.
Pop-outs in (B) and (D) depict the transposon insertion sites relative to experimentally determined Transcription
Start Sites(127, 128) (TSS). The insertion in (D) interrupts a TANNNT Pribnow Box (blue), destroying the Rv3916c
promoter. Error bars are shown when there was a deviation in the calculated MIC90 across triplicates.

To interpret how the interrupted genes might contribute to PZA resistance, we inspected the

structural and functional data available from our I-TASSER results (Figs. 8 & 9). PZA is a structural

analog of nicotinamide(96), suggesting the putative nicotinamide binding domain of Rv2705c

(Fig. 7A) may interact directly with PZA or POA. While it remains difficult to confidently annotate

Rv2706c (Fig. 7B), considering its position immediately upstream of Rv2705c it may alter PZA

sensitivity by influencing expression of Rv2705c.

Rv3256c structurally resembles multiple phosphosugar isomerases—particularly

phosphoglucose (PGI) and phosphomannose (PMI)—and Glutamine-fructose-6-phosphate

transaminases (GlmS). Rv3256c has neither the conserved residues essential for PMI/PGI catalysis

(97) (Fig. 8A,C) nor the glutamine amidotransferase domain required for GlmS activity (98) (Fig.

8B,D), effectively ruling out these functions. The common structural feature among these

functionally disparate matches is a sugar isomerase (SIS) domain (Fig. 8). SIS domain is a

phosphosugar-binding module(99), implicating Rv3256c in phosphosugar metabolism or its

regulation. Rv3256c lacks the HTH domain common to RpiR-like SIS domain proteins(99) that

regulate phosphosugar metabolism genes, refuting the possibility of a RpiR-like transcriptional

regulatory function. Flanking Rv3256c (Fig 7C), however, are mannose donor biosynthesis

genes—Rv3255c (a PMI) and Rv3257c (a phosphomannomutase). In *M smegmatis*, Rv3256c

overexpression decreased cell-surface mannosylation(100), consistent with a role in regulating

phosphosugar metabolism, though a specific molecular function for Rv3256c remains unclear.

Presumably, the role played by Rv3256c in phosphosugar metabolism is disrupted in the

Rv3256c::Tn mutant. This disruption may alter composition of acyl-CoA/CoA pools (e.g., through

disrupting/promoting acylation of cell wall constituents) or the metabolic and cell wall

restructuring response under CoA depletion(93) induced by PZA treatment.

601
Fig. 8. Structural analysis informs specific functional hypotheses for the basis of PZA resistance in the
***Rv3256c::Tn* mutant. (A-D)** Structural analysis identifies Rv3256c as a sugar isomerase (SIS) domain-containing
protein likely involved in phosphosugar metabolism or its regulation. InterPro functional domains are displayed for
the two strongest structural matches of Rv3256c, (**A**) *P. aerophilum* bifunctional phosphoglucose/phosphomannose
isomerase (*P. aer* PGI/PMI) and **(B)** *E. coli* glucosamine-6-P synthase (*E. col* GlmS). The InterPro domains labelled in
A and B are mapped onto the 3D structures of (**C**) *P. aer* PGI/PMI and (**D**) *E. col* GlmS. Rv3256c (charcoal) modeled
protein structure is optimally superposed to each of its matches. Rv3256c is structurally homologous to the SIS
domains of *E. col* GlmS and *P. aer* PGI/PMI and exhibits the alpha-beta-alpha sandwich fold of SIS(129). The pop-
out in A (**<5Å between residues) and labelled residue in C show the threonine residue essential for isomerase
activity in *P. aer* PGI/PMI (Thr[291]) and other PMI homologs. The Thr[291]-containing region appears to be absent from
Rv3256c. Likewise, Rv3256 lacks a glutamine amidotransferase domain homologous to *E. col* GlmS. From this
structural evidence, we conclude that Rv3256c is a SIS domain protein putatively involved in phosphosugar
metabolism and/or its regulation. Structural images were rendered in PyMol. Structurally homologous sequence
alignments from TM-align(22).

Rv3916c structurally resembles numerous Gcn5-related N-acetyltransferase (GNAT) proteins

and exhibits structural, topological, and local features characteristic of GNAT enzymes (Fig. 9).

Perhaps most tellingly, Rv3916c has a predicted acetyl-CoA binding site consistent with known

GNAT enzymes (Fig. 9B,C). While the acyl donor and substrate of Rv3916 remain unclear, the

conservation of these structural features involved acyl-CoA interaction in functionally

characterized GNAT superfamily proteins strongly suggest that Rv3916 is a GNAT N-

acetyltransferase. This putative function implicates the $_P$Rv3916c::Tn mutant in the CoA-depletion

model of POA action. Destruction of the Rv3916c promoter would reduce its expression, in turn

altering acyl-CoA pool modulation. Structure-function insights gleaned from these structural

models inform specific functional hypotheses for these mutants' role in PZA resistance and

demonstrate how the provided structural data can enrich the interpretation of large-scale

screens and generate specific functional hypotheses.

Fig. 9. Structural analysis of transposon mutants refines functional hypotheses for their role in PZA resistance. (A-C) Structural analysis supports Rv3916c as a General control non-repressible 5 (GCN5)-related *N*-acetyltransferases (GNAT). Rv3916c matches exclusively comprised GNAT enzymes with low sequence similarity, typical among homologous GNAT enzymes (130). **(A)** Rv3916c (orange) superposed with its closest structural match (PDB 2a4n, charcoal), *Enterococcus faecium* aminoglycoside 6'-N-acetyltransferase (*E. faec* AAC(6')-Ii). (**B**) *M. tuberculosis* Rv3916c structural model (top) and secondary structure topology (bottom), colored according to the four core conserved GNAT folds (wheat = poorly conserved secondary structure element) and secondarily (tones of the primary colors) by secondary structure elements in *E. faec* AAC(6')-Ii (as defined by (131)) with which Rv3916c structurally aligns. All secondary structure elements of *E. faec* AAC(6')-Ii are present in Rv3916c, with two gratuitous alpha-helices in its C-terminal arm (which is not well-conserved among GNAT enzymes(130)). The distinctive β-bulge (pop-out) within the β4 strand (red, bulge residues colored blue) characteristic of GNAT enzymes is present in Rv3916c, diverting β4 away from β5 to create the chasm where the acetyl-coenzyme A (yellow) is predicted to bind (**C**) Primary sequence of Rv3916c colored according to the scheme described in (B)**.** Asterisks mark predicted acetyl-CoA binding residues (black) and residues structurally aligning to known CoA-interacting residues (blue). All known CoA-interacting residues from *E. faec* AAC(6')-Ii are conserved in Rv3916c and all predicted acetyl-CoA binding sites coincide or directly flank demonstrated sites of CoA interaction. The presence of the features in Rv3916c suggest it

is a GNAT enzyme. GNAT enzymes catalyze transfer of an acyl moiety from an acyl-CoA to various substrates (130)
making Rv3916c a probable acyl-CoA acetyltransferase and implicating $_p$Rv3916c::Tn in CoA pool modulation model
of PZA resistance. **(D)** Summary of functional hypotheses for PZA resistance-conferring transposon mutants. All
structural images were rendered in PyMol. Structurally homologous sequence alignments are based on TM-
align(22).

Discussion

Functional genome annotation is critical for interpreting the deluge of omics data generated by

emerging high throughput technologies. Here, we devised procedures to systematically curate

annotations from published literature and infer putative function through structure-based

function inference and applied them to annotate the *M. tuberculosis* virulent type strain and

primary reference genome, H37Rv. We curated annotations for hundreds of proteins with

published functions lacking from common resources (Table 2), a quarter of which were absent

from all five annotation resources examined, highlighting the importance of community-specific

manual curation. To complement these manually curated annotations, we built a structural

modeling and functional inference pipeline (Figs. 1-4), calibrated it to include confident

annotations (Fig. 1), orthogonally validated it with established remote homology detection

methods (Fig. 3). Through structure-function inference we annotated hundreds of genes (Fig. 5),

including dozens of potential transport proteins, resistance genes, and virulence factors (Dataset

1).

Elucidating the determinants of *M. tuberculosis'* survival under drug pressure and within the

context of infection are chief objectives of tuberculosis research. Integrating this updated

functional annotation with new (Figs. 7-9) and published (Tables 5 & 6) functional screens

showed it can aid understanding the genetic basis of *M. tuberculosis'* resistance to drug pressure

and infection-like conditions. The structural models provided a rational basis for functional hypotheses of the molecular basis of resistance for four novel PZA-resistant strains with mutations in otherwise unannotated genes (Figs. 7-9). In particular, the Rv3256c and Rv3916c mutants implicate CoA homeostasis in PZA resistance (Figs. 8 & 9), linking them to the CoA depletion model of PZA mode of action described recently for other PZA-R mutants (93). The functional interpretation of TnSeq mutants afforded by this resource informs hypotheses for the mechanistic basis of these mutants' PZA resistance for investigation in future work.

Our systematic approach to manual literature curation has limitations. First is the time and attention from researchers with specialized knowledge required for manual literature curation. To mitigate this limitation in the future, contributions from the TB research community can be submitted, and will be incorporated with standardized criteria and structured ontologies(24, 28). A second limitation is the inevitable subjectivity of the curator. We addressed this by requiring two curators review each paper independently and provided explicit guidelines for what evidence warrants annotation with the degree of confidence connoted systematically by qualifying adjectives.

Other limitations arise from the scope of our annotation. First, we only curated functions for 1,725 of over 4,000 ORFs in *M. tuberculosis*. Products that did not meet our criteria for inclusion may have useful functional characterizations excluded by our approach. Second, we only searched for the locus tag during curation. While most publications include locus tag, some do not and therefore some experimental characterizations may remain unannotated. Last, we searched literature back from 2010, as TubercuList updated continuously through March 2013,

and we assumed annotations to that point were captured. However, the absence of dozens of characterizations from all resources suggests some findings prior to 2010 may remain unintegrated. Despite these limitations, the numerous genes we curated that were absent from all frequented annotation sources are now centralized in an single updated annotation that is clear in source and confidence level, in a consistent and extendable format.

Multiple factors contribute to error and bias in resolving protein structure and function. These factors fall unevenly across protein classes and families(101), making them challenging to account for. Considering this while designing our structure-based inference pipeline, we favored simple, interpretable inclusion criteria, coupled with downstream quality assurance measures. Future work more focused on customizing inclusion criteria optimized for features of protein structure or function may improve prediction accuracy. Our simplified approach let us circumvent accounting for these biases explicitly, which would require further methods development and introduce additional bias if not executed carefully.

Our structural approach to functional inference also has limitations. First, it depends on the input sequence. We took amino acid sequences as provided by TubercuList without accounting for the impact of known, uncorrected sequencing errors(102) or corrections to amino acid sequences proposed by UniProt curators. Furthermore, some genes have multiple translation initiation sites, or isoforms(103), but we considered one sequence per gene. Second, our approach compares global, rather than local, structural similarity and can be challenged by functionally diverse folds(104), and proteins with dynamic active sites(105) or context-specific conformation and activity(106). Our empirically-driven inclusion criteria (Fig. 1) and quality

control measures helped to mitigate false positive annotations (Figs. 3 and 4). In future analysis of structural models, emerging methods that capture functional conservation distributed across primary and tertiary structure may identify functionally informative protein features missed by our approach. Promising approaches include direct coupling analysis[107], statistical coupling analysis[108], Bayesian partitioning with pattern selection[109], and structurally interacting pattern residues' inferred significance[110]. Third, proteins from model organisms and humans are overrepresented among crystallized structures on PDB (https://www.rcsb.org/) [111]. This adds bias toward inferring function from these proteins. Finally, the structure-based annotations should be interpreted as tentative, since inclusion criteria required similarity implying >50% ("putative") or >75% ("probable") likelihood of being correct. Structure-based annotations should be viewed accordingly, as well-informed hypotheses rather than established truth.

 Over half of structure models (871/1,711) were low-quality (C-score < -1.5)[21] (Dataset 1). Several phenomena may challenge effective modeling of these under-annotated genes: 1) No proteins of similar folds have been solved; 2) The protein is highly disordered[112]; 3) These are multi-domain proteins that need to be split into individual domains[14]; 4) Sequencing errors; 5) gene coordinate misannotation[102]; 6) Pseudogenization. We suspected reason 3 as a major factor, considering we did not attempt to break up multi-domain polypeptides into their constituent domains[14]. However, the protein length distributions of proteins of high (greater than -1.5) and low (below -1.5) C-scores were similar (*Text S1*, Fig. S5C), which suggests the presence of multiple domains was not a primary cause of poor models. Each of the other reasons

likely contributes to some extent, but reasons 1 and 2 are most troublesome for PE/PPE genes

and other protein classes specific to *Mycobacteria*.

I-TASSER failed to produce models for 14 under-annotated genes (Dataset 1). Six of these

sequences are pseudogenes and the remaining 8 belong to PPE or PE_PGRS gene families, which

are especially prone to sequencing errors and intrinsically hypervariable[113]. Although we

ascribed putative functions for some PE/PPE genes the function of most remains unclear. Far

fewer PE/PPE proteins (20/166, 12.0%) than non-PE/PPE genes (518/1,559, 33.2%) met inclusion

criteria for structure-based annotation (P = 2.13x10^{-9}, odds ratio = 0.276; Fisher's exact). This

likely owes partly to their intrinsic disorder and partly to their specificity to the *M. tuberculosis*

complex[114], which limits the number of homologous structures in PDB with known function,

challenging accurate structural modeling and structure-based functional annotation. Moreover,

PE/PPE and other effector proteins require precise metabolic contexts or immunological cues,

precluding observation of their function *in vitro*. Characterizing function for these genes will

require high-throughput biochemical assays and development of techniques that directly assay

or precisely reconstruct host microenvironments; formidable challenges, indeed. In the

meantime, carefully designed and caveated inferential methods can make valuable surrogates

and streamline candidate prioritization for experimental confirmation or more comprehensive *in*

silico analysis.

Systematically curated literature and structure-derived annotations are available at

https://gitlab.com/LPCDRP/Mtb-H37Rv-annotation. Researchers can file issues to report future

published characterizations and submit merge requests to incorporate future functional

characterizations. These methods can continue to furnish annotations as functional

characterizations are published in the primary literature, structure-function relationships in PDB

expand, *M. tuberculosis* gene product functions are determined, and sequence-structure-

function prediction tools become more resource-efficient.

Author contributions SJM, AE, and FV conceptualized the overarching research goals and

aims; SJM, AE, DG, AMZ, NK, and CKC developed the manual curation protocol; SJM, DG, AMZ,

NK, and CKC manually curated annotations from literature, and independently validated

annotations; SJM, AE, DG, AMZ, NK, and CR developed and tested code for data processing and

analysis; AE and SJM developed the structural comparison workflow; SJM, AE, and ND developed

and executed the structural QC workflow; NAD and ADB conceptualized the PZA resistance

experiments; NAD isolated and characterized the PZA resistant mutants; SJM, AE, DG, AMZ, NAD,

and ND prepared the manuscript; SJM, DG, AMZ, and NAD prepared figures and tables; SJM and

FV managed the project.

Materials and Methods

Additional detail on methods are provided in *Text S1*.

Data availability. We provide final annotations in common machine (GFF3) and human (Dataset

1) readable formats, including EC numbers, GO terms, CATH topologies, and product name

annotations. Annotations in the GFF3 are defined by our inclusion criteria. PDB templates with

structures similar yet below our criteria are provided in Dataset 3 (top 3 PDB templates for each

under-annotated gene) and Dataset 3 (all matches where $TM_{ADJ} > 0.52$ (Equation 2, Text S1)

and/or μ_{geom} > EC3 (putative) threshold). I-TASSER results and model protein structures for 1,711

under-annotated genes are freely accessible at https://tuberculosis.sdsu.edu/structures/H37Rv/

including functional predictions by COFACTOR, predicted ligand binding sites, local secondary

structure confidence (B-factor), and other quality and similarity metrics.

Manual curation protocol. All publications mentioning each of the 1,725 under-annotated

gene were independently evaluated for annotation-worthy functional characterization by two

researchers and quality checked by a third for format and protocol compliance (*Text S1*, Fig. S1).

Qualifying adjectives were defined by evidence quality and systematically assigned to connote

annotation confidence. Notes relevant to function but insufficient to assign product name were

also annotated (*Text S1*, Fig. S1).

Precision benchmarking. We designed procedures and inclusion criteria to maximize precision

(Equation 1) and minimize "Overannotation"(101): Only annotations with 50% or greater

precision were included, regardless of source. Whereas other metrics had applicable precision

benchmarks (*Text S1*), EC number and GO terms did not. We assessed how precision of EC

number and GO term predictions (Equation 2) correlated with similarity metrics. We evaluated

which I-TASSER metrics were most predictive of precision (Equation 1) through logistic

regression (*Text S1*, Fig. S2).

$$precision \ = \ \frac{(TP)}{(TP \ + \ FP)} \tag{1}$$

where TP = "True Positive", and FP = "False Positive.

We gauged sequence similarity by amino acid identity (AA%) and structural similarity by

Template Modeling Score (TM-score). TM-score describes structural similarity from 0 and 1. It

represents the average root-mean squared deviation across all atoms in the predicted structure

with respect to the PDB template model, normalized to remove apparent deviation arising falsely

due to local differences(14, 115) (Methods & Materials), allowing proteins of different length to

be compared(115).

To base inclusion criteria off precision (Equation 1), we regressed against sequence and

structural similarity metrics: amino acid identity, C-score, TM-score (Equation 1, as calculated by

Zhang and Skolnick(115)), and the geometric mean of TM-score and AA% (μ_{geom}) against

precision of EC number assignment in a positive control set of 363 *M. tuberculosis* genes with

known function but unknown structure. Because EC numbers and GO terms encode the same

fundamental information, although GO terms have many false negatives and were relatively

underpowered (*Text S1*, Fig. S2), we included both according to the same criteria: μ_{geom} values

corresponding to 50% ("putative") and 75% ("probable") precision for each tier of specificity (Fig.

1).

$$TM\text{-}score = (1/L_{\mathrm{N}})\sum_{i=1}^{L_{\mathrm{T}}}\frac{1}{(1+d_i^2+d_0^2)} \qquad \{2\}$$

where L_N is protein length, L_T is the length of the aligned residues to the template, d_i is the

distance of the i^{th} pair of residues between two structures after an optimal superposition, and

$d_0 = 1.24\sqrt[3]{L_N - 15} - 1.8$, as described by Zhang *et al.*, normalizes for protein length(21). TM-

score measures the difference between *predicted* structure and known structure of the putative

homolog/analog on PDB. Since we are interested in the similarity between the *true* (unknown)

structure and its putative homolog/analog on PDB, we used an adjusted TM-score, "TM$_{ADJ}$".

TM$_{ADJ}$ subtracts from TM-score the expected difference in TM-score between the modeled

protein structure and its (unknown) true structure (See Equations 2 and 3 and accompanying text

in *Text S1* for additional discussion of this rationale).

Training data selection. We combined 200 randomly selected *M. tuberculosis* protein sequences

with known function with 163 manually annotated under-annotated genes with "probable" or

higher annotation confidence (Dataset 1) to form a set of training genes. We extracted EC

numbers and GO terms that were marked as experimentally verified in UniProt(116) from the 363

training genes.

Annotation inclusion criteria. Structure-inferred annotations comprised product names, GO

terms, EC numbers, CATH topologies, and LBS. We systematically transferred EC numbers and

GO terms according to μ_{geom} thresholds corresponding to 50% and 75% precision (Fig. 1). LBS

and CATH predictions were included according to previous precision benchmarks(21, 117). CATH

annotations were retrieved using the REST API of PDB for structure matches surpassing the TM-

score corresponding to 50% precision, after correcting TM-score for expected modeling error

(S2 Text). Under-annotated genes with quality models (C-score > -1.5) and a TM-score greater

than 0.85 and/or μ_{geom} meeting the inclusion criteria for putative EC 3rd digit (0.374,

corresponding to a precision > 0.5), and further criteria based on aligned portions and method

of UniProt annotation of the PDB template (*Text S1)*. Because transport proteins are more

conserved in structure than in sequence relative to globular proteins(12) we weighted structural

similarity more heavily than AA% in their inclusion criteria: Transport protein annotations were

transferred if (i) greater than 90% of the PDB structure implicated in transport aligned with the

under-annotated gene model, and (ii) structural similarity exceeded the threshold for CATH

topology transfer. All analyses were implemented in R(118).

Product naming protocol. To translate transferred GO terms and EC numbers into product

names we converted GO terms that describe enzymatic activity into EC numbers by searching

EXPaSY ENZYME. Product names were converted from EC numbers (including those derived from

GO terms) using the ENZYME.dat file from the EXPaSY database (*Text S1,* Figs. S3 and S4). GO

terms that mapped to multiple EC numbers, were merged at the most specific level at which they

converged (e.g., 3.2.1.5 and 3.2.2.4 would resolve to 3.2.-.-). When GO terms did not map to an

EC numbers, we translated sufficiently descriptive GO terms into product names (e.g., "DNA

binding transcription factor activity" is sufficiently descriptive whereas "pathogenesis" is not).

Product names for PDB matches lacking GO or EC annotations were determined manually.

Transport proteins were named with lower specificity than their PDB matches (e.g. "transport

protein" instead of "Na+/H+ antiporter"), unless (i) all three strongest PDB matches converged

on a more specific description and (ii) TM-score exceeded 0.85 for at least one of the three, the

name in common between the three strongest matches was transferred. LBS predictions and the

residues predicted to coordinate binding (Dataset 3D) can be interpreted as being at least 60%

likely to be true(23), though most have greater confidence.

For proteins annotated only with structure-based functional inferences that had EC number

annotation modified by HHpred-filtering, or had multiple EC numbers corroborated by HHpred,

the implicated structural homologs were inspected manually, and spurious or infeasible

annotations were pruned. Reasons for pruning EC numbers include cases where one of the implied catalytic functions was exceedingly unlikely (such as eukaryotic proteins with bacterial homologs that had evolved distinct, non-overlapping functions) or there was a clear reason for a false positive (such structural alignment with a multifunctional protein to only one of the functional domains). Additional annotation specificity was added in rare cases, where HHpred results strongly corroborated evidence from structural alignment that alone did not meet inclusion criteria for specific annotation. Rv3433c exemplifies such cases. It was annotated with EC 4.2.1.- and EC 5.-.-.- and both were corroborated by HHpred. Upon inspection, the annotators noted that the top hits from structural alignment and HHpred were a mixture of EC 5.1.99.6 and EC 4.2.1.136 proteins and bifunctional proteins encoding both catalytic functions. The portions aligning to the respective EC functions were mutually exclusive, and Rv3433c was of similar length to characterized bifunctional enzymes including both functions. In this case, EC numbers were updated to full specificity and the product name was changed from "putative hydro-lyase/putative isomerase" to "putative Bifunctional NAD(P)H-hydrate repair enzyme"

Comparison with other databases. To assess the novelty of manual product annotations, we compared our annotation for each under-annotated gene with the corresponding entry on UniProt(116), Mtb Network Portal(9) (which included annotations from TBDB(5)), PATRIC(6), RefSeq(36), BioCyc(119), and KEGG(120). Comparisons were performed programmatically where possible, and systematically otherwise.

Enzyme commission number assignment. We assigned EC numbers to under-annotated genes with experimentally verified enzymatic activity using that assigned by the source article's author

when compliant with IUBMB standards. Otherwise, we manually assigned one using the official

IUBMB database(25).

HHpred filtering of structure-inferred functional annotations. All proteins with functions

assigned solely by structural inference were run through HHpred, searching against the PDB70

and ECOD databases, limiting maximum number of hits to 1000, and using default parameters

for the searches. All HHpred results were filtered, and only hits where Prob > 0.95 were retained

for downstream analysis. Each function assigned to a protein was evaluated separately (e.g. a

bifunctional protein could have one function culled and the second function retained).

Annotations were evaluated differently depending on whether they had a corresponding EC

number or now. All annotations with an EC number assigned were evaluated programmatically

and retained to the degree of EC specificity matched by HHpred hit(s). Annotations without

corresponding EC numbers were evaluated manually, independently, by two curators. Each

curator screened all HHpred hits and evaluated whether HHpred hits supported the assigned

function wholly, entirely, or not at all. In cases where function was partially supported, each

curator submitted a suggested product name change. After evaluating all proteins, the curators

reconciled any disparate assignments. Functions entirely uncorroborated by HHpred that passed

the Ramachandran plot analysis filtering step were subsequently evaluated to determine whether

the structural similarity used to infer function had substantial evidence warranting a transferred

annotation. Original structural inferences were either discarded, retained, or modified at the

discretion of the curators. For an annotation to be accepted, curators verified that model proteins

were not threaded on low-complexity proteins, checking whether regions underlying function of

the structurally solved protein structurally aligned to the protein model being annotated, and for

conservation of any known residues or structural motifs essential for function.

Ramachandran plot analysis. To evaluate structure model quality we computed the fraction of

the residues in "most favored" regions, "additionally allowed" regions, "generously allowed"

regions and "disallowed regions", via the PROCHECK server (121). The .pdb file containing the

atomic coordinates of each model protein structure of interest was uploaded to PROCHECK and

proportions of residues in each regional favorability classification were extracted from the

"results summary" file and collated into tabular format for further analysis. To determine a

threshold for including structures not corroborated by HHpred on the basis of quality protein

structure, we assessed the distributions of "most favored" region residues between proteins

wholly uncorroborated by HHpred and those fully corroborated by HHpred with maximally

specific EC numbers (Fig. 3).

Structural model visualization and annotation. Protein structure models (.pdb files) from I-

TASSER and solved protein structures form The Protein Data Bank were visualized and annotated

with pymol (https://pymol.org/).

Comparison with annotations from recent functional screens. Tables S2 and S4 from a

recently published Transposon mutant functional screen(83) were downloaded and the

intersection of their locus tags and the under-annotated gene set of this study went on for

further analysis. Genes with annotated functions by Bellerose and colleagues(83) were filtered

out. PE/PPE genes and Transcriptional regulatory proteins were also excluded, as the novelty

comparison was determined according to product name, which typically remains generic for

these two classes of proteins even upon updated functional information.

Bacterial strains and growth media. *M. tuberculosis strain* H37Rv was a gift from W.R. Jacobs,

Jr. of the Albert Einstein College of Medicine. Strains were grown in Middlebrook 7H9 medium

(Difco) supplemented with 10% (vol/vol) oleic acid-albumin-dextrose-catalase (OADC) (Difco),

0.2% (vol/vol) glycerol, and 0.05% (vol/vol) tyloxapol.

Characterization of POA resistant strains. Strain H37Rv was mutagenized with the *mariner-*

based transposon(122, 123). Approximately 10^5 independent transposon mutagenized bacilli

were plated on Middlebrook 7H10 medium supplemented with 10% (vol/vol) OADC (Difco), and

0.2% (vol/vol) glycerol with 50 µg/mL of pyrazinoic acid (POA) (Sigma). Resistant mutants were

selected from the plates containing 50 µg/mL POA. The initial isolates were plated on 7H10

medium supplemented with 10% (vol/vol) OADC (Difco), and 0.2% (vol/vol) glycerol either

containing 400 µg/mL POA or no drug after their initial isolation to confirm their POA resistance

prior to the more detailed drug susceptibility testing. (123, 124)Transposon insertion sites were

identified as previously described.

The antimicrobial drug susceptibility was determined by assessing the minimum concentration

of drug that was required to inhibit 90% of growth (MIC$_{90}$) relative to a no drug control. Growth

was assessed by measuring OD$_{600}$ of cultures after 14 days of incubation at 37ºC. Drug

susceptibility testing for PZA and POA was carried out in 7H9 broth supplemented with OADC,

glycerol, and tyloxapol (pH 5.8) as indicated above. INH MIC$_{90}$ determinations were carried out

in medium with the same composition at pH 6.8.

ACKNOWLEDGEMENTS We extend our gratitude to Dr. Vikram Alva for running a large set of

proteins through HHpred in batch mode on our behalf. We thank Sarah Radecke, Derek Conkle-

Gutierrez, and Matthew Onorato for their extensive review of the manuscript and helpful

comments. We would also like to acknowledge Dr. Yusuke Minato for his contributions to PZA

experiments. This work was funded by grants from National Institute of Allergy and Infectious

Diseases (NIAID Grant No. R01AI105185 to FV and R01AI123146 to ADB). SJM, AMZ, DG, AE, NK,

CR, ND, CKC and FV were supported by R01AI105185. SJM was also supported by scholarships

from a National Science Foundation DUE Training Grant to FV (0966391). ADB and NAD were

funded by R01AI123146. NAD was also supported by NHLBI (HL007741). The funding bodies had

no role in the design of the study or in collection, analysis, and interpretation of data or in writing

the manuscript.

CITATIONS
1. Poux S, Arighi CN, Magrane M, Bateman A, Wei C-H, Lu Z, Boutet E, Bye-A-Jee H, Famiglietti ML, Roechert B. On expert curation and scalability: UniProtKB/ Swiss-Prot as a case study.
2. Steenken W, Oatway WH, Petroff SA. 1934. BIOLOGICAL STUDIES OF THE TUBERCLE BACILLUS. J Exp Med 60.
3. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon S V., Eiglmeier K, Gas S, Barry CE, Tekaia F, Badcock K, Basham D, Brown D, Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N, Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L, Oliver K, Osborne J, Quail MA, Rajandream M-A, Rogers J, Rutter S, Seeger K, Skelton J, Squares R, Squares S, Sulston JE, Taylor K, Whitehead S, Barrell BG. 1998. Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. Nature 393:537–544.
4. Camus J-C, Pryor MJ, Médigue C, Cole ST. 2002. Re-annotation of the genome sequence of Mycobacterium tuberculosis H37Rv. Microbiology 148:2967–2973.
5. Reddy TBK, Riley R, Wymore F, Montgomery P, DeCaprio D, Engels R, Gellesch M, Hubble J, Jen D, Jin H, Koehrsen M, Larson L, Mao M, Nitzberg M, Sisk P, Stolte C, Weiner B, White J, Zachariah ZK, Sherlock G, Galagan JE, Ball CA, Schoolnik GK. 2009. TB database: an integrated platform for tuberculosis research. Nucleic Acids Res 37:D499–D508.
6. Gillespie JJ, Wattam AR, Cammer SA, Gabbard JL, Shukla MP, Dalay O, Driscoll T, Hix D, Mane SP, Mao C, Nordberg EK, Scott M, Schulman JR, Snyder EE, Sullivan DE, Wang C, Warren A, Williams KP, Xue T, Seung Yoo H, Zhang C, Zhang Y, Will R, Kenyon RW, Sobral BW. 2011. PATRIC: the Comprehensive Bacterial Bioinformatics Resource with a Focus on Human Pathogenic Species ▽. Infect Immun 79:4286–4298.
7. Garcia BJ, Datta G, Davidson RM, Strong M. 2015. MycoBASE: expanding the functional annotation coverage of mycobacterial genomes. BMC Genomics 16:1102.
8. Schito M, Dolinger DL. 2015. A Collaborative Approach for "ReSeq-ing" Mycobacterium tuberculosis Drug Resistance: Convergence for Drug and Diagnostic Developers. EBioMedicine 2:1262–1265.
9. Ma S, Minch KJ, Rustad TR, Hobbs S, Zhou S-L, Sherman DR, Price ND. 2015. Integrated Modeling of Gene Regulatory and Metabolic Networks in Mycobacterium tuberculosis. PLoS Comput Biol 11:e1004543.
10. Faksri K, Tan JH, Chaiprasert A, Teo Y-Y, Ong RT-H. 2016. Bioinformatics tools and databases for whole genome sequence analysis of Mycobacterium tuberculosis. Infect Genet Evol 45:359–368.
11. Silva D, Almeida PE, Von Groll A, Martin A, Palomino JC. 2011. Efflux as a mechanism for drug resistance in Mycobacterium tuberculosis. FEMS Immunol Med Microbiol 63:1–9.
12. Theobald DL, Miller C. 2010. Membrane transport proteins: surprises in structural sameness. Nat Struct Mol Biol 17:2–3.

13.   Drayman N, Glick Y, Ben-nun-Shaul O, Zer H, Zlotnick A, Gerber D, Schueler-Furman O, Oppenheim A. 2013. Pathogens Use
Structural Mimicry of Native Host Ligands as a Mechanism for Host Receptor Engagement. Cell Host Microbe 14:63–73.
14.   Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. 2015. The I-TASSER Suite: protein structure and function prediction. Nat
Methods 12:7–8.
15.   Rose PW, Prlić A, Altunkaya A, Bi C, Bradley AR, Christie CH, Costanzo L Di, Duarte JM, Dutta S, Feng Z, Green RK, Goodsell
DS, Hudson B, Kalro T, Lowe R, Peisach E, Randle C, Rose AS, Shao C, Tao Y-P, Valasatava Y, Voigt M, Westbrook JD, Woo
J, Yang H, Young JY, Zardecki C, Berman HM, Burley SK. 2017. The RCSB protein data bank: integrative view of protein,
gene and 3D structural information. Nucleic Acids Res 45:D271–D281.
16.   Moult J, Fidelis K, Kryshtafovych A, Rost B, Hubbard T, Tramontano A. 2007. Critical assessment of methods of protein
structure prediction—Round VII. Proteins Struct Funct Bioinforma 69:3–9.
17.   Kryshtafovych A, Krysko O, Daniluk P, Dmytriv Z, Fidelis K. 2009. Protein structure prediction center in CASP8. Proteins
Struct Funct Bioinforma 77:5–9.
18.   Moult J, Fidelis K, Kryshtafovych A, Tramontano A. 2011. Critical assessment of methods of protein structure prediction
(CASP)—round IX. Proteins Struct Funct Bioinforma 79:1–5.
19.   Huang YJ, Mao B, Aramini JM, Montelione GT. 2014. Assessment of template-based protein structure predictions in CASP10.
Proteins Struct Funct Bioinforma 82:43–56.
20.   Moult J, Fidelis K, Kryshtafovych A, Schwede T, Tramontano A. 2016. Critical assessment of methods of protein structure
prediction: Progress and new directions in round XI. Proteins Struct Funct Bioinforma 84:4–14.
21.   Xu J, Zhang Y. 2010. How significant is a protein structure similarity with TM-score = 0.5? Bioinformatics 26:889–895.
22.   Zhang Y, Skolnick J. 2005. TM-align: a protein structure alignment algorithm based on the TM-score. Nucleic Acids Res
33:2302–9.
23.   Zhang C, Freddolino PL, Zhang Y. 2017. COFACTOR: improved protein function prediction by combining structure, sequence
and protein–protein interaction information. Nucleic Acids Res 45:W291–W299.
24.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill
DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene Ontology:
tool for the unification of biology. Nat Genet 25:25–29.
25.   Webb EC, others. 1992. Enzyme nomenclature 1992. Recommendations of the Nomenclature Committee of the International
Union of Biochemistry and Molecular Biology on the Nomenclature and Classification of Enzymes. Academic Press.
26.   Yang J, Roy A, Zhang Y. 2013. BioLiP: a semi-manually curated database for biologically relevant ligand-protein interactions.
Nucleic Acids Res 41:D1096-103.
27.   Hill DP, Davis AP, Richardson JE, Corradi JP, Ringwald M, Eppig JT, Blake JA. 2001. PROGRAM DESCRIPTION: Strategies for
Biological Annotation of Mammalian Systems: Implementing Gene Ontologies in Mouse Genome Informatics. Genomics
74:121–128.
28.   The Gene Ontology Consortium, Gene Ontology Consortium, The Gene Ontology Consortium. 2017. Expansion of the Gene
Ontology knowledgebase and resources. Nucleic Acids Res 45:D331–D338.
29.   Ramakrishnan G, Ochoa-Montaño B, Raghavender US, Mudgal R, Joshi AG, Chandra NR, Sowdhamini R, Blundell TL, Srinivasan
N. 2015. Enriching the annotation of Mycobacterium tuberculosis H37Rv proteome using remote homology detection
approaches: Insights into structure and function. Tuberculosis 95:14–25.
30.   Functional assignment of Mycobacterium tuberculosis proteome revealed by genome-scale fold-recognition.
31.   Al-Khafaji ZM. 2013. In Silico Investigation of Rv Hypothetical Proteins of Virulent Strain Mycobacterium tuberculosis H37Rv.
ndian J Pharm Biol Res 1:81–88.
32.   Doerks T, Noort V van, Minguez P, Bork P. 2012. Annotation of the M. tuberculosis Hypothetical Orfeome: Adding Functional
Information to More than Half of the Uncharacterized Proteins. PLoS One 7:e34302.
33.   Mazandu GK, Mulder NJ. 2012. Using the underlying biological organization of the Mycobacterium tuberculosis functional
network for protein function prediction. Infect Genet Evol 12:922–932.
34.   Sangrador-Vegas A, Mitchell AL, Chang H-Y, Yong S-Y, Finn RD. 2016. GO annotation in InterPro: why stability does not
indicate accuracy in a sea of changing annotations. Database 2016:baw027.
35.   Liberal R, Pinney JW. 2013. Simple topological properties predict functional misannotations in a metabolic network.
Bioinformatics 29.
36.   Tatusova T, Ciufo S, Fedorov B, O'Neill K, Tolstoy I. 2014. RefSeq microbial genomes database: new representation and
annotation strategy. Nucleic Acids Res 42:D553–D559.
37.   Sultana R, Vemula MH, Banerjee S, Guruprasad L. 2013. The PE16 (Rv1430) of Mycobacterium tuberculosis Is an Esterase
Belonging to Serine Hydrolase Superfamily of Proteins. PLoS One https://doi.org/10.1371/journal.pone.0055320.
38.   Padhi A, Naik SK, Sengupta S, Ganguli G, Sonawane A. 2016. Expression of Mycobacterium tuberculosis NLPC/p60 family
protein Rv0024 induce biofilm formation and resistance against cell wall acting anti-tuberculosis drugs in Mycobacterium
smegmatis. Microbes Infect 18:224–236.
39.   Kateete DP, Katabazi FA, Okeng A, Okee M, Musinguzi C, Asiimwe BB, Kyobe S, Asiimwe J, Boom WH, Joloba ML. 2012.
Rhomboids of Mycobacteria: Characterization Using an aarA Mutant of Providencia stuartii and Gene Deletion in
Mycobacterium smegmatis. PLoS One https://doi.org/10.1371/journal.pone.0045741.
40.   Nambi S, Long JE, Mishra BB, Baker R, Murphy KC, Olive AJ, Nguyen HP, Shaffer SA, Sassetti CM. 2015. The Oxidative Stress
Network of Mycobacterium tuberculosis Reveals Coordination between Radical Detoxification Systems. Cell Host Microbe
https://doi.org/10.1016/j.chom.2015.05.008.
41.   Flentie K, Garner AL, Stallings CL. 2016. Mycobacterium tuberculosis transcription machinery: Ready to respond to host
attacks. J Bacteriol.
42.   Primm TP, Andersen SJ, Mizrahi V, Avarbock D, Rubin H, Barry Iii CE. 2000. The Stringent Response of Mycobacterium
tuberculosis Is Required for Long-Term Survival. J Bacteriol 182:4889–4898.

43. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology detection and structure prediction.
Nucleic Acids Res 33:W244.
44. Bhattacharya A, Tejero R, Montelione GT. 2007. Evaluating protein structures determined by structural genomics consortia.
Proteins Struct Funct Genet 66:778–795.
45. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. 1993. PROCHECK: a program to check the stereochemical quality of
protein structures. J Appl Crystallogr 26:283–291.
46. He YX, Huang L, Xue Y, Fei X, Teng Y Bin, Rubin-Pitel SB, Zhao H, Zhou CZ. 2010. Crystal structure and computational
analyses provide insights into the catalytic mechanism of 2,4-diacetylphloroglucinol hydrolase PhlG from Pseudomonas
fluorescens. J Biol Chem 285:4603–4611.
47. Han J, Zhang S, Jia W, Zhang Z, Wang Y, He Y. 2019. Discovery and structural analysis of a phloretin hydrolase from the
opportunistic human pathogen *Mycobacterium abscessus*. FEBS J 286:1959–1971.
48. Brzostek A, Pawelczyk J, Rumijowska-Galewicz A, Dziadek B, Dziadek J. 2009. Mycobacterium tuberculosis is able to
accumulate and utilize cholesterol. J Bacteriol 191:6584–91.
49. Lack NA, Yam KC, Lowe EE, Horsman GP, Owen RL, Sim E, Eltis LD. 2010. Characterization of a carbon-carbon hydrolase
from Mycobacterium tuberculosis involved in cholesterol metabolism. J Biol Chem 285:434–443.
50. Wilburn KM, Fieweger RA, VanderVen BC. 2018. Cholesterol and fatty acids grease the wheels of Mycobacterium
tuberculosis pathogenesis. Pathog Dis. Oxford Academic.
51. Wang R, Yin YJ, Wang F, Li M, Feng J, Zhang HM, Zhang JP, Liu SJ, Chang WR. 2007. Crystal structures and site-directed
mutagenesis of a mycothiol-dependent enzyme reveal a novel folding and molecular basis for mycothiol-mediated
maleylpyruvate isomerization. J Biol Chem 282:16288–16294.
52. Newton GL, Leung SS, Wakabayashi JI, Rawat M, Fahey RC. 2011. The DinB Superfamily Includes Novel Mycothiol, Bacillithiol,
and Glutathione S-Transferases. Biochemistry 50:10751–10760.
53. Liu TT, Zhou NY. 2012. Novel L-cysteine-dependent maleylpyruvate isomerase in the gentisate pathway of Paenibacillus sp.
strain NyZ101. J Bacteriol 194:3987–3994.
54. Slama N, Jamet S, Frigui W, Pawlik A, Bottai D, Laval F, Constant P, Lemassu A, Cam K, Daffé M, Brosch R, Eynard N, Quémard
A. 2015. The changes in mycolic acid structures caused by hadC mutation have a dramatic effect on the virulence of
Mycobacterium tuberculosis. Mol Microbiol 99:n/a-n/a.
55. Singh A, Crossman DK, Mai D, Guidry L, Voskuil MI, Renfrow MB, Steyn AJC. 2009. Mycobacterium tuberculosis WhiB3
Maintains Redox Homeostasis by Regulating Virulence Lipid Anabolism to Modulate Macrophage Response. PLoS Pathog
5:e1000545.
56. Olivella M, Gonzalez A, Pardo L, Deupi X. 2013. Relation between sequence and structure in membrane proteins.
Bioinformatics 29:1589–1592.
57. Shi W, Chen J, Zhang S, Zhang W, Zhang Y. 2018. Identification of novel mutations in LprG (rv1411c), rv0521, rv3630,
rv0010c, ppsC, and cyp128 associated with pyrazinoic acid/pyrazinamide resistance in mycobacterium tuberculosis.
Antimicrob Agents Chemother 62.
58. Costa P, Amaro A, Ferreira AS, Machado D, Albuquerque T, Couto I, Botelho A, Viveiros M, Inácio J. 2014. Rapid identification
of veterinary-relevant Mycobacterium tuberculosis complex species using 16S rDNA, IS6110 and Regions of Difference-
targeted dual-labelled hydrolysis probes. J Microbiol Methods 107:13–22.
59. Strong EJ, Jurcic Smith KL, Saini NK, Ng TW, Porcelli SA, Lee S. 2020. Identification of autophagy-inhibiting factors of
mycobacterium tuberculosis by high-throughput loss-of-function screening. Infect Immun 88.
60. Kania E, Pająk B, O'Prey J, Sierra Gonzalez P, Litwiniuk A, Urbańska K, Ryan KM, Orzechowski A. 2017. Verapamil treatment
induces cytoprotective autophagy by modulating cellular metabolism. FEBS J 284:1370–1387.
61. Abramovitch RB, Rohde KH, Hsu F-F, Russell DG. 2011. {aprABC}: {A} {Mycobacterium} tuberculosis complex-specific locus
that modulates {pH}-driven adaptation to the macrophage phagosome. Mol Microbiol 80:678–694.
62. Agranoff D, Krishna S. 2004. Metal ion transport and regulation in {Mycobacterium} tuberculosis. Front Biosci A J Virtual
Libr 9:2996–3006.
63. Rodrigues L, Cravo P, Viveiros M. 2020. Efflux pump inhibitors as a promising adjunct therapy against drug resistant
tuberculosis: a new strategy to revisit mycobacterial targets and repurpose old drugs. Expert Rev Anti Infect Ther. Taylor
and Francis Ltd.
64. Lin Y, Dong Y, Gao Y, Shi R, Li Y, Zhou X, Liu W, Li G, Qi Y, Wu Y. 2021. Identification of CTL Epitopes on Efflux Pumps of
the ATP-Binding Cassette and the Major Facilitator Superfamily of Mycobacterium tuberculosis. J Immunol Res 2021:1–13.
65. Moult J, Melamud E. 2000. From fold to function. Curr Opin Struct Biol 10:384–389.
66. Evans JC, Mizrahi V. 2015. The application of tetracyclineregulated gene expression systems in the validation of novel drug
targets in Mycobacterium tuberculosis. Front Microbiol 6.
67. Judd JA, Canestrari J, Clark R, Joseph A, Lapierre P, Lasek-Nesselquist E, Mir M, Palumbo M, Smith C, Stone M, Upadhyay A,
Wirth SE, Dedrick RM, Meier CG, Russell DA, Dills A, Dove E, Kester J, Wolf ID, Zhu J, Rubin ER, Fortune S, Hatfull GF, Gray
TA, Wade JT, Derbyshire KM. 2021. A mycobacterial systems resource for the research community. MBio 12:1–15.
68. Lato DF, Golding GB. 2020. Spatial Patterns of Gene Expression in Bacterial Genomes. J Mol Evol 88:510–520.
69. Eisen JA, Heidelberg JF, White O, Salzberg SL. 2000. Evidence for symmetric chromosomal inversions around the replication
origin in bacteria. Genome Biol 1:research0011.1.
70. Repar J, Warnecke T. 2017. Non-Random Inversion Landscapes in Prokaryotic Genomes Are Shaped by Heterogeneous
Selection Pressures. Mol Biol Evol 34:1902–1911.
71. Galperin MY, Kristensen DM, Makarova KS, Wolf YI, Koonin E V. 2019. Microbial genome analysis: the COG approach. Brief
Bioinform 20:1063–1070.
72. Merrikh CN, Merrikh H. 2018. Gene inversion potentiates bacterial evolvability and virulence. Nat Commun 9:4662.
73. Onwueme KC, Vos CJ, Zurita J, Ferreras JA, Quadri LEN. 2005. The dimycocerosate ester polyketide virulence factors of

1113    mycobacteria. Prog Lipid Res 44:259–302.

1114 74. Lee JS, Krause R, Schreiber J, Mollenkopf H-J, Kowall J, Stein R, Jeon B-Y, Kwak J-Y, Song M-K, Patron JP, Jorg S, Roh K, Cho
1115    S-N, Kaufmann SHE. 2008. Mutation in the Transcriptional Regulator PhoP Contributes to Avirulence of Mycobacterium
1116    tuberculosis H37Ra Strain. Cell Host Microbe 3:97–103.

1117 75. Elghraoui A, Modlin SJ, Valafar F. 2017. SMRT genome assembly corrects reference errors, resolving the genetic basis of
1118    virulence in Mycobacterium tuberculosis. BMC Genomics 18:302.

1119 76. Passemar C, Arbués A, Malaga W, Mercier I, Moreau F, Lepourry L, Neyrolles O, Guilhot C, Astarie-Dequeker C. 2014. Multiple
1120    deletions in the polyketide synthase gene repertoire of Mycobacterium tuberculosis reveal functional overlap of cell envelope
1121    lipids in host–pathogen interactions. Cell Microbiol 16:195–213.

1122 77. Minnikin DE, Dobson G, Hutchinson IG. 1983. Characterization of phthiocerol dimycocerosates from Mycobacterium
1123    tuberculosis. Biochim Biophys Acta - Lipids Lipid Metab 753:445–449.

1124 78. Mann FM, Xu M, Davenport EK, Peters RJ. 2012. Functional characterization and evolution of the isotuberculosinol operon
1125    in Mycobacterium tuberculosis and related Mycobacteria. Front Microbiol 3.

1126 79. Mann FM, Prisic S, Hu H, Xu M, Coates RM, Peters RJ. 2009. Characterization and Inhibition of a Class II Diterpene Cyclase
1127    from Mycobacterium tuberculosis. J Biol Chem 284:23574–23579.

1128 80. Mann FM, Peters RJ. 2012. Isotuberculosinol: the unusual case of an immunomodulatory diterpenoid from Mycobacterium
1129    tuberculosis. Medchemcomm 3:899–904.

1130 81. Sieniawska E, Swatko-Ossor M, Sawicki R, Skalicka-Woźniak K, Ginalska G. 2017. Natural Terpenes Influence the Activity of
1131    Antibiotics against Isolated Mycobacterium tuberculosis. Med Princ Pract Int J Kuwait Univ Heal Sci Cent 26:108–112.

1132 82. Layre E, Lee HJ, Young DC, Martinot AJ, Buter J, Minnaard AJ, Annand JW, Fortune SM, Snider BB, Matsunaga I, Rubin EJ,
1133    Alber T, Moody DB. 2014. Molecular profiling of Mycobacterium tuberculosis identifies tuberculosinyl nucleoside products
1134    of the virulence-associated enzyme Rv3378c. Proc Natl Acad Sci 111:2978–2983.

1135 83. Bellerose MM, Proulx MK, Smith CM, Baker RE, Ioerger TR, Sassetti CM. 2020. Distinct Bacterial Pathways Influence the
1136    Efficacy of Antibiotics against Mycobacterium tuberculosis. mSystems 5.

1137 84. Jansen RS, Mandyoli L, Hughes R, Wakabayashi S, Pinkham JT, Selbach B, Guinn KM, Rubin EJ, Sacchettini JC, Rhee KY. 2020.
1138    Aspartate aminotransferase Rv3722c governs aspartate-dependent nitrogen metabolism in Mycobacterium tuberculosis.
1139    Nat Commun 11:1–13.

1140 85. Zhang L, Hendrickson RC, Meikle V, Lefkowitz EJ, Ioerger TR, Niederweis M. 2020. Comprehensive analysis of iron utilization
1141    by Mycobacterium tuberculosis. PLoS Pathog 16:e1008337.

1142 86. Scorpio A, Zhang Y. 1996. Mutations in pncA, a gene encoding pyrazinamidase/nicotinamidase, cause resistance to the
1143    antituberculous drug pyrazinamide in tubercle bacillus. Nat Med 2:662–667.

1144 87. Zhang Y, Wade MM, Scorpio A, Zhang H, Sun Z. 2003. Mode of action of pyrazinamide: Disruption of Mycobacterium
1145    tuberculosis membrane transport and energetics by pyrazinoic acid. J Antimicrob Chemother 52:790–795.

1146 88. Zimhony O, Cox JS, Welch JT, Vilchèze C, Jacobs WR. 2000. Pyrazinamide inhibits the eukaryotic-like fatty acid synthetase I
1147    (FASI) of Mycobacterium tuberculosis. Nat Med 6:1043–1047.

1148 89. Shi W, Zhang X, Jiang X, Yuan H, Lee JS, Barry CE, Wang H, Zhang W, Zhang Y. 2011. Pyrazinamide inhibits trans-translation
1149    in Mycobacterium tuberculosis. Science (80-) 333:1630–1632.

1150 90. Boshoff HI, Mizrahi V, Iii CEB, Barry CE. 2002. Effects of Pyrazinamide on Fatty Acid Synthesis by Whole Mycobacterial Cells
1151    and Purified Fatty Acid Synthase I. J Bacteriol 184:2167–2172.

1152 91. Peterson ND, Rosen BC, Dillon NA, Baughn AD. 2015. Uncoupling environmental pH and intrabacterial acidification from
1153    pyrazinamide susceptibility in Mycobacterium tuberculosis. Antimicrob Agents Chemother 59:7320–7326.

1154 92. Dillon NA, Peterson ND, Feaga HA, Keiler KC, Baughn AD. 2017. Anti-tubercular Activity of Pyrazinamide is Independent of
1155    trans-Translation and RpsA. Sci Rep 7:6135.

1156 93. Gopal P, Yee M, Sarathy JPJ, Liang Low J, Sarathy JPJ, Kaya F, Dartois VV, Gengenbacher M, Dick T, Low JL, Sarathy JPJ, Kaya
1157    F, Dartois VV, Gengenbacher M, Dick T. 2016. Pyrazinamide resistance is caused by two distinct mechanisms: Prevention of
1158    coenzyme a depletion and loss of virulence factor synthesis. ACS Infect Dis 2:616–626.

1159 94. Anthony RM, den Hertog AL, van Soolingen D. 2018. 'Happy the man, who, studying nature's laws, Thro'' known effects can
1160    trace the secret cause.'† Do we have enough pieces to solve the pyrazinamide puzzle?' J Antimicrob Chemother 73:1750–
1161    1754.

1162 95. Dillon NA, Peterson ND, Rosen BCBC, Baughn AD. 2014. Pantothenate and pantetheine antagonize the antitubercular
1163    activity of pyrazinamide. Antimicrob Agents Chemother 58:7258–7263.

1164 96. Zhang Y, Mitchison D. 2003. The curious characteristics of pyrazinamide: A review. Int J Tuberc Lung Dis 7:6–21.

1165 97. Swan MK, Hansen T, Schönheit P, Davies C. 2004. Structural basis for phosphomannose isomerase activity in
1166    phosphoglucose isomerase from Pyrobaculum aerophilum: A subtle difference between distantly related enzymes.
1167    Biochemistry 43:14088–14095.

1168 98. Mouilleron S, Badet-Denisot MA, Golinelli-Pimpaneau B. 2006. Glutamine binding opens the ammonia channel and activates
1169    glucosamine-6P synthase. J Biol Chem 281:4404–4412.

1170 99. Bateman A. 1999. The SIS domain: A phosphosugar-binding domain. Trends Biochem Sci 24:94–95.

1171 100. Keiser TL, Schlesinger LS, King S, Munson R, Seveau S. 2014. Biosynthesis of mannose-containing cell wall components
1172    important in Mycobacterium.

1173 101. Schnoes AM, Brown SD, Dodevski I, Babbitt PC. 2009. Annotation Error in Public Databases: Misannotation of Molecular
1174    Function in Enzyme Superfamilies. PLoS Comput Biol 5.

1175 102. Ioerger TR, Feng Y, Ganesula K, Chen X, Dobos KM, Fortune S, Jacobs WR, Mizrahi V, Parish T, Rubin E, Sassetti C, Sacchettini
1176    JC. 2010. Variation among Genome Sequences of H37Rv Strains of Mycobacterium tuberculosis from Multiple Laboratories.
1177    J Bacteriol 192:3645–3653.

1178 103. Shell SS, Wang J, Lapierre P, Mir M, Chase MR, Pyle MM, Gawande R, Ahmad R, Sarracino DA, Ioerger TR, Fortune SM,

Derbyshire KM, Wade JT, Gray TA. 2015. Leaderless Transcripts and Small Proteins Are Common Features of the
Mycobacterial Translational Landscape. PLoS Genet 11.

104.	Nagano N, Orengo CA, Thornton JM. 2002. One fold with many functions: the evolutionary relationships between TIM barrel
families based on their sequences, structures and functions. J Mol Biol 321:741–765.

105.	Murphy GS, Greisman JB, Hecht MH. 2016. De Novo Proteins with Life-Sustaining Functions Are Structurally Dynamic. J Mol
Biol 428:399–411.

106.	Miskei M, Gregus A, Sharma R, Duro N, Zsolyomi F, Fuxreiter M. 2017. Fuzziness enables context dependence of protein
interactions. FEBS Lett 591:2682–2695.

107.	Morcos F, Pagnani A, Lunt B, Bertolino A, Marks DS, Sander C, Zecchina R, Onuchic JN, Hwa T, Weigt M. 2011. Direct-
coupling analysis of residue coevolution captures native contacts across many protein families. Proc Natl Acad Sci U S A
108:E1293–E1301.

108.	Lockless SW, Ranganathan R. 1999. Evolutionarily conserved pathways of energetic connectivity in protein families. Science
(80-) 286:295–299.

109.	Neuwald AF. 2014. A Bayesian Sampler for Optimization of Protein Domain Hierarchies. J Comput Biol 21:269–286.

110.	Neuwald AF, Aravind L, Altschul SF. 2018. Inferring joint sequence-structural determinants of protein functional specificity.
Elife 7.

111.	PDB RCSB. PDB Data Distribution by Resolution.

112.	Jensen MR, Zweckstetter M, Huang J, Blackledge M. 2014. Exploring Free-Energy Landscapes of Intrinsically Disordered
Proteins at Atomic Resolution Using NMR Spectroscopy. Chem Rev 114:6632–6660.

113.	Fishbein S, van Wyk N, Warren RM, Sampson SL. 2015. Phylogeny to function: PE/PPE protein evolution and impact on
Mycobacterium tuberculosis pathogenicity. Mol Microbiol 96:901–916.

114.	McEvoy CREE, Cloete R, Müller B, Schürch AC, Helden PD van, Gagneux S, Warren RM, Pittius NCG van, van Helden PD,
Gagneux S, Warren RM, Gey van Pittius NC. 2012. Comparative Analysis of Mycobacterium tuberculosis pe and ppe Genes
Reveals High Sequence Variation and an Apparent Absence of Selective Constraints. PLoS One 7:e30593.

115.	Zhang Y, Skolnick J. 2004. Scoring function for automated assessment of protein structure template quality. Proteins Struct
Funct Bioinforma 57:702–710.

116.	UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic Acids Res 45:D158–D169.

117.	Roy A, Yang J, Zhang Y. 2012. COFACTOR: an accurate comparative algorithm for structure-based protein function
annotation. Nucleic Acids Res 40:W471–W477.

118.	R Development Core Team. 2018. A Language and Environment for Statistical Computing. R Found Stat Comput. Vienna,
Austria.

119.	Karp PD, Billington R, Caspi R, Fulcher CA, Latendresse M, Kothari A, Keseler IM, Krummenacker M, Midford PE, Ong Q, Ong
WK, Paley SM, Subhraveti P. 2017. The BioCyc collection of microbial genomes and metabolic pathways. Brief Bioinform
28:1–6.

120.	Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and
drugs. Nucleic Acids Res 45.

121.	Laskowski RA, MacArthur MW, Thornton JM. 2012. *PROCHECK*: validation of protein-structure coordinates, p. 684–687.
In . American Cancer Society.

122.	Rubin EJ, Akerley BJ, Novik VN, Lampe DJ, Husson RN, Mekalanos JJ. 1999. In vivo transposition of mariner-based elements
in enteric bacteria and mycobacteria. Proc Natl Acad Sci 96:1645–1650.

123.	Lee S, Kriakov J, Vilcheze C, Dai Z, Hatfull GF, Jacobs WR, J.F. C, J. M, D. M, N.R. P, W. B, V. K, J. K, L. K, S. B, J. K, J.G. L, W.R.
J, R.W. H, G.F. H. 2004. Bxz1, a new generalized transducing phage for mycobacteria. FEMS Microbiol Lett 241:271–276.

124.	Baughn AD, Deng J, Vilcheze C, Riestra A, Welch JT, Jacobs WR, Zimhony O, Vilchèze C, Riestra A, Welch JT, Jacobs WR,
Zimhony O. 2010. Mutually exclusive genotypes for pyrazinamide and 5-chloropyrazinamide resistance reveal a potential
resistance-proofing strategy. Antimicrob Agents Chemother 54:5323–5328.

125.	Lew JM, Kapopoulou A, Jones LM, Cole ST. 2011. TubercuList--10 years after. Tuberculosis (Edinb) 91:1–7.

126.	Chen J, Guo M, Wang X, Liu B. A comprehensive review and comparison of different computational methods for protein
remote homology detection. Brief Bioinform https://doi.org/10.1093/bib/bbw108.

127.	Cortes T, Schubert OTT, Rose G, Arnvig KBB, Comas I, Aebersold R, Young DBB. 2013. Genome-wide Mapping of
Transcriptional Start Sites Defines an Extensive Leaderless Transcriptome in Mycobacterium tuberculosis. Cell Rep 5:1121–
1131.

128.	Shell SS, Wang J, Lapierre P, Mir M, Chase MR, Pyle MM, Gawande R, Ahmad R, Sarracino DA, Ioerger TR, Fortune SM,
Derbyshire KM, Wade JT, Gray TA. 2015. Leaderless Transcripts and Small Proteins Are Common Features of the
Mycobacterial Translational Landscape. PLoS Genet 11.

129.	Teplyakov A, Obmolova G, Badet-Denisot MA, Badet B, Polikarpov I. 1998. Involvement of the C terminus in intramolecular
nitrogen channeling in glucosamine 6-phosphate synthase: Evidence from a 1.6 Å crystal structure of the isomerase domain.
Structure 6:1047–1055.

130.	Salah Ud-Din A, Tikhomirova A, Roujeinikova A. 2016. Structure and Functional Diversity of GCN5-Related N-
Acetyltransferases (GNAT). Int J Mol Sci 17:1018.

131.	Wybenga-Groot LE, Draker K ann, Wright GD, Berghuis AM. 1999. Crystal structure of an aminoglycoside 6'-N-
acetyltransferase: Defining the GCN5-related N-acetyltransferase superfamily fold. Structure 7:497–507.

132.	Reddy TBK, Riley R, Wymore F, Montgomery P, DeCaprio D, Engels R, Gellesch M, Hubble J, Jen D, Jin H, Koehrsen M, Larson
L, Mao M, Nitzberg M, Sisk P, Stolte C, Weiner B, White J, Zachariah ZK, Sherlock G, Galagan JE, Ball CA, Schoolnik GK.
2009. TB database: an integrated platform for tuberculosis research. Nucleic Acids Res 37:D499–D508.

1245

Figures, Tables, and Supplemental material descriptions

Tables

Table 1. Comparison among frequented annotation resources. "Functional assignments" refer to annotations that describe protein function, and excluded hypothetical, unknown/uncharacterized, and PE/PPE family proteins. Counts reflect database content on May 17, 2017 for RefSeq(36) https://www.ncbi.nlm.nih.gov/refseq/, PATRIC(6) https://www.patricbrc.org/,and Mtb Network Portal(9) http://networks.systemsbiology.net/mtb/ and June 23, 2017 for KEGG(120) https://www.kegg.jp/kegg/genome/pathogen.html and UniProt(116) https://www.uniprot.org/uniprot/ . The number of CDSs in KEGG is reported as 3906 because they include only protein coding genes. The source of annotations for *M. tuberculosis* protein-coding genes in KEGG is TubercuList(125).

Metric	TubercuList	PATRIC	RefSeq	Mtb Network Portal	UniProt	KEGG	BioCyc
Coding sequences	4038	4367	3989	4038	3997	3906	4031
Proteins with functional assignments	2815	3007	2341	2853	2906	1750	2571
Hypothetical proteins	1223	1360	1648	1185	1091	2156	1460
Proteins with ≥ 1 GO term	2629	969	0	2460	3305	0	3557
Proteins with EC number(s) assigned	1293	1074	1081	1003	1138	1050	1018

1256
1257

1258 **Table 2. Genes annotated through systematic manual curation that expand upon annotations from major**
1259 **databases**. Annotations are separated into completely novel, those with similar annotations but with greater
1260 specificity, and those with an additional, orthogonal annotation compared to evaluated databases (*Text S1*, Fig. S4A).
1261 Pubmed IDs (PMID) from which annotations for each product were derived are included. Members of the PE/PPE
1262 family are indicated by asterisk.

Gene	product	PMID	TubercuList	UniProt	Mtb Net Portal
			Novel		
Rv0309	Adhesin/Putative L,D-transpeptidase	23922800; 23889607; 26201501	Possible conserved exported protein	Possible conserved exported protein	
Rv0394c	Hyaluronidase/ Chondrosulfatase	23465892	Possible secreted protein	Possible secreted protein	Possible membrane protein
Rv0431	Probable vesiculogenesis/immune response regulator	24248369; 21170273; 17436267; 26324094; 27765619	Putative tuberculin related peptide	Putative tuberculin related peptide	Tuberculin related peptide
Rv1430*	Esterase	23383323	PE family protein PE16	PE family protein PE16	PE family protein
Rv1993c	Putative chaperone	21925112	Conserved protein	Uncharacterized protein	
Rv2345	Probable phosphatase	25782739	Possible conserved transmembrane protein	UPF0603 protein	Possible membrane protein
Rv2923c	Probable Osmotically induced bacterial protein C (OsmC, a probable peroxide reductase)	22088319	Conserved protein	Uncharacterized protein	
Rv2954c	Probable Methyltransferase	23536839	Hypothetical protein	Uncharacterized protein	
Rv2969c	Periplasmic disulfide-bond forming (Dsb) Enzyme	24100317; 18539140	Possible conserved membrane or secreted protein	Membrane protein (Possible conserved membrane or secreted protein)	POSSIBLE CONSERVED MEMBRANE OR SECRETED PROTEIN
Rv3528c	Probable serine hydrolase	26853625	Unknown protein	Uncharacterized protein	
			Greater specificity		
Rv0059	Probable toxin DarT/Probable DNA ADP-ribosyl transferase	27939941	Hypothetical protein	Uncharacterized protein	
Rv0060	Probable antitoxin DarG/Probable DNA ADP-ribosyl glycohydrolase	27939941	Conserved hypothetical protein	Uncharacterized protein	ADP-ribose 1-phosphate phophatase related protein
Rv1337	Probable rhomboid protease/integral membrane protein	19165721; 23029216	Probable integral membrane protein	Uncharacterized protein	Rhomboid membrane family protein
Rv1357c	Cyclic diguanylate phosphodiesterase	21151497	Conserved hypothetical protein	Uncharacterized protein	Sensory box/GGDEF family protein
Rv1566c	Probable non-catalytic peptidoglycan binding RipD protein/probable antigen	24107184; 26481294	Possible Inv protein	Possible Inv protein	Invasion protein
Rv2024c	Restriction enzyme/m-6-adenine DNA methyltransferase (Mycobacterial adenine methyltransferase B "MamB")	26704977	Conserved hypothetical protein	Uncharacterized protein	putative helicase

1263

Rv2695	Probable serine hydrolase	26853625	Conserved hypothetical alanine rich protein	Conserved hypothetical alanine rich protein	
Rv2991	Probable flavin/deazaflavin oxidoreductase	26434506	Conserved protein	Conserved protein (F420-dependent protein)	
Rv3036c	Esterase	25224799	Probable conserved secreted protein TB22.2	Probable conserved secreted protein TB22.2	Possible membrane protein
Rv3354	protein kinase	25139900	Conserved hypothetical protein	Lipoprotein	Possible lipoprotein LprJ

<div align="center">Orthogonal annotation</div>

1264

Rv0256c *	B cell Antigen/probable INOS promoter binding protein	23827809; 28071726	PPE family protein PPE2	Uncharacterized PPE family protein PPE2	Predicted cobalt transporter in Mycobacteria
Rv2204c	probable serine hydrolase	26298037; 26853625; 26536359	Conserved protein	Protein Rv2204c	probable iron binding protein from the HesB_IscA_SufA family
Rv3779	polyprenylphosphomannosyl synthase/galactosaminyltransferase	21030587; 19717608	Probable conserved transmembrane protein alanine and leucine rich	Membrane protein (Probable conserved transmembrane protein alanine and leucine rich)	

1265
1266

1267 **Table 3.** Functional annotation of HHpred-uncorroborated enzymes following manual structural analysis.

Rv number	PDB ID	AA %	Inferred EC	Updated EC	Change	Recommended product name	Product name (Mycobrowser)
Rv0036c	2nsg	14%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved protein
Rv0738	2nsg	17%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved protein
Rv1632c	2p12	59%	3.6.1.-	3.6.1.-	Verified	putative cytidylyl-2-hydroxypropylphosphonate hydrolase	Hypothetical protein
Rv1727	2nsg	12%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein
Rv1734c	3l60	26%	1.2.4.-	2.3.-.-	Changed function	putative acyltransferase	Conserved hypothetical protein
Rv1775	3hwp	29%	3.-.-.-	3.7.1.-	Increased specificity	putative 3-oxo-carboxylic acid hydrolase	Conserved hypothetical protein
Rv2036	2nsg	15%	5.2.1.-	None	Reduced specificity	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein
Rv2968c	3kp9	21%	1.1.4.-	1.17.4.-	Nomenclature change	putative Vitamin K epoxide reductase	Probable conserved integral membrane protein
Rv3224B	1dbx	38%	3.1.1.-	None	Reduced specificity	putative trans-editing enzyme	Conserved hypothetical protein

1268

Table 4. Novel annotations transferred through structural similarity despite low sequence similarity. Selected
proteins with modelled structures highly similar to solved PDB structures of known function. Sequence similarities
range in the "twilight zone" of sequence similarity, below which remote homology is undetectable by sequence
similarity(126). TM$_{ADJ}$ above 0.52 indicates that template and under-annotated gene share structural folds.
Annotations from UniProt, Mtb systems Portal, and TubercuList are shown, along with the highest error-adjusted
structural similarity match, its identifier ("PDB"), and final product annotation. "Affirmatory" corroborate the
annotations in UniProt or Mtb Network Portal. "Novel" are entirely novel annotations to those in UniProt and Mtb
Network Portal, while "More specific" are in accord with annotations in other databases but describe product
function in greater detail.

Rv number	Top I-TASSER hit	AA%	TM$_{ADJ}$	PDB	Final annotation	MycoBrowser	UniProt	Mtb Net Portal	Type
Rv1139c	Integral membrane Methyltransferase	18	0.86	4a2n	putative integral Membrane Methyltransferase	Conserved hypothetical Membrane protein	Conserved hypothetical membrane protein (Membrane protein)	None	Novel
Rv1766c	Copper-sensing transcriptional Repressor CsoR	29	0.84	4m1p	Putative Transcription factor	conserved protein	Conserved protein	None	Novel
Rv3192c	5,10-methylenetetrahydro Methanopterin Reductase	16	0.83	1z69	Putative Monooxygenase	Conserved hypothetical alanine and Proline-rich protein	Conserved hypothetical Alanine and Proline-rich protein	Oxidoreductase	More specific
Rv2141c	M20 family Metallo peptidase	20	0.82	2pok	putative linear amide hydrolase	conserved protein	conserved protein	FIG016551: Putative peptidase	Affirmatory
Rv1775	2,4-diacetylphloroglucinol hydrolase	29	0.82	3hwp	putative 3-oxo-carboxylic acid hydrolase	Conserved hypothetical protein	Uncharacterized protein	None	Novel
Rv0052c	Isonitrile hydratase	33	0.81	3noo	putative hydro-lyase/putative deglycase	Conserved protein	Conserved protein	ThiJ/PfpI Family protein	Novel
Rv2036	mycothiol-dependent maleylpyruvate isomerase	15	0.73	2nsg	putative thiol-dependent DinB-like metalloenzyme	Conserved hypothetical protein	DinB family protein	None	More specific

1278
1279

Rv number	Product annotation	AA (%)	PDB macromolecule name	PDB ID	TM-score	TM_{ADJ}
Rv1085c	putative membrane transporter receptor protein	0.073	Chloride pumping rhodopsin	5b2nA	0.71	0.59
		0.058	Sodium pumping rhodopsin	4xtlA	0.71	0.59
Rv1462	putative transporter	0.173	ABC transporter, ATP-binding protein	4dn7A	0.79	0.67
Rv1510	putative Na+/H+ antiporter drug efflux protein	0.104	Putative drug/sodium antiporter	4z3nA	0.89	0.60
		0.088	Multi antimicrobial extrusion protein (Na(+)/drug antiporter) MATE-like MDR efflux pump	3mktA	0.84	0.54
Rv1680	putative phosphonate transporter component	0.151	PhnD, subunit of alkylphosphonate ABC transporter	3p7iA	0.91	0.77
		0.165	Binding protein component of ABC phosphonate transporter	3n5lA	0.89	0.76
Rv2325c	putative transport protein	0.165	Putative cobalt ABC transporter, permease protein	5d3mD	0.87	0.60
		0.215	Energy-coupling factor transporter transmembrane protein EcfT	4huqT	0.81	0.54
Rv2508c	putative MFS membrane transporter	0.104	Solute carrier family 2, facilitated glucose transporter member 3	5c65A	0.79	0.63
		0.119	D-xylose-proton symporter	4gbyA	0.78	0.63
Rv3630	putative Na+/H+ antiporter drug efflux protein	0.087	Multi antimicrobial extrusion protein (Na(+)/drug antiporter) MATE-like MDR efflux pump	3mktA	0.84	0.63
		0.108	Putative drug/sodium antiporter	4z3nA	0.83	0.62

1285

Table 6. Updated annotations add functional knowledge to genes required for optimal fitness during TB infection. Source annotation is the annotation listed by Bellerose and colleagues(83) and new annotation derived from the current project. Protein data Bank identifiers (PDB ID) of the protein structures matching H37Rv protein models are listed for structure-based annotations. PubMed IDs are listed for the papers from which functional annotations were manually curated.

Rv number	Source annotation	New annotation	PubMed ID	PDB ID
Literature annotations				
Rv1205	hypothetical protein	riboside monophosphate phosphoribohydrolase	25728768	
Rv2018	hypothetical protein	probable antitoxin VapB/Antigen	28066388, 23140854	
Rv2272	transmembrane protein	Probable gamma delta T-cell activator	23389928	
Rv2525c	tat pathway signal sequence	probable peptidoglycan hydrolase	16952959, 25869294, 25260828	
Rv2923c	hypothetical protein	probable Osmotically induced bacterial protein C (OsmC, a probable peroxide reductase)	22088319	
Rv3632	membrane protein	Putative flippase	21030587	
Rv3763	lipoprotein LpqH	adhesin/antigen LpqH	16098710, 12594264	
Rv3788	hypothetical protein	probable secondary channel binding-factor of RNA polymerase	22194445	
Literature & Structural inference				
Rv0191	integral membrane protein	Putative efflux pump	25690361, 12520088, 22132058	5c65, 4gby
Rv1433	exported protein	Probable l,d-transpeptidase LdtMt3	24041897	3tur, 3vae, 4jmn
Rv1769	hypothetical protein	T-cell antigen/putative aldehyde-lyase	26853625, 15102765	4v15
Rv3722c	hypothetical protein	probable serine hydrolase, probable aspartate transaminase		3ppl
Structural inference				
Rv0047c	hypothetical protein	Putative transcriptional regulator		1yg2a, 3l9f
Rv0259c	hypothetical protein	Putative lyase		2jh3, 4ccs
Rv0323c	hypothetical protein	Putative hydro-lyase/Putative linear amide hydrolase		1q7t, 5cgz
Rv0449c	hypothetical protein	Putative oxidoreductase		2ive, 1sez, 3nks, 3i6d, 3lov
Rv0767c	hypothetical protein	putative transcription factor		3mnl
Rv1085c	hemolysin-like protein	putative membrane transporter receptor protein		5b2nA, 4xtlA
Rv2052c	hypothetical protein	putative endodeoxyribonuclease		3igh
Rv2160A	hypothetical protein	putative transcription factor		2hyj
Rv3226c	hypothetical protein	putative peptidase		2icu
Rv3433c	hypothetical protein	putative hydro-lyase/putative isomerase		2ax3
Rv3719	hypothetical protein	putative amide-bond oxidoreductase		3dq0, 2exr, 4o95

1293 **Table 7. Updated annotation enriches functional interpretation of under-annotated genes affecting drug**
1294 **sensitivity.** Source annotation is the annotation listed by Bellerose and colleagues(83). Enriched conditions are the
1295 drugs' exposure under which differential mutant abundance was observed. Sources of updated annotation are listed
1296 in the evidence column. INH=Isoniazid; RIF=Rifampicin; EMB=Ethambutol; PZA=Pyrazinamide; HRZE=Combination
1297 regimen of INH, RIF, PZA, and EMB. "(+)" indicates enrichment observed at multiple time points.

Rv number	Source annotation	New annotation	Enriched condition(s)	Evidence
Rv0998	hypothetical protein	cAMP-dependent lysine acetyltransferase	EMB, RIF, HRZE	23553634
Rv1205	hypothetical protein	riboside monophosphate phosphoribohydrolase	EMB (+)	25728768
Rv0767c	hypothetical protein	putative transcription factor	INH	3mnl, 3bjb (PDB ID)
Rv3131	hypothetical protein	putative nitroreductase	INH	27094446; 28261197
Rv2140c	hypothetical protein	Phosphatidylethanolamine-binding protein TB18.6	INH	23907008; 27895634; 26238929
Rv2061c	hypothetical protein	probable serine hydrolase	INH	26853625; 26536359
Rv3267	hypothetical protein	probable peptidoglycan-arabinogalactan ligase	INH	27486192
Rv1770	hypothetical protein	probable serine hydrolase	PZA	26853625
Rv3005c	hypothetical protein	Probable membrane oxidoreductase component (MRC) DoxX	RIF	26067605
Rv1184c	exported protein	mycoacyltransferase	RIF	25331437; 25124040
Rv3036c	secreted protein	secreted esterase	RIF, INH	25224799
Rv2190c	hypothetical protein	peptidoglycan peptidase RipC/Antigen	RIF	24843173; 22952680; 28241799
Rv0079	hypothetical protein	Putative Dormancy Associated Translation Inhibitor (DATIN)	RIF (+)	22719925; 23819907; 28261197

1298

1299 **Figures**

1300
Fig. 1. Determining similarity thresholds for annotation inclusion criteria. Precision of EC number as regressed
against the geometric mean of TM-score and AA% ("μ_{geom}") for each specificity tier. Horizontal lines define (50%
and 75%) thresholds, the points where precision intersects with regression lines for each EC specificity curve
(labelled). Circles at the bottom and top are individual data points (incorrect = 0 and correct = 1, y-axis: precision,
x-axis: μ_{geom}). Circles are rendered at 10% opacity to visualize observation density. Only templates with AA% < 40%
were included.

1307

AA sequences

modeled structure solved structures

similarity metrics
- AA% identity
- TM-score

Observed TM-score/AA%
relationship with precision

Posterior probability of matching
CATH given TM-score[1]

ID under-annotated genes

Similarity to solved structures

extract EC, CATH, & GO from PDB entries

I-TASSER
Protein Structure & Function Predictions

manual curation   official enzyme nomenclature

experimentally determined GO & EC annotations

UniProt

Relate precision to similarity metrics

set similarity score-based inclusion criteria

Updated H37Rv reference genome annotation

locus_tag   Rv3722c
product   Probable aspartate transaminase
EC_number   2.6.1.1
note   Similar predicted tertiary structure to PDB:3pplA
inference   ab initio prediction:I-TASSER:5.1
inference   alignment:TM-align:PDB:3pplA
go_function   transaminase activity|GO:0008483|ISS

Protein model quality control

HHPred
MPI
Bioinformatics
Toolkit

Current annotation
locus_tag   Rv3722c
productConserved protein
noteFunctionalCategory: conserved hypotheticals

integrate new annotations

Tertiary Structure models

Gene products with known
EC and GO as positive control

Orthogonal validation and
Quality control

Structure-based
functional annotation

Fig. 2. Information flow for producing annotations from structural similarity. The flow of information and procedures for acquiring, processing, filtering, and representing information, running from retrieval of amino acid sequences to the final updated H37Rv annotation. Some details are omitted for clarity. The 1,725 amino acid sequences were retrieved from TubercuList and run through a local installation of I-TASSER v5.1. 1,711 of 1,725 Amino Acid sequences had models generated successfully. Comparison metrics for sequence (Amino acid identity) and Structure (TM-score) were extracted from I-TASSER output. To set criteria for annotation transfer, precision (Equation 1) of GO Term and EC number concordance between similar matches on PDB and true function of 363 positive controls with GO terms and EC numbers of known function were regressed against extracted similarity metrics to generate a curve relating the geometric mean of TM-score and Amino acid similarity to precision. These informed inclusion thresholds for transferring GO and EC annotations from structures on PDB similar to the 1,711 modeled structures. CATH Topology folds were transferred according to a previous precision curve based on TM-score. This threshold was also used for inclusion of proteins classes that vary in sequence more than structure (e.g. transporters) and as criteria for transferring annotations from structures that were not annotated with EC numbers or GO terms. Annotations derived only from structure were passed through orthogonal validation and manual structure analysis for verification that transferred annotations were reasonable. All annotations were programmatically collated into an updated H37Rv reference genome annotation.

Fig. 3. Orthogonal validation and quality assurance for structure-to-function inference. (A) Quality assurance and validation protocol assignment and (B) Decision workflow for retaining functional annotations inferred from structural similarity to proteins of solved structure and known function. (C) Heat map depicting fraction of EC number inferences corroborated by HHpred at each level of specificity. Fraction denominator is binned according to the number of E.C digits annotated (x-axis). (D,E) Structure quality assurance. Distribution of fraction of residues in "most favorable" region of Ramachandran plot prior to (D) and following application of a heuristic threshold (E) to discard biophysically improbable structural models.

Protein	Length (AA)	TM_{ADJ}	Seq. identity	Predicted Ligands	Metal-coordinating triad	Speculated function(s)	Evidence
Rv0036c	257	0.63	14%	L-Glutamic acid; Ni^{2+}	His43, His142, Val146	Intraphagosomal stress response; detoxification	Increased abundance in intraphagosomally-grown bacilli
Rv0738	182	0.75	17%	Zn^{2+}	His43, Asp112, His116	Hypoxic stress response; detoxification	More abundant in biofilm than in planktonic bacilli
Rv1727	189	0.69	12%	Fe^{3+}	His43, Ser121, His125	Iron starvation stress response	Co-transcribed transferrin-sequestering protein (Rv1872c) under carbon limitation
Rv2036	213	0.74	15%	Zn^{2+}	His52, Asp117, His121	Hypoxic stress response; detoxification	Highly correlated (R=0.72) expression with co-operonic transcriptional activator of hypoxic response (DosR) and virulence (PhoP) master regulators

Fig. 4. Manual structural analysis refines functional annotations uncorroborated by HHpred. (A-B) Conservation of structure- and sequence-features essential for C—C bond hydrolysis supports the inferred hydrolase function of Rv1775. **(A)** Structural alignment of modeled Rv1775 and its closest structural match (PDB 3hpw), a 2,4-diacetyl-phloroglucinolhydrolase of *Pseudomonas fluorescens* (*Pf*PhlG), The structures are superposed (top). Zoomed and re-oriented images of *Pf*PhlG Zinc-coordinating (box on left) and catalytic (pop-out) residues superposed with analogous *Mt*Rv1775 residues. **(B)** Comparison of functional and structural features between *Mt*Rv1775 and a putative PhlG homolog of *M. absecssus* (*Ma*PhlG), phloretin hydrolase, which catalyzes C—C bond hydrolysis of a different substrate. Comparison carried out in a similar scheme as in **A**. Superposition of the putative

homologs, color-annotated with conserved residues essential for phloroglucol moiety recognition and for phloretin substrate specificity in *Ma*PhlG(47). The structural similarity and conserved Zinc-coordinating and catalytic residues affirm Rv1775 as a *bona fide* C—C hydrolase, potentially with a substrate that includes a phloroglucol moiety, but likely not phloretin. Conservation of structure- and sequence-features characteristic of DinB-like metalloenzymes exemplified by structural homology of Rv2036 and a mycothiol-dependent maleylpyruvate isomerase from *Corynebacterium glutamicum* (*C. glu* MDMPI) (**C-G**). (**C**) Superposition of Rv2036 structure model and *C. glu* MDMPI (PDB: 2nsf). Conserved Zn^{2+}-coordinating (pink) and catalytic residues are highlighted (green). (**D**) Highly conserved residues Arg^{222} (C-terminal domain, Arg^{193} in Rv2036) and Asp^{151} (N-terminal domain, in Asp^{124} Rv2036) are in close proximity (4.22Å), suggesting conservation of their proposed role as inter-domain protein stabilizers(51) (**E**) Spatial conservation of Zn^{2+}-coordinating residues of the catalytic triad (Asp and Glu are observed interchangeably) are consistent with conserved catalytic function. (**F**) Surface hydrophobicity of Rv2036 model and 2nsf shows that the hydrophilic core proposed to underlie MDMPI catalysis(51) is relatively conserved. (**G**) Structure-based sequence alignment of Rv2036 and *C. glu* MDMPI with conserved residues were manually annotated according to prior work(51). (**H**) Summary of relevant genomic context potentially informative of function, protein similarity metrics between putative *M. tub* MDMPI homologs and *C. glu* MDPMI, and predicted protein features. All structural images were rendered in PyMol. Structurally homologous sequence alignments are based on TM-align(22) (****<5Å** between residues; *<10Å between residues).

Fig. 5. An updated H37Rv functional annotation. (**A-C**) Circos plots illustrating annotation coverage prior to the annotation effort (left) and following it (right), colored according to annotation status. In plots A and C, all 4,031 CDSs are represented as segments of equal width whereas (B) segments the ring into only the 1,725 under-annotated genes. Black genes reflect what was on TubercuList, are considered "annotated", and are mutually exclusive from the 1,725 under-annotated genes (white). (B) shows only the 1,725 under-annotated genes, whereas A and C include all 4,031 original CDS. Inside the circos rings are stacked bar charts with genes in 100kb bins according to gene start position. The terminal-proximal (± 250 kilobases) region is marked with dashed lines and labelled (*ter*). (**D**) Cumulative number of genes annotated, by annotation type. LBS = Ligand binding site. Percentages refer to under-annotated genes annotated/1,725 initial under-annotated genes. Genes are binned into mutually exclusive categories hierarchically: manually curated product name > structure-derived > literature notes > CATH > LBS. Manually curated and literature notes categories are combined as "Literature-curated" in the visualization. For the purposes of these counts, functional notes from publications implicating many proteins but not clearly establishing function were not counted (e.g. PMIDs 26536359, 21085642, and 22485162).

1375
Fig. 6. Functional annotations across *M. tuberculosis* metabolism. Annotated EC numbers for manually curated
and structure-inferred products were compared with annotations for each under-annotated gene in popular
databases **(A)** Set analysis of under-annotated genes (UAG) with an EC number assigned in this study compared to
popular databases. **(B)** Novelty of EC numbers for UAG annotated in this study with respect to popular databases.
(**C & D**) Distribution of EC numbers annotated across KEGG (**C**) Subsystems and (**D**) Pathways. Generic KEGG
subsystems are depicted. All pathways with at least three genes have the number of EC numbers displayed. For
subsystems with no pathways with three or more genes, the highest total pathway is displayed.

Fig. 7. Annotation of genes involved in pyrazinamide resistance. A library of *M. tuberculosis* H37Rv transposon insertion mutants was used to select for strains that were resistant to POA. The transposon insertion sites were mapped and the strains were characterized for their susceptibility to PZA, POA, and INH in comparison with wild-type H37Rv (MIC$_{90}$: 50 μg/mL PZA, 200 μg/mL POA, 0.0625 μg/mL INH). (A) H37Rv *Rv2705c::Tn*. (B) H37Rv *Rv2706c::Tn*. (C) H37Rv *Rv3256c::Tn*. (D) H37Rv *Rv3916c::Tn*. Error bar depicts standard deviation across triplicates. Pop-outs in (B) and (D) depict the transposon insertion sites relative to experimentally determined Transcription Start Sites(127, 128) (TSS). The insertion in (D) interrupts a TANNNT Pribnow Box (blue), destroying the Rv3916c promoter. Error bars are shown when there was a deviation in the calculated MIC90 across triplicates.

1392
Fig. 8. Structural analysis informs specific functional hypotheses for the basis of PZA resistance in the
***Rv3256c::Tn* mutant. (A-D)** Structural analysis identifies Rv3256c as a sugar isomerase (SIS) domain-containing
protein likely involved in phosphosugar metabolism or its regulation. InterPro functional domains are displayed for
the two strongest structural matches of Rv3256c, **(A)** *P. aerophilum* bifunctional phosphoglucose/phosphomannose
isomerase (*P. aer* PGI/PMI) and **(B)** *E. coli* glucosamine-6-P synthase (*E. col* GlmS). The InterPro domains labelled in
A and B are mapped onto the 3D structures of **(C)** *P. aer* PGI/PMI and **(D)** *E. col* GlmS. Rv3256c (charcoal) modeled
protein structure is optimally superposed to each of its matches. Rv3256c is structurally homologous to the SIS
domains of *E. col* GlmS and *P. aer* PGI/PMI and exhibits the alpha-beta-alpha sandwich fold of SIS(129). The pop-
out in A (**<5Å between residues) and labelled residue in C show the threonine residue essential for isomerase
activity in *P. aer* PGI/PMI (Thr^{291}) and other PMI homologs. The Thr^{291}-containing region appears to be absent from
Rv3256c. Likewise, Rv3256 lacks a glutamine amidotransferase domain homologous to *E. col* GlmS. From this
structural evidence, we conclude that Rv3256c is a SIS domain protein putatively involved in phosphosugar
metabolism and/or its regulation. Structural images were rendered in PyMol. Structurally homologous sequence
alignments from TM-align(22).
1407

Fig. 9. Structural analysis of transposon mutants refines functional hypotheses for their role in PZA resistance. (A-C) Structural analysis supports Rv3916c as a General control non-repressible 5 (GCN5)-related *N*-acetyltransferases (GNAT). Rv3916c matches exclusively comprised GNAT enzymes with low sequence similarity, typical among homologous GNAT enzymes (130). **(A)** Rv3916c (orange) superposed with its closest structural match (PDB 2a4n, charcoal), *Enterococcus faecium* aminoglycoside 6′-N-acetyltransferase (*E. faec* AAC(6′)-Ii). **(B)** *M. tuberculosis* Rv3916c structural model (top) and secondary structure topology (bottom), colored according to the four core conserved GNAT folds (wheat = poorly conserved secondary structure element) and secondarily (tones of the primary colors) by secondary structure elements in *E. faec* AAC(6′)-Ii (as defined by (131)) with which Rv3916c structurally aligns. All secondary structure elements of *E. faec* AAC(6′)-Ii are present in Rv3916c, with two gratuitous alpha-helices in its C-terminal arm (which is not well-conserved among GNAT enzymes(130)). The distinctive β-bulge (pop-out) within the β4 strand (red, bulge residues colored blue) characteristic of GNAT enzymes is present in Rv3916c, diverting β4 away from β5 to create the chasm where the acetyl-coenzyme A (yellow) is predicted to bind **(C)** Primary sequence of Rv3916c colored according to the scheme described in (B)**.** Asterisks mark predicted acetyl-CoA binding residues (black) and residues structurally aligning to known CoA-interacting residues (blue). All known CoA-interacting residues from *E. faec* AAC(6′)-Ii are conserved in Rv3916c and all predicted acetyl-CoA binding sites coincide or directly flank demonstrated sites of CoA interaction. The presence of the features in Rv3916c suggest it

1425 is a GNAT enzyme. GNAT enzymes catalyze transfer of an acyl moiety from an acyl-CoA to various substrates (130)
1426 making Rv3916c a probable acyl-CoA acetyltransferase and implicating $_p$Rv3916c::Tn in CoA pool modulation model
1427 of PZA resistance. **(D)** Summary of functional hypotheses for PZA resistance-conferring transposon mutants. All
1428 structural images were rendered in PyMol. Structurally homologous sequence alignments are based on TM-
1429 align(22).
1430
1431 **Supplementary Figures**

1432
Fig. S1. Information flow for producing annotations from literature curation. An initial extraction of the
existing annotation every "conserved hypothetical" and "unknown" protein from TubercuList totaled 1,057
unannotated protein-coding genes. Additionally, a 668 "ambiguous set" was manually determined from the
annotations on TubercuList, and these annotations were extracted and combined with the 1,057 hypothetical and
unknown proteins to give a total of 1,725 under-annotated genes. These genes were then searched in Google
Scholar, and pertinent articles were analyzed for annotation information, which was recorded in NCBI's Table File
Format (.tbl extension) for each gene, one file per gene. Every gene annotated with a novel product was compared
to annotation in other databases (Results). Decisions were made based on the criteria depicted above.

1441
Fig. S2. Relating precision to similarity metrics. A-D display binomial regressions on similarity metrics. **(A)**
Precision of GO term prediction as a function of C-scores from COFACTOR. Precision of GO term predictions are
plotted as a function of C-score for each of the three ontologies (Biological process, BP; Cellular Component, CC;
Molecular Function). The circles at the bottom and top of each plot are individual data points (representing 0 and 1
for incorrect and correct, respectively at a particular value). The circles are rendered at 10% opacity to visually depict
observation density. For benchmarking, only templates with C-scores above -1.5 were included, as structural
predictions with lower confidence are unlikely to reflect correct protein topology(21). **(B)** TM-score and amino acid
sequence identity (AA%) colored by correctness in the sample data. Dots' position indicates structural similarity (TM-
score, y-axis) and AA% (x-axis) between modeled structure and PDB entry, and their color indicates concordant (red)
or discordant (blue) EC number, to the specificity indicated by the pane label. **(C)** Function of EC number precision
using C-scores from the structure-to-function platform COFACTOR(23). Horizontal lines indicate the cutoffs used to
set the thresholds for hierarchical incorporation. **(D)** Similar to **A** & **C**, but using the geometric mean of Amino acid
identity and TM-score. This is the same data plotted in Figure 1 of the main text, and the model we selected for
determining inclusion criteria.
1456

1457
Fig. S3. Structure-based annotation inclusion protocols. A) Enzyme Commission (EC) number inclusion
protocol. The flow of processes and decisions each protein model was subjected to for determining EC based
annotations from structural homologs. Most processes and decisions were implemented in a fully automated
manner, but some corner cases had to be resolved manually. These manual cases were handled algorithmically, or
by previously established procedures where possible. For example, when EC numbers had to be assigned manually,
the procedures put forth by IUBMB were consulted and followed directly(25). **B) Gene Ontology (GO) term**
inclusion protocol. The flow of processes and decisions each modeled protein structure was subjected to for

determining GO based annotations from structural homologs/analogs. All processes and decisions were implemented in a fully automated manner, up until product assignment, those of which did not map to EC number had to be resolved manually.

Fig. S4. Comparison of annotations to existing databases. (A) To assess the novelty of manual product annotations, we compared each to their counterparts on UniProt(116), Mtb Network Portal(9) (which included annotations from TBDB(132)), PATRIC(6), RefSeq(36), BioCyc(119), and KEGG(120). We obtained the UniProt, Mtb Network Portal, and RefSeq annotations for each of our genes with new product annotation programmatically. For each under-annotated gene, we determined whether the database annotations agreed or disagreed with our manually curated annotations. For annotations that agreed, we recorded which source had the more descriptive annotation. Annotation from our literature curation absent in the other databases were considered novel gene annotations. If the annotations disagreed, we considered our annotation a candidate for additional gene product annotation, since both our annotation and those in other databases may describe true functions (bifunctional/moonlighting proteins). Existence of functional annotations for these genes were tallied for each database to assess their comprehensiveness and identify discrepancies between them. Furthermore, genes unannotated in any of the listed databases, but with annotations assigned in this study, were identified and enumerated. EC number assignments were also compared among the databases (Results). Of these databases, BioCyc and UniProt are the most comprehensive for GO term annotations, while UniProt and Mtb Network Portal have the fewest hypothetical proteins (Table 1). **(B-C) EC number annotation in the manual curation effort compared to widely used databases**. Manually curated annotations were compared with those in the databases in Table 1 to identify the presence or absence of EC number in GUF annotations. (**B**) Set of under-annotated genes annotated with an EC number in each of the five databases compared. The non-overlapping segments indicate the number of under-annotated genes annotated uniquely in that database. **(C)** Set of unique EC numbers across under-annotated for each database.

1491
**Fig. S5. Relationships between model quality, annotation specificity, and protein features. (A) Distribution
of model quality (C-score) for levels of annotation confidence and specificity.** For each of the 1,711 protein
models, the top three most similar PDB entries are included. Despite not explicitly accounting for differential degrees
of error in C-score across proteins modeled, annotations were transferred higher quality models were provided EC
thresholds correlated positively to model quality. (**B) Annotation classification as a function of amino acid
identity and TM-score.** Each dot represents a relationship between a structure in PDB and a predicted structure of
one of the 1,711 proteins for which structural models were able to be generated. Colors indicate which classification
the relationship qualified for based on our inclusion criteria. Shades running from blue to green depict progressively
stricter thresholds for EC inclusion tiers and confidence as determined by μ_{geom}:precision relation (Fig. 1). Red dots
indicate that similarity between the structure model/PDB structure was insufficient to qualify for an EC or GO
annotation, but met the TM-score$_{ADJ}$ criteria for transfer of CATH topology (blue and green points may have also
qualified for CATH transfer). Black dots indicate that though the raw TM-score would qualify the relationship for
transfer of EC/GO annotation, the model was of insufficient quality (C-score < -1.5), and the annotation was
therefore not transferred. The sharp transitions in the distribution reflect the hierarchical implementation of inclusion
criteria for functional annotations (EC number and GO terms). **(C) Protein length distributes similarly in models
in high- and low-quality structural models.** Density in the distribution of protein length for model above (red)
and below (yellow) the threshold for protein structure model quality (C-score > -1.5)(14, 21). **(D) Most frequently
annotated Cellular Component Gene Ontology (GO) terms**. Cellular component ontology GO term describe
proteins by the cellular location where they perform their function (28). GO terms were incorporated as described
in the flow diagram of Fig. S3B. Only those terms with ten or more occurrences are plotted, and no attempt was
made to collapse child ontologies into parents. GO terms implicating function specific to eukaryotic cellular
components are shown in red.

1514
1515 **Supplemental material descriptions**
1516
1517 **Dataset 1:** (A) **Under-annotated genes.** List of all under-annotated genes analyzed in the study, their reason for
1518 inclusion, and product annotation on TubercuList. (B) **Annotation summary**. A comprehensive annotation overview
1519 for all 1,725 under-annotated genes. For each under-annotated gene, annotations according to TubercuList, manual
1520 curation, structurally similar proteins from I-TASSER, CATH structural topologies, and Ligand Binding sites are listed,

along with structural model quality score (C-score) and classification (poor, strong, no model). (C) **Novel manual annotations**. All genes with manually curated product annotations novel to TubercuList, UniProt, and Mtb Network Portal. Annotations from each source are enumerated, along with PubMed IDs for each publication from which manually curated annotations were derived. (D) **Training set genes.** The list of genes selected randomly among proteins with experimentally-characterized, reviewed functional annotations on from UniProt (n=200) and manually curated annotations with specific functions (n=163) that were used as the training set to determine the putative and probable thresholds for transferring functional annotations between structurally similar proteins. **(E) Manually annotated gene products.** All genes with manually curated annotations from the literature in this work, their product annotations, level of confidence, and the sources from which the annotations were curated. (F) **Still lacking annotation.** All genes lacking LBS, CATH, functionally informative literature notes, or product annotations. Contains information about alternative databases with functional information and structure-quality (C-score > -1.5) that may provide leads to functional characterization or hypothesis generation. (G) **Syntenic unknown clusters.** Clusters of ≥3 consecutive genes lacking annotation have their potential roles listed, based on gene neighborhood. (H) **Chromosomal bias.** All genes in TubercuList and their annotation status, functional categories, and chromosomal location data (e.g. lagging vs. leading, terminus-proximal, etc.) used for enrichment analyses to identify orientation and spatial biases for uncharacterized genes. (I) **EC assignments.** All Enzyme commission numbers assigned to under-annotated genes in the study. (J) **KEGG mappings.** Generic KEGG pathway and subsystems mappings for all fully specified EC numbers (all four digits) annotated in this study. Many EC numbers map to multiple pathways/subsystems.

Dataset 2. Comparison of manually curated annotations to annotations in UniProt. (A-D) Contains results of comparison of 283 manually annotated under-annotated genes to UniProt annotations to identify instances where:
1. UniProt annotations and annotations from this study were discordant.
2. UniProt annotations and annotations from this study were concordant.
3. UniProt annotation was more informative of the function of the under-annotated genes.
4. Annotation from this study was more informative of the function of the under-annotated genes or
5. UniProt does not have a function annotation for the under-annotated genes.

We then distinguished genes annotated solely as antigens from those with other annotations to identify instances where the under-annotated genes had non-antigenic other functions associated with it in UniProt or the manual curation. (E) **Comparison of annotations manual annotations novel to UniProt to Mtb Network Portal, PATRIC, and RefSeq.** Results of comparing 173 under-annotated genes with manually annotated products absent from UniProt to the Mtb Network Portal, PATRIC, and RefSeq databases. Genes novelty are indicated by cell color. Yellow: Genes given a novel annotation in one or more databases where other databases lacked or had ambiguous annotation. For example, Rv0394c is annotated as Hyaluronidase/Chondrosulfatase only in our manual annotation; it is annotated ambiguously ("membrane protein" or "hypothetical protein") in other databases, so only the annotation in the "Our manual annotation" column is highlighted. Magenta: Genes given an annotation more informative or of greater confidence annotation compared to other analyzed databases. For example, Rv3632 is annotated as a flippase in our manual annotation; which is more specific than annotations of "membrane protein" or "cell division protein" in other databases. Lime Green: Gene annotations that matched between our manual annotation and other database annotations. Cyan: Gene annotations that were distinct in our manual annotation from one or more annotations in other databases such that it is possible our annotation and the other annotations could both describe the gene's functional product. For example, Rv0020c is annotated as a Probable Cell wall synthesis protein with FHA domain, FhaA in our database and Mtb Network portal's database, but these annotations differ from PATRIC's annexin VII annotation, so the annotations in all three of these columns are highlighted. Empty rows indicate that a database did not have an annotation for the protein. This sheet can be converted to html format using Microsoft Excel/Open office/Libreoffice or an online file conversion tool for ease of programmatically reproducing counts. PATRIC comments are included in the analysis for comparison, but mostly contain putative annotations from Phyre2 ab initio prediction software, which are considered weaker than the experimental evidence found by manual curation.

Dataset 3. Structure-based annotations. (A) **Strong protein-PDB matches.** All matches between predicted 3D structure of proteins encoded by under-annotated genes and PDB structure entries. All matches that either exceeded the geometric mean cutoff for inclusion in annotation as "putative" (> 50% observed probability of correct annotation), or with an adjusted TM-score > 0.52 are included. PDB IDs, macromolecular names, GO terms of the PDB match, are included, as well as statistic of model quality and match strength. (B) **Top PDB matches.** For all proteins with a model generated, the top 3 PDB entry matches are listed, irrespective of similarity metrics. Similarity metrics are listed, along with information about the matching PDB entry. (C) **CATH families.** All CATH families for all structure-PDB entry matches with adjusted TM-score greater than 0.52. The coordinates annotated with each CATH are listed, along with similarity metrics and model quality (C-score) (D) **Ligand Binding Sites.** All predicted Ligand Binding Sites for which C-score$_{LBS}$ ≥0.6. (E) **Transport-like PDB hits.** All matches according to criteria for transporters in methods are included. Information about the matching PDB template (E.C. number, GO terms, macromolecular name) are included, as well as similarity metrics between 3D structure models and PDB entries. (F) **Corroborated transport proteins.** The top two Transport-like protein hits (ranked by TM$_{ADJ}$) that passed the structure-inferred quality control pipeline. Sheets G-K contain orthogonal validation and manual structure analysis quality assurance steps. (G) **HHpred EC concordance.** EC numbers for each enzymatic annotations from proteins with functions inferred by structure alone. Left columns are the structure-inferred EC digits, and the rightmost four are those that were corroborated by HHpred hits. (H) **HHpred results.** Parsed HHpred output for all proteins with non-enzymatic annotations, alongside their structure-inferred product name. (I) **Consensus non-EC HHpred.** Sheet containing final reconciled product name corroborations, refutations, and modifications based on two-reviewer consensus of match/mismatch between HHpred hits and structure-inferred product annotations. (J) **Ramachandran filtering.** Fraction of residues in most favored, additional allowed, generously allowed, and disallowed regions for the proteins uncorroborated by HHpred, and some fully corroborated as positive controls used to build the threshold to define inclusion criteria to go on to manual structural analysis. (K) **Manual structure inspection.** Notes and results of manual verification of structure quality, structure-to-function mapping, and conservation of key structural features and residues for HHpred-uncorroborated proteins whose model structure passed Ramachandran filtering. (L) **CATH topologies.** Most commonly transferred CATH topologies and their associated functions. (M) **TetR matches.** Genes with PDB matches with Tetracyline Repressor 2 (TetR) CATH topologies. Only the highest TM$_{ADJ}$ is shown. Amino acid coordinates annotated with the CATH topology are indicated (AA coords), along with the PDB template, model quality (C-Score), Amino acid identity to the PDB match (Identity), and raw TM-score (TM-Score). (N) **Mycobacterial core genes.** Genes identified recently as part of the mycobacterial core genome(67) and annotated as hypothetical proteins (according to Table S3 of the recent work) are listed, along with their updated annotation and its source(s).

Text S1. Further details of approach, methods, and results.

Supplemental annotation file. Full machine-readable H37Rv annotation file in GFF format. Contains all annotations, notes, and evidence from the study