Chicken uric acid elimination via the uric acid transporters BCRP and MRP4 in the liver, kidneys, and intestines

Xuedong Ding, Manman Li, Shoufa Qian, Yuying Ma, Tianyi Fang, Xinlu Li, Huan Liu, Shibin Feng, Yu Li, Xichun Wang, Jinchun Li, and Jinjie Wu*

College of Animal Science and Technology, Clinical Veterinary Medicine, Anhui Agricultural University, 130 West Changjiang Road, Hefei 230036, China

Running title: Chicken uric acid elimination via BCRP and MRP4

*Corresponding author: Jinjie Wu. Tel: +86-13505514926; Email: wjj@ahau.edu.cn.

Key words: uric acid, BCRP, MRP4, chicken

Summary statement: The uric transporters BCRP and MRP4 in different tissues may exhibit the interaction to maintain the serum uric acid levels.

Abstract

Breast cancer resistance protein (BCRP) and multidrug resistance protein 4 (MRP4) are involved in uric acid excretion in humans and mice. Despite evidence suggesting that chicken renal proximal tubular epithelial cells participate in uric acid secretion, the roles of BCRP and MRP4 in chickens remain unclear. This study evaluated the relationship between chicken BCRP and MRP4 expression and renal function in the liver, kidneys, and intestines. Sixty 20-day-old Isa brown laying hens were randomly divided into four groups: a control group (NC) and groups provided with sulfonamide-treated drinking water (SD), a diet supplemented with fishmeal (FM), and an intraperitoneal injection of uric acid (IU). Serum uric acid, creatinine, and blood urea nitrogen (BUN) levels were significantly higher in the SD and IU groups than in the NC group. BCRP and MRP4 levels in the SD and IU groups were significantly increased in the kidneys and ileum and decreased in the liver. In the FM group, BCRP and MRP4 were significantly increased in the kidneys and slightly increased in the ileum. These results demonstrate that chicken BCRP and MRP4 are involved in renal and intestinal uric acid excretion. When renal function is impaired, serum uric acid increased and BCRP and MRP4 in the liver, kidneys, and ileum exhibit compensatory increases; when renal function is normal, serum uric acid changes have no effect on ileum BCRP and MRP4 expression. Therefore, this study may provide the references to the uric acid regulation in human.

Introduction

Uric acid (urate) is the final product of purine metabolism. Although in some cases, dietary (Zhang et al.,2016), genetic (Maalouf,2011; Petru et al.,2016), or disease-related (Kumar et al.,2017) uric acid overproduction is the basis of hyperuricemia, the main cause, in fact, of hyperuricemia is reduced uric acid excretion (Oka et al.,2014; Tsai et al.,2017). In an analysis of uric acid metabolism in 65 patients with hyperuricemia, six (9.2%) exhibited an overproduction phenotype, 52 (80.0%) exhibited an underexcretion phenotype, and seven (10.8%) exhibited a mixed phenotype (Oka et al.,2014). The kidney is the main organ responsible for uric acid

excretion, accounting for approximately two-thirds of the total uric excretion in the

body; the remaining one-third mainly involves the intestines (Maesaka and

Fishbane,1998). This process involves several uric acid transporters; breast cancer

resistance protein (BCRP; ABCG2) and multidrug resistance protein 4 (MRP4;

ABCC4) are the major proteins involved in uric acid excretion and expression in the

human liver, kidneys, and intestines (Maliepaard et al.,2001; Rius et al.,2003; Van

Aubel RA et al.,2005; Huls et al.,2008). BCRP is a high-capacity uric acid transporter

that physiologically mediates renal and extra-renal (intestinal) uric acid excretion; its

dysfunction leads to hyperuricemia (Matsuo et al.,2014). Extensive data indicates that

BCRP plays an important role in intestinal uric acid excretion in mice and humans

(Hosomi et al.,2012). Renal uric acid excretion is significantly reduced after

nephrectomy in mice, while serum uric acid does not change and ileum BCRP

expression is significantly increased (Yano et al.,2014). Therefore, alterations in

intestinal BCRP may serve as a compensatory mechanism. When renal uric acid

excretion is reduced, intestinal uric acid excretion is increased to maintain the uric

balance.

Many studies have examined the functions of human, rat, and mouse BCRP in the

kidneys and intestines; however, relatively few studies have evaluated MRP4. Similar

to BCRP, MRP4 is a uric acid unidirectional efflux pump with multiple allosteric

substrate binding sites, which is expressed in the apical membrane of human renal

proximal tubules (van Aubel RA et al.,2002). It is responsible for uric acid excretion

by transporting uric acid from tubular epithelial cells into renal tubule lumens. MRP4

is also expressed in the basal membrane of human hepatocytes and is involved in the

transport of uric acid in the liver (Russel et al.,2008). In HEK293 cells, MRP4 can

transport uric acid concurrently with cAMP or cGMP and uric acid excretion

increases with the overexpression of MRP4 (Van Aubel RA et al.,2005).

Uricase in the mouse liver can convert uric acid into allantoin; however, human and

chicken livers lack uricase (Wu et al.,1989). Accordingly, uric acid metabolism in

humans occurs via a different mechanism than in mice. Therefore, the chicken may

constitute a more useful model than mice for studying human uric acid transporters. In

vitro studies have demonstrated the presence of active urate secretion in chicken renal

proximal tubular epithelial cells (cPTCs) and this may be related to multiple uric acid

transporters (Dudas et al.,2005). Subsequently, Bataille et al. (Bataille et al.,2008)

showed that BCRP and MRP4 are expressed in cPTCs and that uric acid secretion is

reduced by 60–70% in response to a 75% reduction in chicken *Mrp4* expression by

short hairpin RNA interference. The net transepithelial transport of uric acid decreases

when *BCRP* is knocked down (Bataille et al.,2011), though the change is not

significant, indicating that MRP4 is the main route for urate secretion in chicken

proximal tubules. However, the roles of BCRP and MRP4 in chicken uric acid

excretion remain unclear. Therefore, the aim of this study was to investigate the

relationship between serum uric acid levels and BCRP and MRP4 levels in the liver,

kidneys, and intestines and to evaluate kidney and extrarenal uric acid excretion in

chickens. These findings may lay the foundation for the treatment and prevention of

hyperuricemia.

Materials and methods

Animal grouping and Treatment

Seventy female 1-day-old Isa brown laying hens were purchased from Anhui Poultry

Industry Co., Ltd. (China), all chickens were reared in cages and allowed ad libitum

consumption of feed and water. The room temperature was 25~30°C. The diets

compositions were arranged based on National Research Council (1994)

recommended requirements and diet containing 204.3 g kg^{-1} of crude protein, 11.5 g

kg^{-1} of calcium, 4.2 g kg^{-1} of phosphorus and 12.11 MJ kg^{-1} of ME. On the 20st day,

sixty healthy chicken were adopted and randomly divided into four groups (n = 15 per

group, weight 189.3 ±13.8 g). The normal control group (NC) was fed the basal diet;

the sulfonamide drug group (SD) was fed the basal diet, and the sulfamonomethoxine

sodium soluble powder was added to the drinking water (8 mg L^{-1} d^{-1}); the fish meal

group (FM) was added 16% fishmeal in the basal diet (crude protein 27.6%); the

injection uric acid group (IU) was fed on the basal diet and received uric acid 250 mg

kg^{-1} d^{-1} (suspended in 0.5% CMC-Na solution) by intraperitoneally injected. The

experiment was lasted for 3 weeks, on the 41st day, the blood were collected from

jugular vein after fasting for 12 h. After clot for approximately 30 min at room

temperature, the blood was centrifuged at 3,500 g min^{-1} for 10 min to obtain serum.

The collected serum was stored at -20°C. Finally all chickens were killed by

decapitation. The liver, kidney, jejunum, and ileum were collected and stored in 4%

paraformaldehyde and liquid nitrogen for testing. All experimental procedures for the

care and use of animals in the present study were approved by the Animal Care

Committee of Anhui Agricultural University.

Instruments and Reagents

Automated biochemical analyzer (Beckman AU680, USA), Electrophoresis apparatus

(Tanon, China), Cryogenic centrifuge (TGL-18R, China), Microscope (Olympus

CX31, Japan), ChemiDoc Imaging System (Bio-Rad, USA), RNA Concentration

analyzer (NanoVue plus, Thermo, USA), Real-time Quantitative PCR (Thermo, USA),

RIPA cell lysate (Biosharp, China), Protein phosphatase inhibitor (Solarbio, China),

BCA protein concentration assay kit (Biosharp, China), Sulfamonomethoxine Sodium

soluble powder (Shijiazhuang City Hengxin Pharmaceutical Co., Ltd. China), Uric

acid (BioXtra, ≥ 99% HPLC, Sigma, USA), Sodium carboxymethyl cellulose

(CMC-Na, Solarbio, China), Defatted fish meal (Crude protein ≥65%, China).

Serum uric acid, creatinine and blood urea nitrogen (BUN) Levels

The serum uric acid, creatinine and BUN levels was measured using an automatic

biochemical analyzer.

***Real-time Quantitative PCR* (qPCR)**

Total RNA was extracted from liver, kidney, jejunum, and ileum (100 mg). And 500

ng total RNA was reverse transcribed from each sample. QPCR reaction program was

as follows.: 95°C, 02 min; 95°C, 15 s; 60°C, 1 min (40 cycles); 60°C, 30 s; 60°C ~

95°C, 0.2°C s^{-1}; 20°C, 10 s. The chicken *BCRP* and *MRP4* qPCR primers are shown

in Table 1.

Western Blotting

Total protein was extracted from liver, kidney, jejunum, and ileum (100mg) for

western blotting analysis of BCRP, MRP4 and β-actin. Immunoblotting was assayed

using anti-BCRP (cat. no. bs-0662R, Polyclonal, 1:1000, Bioss, China), MRP4 (cat.

no. bs-1422R, Polyclonal, 1:1000, Bioss, China) as well as β-actin (cat. No.

abs137975, Monoclonal, 1:1000, Absin, China) antibodies. The proteins were

visualized using western blotting detection kit (Advansta, USA). The density of bands

was analyzed by Image pro plus 6.0 and normalized to β-actin.

Immunohistochemistry

Tissues of liver, kidney, jejunum and ileum were fixed in 4% paraformaldehyde and

paraffin-embedded sections (5 μm thick). BCRP and MRP4 protein expression were

detection by immunohistochemistry as described by Liu et al (Liu et al.,2015). The

primary antibodies were used in this study: anti-BCRP (1:400) and anti-MRP4 (1:300).

And the secondary antibody was used Goat anti-rabbit IgG(cat. no. AP132P, 1:1,000,

Millipore, USA). Finally the immunolabeled sections were observed under a light

microscope and taken photos.

Statistical Analysis

Statistical analysis was used by IBM SPSS Statistics Version19 (SPSS). Data were

expressed as mean ± standard error of mean (s.e.m.). One-way analysis of variance

with SPSS, Duncan multiple comparison, and *P < 0.05* was statistically significant.

Results

Renal function

As shown in Table 2, compared with the control group, serum uric acid levels were

higher ($p = 0.01$) and creatinine and BUN levels were significantly higher ($p < 0.01$)

in the sulfonamide-treated drinking water (SD) group. Serum uric acid, creatinine, and

BUN levels were significantly higher in the intraperitoneal injection of uric acid (IU)

group than in the control group ($p < 0.01$); however, there was no significant

difference between the diet supplemented with fishmeal (FM) group and the control

group.

BCRP and MRP4 mRNA and protein expression in normal chickens

Real-time quantitative PCR and western blotting were used to detect the distributions

of BCRP and MRP4 in the liver, kidneys, jejunum, and ileum of normal chickens (Fig.

1). The results showed that chicken BCRP (Fig. 1A) was highly expressed in the

jejunum and ileum ($p < 0.01$), lowly expressed in the liver, and minimally expressed

in the kidneys ($p < 0.01$). MRP4 expression was similar to that of BCRP (Fig. 1B); its

expression levels in the liver and kidneys were lower than in the jejunum and ileum (p

< 0.01), but did not differ significantly between the liver and kidneys. In addition,

relative expression analyses showed that BCRP and MRP4 were mainly expressed in

the jejunum and ileum. BCRP levels were higher than MRP4 levels in the jejunum,

ileum, and liver and BCRP and MRP4 levels did not differ significantly in the kidneys

(Fig. 1C).

Immunohistochemical staining showed that BCRP and MRP4 were expressed in the

liver cells, renal apical membrane, intestinal smooth muscle, and intestinal villi (Fig. 2

and Fig. 3). Moreover, MRP4 levels in these four tissues were lower than BCRP

levels.

BCRP and MRP4 expression in various treatment groups

BCRP and *MRP4* mRNA levels in the liver, kidneys, jejunum, and ileum of chickens

in each treatment group were evaluated by qPCR. *BCRP* and *MRP4* levels showed

similar trends. As shown in Fig. 4, compared with the control group, *BCRP* and *MRP4*

levels were significantly higher ($p < 0.05$, $p < 0.01$) in the ileum and slightly higher

in the kidneys of the SD group. *BCRP* and *MRP4* levels were significantly increased

in the kidneys and ileum of the IU group. Additionally, in the SD and IU groups,

BCRP and *MRP4* expression levels were significantly decreased in the liver ($p < 0.05$,

$p < 0.05$). In the FM group, *BCRP* and *MRP4* expression levels in the kidneys were

significantly increased ($p < 0.05$, $p < 0.05$), the levels in the ileum were slightly

increased, and there was no obvious difference in the liver. Moreover, *BCRP* and

MRP4 expression levels in the jejunum of the three experimental groups showed a

decreasing trend, though the difference was not significant.

Western blotting results showed that BCRP and MRP4 expression levels in the liver,

kidneys, jejunum, and ileum of each group were consistent with the mRNA

expression levels (Fig. 5). In the SD group, BCRP and MRP4 protein levels were

slightly increased in the kidneys and significantly increased in the ileum ($p < 0.05$, p

< 0.01). In the IU group, BCRP and MRP4 protein levels were significantly increased

in the kidneys and ileum. In the SD and IU groups, BCRP and MRP4 levels were

decreased in the liver. In the FM group, BCRP and MRP4 protein levels were

significantly increased in the kidneys ($p < 0.01$, $p < 0.05$) and slightly increased in the

ileum, while there was no significant difference in the liver levels compared to the

control group. BCRP and MRP4 protein levels in the jejunum of the three

experimental groups did not differ significantly.

Finally, we used immunohistochemistry to evaluate BCRP and MRP4 protein

expression levels in the liver, kidneys, jejunum, and ileum in each treatment group. As

shown in Fig. 6 and Fig. 7, BCRP and MRP4 were more highly expressed in the

kidneys and ileum of the three experimental groups than in the control group, whereas

jejunum BCRP and MRP4 levels were similar to those of the control group. Liver

BCRP and MRP4 protein expression levels were lower in the SD and IU groups than

in the control group, while in the FM group, liver expression levels were similar to

those in the control group.

Dissussion

BCRP and MRP4 are uric acid transporters present in various organs, such as the

human liver, kidneys, and intestines, which can be expressed in heterogeneous

systems for uric acid transport (Dudas et al.,2005; Van Aubel RA et al.,2005;

Woodward et al.,2009). BCRP and MRP4 are involved in human and mouse uric acid

excretion and in vitro studies have revealed that BCRP and MRP4 are expressed in

cPTCs (Bataille et al.,2008). However, their roles in the chicken uric acid transport

system remain unclear. The results of this study showed that BCRP and MRP4 are

highly expressed in the jejunum and ileum of chickens, with low expression in the

liver and kidney and minimal expression of BCRP in the kidneys. Several BCRP

localization studies have reported relatively high expression in rat and mouse kidneys

as well as in the small intestine, especially in the ileum (Tanaka et al.,2005), while in

humans, the apical membrane of hepatocytes, colonic epithelial cells, and placental

syncytium trophoblasts exhibit relatively high expression (Doyle et al.,1998;

Maliepaard et al.,2001). MRP4 is most highly expressed in the human kidneys,

followed by the liver and intestines (Hilgendorf et al.,2007). However, in mice, MRP4

levels are significantly higher in the kidneys than in the liver and intestines, and

female liver and kidney expression levels are significantly higher than in male mice

(Maher et al.,2005). These results indicate that although BCRP and MRP4 can be

expressed heterologously, their tissue distributions differ among species and these

differences may be related to species-specific mechanisms of uric acid metabolism.

Previous studies have shown that endogenous uric acid in mice is secreted directly

from the blood into the intestinal lumen of all bowel segments (Xu et al.,2016; Yun et

al.,2017). Ileum secretion is approximately 3-fold and 2-fold higher than jejunum and

colon secretion, respectively (Hosomi et al.,2012). These findings indicate that the

mouse ileum is the main site of intestinal uric acid secretion (Yun et al.,2017). The

results of this study demonstrate that chicken BCRP and MRP4 are mainly expressed

in the jejunum and ileum, with higher expression in the ileum; accordingly, the roles

of BCRP and MRP4 in ileum uric acid excretion may be particularly important.

The kidney has been recognized as the main regulator of serum uric acid and the

excretion of renal uric acid is determined by the balance of urate reabsorption and

re-secretion (Bobulescu and Moe,2012). In humans, approximately 70% of uric is

secreted into the urine through the renal tubules (Maesaka and Fishbane,1998). BCRP

and MRP4 are critical for uric acid secretion in human and mouse kidneys

(Sakurai,2013). A high-protein diet can increase chicken serum uric acid levels (Guo

et al.,2005). However, in this study, the high-protein diet FM group did not

demonstrate an increase in serum uric acid, creatinine, or BUN levels, while renal

BCRP and MRP4 expression increased significantly and ileum expression increased

slightly. These findings indicate that when renal function is normal, the kidney is the

main site of uric acid clearance and that renal BCRP and MRP4 are involved in renal

uric acid excretion.

In this study, serum uric acid, creatinine, and BUN levels were significantly

increased in the SD and IU groups compared with the control group. In the SD group,

sulfonamide crystallization may have blocked the renal tubules and caused renal

damage (Mustafa et al.,2014), thereby reducing uric acid excretion and increasing

serum uric acid. In the IU group, intraperitoneal injection of uric acid not only raised

serum uric acid levels, but also caused renal damage (Braga et al.,2017; Romi et

al.,2017). Mouse studies have shown that the ileum plays an important role in ileum

uric acid clearance during kidney injury (Yano et al.,2014; Nagura et al.,2016).

Similarly, our results showed that chicken serum uric acid increased when serum

creatinine and BUN levels were elevated in the SD and IU groups. In addition, ileum

BCRP and MRP4 protein and gene expression levels were significantly increased.

These results suggest that chicken kidney and intestine BCRP and MRP4 are involved

in uric acid clearance and that when kidney function is impaired, uric acid excretion in

the ileum can provide a compensatory mechanism by increasing BCRP and MRP4

expression. In addition, BCRP and MRP4 levels in the jejunum were slightly lower in

the three experimental groups compared with expression in the control group. The

mechanisms underlying these differences should be evaluated in future studies.

Serum uric acid levels in the SD and IU groups were significantly higher than those

in the control group, while liver BCRP and MRP4 expression levels were significantly

lower. In addition, serum uric acid levels in the FM group were decreased and liver

BCRP and MRP4 expression levels were slightly increased. These results indicate that

changes in liver BCRP and MRP4 expression contradict the changes in serum uric

acid levels. Previous studies have shown that BCRP and MRP4 are expressed as uric

acid efflux proteins in the basolateral membrane of hepatocytes (Rius et al.,2003). In

this study, immunohistochemical staining showed that BCRP and MRP4 were also

expressed in blood vessels, indicating that they may participate in liver uric acid entry

into the blood circulation. The decrease in liver BCRP and MRP4 expression with

increasing serum uric acid levels may reduce serum uric acid levels.

Several studies have proposed a "Remote Sensing and Signaling Hypothesis" (Ahn

and Nigam,2009; Wu et al.,2011; Nigam,2015), suggesting that some uric transporters,

such as BCRP and MRP4, present in different tissues are part of an inter-organ and

inter-organismal communication network that maintains uric acid levels in the case of

kidney or other organ injury (Bhatnagar et al.,2016). Our study supports the

hypothesis that the uric acid transporters BCRP and MRP4 are involved in the

regulation of serum uric acid levels in the liver, kidneys, and intestines.

This study had some limitations; the mechanisms underlying the interaction

between changes in serum uric acid levels and liver, kidney, and intestinal BCRP and

MRP4 levels remain unclear. In addition, an interaction between BCRP and MRP4

may exist and further studies are needed to evaluate this.

Conclusions

Our results show that chicken BCRP and MRP4 participate in renal and intestinal

uric acid excretion. When renal function is impaired, BCRP and MRP4 expression in

the kidneys and ileum exhibit compensatory increases; however, when renal function

is normal, changes in serum uric acid levels have no effect on ileum BCRP and MRP4

levels. Importantly, inter-organ communication between uric transporters in different

tissues during uric regulation remains unclear and this coordination should be

investigated in future studies.

Acknowledgments

We are grateful to the animal hospital of Anhui Agricultural University. We wish to

thank anonymous reviewers for their kind advice.

Competing Interests

No conflicts of interest are declared by the authors.

Funding

This work was supported by grants from the National Key Research and

Development Program of China [2016YFD0501205] and the Key Research and

Development Program of Anhui Province of China [1804a07020135].

References

Ahn, S.Y. and Nigam, S.K. (2009). Toward a systems level understanding of organic
anion and other multispecific drug transporters: a remote sensing and signaling
hypothesis. *Mol. Pharmacol.* **76**,481-490.

Bataille, A.M., Goldmeyer, J. and Renfro, J.L. (2008). Avian renal proximal tubule
epithelium urate secretion is mediated by Mrp4. *Am. J. Physiol. Regul. Integr.*
Comp. Physiol. **295**,R2024-2033.

Bataille, A.M., Maffeo, C.L. and Renfro, J.L. (2011). Avian renal proximal tubule
urate secretion is inhibited by cellular stress-induced AMP-activated protein
kinase. *Am. J. Physiol. Renal Physiol.* **300**,F1327-1338.

Bhatnagar, V., Richard, E.L., Wu, W., Nievergelt, C.M., Lipkowitz, M.S., Jeff, J.,
Maihofer, A.X. and Nigam, S.K. (2016). Analysis of ABCG2 and other urate
transporters in uric acid homeostasis in chronic kidney disease: potential role
of remote sensing and signaling. *Clin Kidney J* **9**,444-453.

Bobulescu, I.A. and Moe, O.W. (2012). Renal transport of uric acid: evolving
concepts and uncertainties. *Adv Chronic Kidney Dis* **19**,358-371.

Braga, T.T., Forni, M.F., Correa-Costa, M., Ramos, R.N., Barbuto, J.A., Branco,
P., Castoldi, A., Hiyane, M.I., Davanso, M.R., Latz, E., et al. (2017).
Soluble Uric Acid Activates the NLRP3 Inflammasome. *Sci Rep* **7**,39884.
Doyle, L.A., Yang, W., Abruzzo, L.V., Krogmann, T., Gao, Y., Rishi, A.K. and
Ross, D.D. (1998). A multidrug resistance transporter from human MCF-7
breast cancer cells. *Proc. Natl. Acad. Sci. U.S.A.* **95**,15665-15670.
Dudas, P.L., Pelis, R.M., Braun, E.J. and Renfro, J.L. (2005). Transepithelial urate
transport by avian renal proximal tubule epithelium in primary culture. *J. Exp.*
Biol. **208**,4305-4315.
Guo, X., Huang, K. and Tang, J. (2005). Clinicopathology of gout in growing layers
induced by high calcium and high protein diets. *Br. Poult. Sci.* **46**,641-646.
Hilgendorf, C., Ahlin, G., Seithel, A., Artursson, P., Ungell, A.L. and Karlsson, J.
(2007). Expression of thirty-six drug transporter genes in human intestine,
liver, kidney, and organotypic cell lines. *Drug Metab. Dispos.* **35**,1333-1340.
Hosomi, A., Nakanishi, T., Fujita, T. and Tamai, I. (2012). Extra-renal elimination
of uric acid via intestinal efflux transporter BCRP/ABCG2. *PLoS ONE*
7,e30456.
Huls, M., Brown, C.D., Windass, A.S., Sayer, R., van den Heuvel JJ, Heemskerk,
S., Russel, F.G. and Masereeuw, R. (2008). The breast cancer resistance
protein transporter ABCG2 is expressed in the human kidney proximal tubule
apical membrane. *Kidney Int.* **73**,220-225.
Kumar, R., Joshi, G., Kler, H., Kalra, S., Kaur, M. and Arya, R. (2017). Toward an
Understanding of Structural Insights of Xanthine and Aldehyde Oxidases: An
Overview of their Inhibitors and Role in Various Diseases. *Med Res Rev* .
Liu, Y.W., Sun, W.F., Zhang, X.X., Li, J. and Zhang, H.H. (2015). Compound
Tufuling Granules ([characters: see text]) regulate glucose transporter 9
expression in kidney to influence serum uric acid level in hyperuricemia mice.
Chin J Integr Med **21**,823-829.
Maalouf, N.M. (2011). Metabolic syndrome and the genesis of uric acid stones. *J Ren*
Nutr **21**,128-131.
Maesaka, J.K. and Fishbane, S. (1998). Regulation of renal urate excretion: a
critical review. *Am. J. Kidney Dis.* **32**,917-933.
Maher, J.M., Slitt, A.L., Cherrington, N.J., Cheng, X. and Klaassen, C.D. (2005).
Tissue distribution and hepatic and renal ontogeny of the multidrug
resistance-associated protein (Mrp) family in mice. *Drug Metab. Dispos.*
33,947-955.
Maliepaard, M., Scheffer, G.L., Faneyte, I.F., van Gastelen MA, Pijnenborg, A.C.,
Schinkel, A.H., van De Vijver MJ, Scheper, R.J. and Schellens, J.H. (2001).
Subcellular localization and distribution of the breast cancer resistance protein
transporter in normal human tissues. *Cancer Res.* **61**,3458-3464.
Matsuo, H., Nakayama, A., Sakiyama, M., Chiba, T., Shimizu, S., Kawamura, Y.,
Nakashima, H., Nakamura, T., Takada, Y., Oikawa, Y., et al. (2014).
ABCG2 dysfunction causes hyperuricemia due to both renal urate
underexcretion and renal urate overload. *Sci Rep* **4**,3755.

Mustafa, S., Alsughayer, A., Elgazzar, A., Elassar, A. and Al, S.F. (2014). Effect of
sulfa drugs on kidney function and renal scintigraphy. *Nephrology (Carlton)*
19,210-216.

Nagura, M., Tamura, Y., Kumagai, T., Hosoyamada, M. and Uchida, S. (2016).
Uric acid metabolism of kidney and intestine in a rat model of chronic kidney
disease. *Nucleosides Nucleotides Nucleic Acids* **35**,550-558.

Nigam, S.K. (2015). What do drug transporters really do. *Nat Rev Drug Discov*
14,29-44.

Oka, Y., Tashiro, H., Sirasaki, R., Yamamoto, T., Akiyama, N., Kawasugi, K.,
Shirafuji, N. and Fujimori, S. (2014). Hyperuricemia in hematologic
malignancies is caused by an insufficient urinary excretion. *Nucleosides*
Nucleotides Nucleic Acids **33**,434-438.

Petru, L., Pavelcova, K., Sebesta, I. and Stiburkova, B. (2016). Genetic
background of uric acid metabolism in a patient with severe chronic
tophaceous gout. *Clin. Chim. Acta* **460**,46-49.

Rius, M., Nies, A.T., Hummel-Eisenbeiss, J., Jedlitschky, G. and Keppler, D.
(2003). Cotransport of reduced glutathione with bile salts by MRP4 (ABCC4)
localized to the basolateral hepatocyte membrane. *Hepatology* **38**,374-384.

Romi, M.M., Arfian, N., Tranggono, U., WAW, S. and DCR, S. (2017). Uric acid
causes kidney injury through inducing fibroblast expansion, Endothelin-1
expression, and inflammation. *BMC Nephrol* **18**,326.

Russel, F.G., Koenderink, J.B. and Masereeuw, R. (2008). Multidrug resistance
protein 4 (MRP4/ABCC4): a versatile efflux transporter for drugs and
signalling molecules. *Trends Pharmacol. Sci.* **29**,200-207.

Sakurai, H. (2013). Urate transporters in the genomic era. *Curr. Opin. Nephrol.*
Hypertens. **22**,545-550.

Tanaka, Y., Slitt, A.L., Leazer, T.M., Maher, J.M. and Klaassen, C.D. (2005).
Tissue distribution and hormonal regulation of the breast cancer resistance
protein (Bcrp/Abcg2) in rats and mice. *Biochem. Biophys. Res. Commun.*
326,181-187.

Tsai, C.W., Lin, S.Y., Kuo, C.C. and Huang, C.C. (2017). Serum Uric Acid and
Progression of Kidney Disease: A Longitudinal Analysis and Mini-Review.
PLoS ONE **12**,e0170393.

Van Aubel RA, Smeets, P.H., van den Heuvel JJ and Russel, F.G. (2005). Human
organic anion transporter MRP4 (ABCC4) is an efflux pump for the purine
end metabolite urate with multiple allosteric substrate binding sites. *Am. J.*
Physiol. Renal Physiol. **288**,F327-333.

van Aubel RA, Smeets, P.H., Peters, J.G., Bindels, R.J. and Russel, F.G. (2002).
The MRP4/ABCC4 gene encodes a novel apical organic anion transporter in
human kidney proximal tubules: putative efflux pump for urinary cAMP and
cGMP. *J. Am. Soc. Nephrol.* **13**,595-603.

Woodward, O.M., Köttgen, A., Coresh, J., Boerwinkle, E., Guggino, W.B. and
Köttgen, M. (2009). Identification of a urate transporter, ABCG2, with a
common functional polymorphism causing gout. *Proc. Natl. Acad. Sci. U.S.A.*

106,10338-10342.

Wu, W., Dnyanmote, A.V. and Nigam, S.K. (2011). Remote communication through
solute carriers and ATP binding cassette drug transporter pathways: an update
on the remote sensing and signaling hypothesis. *Mol. Pharmacol.* **79**,795-805.

Wu, X.W., Lee, C.C., Muzny, D.M. and Caskey, C.T. (1989). Urate oxidase:
primary structure and evolutionary implications. *Proc. Natl. Acad. Sci. U.S.A.*
86,9412-9416.

Xu, X., Li, C., Zhou, P. and Jiang, T. (2016). Uric acid transporters hiding in the
intestine. *Pharm Biol* **54**,3151-3155.

Yano, H., Tamura, Y., Kobayashi, K., Tanemoto, M. and Uchida, S. (2014). Uric
acid transporter ABCG2 is increased in the intestine of the 5/6 nephrectomy
rat model of chronic kidney disease. *Clin. Exp. Nephrol.* **18**,50-55.

Yun, Y., Yin, H., Gao, Z., Li, Y., Gao, T., Duan, J., Yang, R., Dong, X., Zhang, L.
and Duan, W. (2017). Intestinal tract is an important organ for lowering
serum uric acid in rats. *PLoS ONE* **12**,e0190194.

Zhang, M., Gao, Y., Wang, X., Liu, W., Zhang, Y. and Huang, G. (2016).
Comparison of the effect of high fruit and soybean products diet and standard
diet interventions on serum uric acid in asymptomatic hyperuricemia adults:
an open randomized controlled trial. *Int J Food Sci Nutr* **67**,335-343.

    Figure Legends

Fig. 1: *BCRP* and *MRP4* **expression in the liver, kidneys, jejunum, and ileum of**

normal chickens.

A and B: Real-time quantitative PCR of *BCRP* and *MRP4* in the liver, kidneys,

jejunum, and ileum of normal chicken (N = 5); compared with the liver, $^*P<0.05$,

$^{**}P<0.01$. C: The relative expression of *BCRP* and *MRP4* in the liver, kidneys,

jejunum, and ileum of normal chicken (N = 5). The relative expression levels were

normalized to 18S expression levels. The expression levels of *BCRP* in the liver were

set to 1. Compared with *BCRP* in the liver, $^*P<0.05$, $^{**}P<0.01$; compared with MRP4

in the liver, $^\#P<0.05$, $^{\#\#}P<0.01$. D: BCRP and MRP4 protein expression in the liver,

kidneys, jejunum, and ileum of normal chickens by western blotting (N = 3).

Compared with the liver, $^*P<0.05$, $^{**}P<0.01$. All data are means ± s.e.m.

Fig. 2: **BCRP protein expression in the liver, kidneys, jejunum, and ileum of**

normal chickens, as determined by immunohistochemistry.

A and B: Liver; C and D: Kidneys; E and F: Jejunum; G and H: Ileum. A, C, E, F, G,

and H: scale bar = 100 μm; B and D: scale bar = 20 μm.

Fig. 3: **MRP4 protein expression in the liver, kidney, jejunum, and ileum of**

normal chickens, as determined by immunohistochemistry.

A and B: Liver; C and D: Kidney; E and F: Jejunum; G and H: Ileum. A, C, E, F, G,

and H: scale bar = 100 μm; B and D: scale bar = 20 μm.

Fig. 4: **Relative BCRP and MRP4 mRNA levels in the liver, kidneys, jejunum,**

and ileum of normal chickens, as determined by real-time quantitative PCR.

Compared with the NC group, $^*P<0.05$, $^{**}P<0.01$. NC: normal control group; SD:

sulfonamide group; FM: fish meal group; IU: injection uric acid group. A and B:

BCRP and *MRP4* mRNA expression in the liver; C and D: *BCRP* and *MRP4* mRNA

expression in the kidney; E and F: *BCRP* and *MRP4* mRNA expression in the jejunum;

G and H: *BCRP* and *MRP4* mRNA expression in the ileum. All data are means ± s.e.m;

N = 5 samples per treatment.

Fig. 5: **BCRP and MRP4 protein expression in the liver, kidneys, jejunum, and ileum in each group of chickensas, as determined by western blotting.**

Compared with the NC group, $^*P<0.05$, $^{**}P<0.01$. NC: normal control group; SD: sulfonamide group; FM: fish meal group; IU: injection uric acid group. A: Protein expression of BCRP and MRP4 in the liver; B: Protein expression of BCRP and MRP4 in the kidney; C: Protein expression of BCRP and MRP4 in the jejunum; D: Protein expression of BCRP and MRP4 in the ileum. All data are means ± s.e.m; N = 3 samples per treatment.

Fig. 6. BCRP protein expression in the liver, kidneys, jejunum, and ileum in each
group of chickens based on immunohistochemistry.

NC: normal control group; SD: sulfonamide group; FM: fish meal group; IU: injection uric acid group. A to H: scale bar = 20 μm; I to P: scale bar = 200 μm.

Fig. 7. MRP4 protein expression in the liver, kidney, jejunum, and ileum in each

group of chickens based on immunohistochemistry.

SD: sulfonamide group; FM: fish meal group; IU: injection uric acid group. A to H:

scale bar = 20 μm; I to P: scale bar = 200 μm.

498 **Tables**

499 **Table 1. Real-Time Quantitative PCR Primer Sequence**

Gene	Primer Sequence (5'–3')	Length	Accession No.
BCRP-F	CAGCAAGCAAGGAAGATCAC		
		129 bp	NM_001328490.1
BCRP-R	GGCTGGAGTTGAGATACTTC		
MRP4-F	TAGTGTTGGTCAGAGACAGC		
		167 bp	NM_001030819.1
MRP4-R	GTGCAATGGTCAGAACTGTG		
18S-F	CGGCGACGACCCATTCGAAC		
		99bp	M_59389.1
18S-R	GAATCGAACCCTGATTCCCCGTC		

500 **Table 2. Serum uric acid levels and renal function in each treatment group**

Groups	Uric (μmol/L)	Creatinine (μmol/L)	Bun (mmol/L)
NC	115.7±7.3	2.79±0.14	0.22±0.01
SD	141.7±6.2[*]	3.66±0.24[**]	0.33±0.03[**]
FM	112.9±7.8	2.97±0.15	0.22±0.02
IU	152.9±6.1[**]	3.79±0.19[**]	0.28±0.01[**]

501 Compared with the NC group, [*]$P<0.05$, [**]$P<0.01$. NC: normal control group; SD:

502 sulfonamide group; FM: fish meal group; IU: injection uric acid group. All data are

503 means ± s.e.m; N = 10 samples per treatment.