Dynamic expression of Id3 defines the stepwise differentiation of tissue-resident
regulatory T cells

Jenna M. Sullivan[1,2], Barbara Höllbacher[1] and Daniel J. Campbell[1,2]*

[1]Immunology Program, Benaroya Research Institute, Seattle, WA 98101

[2]Department of Immunology, University of Washington School of Medicine, Seattle, WA 98195

*Corresponding Author: Daniel J. Campbell, Benaroya Research Institute, 1201 Ninth Avenue,

Seattle, WA 98101-2795. E-mail address: campbell@benaroyaresearch.org

14 Running Title: Dynamic Id3 expression in development of tissue Tregs

17 Funding: This work was supported by grants to DJC from the National Institutes of Health

18 (AI067750, AI124693). JMS was supported by NIH-NIAID T32 (AI106677, UW Immunology).

<u>Abstract</u>

Foxp3$^+$ regulatory T (T$_R$) cells are phenotypically and functionally diverse, and broadly distributed in lymphoid and non-lymphoid tissues. However, the pathways guiding the differentiation of tissue-resident T$_R$ populations have not been well defined. By regulating E-protein function, Id3 controls the differentiation of CD8$^+$ effector T cells and is essential for T$_R$ maintenance and function. We show that dynamic expression of Id3 helps define three distinct T$_R$ populations, Id3$^+$CD62L^{hi}CD44^{lo} central (c)T$_R$, Id3$^+$CD62L^{lo}CD44^{hi} effector (e)T$_R$ and Id3$^-$ eT$_R$. Adoptive transfer experiments and transcriptome analyses support a stepwise model of differentiation from Id3$^+$ cT$_R$ to Id3$^+$ eT$_R$ to Id3$^-$ eT$_R$. Furthermore, Id3$^-$ eT$_R$ have high expression of functional inhibitory markers and a transcriptional signature of tissue-resident T$_R$. Accordingly, Id3$^-$ eT$_R$ are highly enriched in non-lymphoid organs, but virtually absent from blood and lymph. Thus, we propose that tissue-resident T$_R$ develop in a multi-step process associated with Id3 downregulation.

<u>Introduction</u>

Several recent studies have highlighted the phenotypic and functional heterogeneity of

regulatory T (T$_R$) cells during both steady state and inflammation (1-4). We and others have

shown that at steady state in lymphoid organs T$_R$ can be broadly divided by expression of CD44

and CD62L into distinct subsets which differ in their localization, dependence on IL-2, and

extent of PI3K signaling (2, 5, 6). Moreover, CD44^{hi}CD62L^{lo} effector (e)T$_R$ display diverse

expression of transcription factors and chemokine receptors that promote their migration to

inflamed tissues and their response to different types of inflammatory signals (1, 7). Accordingly,

T$_R$ found in nonlymphoid tissues have a distinct molecular profile that includes high expression

of Gata3 and ST2 (the IL-33R), and are functionally equipped to suppress inflammation at

barrier sites (8, 9). Although these data highlight the anatomical, functional and molecular

diversity of T$_R$, the pathways by which these T$_R$ populations differentiate have not been

completely defined.

The inhibitors of DNA binding (Id) proteins have been extensively studied in lymphocyte

development (10, 11). Studies of CD8$^+$ effector T cells revealed that Id2 and Id3 are powerful

transcriptional regulators of differentiation that are dynamically regulated during T cell activation

and effector/memory T cell differentiation (12, 13). Through their regulation of E protein function,

Id2 and Id3 help to control expression of genes essential for CD8$^+$ effector cell differentiation

and survival such as *Tcf7*, *Tbx21*, *Bcl2* and *Klrg1* (14, 15). Although less well studied, Id

proteins have been shown to have essential roles in CD4$^+$ T cell function. For instance, Id2 and

Id3 are essential for T$_R$ maintenance and function, with T$_R$ lacking both Id2 and Id3 having

impaired proliferation and survival (16). In T$_R$, Id3 helps to stabilize Foxp3 through restriction of

the E protein E47 and its downstream targets Spi-B and SOCS3 (17). However, Id3 expression

is not uniform in T$_R$, and distinct populations of Id3$^+$ and Id3$^-$ have been identified (16, 18). In

this study, we show that Id3 is dynamically regulated in T$_R$, and that progressive loss of Id3

correlates with the stepwise differentiation of a highly-functional T_R population localized primarily

in non-lymphoid tissues.

<u>Materials and Methods</u>

<u>Mice</u>

C57BL/6, RAG1-deficient and Foxp3-mRFP mice were purchased from The Jackson

Laboratory. Id3-GFP mice were a gift from Ananda Goldrath (UCSD, La Jolla, California) and

have been previously described (12, 14). Mice were bred and housed under the approval of the

Institutional Animal Care and Use Committee of the Benaroya Research Institute.

<u>Cell Isolation</u>

Unless noted below, single cell suspensions isolated from tissues using manual disruption. PEC

isolated by injecting sterile PBS into peritoneal cavity of euthanized mice, gentle disruption to

dislodge cells and collection of injected PBS. IEL and LPL were isolated from pooled large and

small intestine as previous described (19). Lymphocytes further purified by resuspension in 44%

Percoll^{TM} (GE Healthcare) layered over 67% Percoll^{TM} and spun at 2,800rpm for 20 mins. Lung

and fat were finely minced, digested with 0.26U/mL Liberase TM (Roche) and 10U/mL DNAse

(Sigma) for 1 hr at 37°C and filtered. For skin tissue, ears were processed as above with

0.14U/mL Liberase TM and 10U/mL DNAse. For lymph collection, mice were fed 20mL/kg 'Half

and Half' by oral gavage and sacrificed 2-3 hrs later. Lymph collected from the cisterna chyli

was directly stained for flow cytometry.

<u>Flow cytometry</u>

Single cells suspensions were stained with fixable Viability Dye eFluor 780 (eBiosciences) in

PBS for 10 min at RT. Cells were stained with directly conjugated Abs in PBS with 0.5% BCS

for 20 min at 4°C. Abs purchased from BioLegend: CD4 (RM4-5), TCRβ (H57-597), CD44 (IM7),

CD62L (MEL-14), CD25 (PC61), ICOS (C398.4A), TIGIT (1G9), CTLA4 (UC10-4B9), GITR

(DTA.1), CD69 (53-7.3). Abs purchased from eBiosciences: KLRG1 (2F1) and CD103 (2E7).

Intracellular stains were performed using a FixPerm Kit (eBiosciences). Data acquired on an

LSR II (BD Biosciences) and analyzed using FlowJo software (TreeStar).

In vitro assays

CD4$^+$ T cells isolated from spleen and LN using CD4 microbeads (Miltenyi). 1x10^6 T cells

cultured with platebound α-CD3 (2C11) and α-CD28 (37.51) from BioXcell at 1μg/mL each for

48 or 66 hrs. Inhibitors purchased and used as follows: ZSTK474 (1μM, Sigma), Rapamycin

(10nM, Selleckchem), NFAT inhibitor (10μM, Tocoris), Mek inhibitor PD0325901 (100nM,

Peprotech) and Erk inhibitor FR180204 (10μM, Tocoris). *In vitro* T$_R$ suppression assays were

performed as previous described (20). Chemotaxis assay performed as previously described

(21).

In vivo T$_R$ transfer

Sorted T$_R$ isolated from spleen and LN as described for RNA-seq. 100,000 sorted cells were

injected retro-orbitally into RAG1-deficient hosts. Spleen, LN and blood of recipient mice

collected two wks later and analyzed by flow cytometry.

Statistical Analysis

The *p* values were calculated by Prism software (GraphPad) using either an unpaired Student's

t test or one way ANOVA as indicated. Values less than 0.05 were considered significant.

RNA-seq

CD4$^+$ T cells were isolated using CD4 microbreads (Miltenyi) from either peripheral LNs or

spleens of 3 littermate Id3-GFP x Foxp3-mRFP mice. Cells were sorted based on viability, CD4,

CD44, CD62L, Id3-GFP and Foxp3-mRFP expression on a FACs Aria II (BD Biosciences). 500

cells were sorted directly into SMART-Seq v4 Ultra Low Input RNA Kit (Takara) lysis buffer and

protocol followed to produce cDNA. Library construction was performed using a modified

protocol of the NexteraXT DNA sample preparation kit (Illumina). Dual-index, single-read

sequencing of pooled libraries was run on a HiSeq2500 sequencer (Illumina) with 58-base

reads and an average depth of 8.6Mio reads per library. Base-calling and demultiplexing were

performed automatically on BaseSpace (Illumina) to generate FASTQ files.

Results/Discussion

Id3 is dynamically expressed in T_R and regulated by TCR signaling

To examine Id3 expression in T_R we generated Id3-GFP x Foxp3-mRFP double reporter

mice. In agreement with previously published reports, most CD4$^+$ T cells in spleen and lymph

nodes (LNs) were Id3$^+$ (16, 18). However, we noted a subset of Id3$^-$ cells in both Foxp3$^+$ T_R and

Foxp3$^-$ conventional CD4$^+$ T cell populations (Fig 1A). Within T_R, the Id3$^-$ population fell

exclusively within the CD44^{hi} CD62L^{lo} eT_R compartment (5), whereas Id3$^+$ T_R were found in both

the CD44^{lo} CD62L^{hi} central (c)T_R and eT_R compartments (Fig 1B). Thus, in secondary lymphoid

organs (SLOs) T_R can be divided into three distinct subsets based on Id3, CD62L and CD44

expression, Id3$^+$ cT_R, Id3$^+$ eT_R or Id3$^-$ eT_R. Within the Id3$^+$ T_R populations Id3$^+$ cT_R had higher

Id3 expression than Id3$^+$ eT_R measured by GFP-MFI (Fig 1B), leading us to hypothesize that T_R

may downregulate Id3 as they transition during their activation and differentiation from cT_R into

eT_R. Indeed, Id3 expression can be downregulated by TCR signaling (16), and we observed a

loss of Id3 expression in the T_R compartment correlating with the strength of TCR stimulation

when cells were activated with different concentrations of platebound αCD3/28 (Sup Fig 1).

Furthermore, inhibiting either the MAP-kinase/Erk or PI3-kinase/mTOR signaling pathways

blocked Id3 downregulation in T_R (Fig 1C), consistent with reports that eT_R development

requires TCR stimulation, mTOR signaling and the PI3-kinase-dependent inactivation of the

transcription factor Foxo1 (5, 6, 22).

To more precisely define the developmental relationship between these T_R subsets we

utilized an adaptive transfer model in which T_R stimulation and expansion depends on

TCR:MHCII interactions (23). For this we sorted Id3$^+$ cT_R, Id3$^+$ eT_R or Id3$^-$ eT_R from spleen and

LNs of reporter mice and transferred individual T_R populations into RAG1-deficient animals, and

evaluated the phenotype and expansion of transferred T_R after 2 weeks. Importantly, we did not

observe any difference in the extent of Foxp3 expression between the T_R populations upon their

recovery, which varied between ~30-80% in different experiments (not shown). Transferred Id3$^+$

cT$_R$ gave rise to all three subsets, with some cells retaining Id3 and CD62L expression but the

majority converting into Id3$^-$ eT$_R$ (Fig 1D). The bulk of Id3$^+$ eT$_R$ downregulated Id3, with no cells

regaining CD62L expression, whereas Id3$^-$ eT$_R$ did not give rise to either of the other

populations, indicating that these cells are likely a terminal differentiated population. Thus, Id3$^-$

eT$_R$ appear to develop from Id3$^+$ cT$_R$ in a stepwise manner during activation, with Id3$^+$ eT$_R$

acting as an intermediate population.

<u>Id3$^-$ eT$_R$ express inhibitory markers and are highly suppressive</u>

To explore the phenotypic and functional differences between these three subsets of T$_R$,

we assessed expression of the T$_R$-associated surface markers on each population of splenic T$_R$.

In agreement with previously published data, cT$_R$ and eT$_R$ showed distinct phenotypes, with eT$_R$

having lower expression of the high affinity IL-2 receptor component CD25, but higher levels of

the activation and functional surface markers ICOS, KLRG1, TIGIT, GITR and CTLA4 (Fig 2A)

(5, 6). Moreover, within the eT$_R$ compartment the Id3$^-$ T$_R$ had higher expression of activation and

inhibitory molecules than their Id3$^+$ T$_R$ counterparts, but had the lowest expression of CD25. As

our lab previously described (5), elevated expression of ICOS and diminished expression of

CD25 suggests that Id3$^-$ eT$_R$ are less dependent on IL-2 for their homeostatic maintenance, and

instead likely rely on continued ICOS signaling. Additionally, increased expression of these T$_R$

functional molecules correlated with enhanced *in vitro* suppressive activity of Id3$^-$ eT$_R$ compared

with either Id3$^+$ cT$_R$ or Id3$^+$ eT$_R$ (Fig 2B).

<u>Transcriptional profiling highlights the stepwise differentiation of Id3$^-$ eT$_R$</u>

To identify and compare their unique transcriptional profiles, we performed RNA-seq on

sorted Id3$^+$ cT$_R$, Id3$^+$ eT$_R$ and Id3$^-$ eT$_R$ from spleen or LNs of Id3-GFP x Foxp3-mRFP reporter

mice. Although there was little difference between LN and spleen samples, principle component

analysis (PCA) showed that each of the three T$_R$ populations were transcriptionally distinct (Fig

3A), and accordingly we identified 1,672 significantly differentially expressed (DE) genes

between the three T_R populations (Fig 3B, Supplemental Table 1). The largest differences were

found between Id3$^-$ eT$_R$ and Id3$^+$ cT$_R$ with 1,471 DE genes, whereas only 474 genes differed

between Id3$^+$ and Id3$^-$ eT$_R$ (Fig 3B). Interestingly among the DE genes we observed a reciprocal

increase in Id2 expression as T_R lose Id3 (Fig 3C). This differential Id expression in T_R is similar

to what occurs in CD8$^+$ T cells, which upregulate Id2 and downregulate Id3 while becoming

activated and gaining effector function (12, 13). This suggests that as T_R downregulate Id3, the

closely related Id2 may take over some of its functions while also driving a unique E-protein-

dependent signature promoting eT$_R$ development. Consistent with the stepwise differentiation

model we propose for these populations, examination of the 300 most DE genes across the

three T_R subsets (based on highest F-value) showed that both up and downregulated genes

were generally expressed in a gradient fashion, with expression in Id3$^+$ eT$_R$ falling between that

of Id3$^+$ cT$_R$ and Id3$^-$ eT$_R$ (Fig 3D-E). Moreover, in accordance with our prior phenotypic analysis

and their enhanced suppressive activity, Id3$^-$ eT$_R$ showed elevated expression of known T_R

function genes, including *Il10, Ctla4, Pdcd1, Tnfrsf4, Lag3 and Ebi3* (Fig 3F). To further validate

our RNA-seq results, we confirmed differential expression of several T_R-associated genes by

flow cytometry (Supplemental Fig 2). Gene Ontology (GO) term enrichment analysis of genes

DE between Id3$^+$ eT$_R$ and Id3$^-$ eT$_R$ identified specific molecular pathways altered between these

closely related populations (Fig 3G). The top six enriched categories all related to cytokine and

chemokine receptor signaling, with Id3$^-$ eT$_R$ expressing high levels of receptors indicating that

they can tune their activity in response to key inflammatory cytokines such as IL-1, IL-18, IL-23,

and IL-25 (Fig 3H). Moreover, among the DE chemokine receptors, Id3$^-$ eT$_R$ had the highest

expression and subsequent responsiveness to chemokines that promote lymphocyte migration

to inflamed tissues, such as *Ccr4 and Cxcr3* (Fig 3H-I) (1) . Thus, Id3$^-$ eT$_R$ have a unique

molecular profile indicative of their development from Id3$^+$ T_R precursors, their elevated

suppressive function and altered migratory capacity.

<u>Id3$^-$ T_R are enriched and resident in non-lymphoid tissues</u>

In contrast to their elevated expression of inflammatory chemokine receptors, Id3$^-$ eT$_R$

had the lowest expression of *Ccr7* and *S1pr1* which function together to promote T$_R$

recirculation through SLOs (24) (Fig 3H). Additionally, expression of CD103 and CD69, which

together act to retain tissue-resident memory T($_{RM}$) cells in non-lymphoid sites, was strongly

enriched in Id3$^-$ eT$_R$, suggesting these cells may be tissue-resident (Fig 4A) (25, 26). Indeed,

utilizing published gene signatures of CD8$^+$ T$_{RM}$ or circulating memory T cells (27), we found

that the CD8$^+$ T$_{RM}$ signature gene set was enriched in Id3$^-$ eT$_R$ compared to either Id3$^+$ cT$_R$ or

Id3$^+$ eT$_R$, whereas the circulating memory gene set was enriched in the Id3$^+$ T$_R$ populations (Fig

4B). Several groups have recently identified the ST2 (IL-33R)-Gata3 axis as a key determinant

of T$_R$ residency and function in non-lymphoid tissues (8, 9, 28). Accordingly, Id3$^-$ eT$_R$ had the

highest expression of all positively associated tissue T$_R$ genes such as *Il1rl1* (ST2), *Gata3*,

Areg, Irf4 and Rora, whereas Id3$^+$ cT$_R$ had the lowest expression of these genes but high

expression of negative regulators of tissue residence such as *Tcf7, Klf2* and *Lef1,* and Id3$^+$ eT$_R$

displayed intermediate expression for all of these genes (Fig 4C). Consistent with their T$_{RM}$-like

transcriptional signature, Id3$^-$ eT$_R$ were a minority of T$_R$ in LNs and spleen, but their frequency

dramatically increased in nonlymphoid tissues, where they highly expressed the T$_{RM}$ surface

markers CD103 and CD69 (Fig 4D-F). Indeed, tissues such as the fat and skin contained almost

exclusively Id3$^-$ eT$_R$. Of particular note Id3$^-$ eT$_R$ were rarest in the lymph and blood, indicating

that Id3$^-$ eT$_R$ do not actively recirculate, but instead are retained as tissue-resident cells in non-

lymphoid organs.

Despite significant interest in tissue-resident T$_R$, the mechanisms regulating their

differentiation and distribution are still poorly defined. Our data show that T$_R$ can be subdivided

based on Id3 expression and known markers of cT$_R$ and eT$_R$ into distinct populations, and

together our phenotypic, functional, transcriptional and transfer analyses strongly support a

stepwise differentiation model in which T$_R$ downregulate Id3 as they progressively gain effector

function, tissue homing capacity and residency in non-lymphoid organs. Similarly, Li *et al.*

recently proposed a stepwise model for the differentiation of T_R in adipose tissue in which

activation in the spleen allowed T_R to migrate into the adipose tissue, where IL-33 signaling

drove their terminal differentiation and functional specialization (28). High expression of co-

stimulatory receptors such as ICOS and receptors for inflammatory cytokines would also allow

Id3$^-$ eT_R to respond to inflammatory signals that enhance Foxp3-mediated transcriptional

repression and promote eT_R differentiation and function (29), in part through activation of

mTORC1 signaling (22). Although the direct role of Id3 downregulation in the functional

differentiation of T_R is still not established, our data highlight the complexity of tissue T_R

development, and identify novel molecular pathways that are modulated during their stepwise

differentiation.

Acknowledgements:

We thank A. Goldrath for Id3-GFP mice. K. Arumuganathan and T. Nguyen for help with flow

cytometry and cell sorting. V. Gersuk and the BRI Genomics Core for running RNA-seq

samples. And members of the Campbell lab for helpful discussions.

References

1. Campbell, D. J. 2015. Control of Regulatory T Cell Migration, Function, and Homeostasis. *J Immunol* 195: 2507-2513.

2. S., S. K., S. Shivani, S. J. Michael, and C. D. J. 2014. Regulatory T-cell homeostasis: steady-state maintenance and modulation during inflammation. *Immunological Reviews* 259: 40-59.

3. Dominguez-Villar, M., and D. A. Hafler. 2018. Regulatory T cells in autoimmune disease. *Nat Immunol*.

4. Josefowicz, S. Z., L. F. Lu, and A. Y. Rudensky. 2012. Regulatory T cells: mechanisms of differentiation and function. *Annu Rev Immunol* 30: 531-564.

5. Smigiel, K. S., E. Richards, S. Srivastava, K. R. Thomas, J. C. Dudda, K. D. Klonowski, and D. J. Campbell. 2014. CCR7 provides localized access to IL-2 and defines homeostatically distinct regulatory T cell subsets. *J Exp Med* 211: 121-136.

6. Luo, C. T., W. Liao, S. Dadi, A. Toure, and M. O. Li. 2016. Graded Foxo1 activity in Treg cells differentiates tumour immunity from spontaneous autoimmunity. *Nature* 529: 532-536.

7. Gratz, I. K., and D. J. Campbell. 2014. Organ-specific and memory treg cells: specificity, development, function, and maintenance. *Front Immunol* 5: 333.

8. Delacher, M., C. D. Imbusch, D. Weichenhan, A. Breiling, A. Hotz-Wagenblatt, U. Trager, A. C. Hofer, D. Kagebein, Q. Wang, F. Frauhammer, J. P. Mallm, K. Bauer, C. Herrmann, P. A. Lang, B. Brors, C. Plass, and M. Feuerer. 2017. Genome-wide DNA-methylation landscape defines specialization of regulatory T cells in tissues. *Nat Immunol* 18: 1160-1172.

9. Wohlfert, E. A., J. R. Grainger, N. Bouladoux, J. E. Konkel, G. Oldenhove, C. H. Ribeiro, J. A. Hall, R. Yagi, S. Naik, R. Bhairavabhotla, W. E. Paul, R. Bosselut, G. Wei, K. Zhao, M. Oukka, J. Zhu, and Y. Belkaid. 2011. GATA3 controls Foxp3(+) regulatory T cell fate during inflammation in mice. *J Clin Invest* 121: 4503-4515.

10. Kee, B. L. 2009. E and ID proteins branch out. *Nat Rev Immunol* 9: 175-184.

11. Quong, M. W., W. J. Romanow, and C. Murre. 2002. E protein function in lymphocyte development. *Annu Rev Immunol* 20: 301-322.

12. Yang, C. Y., J. A. Best, J. Knell, E. Yang, A. D. Sheridan, A. K. Jesionek, H. S. Li, R. R. Rivera, K. C. Lind, L. M. D'Cruz, S. S. Watowich, C. Murre, and A. W. Goldrath. 2011. The transcriptional regulators Id2 and Id3 control the formation of distinct memory CD8+ T cell subsets. *Nat Immunol* 12: 1221-1229.

13. Omilusik, K. D., L. A. Shaw, and A. W. Goldrath. 2013. Remembering one's ID/E-ntity: E/ID protein regulation of T cell memory. *Curr Opin Immunol* 25: 660-666.

14. Miyazaki, M., R. R. Rivera, K. Miyazaki, Y. C. Lin, Y. Agata, and C. Murre. 2011. The opposing roles of the transcription factor E2A and its antagonist Id3 that orchestrate and enforce the naive fate of T cells. *Nat Immunol* 12: 992-1001.

15. Omilusik, K. D., M. S. Nadjsombati, L. A. Shaw, B. Yu, J. J. Milner, and A. W. Goldrath. 2018. Sustained Id2 regulation of E proteins is required for terminal differentiation of effector CD8(+) T cells. *J Exp Med* 215: 773-783.

16. Miyazaki, M., K. Miyazaki, S. Chen, M. Itoi, M. Miller, L. F. Lu, N. Varki, A. N. Chang, D. H. Broide, and C. Murre. 2014. Id2 and Id3 maintain the regulatory T cell pool to suppress inflammatory disease. *Nat Immunol* 15: 767-776.

17. Rauch, K. S., M. Hils, E. Lupar, S. Minguet, M. Sigvardsson, M. E. Rottenberg, A. Izcue, C. Schachtrup, and K. Schachtrup. 2016. Id3 Maintains Foxp3 Expression in Regulatory T Cells by Controlling a Transcriptional Network of E47, Spi-B, and SOCS3. *Cell Rep* 17: 2827-2836.

18. Rauch, K. S., M. Hils, A. J. Menner, M. Sigvardsson, S. Minguet, P. Aichele, C. Schachtrup, and K.
Schachtrup. 2017. Regulatory T cells characterized by low Id3 expression are highly suppressive
and accumulate during chronic infection. *Oncotarget* 8: 102835-102851.
19. Couter, C. J., and N. K. Surana. 2016. Isolation and Flow Cytometric Characterization of Murine
Small Intestinal Lymphocytes. *JoVE*: e54114.
20. Srivastava, S., M. A. Koch, M. Pepper, and D. J. Campbell. 2014. Type I interferons directly inhibit
regulatory T cells to allow optimal antiviral T cell responses during acute LCMV infection. *J Exp*
Med 211: 961-974.
21. Koch, M. A., G. Tucker-Heard, N. R. Perdue, J. R. Killebrew, K. B. Urdahl, and D. J. Campbell.
2009. The transcription factor T-bet controls regulatory T cell homeostasis and function during
type 1 inflammation. *Nat Immunol* 10: 595-602.
22. Sun, I. H., M. H. Oh, L. Zhao, C. H. Patel, M. L. Arwood, W. Xu, A. J. Tam, R. L. Blosser, J. Wen, and
J. D. Powell. 2018. mTOR Complex 1 Signaling Regulates the Generation and Function of Central
and Effector Foxp3(+) Regulatory T Cells. *J Immunol*.
23. Gavin, M. A., S. R. Clarke, E. Negrou, A. Gallegos, and A. Rudensky. 2002. Homeostasis and
anergy of CD4(+)CD25(+) suppressor T cells in vivo. *Nat Immunol* 3: 33-41.
24. Lee, J. H., S. G. Kang, and C. H. Kim. 2006. FoxP3+ T Cells Undergo Conventional First Switch to
Lymphoid Tissue Homing Receptors in Thymus but Accelerated Second Switch to Nonlymphoid
Tissue Homing Receptors in Secondary Lymphoid Tissues. *The Journal of Immunology* 178: 301-
311.
25. Mackay, L. K., A. Rahimpour, J. Z. Ma, N. Collins, A. T. Stock, M. L. Hafon, J. Vega-Ramos, P.
Lauzurica, S. N. Mueller, T. Stefanovic, D. C. Tscharke, W. R. Heath, M. Inouye, F. R. Carbone, and
T. Gebhardt. 2013. The developmental pathway for CD103(+)CD8+ tissue-resident memory T
cells of skin. *Nat Immunol* 14: 1294-1301.
26. Schenkel, J. M., and D. Masopust. 2014. Tissue-resident memory T cells. *Immunity* 41: 886-897.
27. Milner, J. J., C. Toma, B. Yu, K. Zhang, K. Omilusik, A. T. Phan, D. Wang, A. J. Getzler, T. Nguyen, S.
Crotty, W. Wang, M. E. Pipkin, and A. W. Goldrath. 2017. Runx3 programs CD8(+) T cell
residency in non-lymphoid tissues and tumours. *Nature* 552: 253-257.
28. Li, C., J. R. DiSpirito, D. Zemmour, R. G. Spallanzani, W. Kuswanto, C. Benoist, and D. Mathis.
2018. TCR Transgenic Mice Reveal Stepwise, Multi-site Acquisition of the Distinctive Fat-Treg
Phenotype. *Cell*.
29. Arvey, A., J. van der Veeken, R. M. Samstein, Y. Feng, J. A. Stamatoyannopoulos, and A. Y.
Rudensky. 2014. Inflammation-induced repression of chromatin bound by the transcription
factor Foxp3 in regulatory T cells. *Nat Immunol* 15: 580-587.

Figure Legends

<u>Figure 1: Id3 is dynamically expressed in T_R and regulated by TCR signaling</u>

A) Representative flow cytometry plots of Id3-GFP and Foxp3-RFP expression by gated splenic

$TCR\beta^+$ $CD4^+$ T cells. B) Representative flow cytometry analysis of Id3-GFP expression by

splenic $CD44^{lo}CD62L^{hi}$ cT_R and $CD44^{hi}CD62L^{lo}$ eT_R gated as indicated. (Bottom left) Graphical

analysis of gMFI of Id3-GFP in each of the three gated T_R populations. C) Representative flow

cytometry plots of Id3-GFP expression by gated splenic $TCR\beta^+$ $CD4^+$ $Foxp3^+$ T_R 66 hrs after

stimulation of purified $CD4^+$ T cells in the presence or absence of the idicated inhibitors,

representative of 2 independent expts. D) Representative flow cytometry plots and graphical

analysis of CD44 and Id3-GFP expression by gated $TCR\beta^+CD4^+Foxp3^+$ T_R recovered from the

LNs of RAG1-deficient mice 2 weeks after transfer of the indicated T_R population, summary of 3

independent expts, 3-5 mice per group total. Significance determined by one way ANOVA with

Tukey's post-test for pairwise comparisons, $*p < 0.05$, $**p < 0.01$, $****p < 0.001$

<u>Figure 2: $Id3^-$ eT_R express inhibitory markers and are highly suppressive</u>

A) (Top) Representative flow cytometry histograms. (Bottom) Graphical analysis of expression

of the indicated markers by gated splenic T_R populations. B) (Top) Representative flow

cytometry analysis of CPD dilution by $CD4^+Foxp3^-$ effector T cells stimulated with or without the

indicated T_R populations. (Bottom) Graphical analysis of suppression by each of the indicated

populations, n=3. Significance determined by one way ANOVA with Tukey's post-test for

pairwise comparisons, $*p < 0.05$, $**p < 0.01$, $****p < 0.001$

<u>Figure 3: Transcriptional profiling highlights the stepwise differentiation of $Id3^-$ eT_R</u>

A) PCA of RNA-seq data from LN and splenic $Id3^+$ cT_R, $Id3^+$ eT_R and $Id3^-$ eT_R populations sorted

from three individual mice. B) Bar graphs showing the number of differentially expressed genes

(adj.p.value < 0.05 and log2FC >1) for each of the indicated pairwise comparisons. C) Graphical

analysis of normalized transcript reads for Id2 or Id3 from RNA-seq data. D) The 300 most

350 differentially expressed genes (determined by F value) were split into the upregulated (red) and

351 downregulated (blue) fractions based expression in Id3⁻ eT$_R$ vs Id3⁺ cT$_R$. Graph shows the mean

352 log2FC compared to Id3⁺ cT$_R$ for both eT$_R$ populations. Error bars and shaded area represent 1x

353 SD. E) Heatmap and hierarchical clustering of splenic RNA-seq samples based on the 300 most

354 variably expressed genes. F) Heatmap and hierarchical clustering of splenic RNA-seq samples

355 based on T$_R$ signature genes identified in reference (8). G) GO term enrichment analysis for DE

356 genes of Id3⁻ eT$_R$ vs. Id3⁺ eT$_R$. Dashed lines represent adjusted p values of 0.05 and 0.01. H)

357 Heatmap and hierarchical clustering of splenic RNA-seq samples based on DE genes found in

358 the top 6 GO functional categories enriched in the comparison of Id3⁺ and Id3⁻ eT$_R$. I) Graphical

359 analysis of chemotaxis assay.

360 <u>Figure 4: Id3⁻ eT$_R$ are enriched and resident in non-lymphoid tissues</u>

361 A) Representative flow cytometry histograms and graphical analysis of expression of the CD69

362 and CD103 by gated splenic T$_R$ populations. B) Enrichment of CD8⁺ T$_{RM}$ (red) or circulating T

363 cell (blue) gene sets along ranked lists of the indicated pairwise comparisons. C) Heatmap and

364 hierarchical clustering of splenic RNA-seq samples based on ST2⁺ tissue T$_R$ associated genes.

365 D) Representative flow cytometry plots of CD103 and CD69 expression by gated TCRβ⁺ CD4⁺

366 Foxp3⁺ T$_R$ in the indicated tissues. E) Graphical analysis of Id3⁻ eT$_R$ frequency among total T$_R$ in

367 various tissues. F) Representative histograms of GFP expression from various tissues of Id3-

368 GFP x Foxp3-mRFP mice, gated on TCRβ⁺ CD4⁺ Foxp3⁺ T$_R$.

369 <u>Supplementary Figure 1: Loss of Id3 expression correlates with strength of TCR signaling</u>

370 Representative flow cytometry plots of Id3-GFP expression by gated splenic TCRβ⁺ CD4⁺

371 Foxp3⁺ T$_R$ 48 hours after stimulation of purified CD4⁺ T cells with 1µg/mL α-CD28 and indicated

372 titration of α-CD3, summary of 2 independent expts.

373 <u>Supplementary Figure 2: Flow cytometry validation of RNA-seq targets</u>

374   A) Representative flow cytometry histograms. B) Graphical analysis of expression of the

375   indicated markers by gated splenic T_R populations. Significance determined by one way ANOVA

376   with Tukey's post-test for pairwise comparisons, *p <0.05, **p < 0.01, ****p < 0.001

Id3 is dynamically expressed in T$_R$ and regulated by TCR signaling

Id3⁻eT$_R$ express inhibitory markers and are highly suppressive

Transcriptional profiling highlights the stepwise differentiation of Id3⁻ eT_R

Id3⁻eT_R are enriched and resident in non-lymphoid tissues