1 <u>Multi-Method Characterisation of the Human Circulating Microbiome</u>

3 **E. Whittle**[1], M.O. Leonard[2], R. Harrison[1], T.W. Gant[2] and **D.P Tonge**[1]

5 [1]School of Life Sciences, Faculty of Natural Sciences, Keele University, ST5 5BG [2]
6 Centre for Radiation, Chemical and Environmental Hazards, Public Health
7 England, OX11 0RQ

Abstract

The term microbiome describes the genetic material encoding the various microbial
populations that inhabit our body. Whilst colonisation of various body niches (e.g. the
gut) by dynamic communities of microorganisms is now universally accepted, the
existence of microbial populations in other "classically sterile" locations, including the
blood, is a relatively new concept. The presence of bacteria-specific DNA in the
blood has been reported in the literature for some time, yet the true origin of this is
still the subject of much deliberation. The aim of this study was to provide a
comprehensive description of the human blood microbiome using a range of
complementary molecular and classical microbiological techniques. DNA-level
analyses involved the amplification and sequencing of the 16S rDNA gene. RNA-
level analyses were based upon the *de novo* assembly of unmapped mRNA reads.
Molecular studies were complemented by viability data from classical aerobic and
anaerobic microbial culture experiments. At the phylum level, the blood microbiome
was predominated by Proteobacteria, Actinobacteria, Firmicutes and Bacteroidetes.
The key phyla detected were consistent irrespective of molecular method (DNA vs
RNA), and consistent with the results of other published studies. *In silico* comparison
of our data with that of the Human Microbiome Project revealed that members of the
blood microbiome were most likely to have originated from the oral or skin
communities. To our surprise, aerobic and anaerobic cultures were positive in eight
of out the ten donor samples investigated, and we reflect upon their source. Our data
provide further evidence of a core blood microbiome, and provide insight into the
potential source of the bacterial DNA / RNA detected in the blood. Further, data
reveal the importance of robust experimental procedures, and identify areas for
future consideration.

Background

The term "microbiome" describes the genetic material encoding the various microbial
populations that inhabit our body. In contrast, the term "microbiota" refers to the
viable organisms that comprise these communities. Cell for cell, bacterial cells
outnumber our own cells by a factor of ten, and the bacterial microbiome encodes for
one hundred-fold more genes. The microbiome undertakes essential biological
processes and thus it is unsurprising that a number of disease states are associated
with changes in microbiome composition, termed "dysbiosis". Whilst the colonisation
of specific body sites in contact with the external environment (such as the
gastrointestinal tract, skin and vagina) by microorganisms is both well-described and
universally accepted [1], the existence of microbial populations in other "classically
sterile" locations, including the blood, is a relatively new concept.

The presence of bacteria-specific DNA in the blood has been reported in the literature for some time, yet the true origin of this is still the subject of much deliberation. Mounting evidence supports the existence of a blood microbiome (*specifically, the presence of bacterial genetic material*) in man [2-11] and various other species including rodents, cats, chickens and cows [12-15] as determined by amplification and sequencing of the bacterial 16S rRNA gene, or via whole genome sequencing. Such studies report the existence of bacteria-derived genetic material (DNA) within the circulation, but do not provide evidence for the presence of viable organisms.

Convention tells us that the blood is sterile in health, and bacteraemia, even at 1 – 10 bacterial cells per millilitre whole blood, is potentially life threatening. Despite this, several studies have presented evidence of bacteria or bacteria-like structures within the circulation in the absence of overt disease. It should be noted however that Martel and colleagues report that the bacteria-like particles often described following a range of imaging techniques represent non-living membrane vesicles and protein aggregates derived from the blood itself [16]. McLaughlin surveyed the blood of 25 healthy donors and observed the presence of pleomorphic bacteria using dark-field microscopy, electron microscopy, polymerase chain reaction and fluorescent *in situ* hybridisation, in all samples analysed [17]. Further, Potgieter and colleagues described the presence of blood-cell associated bacteria in a range of blood preparations using electron microscopic techniques [8]. Significantly, Damgaard and colleagues found viable (*culturable*) bacteria in 62% of blood donations from donors with no overt disease [18]. This finding is plausible given that various daily activities including chewing, tooth brushing and flossing result in the translocation of oral bacteria into the bloodstream [19-22] however one would expect such organisms to be rapidly targeted and removed with by the immune system in healthy individuals. Kell, D and colleagues provide a detailed hypothesis for the existence of the blood microbiome [4,8] and suggest that it is likely composed of organisms (or parts thereof) that translocate to the circulation from their usual place of habitation (classical niches such as the gastrointestinal tract, oral cavity, skin, vagina) – a process termed atopobiosis. They further describe how this could occur via well-described physiological processes including dendritic cells processes, via micro-fold cells, and in disease, via dysfunctional epithelial junctions. This explanation is supported by studies demonstrating a correlation between gut microbiota dysbiosis and altered microbial signatures detected in the blood [23-25], suggesting that the observed disease-associated blood microbiota is a consequence of increased bacterial translocation across the gut barrier. Furthermore, characterisation of the microbial populations in the coronary artery tissues by Lehtiniemi and colleagues [26] identified known members of the oral microbiota, suggesting that bacteria had translocated from the oral cavities into the bloodstream, potentially as a result of damage caused by tooth brushing or by leakage across the mucosal surfaces.

Various disease states are associated with blood microbiome dysbiosis [2,6,24,25,27], and these changes are likely reflective of dysbiosis at a distant site(s) with well-characterised microbial communities, and the result of subsequent translocation. Limited evidence also suggests that these changes may be disease-specific; Alzheimer's disease for example, has been associated with the detection of mostly coccus microbes, whilst Parkinson's disease has been associated with both coccus

98 and bacillus bacteria [8]. Such changes are of significant scientific and medical interest
99 as they offer opportunities for biomarker and therapeutic target development [25,27].
100
Due to the long-held belief that the bloodstream of healthy individuals is sterile and
since the blood is an unfavourable compartment for the microbes due to its
bacteriostatic and bactericidal components [1,13], it is of principal significance to
understand *whether* and *how* bacteria may persist in it. Accidental contamination
during the collection of blood and or during downstream experimental procedures
has been proposed as an alternative explanation for the existence of the blood
microbiome *per se*. We support this explanation for the detection of *viable* bacteria
within the bloodstream of healthy individuals, however, suggest that this
phenomenon does not adequately explain the existence of the blood microbiome
(*the presence of bacteria-derived nucleic acids*) when one considers the number of
studies that demonstrate significant and apparently disease-specific differences in
the composition of the blood microbiome. Moreover, examination of the bacterial
taxa reported in these studies reveal similar blood microbiota compositions across
the different studies, whereby Proteobacteria dominate (relative abundance values
typically ranging from 85%-90%), and Firmicutes, Actinobacteria, and Bacteroidetes
present to a lesser extent [2,7,25,28]. This suggests the existence of a core blood
microbiome profile that persists independent of study environment or analytical
methodology.
119
Using a range of complementary molecular and classical molecular biology
techniques, here we characterise the human blood microbiome in unparalleled
detail; at the DNA level, we amplify and sequence the 16S rDNA gene, at the
RNA level, we assemble almost 500,000,000 unmapped mRNA reads and map
these to known taxa, and finally, we present viability data from classical
aerobic and anaerobic microbial culture experiments. Our experimental
approach is detailed in Figure 1.
127

128
129
130
131 **Figure 1** – Schematic representation of the multiple-method circulating microbiome
132 characterisation approach implemented herein. NB: Biomarker and mechanistic
133 data are not included within the scope of this publication.
134
135

136 **Materials and Methods**
137
Donor Samples
Atopic asthmatic individuals (n=5) with physician-diagnosed house dust mite allergy,
and gender and age-matched healthy control subjects (n=5) were recruited to the
study via SeraLabs Limited in accordance with the following criteria **(Table 1)**. Whole
blood was drawn, following alcohol cleansing of the skin surface, into EDTA
containing tubes and stored on ice prior to centrifugation at 1000×g to obtain the
plasma component. All samples were analysed anonymously, and the authors
obtained ethical approval (Keele University ERP3) and written informed consent to
utilise the samples for the research reported herein.
147
148 **Table 1** – Donor population characteristics
149

Patient Criteria
• Have a BMI < 30
• Be non-smokers
• Were diagnosed with atopic asthma during childhood
• Have severe/ poorly controlled symptoms
• Are not on any oral steroid treatment
• Must be allergic to house dust mite
• Must not have diabetes, COPD, or hypertension

150
151
DNA-level: Meta-taxonomic Characterisation
The circulating microbiome was investigated at the DNA level by amplification and
sequencing of the bacterial 16S ribosomal gene using oligonucleotide primers that
target variable region 4 [29] **(Table 2)**. Direct amplification of the V4 region was
performed using the Phusion Blood Direct kit (Thermofisher scientific) alongside a
negative control reaction (in which blood was replaced with molecular biology grade
water) that underwent the complete experimental procedure. Amplification was
performed in triplicate as 20µl reactions containing; 1.0 µl plasma, 10µl 2X Phusion
blood PCR buffer, 0.4µl Phusion Blood II DNA polymerase, 1.0 µl of each forward
and reverse oligonucleotide primer (10µM), and 6.6µl of UV-treated molecular
biology grade water. Cycling parameters were as follows; an initial 5 minutes
denaturation step at 98°C followed by 33 cycles of; denaturation (1 second at 98°C),
annealing (5 seconds at 55°C), and extension (15 seconds at 72°C), and a final
extension at 72°C for 7 minutes.
166
Amplicons resulting from triplicate reactions were combined and purified using the
MinElute PCR purification kit (Qiagen) prior to a further 7 cycles of PCR using

AccuPrime *Pfx* SuperMix and a pair of V4 oligonucleotide primers we developed to
incorporate the Illumina XT adapter in preparation for sequencing (**Table 2**). Cycling
parameters were as follows; initial denaturation for 2 minutes at 95°C followed by 7
cycles of; denaturation (15 seconds at 95°C), annealing (30 seconds at 55°C), and
extension (25 seconds at 68°C), and a final extension at 68°C for 10 minutes. PCR
products were purified using the AMPure XP magnetic beads (Agencourt) at a ratio
of 0.8 beads to sample (v/v), eluted in 20μl molecular biology grade water, and
quantified using the Qubit 3.0 high-sensitivity DNA kit. Amplicons were barcoded
using the Nextera DNA library kit, multiplexed, and sequenced using the Ilumina
MiSeq system with a 250bp paired-end read metric. Bioinformatic analysis was
performed using QIIME[30] implemented as part of the Nephele 16S paired-end QIIME
pipeline using closed reference clustering against the SILVA database [31] at a
sequence identity of 99%. All other parameters remained as default.
182
183 **Table 2** – Oligonucleotide primer sequences

Primer Name	Sequence (5' – 3')	Length
V4_F	GTGCCAGCMGCCGCGGTAA	19
V4_R	GGACTACHVGGGTWTCTAAT	20
V4_XT_F	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGTGCCAGCMGCCGCGGTAA	52
V4_XT_R	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGGACTACHVGGGTWTCTAAT	54

184
185
RNA-level: De-novo Assembly of Unmapped RNA Reads
For the purposes of increasing our understanding of the molecular processes that
are deregulated in patients with atopic asthma, we previously performed whole
transcriptome sequencing on RNA extracted from each donor plasma sample (*data
not included within this manuscript*). Here, we hypothesised that the non-mapping
(likely non-human) reads that often result from such analyses would represent
microbial community members that were present in the blood at the time of RNA
extraction. To this end, RNA reads were mapped to the *Homo sapiens* genome
version hg38 using Tophat with default parameters. Reads failing to map to hg38
were identified from the resulting BAM file and reads, in fastq format, extracted using
bedtools bamtofastq. In order to streamline our strategy, a single de novo
transcriptome was produced by concatenating all of the unmapped reads produced
from the entire study, and assembling these using Trinity [32] to form features
representative of candidate non-human genes. To increase computational efficiency,
the total read population was subsampled to a depth of 1M, 10M, 25M, and 100M,
and the number of reads that these features explained computed by building a
bowtie2 index out of each resulting transcriptome, and mapping the unmapped read
population from each sample back to it. The transcriptome assembly with the highest
mapping rate was used for further analysis as follows; (1) Abundance estimation –
the transcriptome was indexed for bowtie2 and the number of reads mapping to each
feature determined for each of our donor samples using RSEM [33]. (2) A matrix of
expression values was produced using the abundance_estimates_to_matrix.pl script
packaged with Trinity [34]. (3) Statistical analysis - a differential expression analysis
was conducted to identify candidate non-human genes that were significantly
differentially expressed between the disease and disease-free donors using edgeR

[35]. (4) Identification - the taxonomic identity of each assembled feature was
determined using Kraken [36] and visualised using Pavian [37].
213
Classical Culture: Bacterial Viability

Classical microbiological culture, using a range of substrates, was carried out to determine whether the human plasma samples contained any viable bacterial cells, i.e. those capable of proliferation. For each sample, 250µl plasma was inoculated into 9ml of brain heart infusion broth and incubated for 5 days at 37^{o}C. The inoculated broth was plated onto agar plates (Columbia agar + 5% horse blood; CLED medium; A.R.I.A + horse blood) and incubated under either aerobic (Columbia blood agar; CLED medium) or anaerobic (A.R.I.A agar) conditions at 37^{o}C for a minimum of three days. Bacterial growth was evaluated by sight, and a selection of individual colonies from each plate were selected for identification by total 16S gene amplification and Sanger sequencing.

Results and Discussion

Donor Characteristics

The donor population were all female and all "never smokers". The asthmatic population were 39.6 years in age (range 19 – 52) with a mean body mass index of 24.4 (range 21.5 – 27.8) and all had physician-diagnosed atopic asthma resulting from house dust mite allergy. The control population were 39.4 years in age (range 23 – 49) with a mean body mass index of 24.3 (range 21.0 – 26.4) and were disease-free. There were no statistically significant differences in age (P = 0.98) or BMI (P = 0.93) between the two groups **(Table 3)**.

Table 3 – Donor Population Characteristics

Donor	Age (years)	Sex	BMI (kg/m^2)	Smoking History	Diagnosis
BRH1017873	52	F	27.8	Never	Atopic Asthma, HDM
BRH1017874	36	F	27.3	Never	Atopic Asthma, HDM
BRH1017875	42	F	23.3	Never	Atopic Asthma, HDM
BRH1017876	19	F	21.5	Never	Atopic Asthma, HDM
BRH1017877	49	F	22.3	Never	Atopic Asthma, HDM
BRH1017878	49	F	26.4	Never	Healthy
BRH1017879	23	F	21	Never	Healthy
BRH1017880	44	F	26.4	Never	Healthy
BRH1017881	49	F	24.9	Never	Healthy
BRH1017882	32	F	22.7	Never	Healthy

DNA-level Circulating Microbiome – Community Structure

The presence of bacterial DNA within the blood of our study cohort was evaluated by amplification and sequencing of the 16S RNA gene variable region 4. A negative experimental control sample mirrored our study samples through the entire experimental procedure upstream of venepuncture including; PCR, bead-based purification, agarose gel electrophoresis, XT-tagging, library preparation and sequencing. Despite this control reaction being PCR-negative, it was nevertheless submitted for sequencing. Using QIIME[30] implemented as part of Nephele, a total of 243, 853 sequencing reads from the amplified V4 region passed quality assessment (mean = 24,385 reads per sample; range = 10, 742 – 35, 701 reads). The results of our taxonomic classification are shown in **(Figure 2 A-D)** at the phylum and genus levels.

Our negative control reaction generated a small number of reads that were identified as the following genera; *Halomonas (6 reads)*, *Corynebacterium (64)*, *Staphylococcus (24)*, *Ralstonia (1726)*, *Stenotrophomonas (460)*, *Pseudomonas (276)*, *Escherichia-Shigella (2420)* and *Ruminococcus* (405), but was overwhelmingly composed of reads mapping to the genus *Serratia (18000)*.

At the phylum level, the blood microbiome was predominated by Proteobacteria (88% of all bacterial DNA in the control population, and 80.9% in the asthmatic

261   population) followed by Actinobacteria (control = 7.8%, asthmatic = 7.1%),
262   Firmicutes (control = 3.5%, asthmatic = 9.2%) and Bacteroidetes (control = 0.1%,
263   asthmatic = 2.2%). These findings mirror previous studies [2,7,25,28] and further support
264   the notion of a core blood microbiome predominated by four key phyla.
265
At the genus level, our blood samples were predominated by the genus
Achromobacter [38] which accounted for 51.1% and 45.3% of the total bacterial DNA
detected in the control and asthma donors respectively. To a lesser extent, the blood
samples also comprised members of the *Pseudomonas* (12.8%, 7.5%) [7,9,18],
Serratia* (0.9%, 11.6%) [39], *Sphingomonas* (3.8%, 5.1%) [7,25], *Staphylococcus* (5.5%,
2.8%) [7,18,40,41], *Corynebacterium* (3.2%, 5.5%) [7,9], and *Acinetobacter* (3.7%, 2.8%) [7]
genera. The genus *Serratia* was excluded from further analysis as the study samples
presented with less reads than did the corresponding negative control reaction and
thus it was considered a contaminant.
275
Whilst all genera found have been previously described in the blood (see
references), the predominance of *Achromobacter,* which is not classically associated
with the blood microbiome, warrants further consideration. Indeed, *Achromobacter*
has been detected abundantly in the lower respiratory tract of healthy mice [42],
humans (HPM airway dataset), and in various respiratory conditions [43,44].
Furthermore, no *Achromobacter* was detected in our control reactions suggesting
that its presence is not the result of experimental contamination.
283
284   Although differential composition of the blood microbiome in response to pathology
285   was not the main focus of this study, we performed principal coordinates analysis
286   and despite the relatively small sample set, this revealed good separation between
287   the two treatment groups based upon their microbiome profile, suggesting that the
288   blood microbiome community was altered in the asthmatic subjects (**Figure 3**).
289
290

291
292
Figure 2 – Relative abundance of the most abundant taxa (>1%) as determined by
amplification and sequencing of the 16S rRNA gene variable region 4. Data are
mean abundance expressed as a percentage of the total bacterial sequence count.
(A) Phylum-level data grouped by condition, (B) Genus-level data grouped by
condition, (C) Phylum-level individual sample data, (D) Genus-level individual
sample data.
299
300

PC2 (19.75 %)

PC1 (37.83 %)

PC3 (17.57 %)

301
302
303 **Figure 3** – Principal coordinates analysis of weighted unifrac distances for control
304 (blue) and asthmatic (red) blood microbiome profiles. Each dot represents an
305 individual sample, and the microbiome of samples that appear more closely together
306 are more similar.

307
308
DNA-level Circulating Microbiome – Likely Origins
In accordance with our hypothesis that the blood microbiome exists as a
consequence of bacterial translocation from other microbiome niches within the
body, we compared the data generated herein with gastrointestinal tract, oral cavity,
and skin data made available by the Human Microbiome Project (HMP). In each
case, the HMP data and our own were combined, weighted UniFrac distances
calculated, and a principal coordinates analysis performed. **Figure 4** demonstrates
that the blood microbiome of our control and asthmatic donors clustered more
closely in PCoA space with the oral cavity and skin HMP data, than it did with the
gastrointestinal tract HMP data. This suggests that the blood microbiome community
is perhaps more likely to result from the translocation of organisms from the oral
cavity and skin, than from organisms that colonise the gastrointestinal tract.

321
Various daily activities including chewing, tooth brushing and flossing have been
shown to result in the translocation of bacteria from the oral cavity into the
bloodstream [19-22]. Further, the skin has a distinct microbial community and is
susceptible to injury, and thus represents a large surface area through which such

organisms may translocate to the bloodstream. It is important here to consider
sources of contamination; venepuncture, the process through which the majority of
blood samples are obtained, is recognised as a cause of transient bacteraemia [45],
and despite the use of preventative measures such as alcohol cleansing of the skin
prior to breaking the surface, there remains the possibility that organisms entered the
sample from the skin via this route.
332
333 One important, but often overlooked limitation of DNA-based microbiome
334 characterisation is that DNA persists post-mortem, even in the presence of harsh
335 environmental conditions. From such analyses it is therefore impossible to confirm
336 whether an organism is present and viable, is present but non-viable, or whether the
337 organism has since left the environment in question yet it's DNA persists.
338
339
340

341
342
Figure 4 – Principal coordinates analysis of weighted UniFrac distances between
variable region 4 16S sequencing data from our donors and the Human Microbiome
Project (A) Gut, (B) Oral Cavity and (C) Skin data. Each dot represents an individual
sample, and the microbiome of samples that appear more closely together are more
similar. In each case, our control donor samples appear in blue, and our asthmatic
donor samples appear in red. Further sample details are provided beneath each
figure, and the number of datasets representing each anatomical location is provided
in brackets.
351

RNA-level Circulating Microbiome – Community Structure

We hypothesise that some of the non-mapping (likely non-human) reads that often result from whole transcriptome analyses (RNA-seq) represent microbial community members that were present (or parts thereof) in the blood at the time of RNA extraction. Furthermore, given the highly labile nature of RNA [46], and the presence of ribonucleases within the circulation [47], we suggest that the detection of bacterial RNA goes further towards confirming at least the recent presence of these organisms when compared with DNA-based approaches. From our previous studies of the circulating transcriptome of our donor community, a total of 439,448,931 paired RNA reads failed to map to the human genome. These reads were used for the following analyses as randomly subsampled populations of 1 million, 10 million, 25 million and 100 million read pairs. Mapping the total read population back to the subsampled populations allowed us to assess how well each subsampled population approximated the starting (entire, ~ 440M read) population. Data revealed only marginal improvements in whole community representation as the subsampled population increased (1M, 10M, 25M, 100M represented 65.05, 66.05, 66.81, 64.24% of the total read population). A subsampled population of 25 million reads (25M) provided an acceptable balance between read representation and computational efficiency and was therefore used for transcriptome assembly. The transcriptome comprised 2050 candidate "non-human genes" with a mean GC content of 53%. Ten-percent of these genes were greater than 517bp, and over half were at least 263bp. Taxonomic identification of each assembled feature was determined using Kraken [36] and this revealed that 729 of the 2050 features were of bacterial origin, and 7 features were from archaea (**Figure 5**).

At the phylum level, the whole transcriptome data was predominated by assembled Proteobacteria sequences (379 sequences, 52%), followed by Firmicutes (143, 19.8%), Actinobacteria (112, 15.5%) and Bacteroidetes (35, 4.8%). In considering the total number of reads mapping to each feature, 379.4M reads mapped to bacterial features (out of a total of 395M reads). Of those reads mapping to bacterial-derived sequences, Proteobacteria (74.9%, 47.0%; Control, Asthma) and Firmicutes (19.5%, 48.0%) predominated with, Actinobacteria (0.01%, 0.04%) and Bacteroidetes (0.05%, 0.008%) present to a much lesser extent. These findings support our DNA-level phylum data and mirror previous studies [2,7,25,28].

At the genus level, the whole transcriptome data was predominated by the genera *Paenibacillus* (17.8%, 44.6%; Control, Asthma reads)[48], *Escherichia* (11.8%, 13.1%) [7,9,40,49], *Acinetobacter* (0.7%, 0.4%) [7], *Pseudomonas* (0.8%, 0.1%) [7,9,18,50] and *Propionibacterium* (0.5%, 0.2%) [9,18,41,50]. With the exception of *Paenibacillus,* all genera detected via *de novo* transcriptome assembly were also present in our DNA-level data and all genera found have been previously described in the blood (see references). The fact that *Paenibacillus* was present within our RNA-level analyses yet absent from our DNA-level data led us to consider whether it could have been introduced as a contaminant during the RNA extraction, library preparation or sequencing procedures. Given that negative control reactions are not routinely conducted for RNA-seq type applications, we were unable to experimentally confirm this. However, we did identify from the literature that this genus has been reported as a common reagent and laboratory contaminant, albeit at the DNA level [51]. Nevertheless there appeared to be little consistency in its presence within our sample set (mean % of reads 39.5 ± 40.3%) and we noted a difference in abundance

402 between our experimental groups despite all preparation procedures being the
403 same. The exact source of this RNA thus remains open to speculation.
404
405

406
407
408
409
Figure 5 – Taxonomic classification of features assembled from a randomly
subsampled population of 25 million unmapped RNA sequencing reads using Trinity,
and identified using Kraken. The numbers present by each taxonomic classification
refer to the number of features classified as such (e.g. 379 assembled features were
identified as Proteobacteria). D – domain, P – phylum, F – family, G – genus, S –
species.
416
417

Classical Culture – Presence of Viable Organisms

The presence of viable, proliferating bacteria in the blood was assessed using growth culture assays as previously described. To our surprise, bacterial cultures were positive for the majority of the blood samples assayed (8 out of 10 samples; 4 control blood samples and 4 asthma blood samples), whilst all negative control plates had no growth as expected. In all instances, bacterial growth was monoculture on microscopy and thus 16S colony PCR (amplifying the entire 16S gene) and Sanger sequencing was conducted on a three independent colonies per plate for identification purposes.

Bacteria were identified using Sanger sequencing following classification with Kraken, and included the following genera; *Staphylococcus* (49 sequences), *Micrococcus* (12), *Kocuria* (6), *Corynebacterium* (6) and *Propionibacterium* (1) (**Figure 6**). These genera belong to the phyla Actinobacteria (*Corynebacterium*, *Kocuria*, *Micrococcus*) and Firmicutes (*Staphylococcus*) and were all represented in our 16S DNA level data (**Figure 4**) and detected within our RNA data. Individual sample data is presented in (**Table 5**). It is noteworthy that, with the exception of Kocuria, all bacteria identified displayed some of the highest total relative abundance scores in the 16S sequencing results; Corynebacterium (4.2%), Kocuria (0.2%), Micrococcus (1.30%), and Staphylococcus at 4.3%.

Due to the long-held belief that the bloodstream of healthy individuals is sterile and since the blood is an unfavourable compartment for the microbes due to its bacteriostatic and bactericidal components; here we consider the likely source of these viable organisms. The skin microbiome is dominated by members of the genera *Corynebacterium*, *Micrococcus*, *Staphylococcus*, and *Propionibacterium*, the proportions of which vary markedly between individuals [52]. Furthermore, several studies report the presence of the genus *Kocuria* on the skin of humans and other mammals [53-55]. We therefore suggest that the organisms detected through our microbial culture experiments most likely originate from the skin. Whilst transient bacteraemia due to a breach of the skin barrier is an accepted occurrence, one would expect such organisms to be rapidly targeted and removed by the immune system. We therefore suggest that the viable organisms detected are the result of venepuncture contamination whereby organisms from the skin are drawn into the vacutainer, contaminating the sample.

456
457
458 **Figure 6** – Taxonomic classification of 16S sequencing reads generated from
459 cultured (viable) bacteria. The numbers present by each taxonomic classification
460 refer to the number of features classified as such (e.g. 49 colonies were identified as
461 Staphylococcus). In all cases, the lowest taxonomic rank is indicated. D – domain, P
462 – phylum, F – family, G – genus, S – species.
463
464

465 **Table 5** – Identification of cultured organisms using 16S colony PCR and Sanger Sequencing. Data are presented at the genus
466 level. The suffix "An" denotes the organism was cultured in an anaerobic environment. Culture Neg denotes no bacterial growth on
467 any of the substrates used.
468

Control Samples					Asthma Samples			
BRH1017878	BRH1017879	BRH1017880	BRH1017881	BRH1017882	BRH1017874	BRH1017875	BRH1017876	BRH1017877
Kocuria rhizophila	Micrococcus luteus	Staphylococcus haemolyticus Staphylococcus epidermidis Propionibacterium acnes	Staphylococcus haemolyticus	Culture Neg	Staphylococcus haemolyticus	Culture Neg	Corynebacterium halotolerans	Staphylococcus epidermidis

Concluding Remarks

478   This study utilised a range of molecular and classical microbiology approaches to characterise the human blood microbiome in unparalleled detail. Our DNA and RNA-based studies revealed a diverse community of bacterial, the main members of which having been described in a range of other studies and therefore providing further evidence of a core blood microbiome. Although disease associated changes in the blood microbiome were not the focus of this study, the fact we identified such changes is encouraging and supports efforts to identify circulating microbiome signatures indicative of disease.

To our surprise, culture based experiments revealed viable bacteria present in eight of our ten study samples. Whilst we attribute their presence to venepuncture-associated contamination of the blood sample (and make recommendations to avoid this in future), the presence of these viable organisms does not undermine our data that support an abundance of bacteria-associated DNA and RNA within the blood, likely present due to translocation. On reflecting upon our experimental approach, we make the following recommendations for future studies;

(1) More attention should be paid to blood collection procedures as any contamination occurring at this stage affects all downstream procedures. In addition to alcohol cleansing of the skin (as performed in this study), we recommend that the first volume drawn is diverted to a secondary tube, and analysed separately. This will allow further investigation of venepuncture-associated contamination.

(2) The inclusion of negative control reactions that are subject to the whole range of experimental procedures, including library preparation and sequencing, is absolutely essential.

(3) Where possible, RNA-seq studies used for unmapped read assembly should include negative control samples that are subject to all of the experimental procedures alongside the study samples. This will allow an appraisal of how significant reagent / laboratory procedure contamination is in these studies. Often the use of RNA-seq derived unmapped reads for microbiome characterisation is a "secondary use", and this is simply not possible.

(4) Seminal studies are still required to satisfactorily investigate the phenomenon of bacterial translocation from well-characterised microbiome niches to the blood.

Acknowledgements

517 This study used the Nephele platform from the National Institute of Allergy and
518 Infectious Diseases (NIAID) Office of Cyber Infrastructure and Computational
519 Biology (OCICB) in Bethesda, MD.

520
521

References

522 **References**
523

1   Markova, N. D. L-form bacteria cohabitants in human blood: significance for health and
diseases. *Discov Med* **23**, 305-313 (2017).
2   Amar, J. *et al.* Blood microbiota dysbiosis is associated with the onset of cardiovascular
events in a large general population: the D.E.S.I.R. study. *PLoS One* **8**, e54461,
doi:10.1371/journal.pone.0054461 (2013).
3   Bhattacharyya, M., Ghosh, T., Shankar, S. & Tomar, N. The conserved phylogeny of blood
microbiome. *Mol Phylogenet Evol* **109**, 404-408, doi:10.1016/j.ympev.2017.02.001 (2017).
4   Kell, D. B. & Pretorius, E. On the translocation of bacteria and their lipopolysaccharides
between blood and peripheral locations in chronic, inflammatory diseases: the central roles
of LPS and LPS-induced cell death. *Integr Biol (Camb)* **7**, 1339-1377, doi:10.1039/c5ib00158g
(2015).
5   Li, Q. *et al.* Identification and Characterization of Blood and Neutrophil-Associated
Microbiomes in Patients with Severe Acute Pancreatitis Using Next-Generation Sequencing.
Front Cell Infect Microbiol **8**, 5, doi:10.3389/fcimb.2018.00005 (2018).
6   Mangul, S. *et al.* Total RNA Sequencing reveals microbial communities in human blood and
disease specific effects. *bioRxiv*, doi:10.1101/057570 (2016).
7   Païssé, S. *et al.* Comprehensive description of blood microbiome from healthy donors
assessed by 16S targeted metagenomic sequencing. *Transfusion* **56**, 1138-1147,
doi:10.1111/trf.13477 (2016).
8   Potgieter, M., Bester, J., Kell, D. B. & Pretorius, E. The dormant blood microbiome in chronic,
inflammatory diseases. *FEMS Microbiol Rev* **39**, 567-591, doi:10.1093/femsre/fuv013 (2015).
9   Dinakaran, V. *et al.* Elevated levels of circulating DNA in cardiovascular disease patients:
metagenomic profiling of microbiome in the circulation. *PLoS One* **9**, e105221,
doi:10.1371/journal.pone.0105221 (2014).
10 Rajendhran, J., Shankar, M., Dinakaran, V., Rathinavel, A. & Gunasekaran, P. Contrasting
circulating microbiome in cardiovascular disease patients and healthy individuals. *Int J*
Cardiol **168**, 5118-5120, doi:10.1016/j.ijcard.2013.07.232 (2013).
11 Nikkari, S., McLaughlin, I. J., Bi, W., Dodge, D. E. & Relman, D. A. Does blood of healthy
subjects contain bacterial ribosomal DNA? *J Clin Microbiol* **39**, 1956-1959,
doi:10.1128/JCM.39.5.1956-1959.2001 (2001).
12 Vientós-Plotts, A. I. *et al.* Dynamic changes of the respiratory microbiota and its relationship
to fecal and blood microbiota in healthy young cats. *PLoS One* **12**, e0173818,
doi:10.1371/journal.pone.0173818 (2017).
13 Mandal, R. K. *et al.* An investigation into blood microbiota and its potential association with
Bacterial Chondronecrosis with Osteomyelitis (BCO) in Broilers. *Sci Rep* **6**, 25882,
doi:10.1038/srep25882 (2016).
14 Sze, M. A. *et al.* Changes in the bacterial microbiota in gut, blood, and lungs following acute
LPS instillation into mice lungs. *PLoS One* **9**, e111228, doi:10.1371/journal.pone.0111228
(2014).
15 Jeon, S. J. *et al.* Blood as a route of transmission of uterine pathogens from the gut to the
uterus in cows. *Microbiome* **5**, 109, doi:10.1186/s40168-017-0328-9 (2017).
16 Martel, J., Wu, C. Y., Huang, P. R., Cheng, W. Y. & Young, J. D. Pleomorphic bacteria-like
structures in human blood represent non-living membrane vesicles and protein particles. *Sci*
Rep **7**, 10650, doi:10.1038/s41598-017-10479-8 (2017).
17 McLaughlin, R. W. *et al.* Are there naturally occurring pleomorphic bacteria in the blood of
healthy humans? *J Clin Microbiol* **40**, 4771-4775 (2002).
18 Damgaard, C. *et al.* Viable bacteria associated with red blood cells and plasma in freshly
drawn blood donations. *PLoS One* **10**, e0120826, doi:10.1371/journal.pone.0120826 (2015).

19   Forner, L., Larsen, T., Kilian, M. & Holmstrup, P. Incidence of bacteremia after chewing, tooth
brushing and scaling in individuals with periodontal inflammation. *J Clin Periodontol* **33**, 401-
407, doi:10.1111/j.1600-051X.2006.00924.x (2006).
20   Horliana, A. C. *et al.* Dissemination of periodontal pathogens in the bloodstream after
periodontal procedures: a systematic review. *PLoS One* **9**, e98271,
doi:10.1371/journal.pone.0098271 (2014).
21   Parahitiyawa, N. B., Jin, L. J., Leung, W. K., Yam, W. C. & Samaranayake, L. P. Microbiology of
odontogenic bacteremia: beyond endocarditis. *Clin Microbiol Rev* **22**, 46-64, Table of
Contents, doi:10.1128/CMR.00028-08 (2009).
22   Lockhart, P. B. *et al.* Bacteremia associated with toothbrushing and dental extraction.
Circulation **117**, 3118-3125, doi:10.1161/CIRCULATIONAHA.107.758524 (2008).
23   Ono, S. *et al.* Detection of microbial DNA in the blood of surgical patients for diagnosing
bacterial translocation. *World J Surg* **29**, 535-539, doi:10.1007/s00268-004-7618-7 (2005).
24   Sato, J. *et al.* Gut dysbiosis and detection of "live gut bacteria" in blood of Japanese patients
with type 2 diabetes. *Diabetes Care* **37**, 2343-2350, doi:10.2337/dc13-2817 (2014).
25   Lelouvier, B. *et al.* Changes in blood microbiota profiles associated with liver fibrosis in obese
patients: A pilot analysis. *Hepatology* **64**, 2015-2027, doi:10.1002/hep.28829 (2016).
26   Lehtiniemi, J., Karhunen, P. J., Goebeler, S., Nikkari, S. & Nikkari, S. T. Identification of
different bacterial DNAs in human coronary arteries. *Eur J Clin Invest* **35**, 13-16,
doi:10.1111/j.1365-2362.2005.01440.x (2005).
27   Ling, Z. *et al.* Blood microbiota as a potential noninvasive diagnostic biomarker for liver
fibrosis in severely obese patients: Choose carefully. *Hepatology* **65**, 1775-1776,
doi:10.1002/hep.28987 (2017).
28   Olde Loohuis, L. M. *et al.* Transcriptome analysis in whole blood reveals increased microbial
diversity in schizophrenia. *Transl Psychiatry* **8**, 96, doi:10.1038/s41398-018-0107-9 (2018).
29   Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data
on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* **79**, 5112-5120,
doi:10.1128/AEM.01043-13 (2013).
30   Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data.
Nat Methods **7**, 335-336, doi:10.1038/nmeth.f.303 (2010).
31   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing
and web-based tools. *Nucleic Acids Res* **41**, D590-596, doi:10.1093/nar/gks1219 (2013).
32   Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity
platform for reference generation and analysis. *Nat Protoc* **8**, 1494-1512,
doi:10.1038/nprot.2013.084 (2013).
33   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323, doi:10.1186/1471-2105-12-323
(2011).
34   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652, doi:10.1038/nbt.1883 (2011).
35   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140,
doi:10.1093/bioinformatics/btp616 (2010).
36   Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification using
exact alignments. *Genome Biol* **15**, R46, doi:10.1186/gb-2014-15-3-r46 (2014).
37   Breitwieser, F. P. & Salzberg, S. L. Pavian: Interactive analysis of metagenomics data for
microbiomics and pathogen identification. *bioRxiv*, doi:10.1101/084715 (2016).
38   Gyarmati, P. *et al.* Metagenomic analysis of bloodstream infections in patients with acute
leukemia and therapy-induced neutropenia. *Sci Rep* **6**, 23532, doi:10.1038/srep23532
(2016).

39   Moriyama, K. *et al.* Polymerase chain reaction detection of bacterial 16S rRNA gene in
human blood. *Microbiol Immunol* **52**, 375-382, doi:10.1111/j.1348-0421.2008.00048.x
(2008).
40   Such, J. *et al.* Detection and identification of bacterial DNA in patients with cirrhosis and
culture-negative, nonneutrocytic ascites. *Hepatology* **36**, 135-141,
doi:10.1053/jhep.2002.33715 (2002).
41   Marques da Silva, R., Lingaas, P. S., Geiran, O., Tronstad, L. & Olsen, I. Multiple bacteria in
aortic aneurysms. *J Vasc Surg* **38**, 1384-1389, doi:10.1016/S0741 (2003).
42   Singh, N., Vats, A., Sharma, A., Arora, A. & Kumar, A. The development of lower respiratory
tract microbiome in mice. *Microbiome* **5**, 61, doi:10.1186/s40168-017-0277-3 (2017).
43   Hogan, D. A. *et al.* Analysis of Lung Microbiota in Bronchoalveolar Lavage, Protected Brush
and Sputum Samples from Subjects with Mild-To-Moderate Cystic Fibrosis Lung Disease.
PLoS One **11**, e0149998, doi:10.1371/journal.pone.0149998 (2016).
44   Segal, L. N., Rom, W. N. & Weiden, M. D. Lung microbiome for clinicians. New discoveries
about bugs in healthy and diseased lungs. *Ann Am Thorac Soc* **11**, 108-116,
doi:10.1513/AnnalsATS.201310-339FR (2014).
45   Depcik-Smith, N. D., Hay, S. N. & Brecher, M. E. Bacterial contamination of blood products:
factors, options, and insights. *J Clin Apher* **16**, 192-201 (2001).
46   Tsui, N. B., Ng, E. K. & Lo, Y. M. Stability of endogenous and added RNA in blood specimens,
serum, and plasma. *Clin Chem* **48**, 1647-1653 (2002).
47   Reddi, K. K. & Holland, J. F. Elevated serum ribonuclease in patients with pancreatic cancer.
Proc Natl Acad Sci U S A **73**, 2308-2310 (1976).
48   Sáez-Nieto, J. A. *et al.* spp. isolated from human and environmental samples in Spain:
detection of 11 new species. *New Microbes New Infect* **19**, 19-27,
doi:10.1016/j.nmni.2017.05.006 (2017).
49   Francés, R. *et al.* A sequential study of serum bacterial DNA in patients with advanced
cirrhosis and ascites. *Hepatology* **39**, 484-491, doi:10.1002/hep.20055 (2004).
50   Bhatt, A. S. *et al.* In search of a candidate pathogen for giant cell arteritis: sequencing-based
characterization of the giant cell arteritis microbiome. *Arthritis Rheumatol* **66**, 1939-1944,
doi:10.1002/art.38631 (2014).
51   Salter, S. J. *et al.* Reagent and laboratory contamination can critically impact sequence-based
microbiome analyses. *BMC Biol* **12**, 87, doi:10.1186/s12915-014-0087-z (2014).
52   Tett, A. *et al.* Unexplored diversity and strain-level structure of the skin microbiome
associated with psoriasis. *NPJ Biofilms Microbiomes* **3**, 14, doi:10.1038/s41522-017-0022-5
(2017).
53   Cosseau, C. *et al.* Proteobacteria from the human skin microbiota: Species-level diversity and
hypotheses. *One Health* **2**, 33-41, doi:10.1016/j.onehlt.2016.02.002 (2016).
54   McIntyre, M. K., Peacock, T. J., Akers, K. S. & Burmeister, D. M. Initial Characterization of the
Pig Skin Bacteriome and Its Effect on In Vitro Models of Wound Healing. *PLoS One* **11**,
e0166176, doi:10.1371/journal.pone.0166176 (2016).
55   Grice, E. A. *et al.* A diversity profile of the human skin microbiota. *Genome Res* **18**, 1043-
1050, doi:10.1101/gr.075549.107 (2008).
56   van Rensburg, J. J. *et al.* The Human Skin Microbiome Associates with the Outcome of and Is
Influenced by Bacterial Infection. *MBio* **6**, e01315-01315, doi:10.1128/mBio.01315-15
(2015).

668