1

The transfer function of the rhesus macaque oculomotor system for small-amplitude slow motion trajectories

Julianne Skinner[1], Antimo Buonocore[2,3], and Ziad M. Hafed[2, 3*]

1. Graduate School of Neural and Behavioural Sciences, International Max Planck Research School, Tübingen University, Tübingen, BW, 72074, Germany
2. Werner Reichardt Centre for Integrative Neuroscience, Tübingen University, Tübingen, BW, 72076, Germany
3. Hertie Institute for Clinical Brain Research, Tübingen University, Tübingen, BW, 72076, Germany

Corresponding Author:

Ziad M. Hafed
Werner Reichardt Centre for Integrative Neuroscience
Otfried-Mueller Str. 25
Tübingen, 72076, Germany
Tel: +49 7071 29-88819
E-mail: ziad.m.hafed@cin.uni-tuebingen.de

30 **The transfer function of the rhesus macaque oculomotor system for**
31 **small-amplitude slow motion trajectories**
32

Julianne Skinner[1], Antimo Buonocore[2,3], and Ziad M. Hafed[2,3]
1. Graduate School of Neural and Behavioural Sciences, International Max Planck
Research School, Tübingen University, Tübingen, BW, 72074, Germany
2. Werner Reichardt Centre for Integrative Neuroscience, Tübingen University,
Tübingen, BW, 72076, Germany
3. Hertie Institute for Clinical Brain Research, Tübingen University, Tübingen, BW,
72076, Germany

40
41

Abstract

Two main types of small eye movements occur during gaze fixation: microsaccades and
slow ocular drifts. While microsaccade generation has been relatively well-studied,
ocular drift control mechanisms are unknown. Here we explored the degree to which
monkey smooth eye movements, on the velocity scale of slow ocular drifts, can be
generated systematically. Two male rhesus macaque monkeys tracked a spot moving
sinusoidally, but slowly, along the horizontal or vertical directions. Maximum target
displacement in the motion trajectory was 30 min arc (0.5 deg), and we varied the
temporal frequency of target motion from 0.1 to 5 Hz. We obtained an oculomotor
"transfer function" by measuring smooth eye velocity gain (relative to target velocity) as
a function of frequency, similar to past work with large-amplitude pursuit. Monkey eye
velocities as slow as those observed during slow ocular drifts were clearly target-motion
driven. Moreover, like with large-amplitude smooth pursuit, eye velocity gain varied with
temporal frequency. However, unlike with large-amplitude pursuit, exhibiting primarily
low-pass behavior, small-amplitude motion tracking was band-pass with the best ocular
movement gain occurring at ~0.8-1 Hz. When oblique directions were tested, we found
that monkey oblique pursuit gain for small-amplitude slow motion trajectories was >1.
Our results provide a catalogue of the control abilities of the monkey oculomotor system
for slow target motions, and they also support the notion that smooth fixational ocular
drifts are controllable. This has implications for neural investigations of drift control and
the image-motion consequences of drifts on visual coding in early visual areas.

Introduction

Small fixational eye movements continuously move the retinal image, even though they never deviate the center of gaze (or fovea) away from objects of interest. Such continuous movements mean that the neural signals representing input images in a variety of early visual areas are continuously modulated, and this has implications for theoretical interpretations of neural variability across trials, as well as population correlations in simultaneously recorded neurons (McFarland et al. 2016; Rucci and Victor 2015). Therefore, understanding the neural control mechanisms for fixational eye movements is an important first step towards understanding the full impact of these eye movements on neural coding, and ultimately visual perception.

Fixational eye movements come in two main flavors: microsaccades, which are rapid gaze shifts that look like big saccades; and smooth ocular drifts, which are slow changes in eye position occurring in between microsaccades (Barlow 1952; Murphy et al. 1975; Nachmias 1961). Even though microsaccades have received a substantial amount of research attention recently (Hafed 2011; Hafed et al. 2015; Krauzlis et al. 2017; Rolfs 2009), drifts remain to be relatively underexplored. For example, the mechanisms for generating ocular drifts are unknown, whereas microsaccade generation has been studied in a variety of different brain areas (Arnstein et al. 2015; Hafed et al. 2009; Hafed and Krauzlis 2012; Peel et al. 2016; Sun et al. 2016; Van Horn and Cullen 2012). Moreover, uncovering detailed neural mechanisms for microsaccade generation in certain key brain areas was instrumental for uncovering important, previously unappreciated, consequences of these eye movements on vision, perception, and even cognition (Chen et al. 2015;

Hafed 2013; 2011; Hafed et al. 2015; Krauzlis et al. 2017; Tian et al. 2016; Veale et al.

2017). Thus, exploring the neural mechanisms for slow ocular drift generation is a

worthwhile effort, especially given the fact that during fixation, microsaccades are brief

events in an otherwise continuous sea of ocular drifts.

Here, towards approaching that ultimate goal, we aimed to characterize the degree to

which the control system for eye position in the macaque monkey brain is able to

generate very slow smooth pursuit eye movements, on the velocity scale of fixational

ocular drifts. In other words, is it possible for monkeys to volitionally generate a slow eye

movement that is as small in amplitude and velocity as slow ocular drifts, but that is

clearly controlled and with a predictable motion trajectory? We designed behavioral

experiments motivated by potential analogies that one can make between slow ocular

drifts and smooth pursuit eye movements (Cunitz 1970; Martins et al. 1985; Nachmias

1961), similar to analogies that one makes between microsaccades and larger saccades

(Hafed 2011; Krauzlis et al. 2017). After all, smooth pursuit eye movements are

(relatively) slow rotations of the eyeball, similar in nature to slow ocular drifts, and the

circuits for generating such smooth pursuit eye movements are well studied (Krauzlis

2004).

We presented macaque monkeys with small-amplitude sinusoidal target trajectories, and

we asked how well their small-amplitude slow eye movements can track such

trajectories. Our approach was to assume that for the frequency ranges that we studied,

the oculomotor system may behave, to a first approximation, like a linear system. This

means that we can present a single frequency and measure the response, and then test

another frequency, and so on. The gain and phase lag of tracking at each frequency can

thus allow estimating an "equivalent" transfer function of the oculomotor system (Ohashi

and Mizukoshi 1991). Previous attempts like this with larger-amplitude (and faster) eye

movements in smooth pursuit were effective, and showed that pursuit tracks very well for

low frequencies (<1 Hz), but that it then behaves relatively poorly with higher

frequencies (exhibiting both lower gain and larger phase lag) (Bahill and McDonald

1983; Collewijn and Tamminga 1984; Fabisch et al. 2009; Martins et al. 1985; Ohashi

and Mizukoshi 1991; Rottach et al. 1996). We were interested in what happens in the

monkey with much smaller and slower eye movements. The key comparison was to see

whether controlled slow eye movements as slow as fixational ocular drifts would be

possible. Critically, we performed our experiments on monkeys in order to demonstrate

that these animals are instrumental for ultimately uncovering the neural control

mechanisms for slow ocular drifts, and also in order to complement earlier human work

on the topic (Cunitz 1970; de Bie and van den Brink 1986; Martins et al. 1985; Murphy et

al. 1975; Nachmias 1961; Wyatt and Pola 1981).

Materials and Methods

Animal preparation and laboratory setup

We recorded eye movements from two male rhesus macaque (macaca mulatta) monkeys

(monkey A and monkey M) aged 6-7 years. We implanted one eye (left for monkey A

and right for monkey M) in each animal with a scleral search coil for eye tracking using

the magnetic induction technique (Fuchs and Robinson 1966; Judge et al. 1980). Surgical

procedures for implantation were similar to those described earlier (Chen and Hafed

2013; Hafed and Ignashchenkova 2013), and the experiments were approved by ethics

committees at the regional governmental offices of the city of Tübingen. These

experiments were in accordance with European Union guidelines on animal research, as

well as the associated implementations of these guidelines in German law.

We used monkeys in this study for two important reasons. First, the monkeys had scleral

search coils implanted, which allowed the most precise measurement of small-amplitude

slow eye movements, including fixational drifts. Video-based eye trackers (used in most

human studies) are less reliable than scleral search coils for measuring slow eye

movements. Second, and more importantly, the monkeys are now being used in

neurophysiological recording experiments, such that direct neural correlates of our

observations here can be identified and disseminated; particularly to complement earlier

human work on similar questions (Cunitz 1970; de Bie and van den Brink 1986; Martins

et al. 1985; Murphy et al. 1975; Nachmias 1961).

During data collection, the animals were seated in a primate chair 73 cm from a CRT

computer monitor in an otherwise dark room. The monitor had a pixel resolution of 34

pixels/deg and a refresh rate of 120 Hz. Stimuli were presented over a uniform gray

background (29.7 Cd/m^2 in Experiment 1, and either 29.7 or 4.4 Cd/m^2 in Experiment 2).

A small white spot (~5 x 5 min arc square) having 86 Cd/m^2 luminance in Experiment 1

and either 86 or 48.1 Cd/m^2 in Experiment 2 was used as the moving target for smooth

pursuit eye movements (see *Behavioral tasks* below).

Graphics on the CRT monitor were presented by a computer running Matlab's

Psychophysics Toolbox (Brainard 1997; Kleiner et al. 2007; Pelli 1997). This computer

in turn received commands from a real-time I/O system from National Instruments

(Austin, USA), which ensured control of the display on a frame-by-frame basis, as well

as real-time monitoring of animal behavior and reward. The system was described

recently (Chen and Hafed 2013; Tian et al. 2018; 2016).

Behavioral tasks

Experiment 1: Temporal frequency series

The monkeys fixated a small white spot for 350-550 ms at trial onset, after which the spot

started moving either horizontally or vertically along a sinusoidal position trajectory of

only 30 min arc (0.5 deg) amplitude. The monkeys had to track this moving target with

their eyes, and target position in deg (along either the horizontal or vertical axis) could be

described by the following equation

$$target\ position = 0.5{\times}\sin{(2\pi ft + \varphi)} \quad (1)$$

where f is the temporal frequency of target trajectory, t is time from motion onset, and ϕ

could be either 0 or π across trials. The temporal frequency, f, of target motion was

chosen randomly from trial to trial from among the following values: 0.1, 0.2, 0.3, 0.4,

0.5, 0.6, 0.8, 1, 2, 3, 4, or 5 Hz. Target motion duration was constant within a session, but

could vary across sessions in the range of 3000-4200 ms, depending on animal

motivation on any one day. In all cases, we had a long enough target motion duration to

ensure that we were analyzing steady-state tracking behavior, even for the smallest values

of f (associated with the slowest target position changes). Horizontal and vertical target

trajectories were collected in different blocks of trials, and we analyzed a total of 867

trials from monkey A and 1392 trials from monkey M. We did not penalize the monkeys

(e.g. by aborting trials) for making catch-up saccades as long as they stayed within a

radius of ~1-1.5 deg around the instantaneous target position.

Experiment 2: Amplitude series

The monkeys performed a similar experiment to that described above, but this time, the

temporal frequency, f, was maintained at 0.5 Hz. Also, we interleaved different target

motion directions, and we varied the amplitude of the target motion. Target motion

trajectory was now described by the following equations

$$horizontal\ target\ position = Amp_h{\times}\sin{(2\pi ft)} \quad (2)$$

$$vertical\ target\ position = Amp_v{\times}\sin{(2\pi ft)} \quad (3)$$

where f was fixed at 0.5 Hz, and Amp_h and Amp_v specified the target motion amplitudes.

For purely horizontal target motions, Amp_h and Amp_v were chosen to result in radial

amplitudes of 0.25, 0.5, 1, or 2 deg. We also introduced two oblique directions: "+45 deg

oblique pursuit" was used to describe the case when Amp_h and Amp_v were equal and

positive (again from among 0.25, 0.5, 1, or 2 deg); "+135 deg oblique pursuit" was used

to describe the case when Amp_v had an opposite sign from Amp_h and Amp_h was negative

(e.g. -0.25 deg for Amp_h and +0.25 deg for Amp_v). In other words, the two oblique

directions both started with upward motion trajectory, but with one first moving

rightward/upward (+45 deg) and the other first moving leftward/upward (+135 deg).

During the sessions, we interleaved all amplitude and direction conditions. We analyzed a

total of 969 trials from monkey A and 2145 trials from monkey M.

Experiment 3: Fixation comparison

For a subset of analyses, we compared eye velocity during tracking of small-amplitude

motion trajectories to eye velocity during fixation. The monkeys simply fixated the same

small spot for 1000-1400 ms before getting rewarded. We analyzed a total of 3160 trials

from monkey A and 5222 trials from monkey M.

Data analysis

Detecting catch-up saccades

We detected catch-up saccades using velocity and acceleration criteria (Chen and Hafed

2013; Hafed et al. 2009; Krauzlis and Miles 1996), and we manually inspected all

movements to correct for misses or false detections. For a subset of the data, we used

instead a novel state-of-the-art machine-learning approach for saccade detection using

convolutional neural networks, which we have developed (Bellet et al. 2018).

Catch-up saccade detection was necessary for performing focused analyses on smooth

pursuit eye movements (see below), but we also analyzed interesting properties of these

saccades themselves. For example, we explored both the frequency of occurrence and

amplitude of these eye movements as a function of either target motion temporal

frequency, f, or amplitude (Amp_h, Amp_v).

Measuring smooth pursuit gain and phase lag

We plotted eye velocity as a function of time for either horizontal or vertical eye

movements. For oblique motion trajectories (Experiment 2), we performed a coordinate

rotation such that one component of eye velocity was along the motion trajectory and the

other was orthogonal, and we plotted eye velocity for the component along the motion

trajectory (in other analyses, we also analyzed the horizontal and vertical components of

oblique eye velocity independently). We then picked a period of steady-state smooth

pursuit execution by excluding the first 1000 ms of eye movement data after target

motion onset (in Experiment 2, we relaxed this to 300 ms). In each velocity trace, we

then excised any interval during which a catch-up saccade was being executed, and we

also removed 10 ms before and 10 ms after each such saccade. The data that were excised

were replaced by Not-a-Number (NaN) labels, such that averages of eye velocity across

trial repetitions of a given condition did not include the large velocity transients

associated with catch-up saccades.

After plotting saccade-free eye velocity from any one trial, we fitted the resulting curve,

using a least-squares fitting algorithm, with a sinusoidal function (of appropriate temporal

frequency for the condition) in which the amplitude and phase values of the sinusoid

were the fitting parameters. This resulted in a population of amplitudes and phases from

the fitting procedure across trial repetitions of a given condition. For example, for 0.5 Hz

horizontal target trajectories in Experiment 1, we could have a population of N fitted

gains or phase lags across trials. We then summarized these N values into the mean gain

or mean phase lag at 0.5 Hz target motion temporal frequency, and with appropriate 95%

confidence intervals. "Gain" was defined as the ratio of the fitted eye velocity amplitude

in a sinusoid divided by the true amplitude of the target velocity sinusoid. For example,

for f Hz horizontal target trajectory with Amp_h deg position amplitude in equation 1 above

(and also equations 2 and 3), the target velocity amplitude was $2\pi f Amp_h$. Similar

procedures were performed for all conditions. In all analyses, we had >25 trials per

condition in each animal (most often, significantly more; e.g. see Fig. 2).

In a subset of analyses (e.g. Fig. 9), we measured eye velocity directly. For example, we

estimated eye velocity during smooth pursuit initiation or during baseline fixation (before

target motion onset). We defined a measurement interval of 50 ms, starting either at -100

ms or +100 ms from motion onset. The earlier interval measured eye velocity during

fixation, whereas the latter interval measured eye velocity during smooth pursuit

initiation.

Spectral analysis of eye positions

For some analyses of the data from Experiment 1, we performed a discrete Fourier

transform decomposition of eye position traces from the different values of f in equation

1. We picked, in each trial, an epoch of steady-state smooth pursuit (i.e. removing the

initial component immediately after motion onset as described above) that was 2500 ms

long. We then applied a Hanning window followed by discrete Fourier transformation to

investigate whether low-gain tracking was still modulated by the temporal frequency of

the target motion. We then plotted the average spectrum across the individual trial

spectra.

In all figures presented in this paper, we designated 95% confidence intervals as error

bars, such that the statistical robustness of our results can be easily assessed.

Results

Band-pass tuning for small-amplitude slow motion tracking in the monkey

Our goal was to systematically characterize the quality of monkey ocular control when

tracking small-amplitude slow motion trajectories. We were motivated by the more

general question of how slow ocular drifts that occur during gaze fixation may be

controlled, and how similar such control may be to the control needed when volitionally

tracking a moving target. We therefore asked two monkeys to pursue a small spot that

moved sinusoidally with an amplitude of only 30 min arc (0.5 deg) and different temporal

frequencies (Experiment 1; Material and Methods). For temporal frequencies of <1 Hz,

the peak velocities of target motion (based on equation 1) were always <3.14 deg/s, and

they were even smaller for even lower frequencies like 0.4 Hz, and also at off-peak-

velocity epochs of tracking. Therefore, the target velocities involved in our experiments

were similar in scale to the velocities with which the eye may drift on its own during

steady fixation (Cherici et al. 2012; Martins et al. 1985).

We found that eye velocity always tracked the temporal frequency of the target, albeit to

varying degrees of success. For example, Fig. 1A shows average saccade-free (Materials

and Methods) eye velocity from monkey M when this monkey tracked a horizontally

moving spot at 0.4 Hz in Experiment 1. Error bars denote 95% confidence intervals

across trials, and the solid blue line shows the true target velocity (based on the derivative

of equation 1). As can be seen, the eye moved sinusoidally at a similar temporal

frequency to the target, but eye velocity gain was quite low; fitting a sinusoid to the eye

velocity data (in the sustained pursuit interval; Materials and Methods; red line in Fig 1A)

showed a peak velocity amplitude in the fit of 0.532 deg/s relative to the true target peak

velocity of 1.26 deg/s, resulting in a gain of only 0.4235. There was also a phase lead of

~90 ms, which amounted to a lead of 0.036 of a full cycle of motion trajectory (3.6% of a

full cycle). When the target temporal frequency was 0.8 Hz instead, eye velocity gain was

significantly better (Fig. 1B), but it then decreased once again for even higher frequencies

(e.g. Fig. 1C; temporal frequency of 2 Hz). Phase lead or lag also matched the gain

changes by progressively shifting towards larger and larger lags, with 0.8 Hz showing

now a minimal phase delay (28 ms, or 0.022 of a full cycle) in tracking (as opposed to a

lead at 0.4 Hz), and the higher frequency showing an even more substantial delay of 35

ms or 0.07 of a full cycle. Therefore, monkey smooth ocular tracking of small-amplitude

slow motion trajectories may be described as being band-pass in nature, unlike the classic

description of smooth pursuit tuning (with much faster target speeds) as being low-pass

(Collewijn and Tamminga 1984; Fabisch et al. 2009; Rottach et al. 1996). This difference

is not due to the use of monkeys in our current study, as opposed to humans in the earlier

ones, because monkeys are indeed capable of high-gain sinusoidal pursuit of foveal spots

when faster target speeds (but similar low temporal frequencies) are used (Hafed et al.

2008; Hafed and Krauzlis 2008). It is very intriguing to us, nonetheless, that very highly

trained human subjects performed significantly better at low temporal frequencies than

our monkeys when face with similar small-amplitude motions (Martins et al. 1985).

Figure 1 Band-pass nature of monkey smooth pursuit of small-amplitude slow motion trajectories.
(**A**) Example tracking behavior from monkey M with 0.4 Hz sinusoidal target motion. The solid blue
sinusoid represents target velocity. The dark blue data plot shows mean eye velocity across trials
(surrounded by 95% confidence interval bands in a fainter color). The red sinusoid is a fit of the eye
velocity data in a sustained interval starting 1000 ms after target motion onset. As can be seen, the eye
tracked the frequency of target motion well, but with a markedly low peak velocity (i.e. low gain). (**B**)
Tracking gain was significantly higher at 0.8 Hz. This panel is formatted identically to **A**. (**C**) For even
higher frequencies, pursuit gain decreased again, as evidenced by the much smaller amplitude of the
sinusoid describing eye velocity relative to that describing target velocity (red and blue sinusoids,
respectively). Note that phase lag also increased (compare the phase of the solid red and blue sinusoids in
each panel). Error bars, when visible, denote 95% confidence intervals.

We confirmed the band-pass nature of small-amplitude slow motion tracking in our two

monkeys, and also with both horizontal and vertical tracking. For each temporal

frequency, f, in equation 1 (Materials and Methods), we estimated the gain of pursuit

(similar to Fig. 1; Materials and Methods) and plotted it for horizontal and vertical

tracking, along with 95% confidence intervals (Fig. 2A). In both monkeys, pursuit gain

peaked at ~0.8-1 Hz, and this happened for both horizontal and vertical tracking,

although vertical tracking always had significantly worse pursuit gain than horizontal

tracking (Rottach et al. 1996). For a frequency of 0.1 Hz, pursuit gain in both monkeys

seemed to increase relative to, say, 0.2 Hz and 0.3 Hz. This was because at 0.1 Hz, the

target was barely moving with a peak speed of only 0.314 deg/s. At such speeds, the

monkeys seemed to rely much more on saccades than slow eye movements to correct eye

350 position errors (e.g. see Fig. 5A), and their own fixational ocular drifts were significantly

351 faster than the target motion (e.g. see Fig. 3A). Therefore, for all frequencies other than

352 0.1 Hz, we observed a band-pass characteristic of monkey smooth pursuit gain for small-

353 amplitude slow motion trajectories.

Figure 2 Band-pass nature of slow motion smooth pursuit in both monkeys. (**A**) Each panel shows the
gain of smooth pursuit (i.e. the ratio of red sinusoid amplitude to blue sinusoid amplitude in Fig. 1) as a
function of target motion frequency. The left panel shows data from monkey M, and the right panel shows
data from monkey A. Error bars denote 95% confidence intervals, and the insets indicate the number of
trials per condition used for analysis. As can be seen, pursuit gain peaked at around 0.8-1 Hz for both
monkeys, and was lower for both lower and higher frequencies. In both animals, pursuit gain seemed high
at 0.1 Hz; at this frequency, the target was barely moving and ocular drift velocities could be substantially
higher than target velocities (Martins et al. 1985). Also, in both animals, pursuit gain was significantly
worse for vertical tracking as opposed to horizontal tracking. (**B**) A similar analysis but now for pursuit
phase (e.g. the phase difference between the red and blue sinusoids of Fig. 1). In monkey M, higher
frequencies were associated with larger phase lags. This dependence of phase lag on frequency was not so
evident in monkey A, for which an overall lag existed even at low frequencies. Note that the phase lag is
displayed here as a fraction of a cycle. Thus, a constant temporal lag would translate into a larger fraction
with increasing frequency (see text). All error bars indicate 95% confidence intervals.

We similarly analyzed pursuit phase lag, reporting it as a fraction of a full cycle (Fig.

2B); that is, a constant temporal delay in phase would mean a larger fraction of a cycle

with increasing frequencies. In monkey M, this was the case: higher target motion

frequencies resulted in progressively more and more pursuit lag when represented as a

fraction of a full cycle. However, this was less so in monkey A, for which there was an

overall lag in pursuit that was independent of frequency. The insets in Fig. 2A show the

total numbers of trials analyzed for each condition.

Therefore, for both horizontal and vertical smooth pursuit eye movements, tracking of

very slow motion trajectories in monkeys is seemingly different from classical

experiments with faster target speeds; for the small-amplitude slow motions, pursuit

efficacy at low frequencies is significantly impaired, and only recovers at ~0.8-1 Hz. For

larger target speeds, evidence from the literature shows that smooth pursuit typically

exhibits low-pass behavior, with high gain at all low frequencies up to ~1 Hz (Collewijn

and Tamminga 1984; Fabisch et al. 2009; Hafed et al. 2008; Hafed and Krauzlis 2008;

Martins et al. 1985; Rottach et al. 1996). As stated above, in highly trained humans, such

a low-pass behavior of smooth pursuit seemed to also still persist for small-amplitude

target trajectories like the ones that we used (Martins et al. 1985); perhaps this difference

from our monkey results is due to extensive training of the humans to avoid making

catch-up saccades.

Controllability of monkey ocular velocities as slow as those during fixational ocular

drifts

Despite the low gain of smooth pursuit at low temporal frequencies in Figs. 1, 2, eye

velocity in our monkeys was still clearly controllable. For example, sinusoidal tracking

was still evident at 0.4 Hz even with the reduced gain (Fig. 1A). As stated earlier, we

were interested in this phenomenon particularly because the velocities with which

tracking occurred at these low temporal frequencies seemed similar to the velocities with

which ocular drifts during fixation normally take place. For example, peak target velocity

at 0.1, 0.2, and 0.3 Hz was 0.314, 0.628, and 0.942 deg/s, respectively, which are all

within the range of eye velocity during fixational ocular drifts (Cherici et al. 2012;

Martins et al. 1985). To confirm this, we collected control fixation data from the same

animals. In these trials, the spot never moved, and the monkeys simply fixated it for

approximately 1000 ms (Materials and Methods). We measured eye velocity during

microsaccade-free fixation epochs, and we related them to the peak eye velocity at each

temporal frequency during tracking (Fig. 3A). Specifically, in Fig. 3A, we plotted the

same data as in Fig. 2A but now as raw measurements of peak eye velocity instead of as

gain values. We then plotted the average velocities observed during gaze fixation

(horizontal lines), again along with 95% confidence intervals. At the lowest and highest

tracking frequencies (e.g. 0.1-0.2 Hz or 4-5 Hz), peak eye velocity during pursuit of slow

motion trajectories was significantly lower than eye velocity during fixation (error bars in

Fig. 3A denote 95% confidence intervals). However, the eye was still well-controlled

because gain was not zero (Fig. 2A). We also further confirmed this by analyzing the

frequency spectrum of eye position traces for different temporal frequencies (particularly

the small ones associated with the slowest peak velocities). We found that there was a

peak in the power spectrum of eye position traces near the frequency with which the

target was moving (Fig. 3B). This means that eye position had a harmonic component

that was close in frequency to that present in the target motion trajectory even at the low

gain values observed in Fig. 2A. Finally, we also inspected example eye position traces

for low (Fig. 3C) or high (Fig. 3D) tracking frequencies, and clear modulation of

saccade-free eye position was still present (e.g. see the gray arrows in Fig. 3C, D).

Therefore, slow ocular movements at velocities similar to or lower than the velocities of

slow fixational ocular drifts are relatively well-controlled in this behavior (Wyatt and

Pola 1981).

Figure 3 Monkey ocular velocities as slow as those during fixational drifts occurred during smooth pursuit of slow motion, but they were still systematically controlled to track the motion trajectory.
(**A**) The curves show the same data as in Fig. 2A, but this time as real measurements of peak eye velocity as opposed to a gain ratio. The solid horizontal lines show average eye velocity during fixation (Materials and Methods), surrounded by 95% confidence intervals (the inset shows an example 1-second fixational eye position trace, demonstrating how ocular drift has substantial non-zero eye velocity even with a stationary fixation spot). Fixational drift velocity was higher than pursuit peak velocities at pursuit frequencies of, say, 0.2 Hz, 0.3 Hz, and 4 Hz. This means that eye velocities as slow as those during ocular drifts are controllable by the central nervous system of the monkey. (**B**) This idea is supported by analyzing the spectral content of eye position traces for different pursuit target frequencies. Even at very low frequencies (e.g. 0.2 Hz and 0.8 Hz) associated with low pursuit velocities (as in **A**), eye position still exhibited a peak in the spectrum at a frequency near target motion. This indicates that eye position was still following target motion (even if at low gain). (**C**, **D**) Sample 1-second eye position traces (shown in gray) from three example trials at 0.2 Hz (**C**) or 3 Hz (**D**), both of which exhibited very weak pursuit gain (**A** and Fig. 2). In both cases, eye position was systematically controlled to follow target trajectory (e.g. see the gray arrows). Thus, monkeys are capable of slow control of their drift velocities in our task.

Dependence of monkey catch-up saccade frequency and amplitude on temporal frequency

Our results so far have focused on smooth eye velocity effects. However, we also

analyzed and catalogued catch-up saccade frequency and amplitude. We found that catch-

up saccades behaved in different ways for temporal frequencies higher or lower than the

frequency associated with peak smooth pursuit gain (~0.8-1 Hz). For example, in Fig.

4A, we plotted the frequency of catch-up saccades as a function of temporal frequency in

Experiment 1. In both animals, catch-up saccade frequency reached a peak at the same

temporal frequency for which smooth pursuit gain was maximum. This suggests that eye

position was continuously adjusted with both smooth pursuit and saccadic eye

movements when overall tracking was particularly effective (i.e. with high gain). Catch-

up saccade frequency then dropped dramatically for higher temporal frequencies, but this

drop was not so dramatic for lower temporal frequencies. This suggests that for the lower

temporal frequencies, when the spot was barely moving, catch-up saccades played a role

more similar to that of fixational microsaccades: they optimized eye position on the target

on average (Guerrasio et al. 2010; Ko et al. 2010; Tian et al. 2018; 2016), especially

when slow eye movements (Fig. 2) were significantly less effective in keeping the eye on

target because of low gain. On the other hand, for very rapid oscillations (high

frequencies), the oculomotor system was unable to keep track of the frequent flips in

target position (even with saccades as opposed to smooth pursuit), and saccades were

therefore more or less random events. Consistent with this, catch-up saccade amplitudes

(Fig. 4B) were always small for all frequencies <1 Hz; on the other hand, catch-up

saccade amplitudes increased for higher frequencies.

Figure 4 Catch-up saccade frequency was highest for frequencies resulting in maximal smooth velocity gain; catch-up saccade amplitude increased for high frequencies. (**A**) For both animals in Experiment 1, we estimated the frequency of catch-up saccades during steady-state smooth pursuit. The highest rate of catch-up saccades occurred at the frequencies associated with maximal smooth velocity gain (Figs. 1-3). For yet higher frequencies, both animals showed a marked decrease in catch-up saccade frequency, suggesting that the reduction in smooth velocity gain associated with these frequencies (Figs. 1-3) was not compensated for by rapid back and forth saccades to try to track the target. (**B**) Instead, catch-up saccades at higher frequencies were apparently random, infrequent movements that were increasingly large in amplitude. These patterns of results (**A** and **B**) were identical for horizontal and vertical pursuit, despite the quantitative difference in smooth velocity gain for these different pursuit directions (Figs. 2-3). All error bars denote 95% confidence intervals.

The above interpretations are also supported by inspecting sample eye position traces

from two example temporal frequencies from Experiment 1 (Fig. 5). Saccades were

frequent at low temporal frequencies, and they kept the eye hovering around target

location, consistent with the role of fixational microsaccades in continuously optimizing

eye position (Guerrasio et al. 2010; Ko et al. 2010; Tian et al. 2018; 2016). On the other

hand, saccades at high temporal frequency were less frequent, large, and very often

increasing eye position error, rather than decreasing it, for substantial amounts of time

(>500 ms).

Figure 5 Monkey catch-up saccades at low temporal frequencies behaved like microsaccades during
fixation. (**A**) Example eye position traces (gray) when tracking a target moving at 0.2 Hz (true target
position is shown by the blue sinusoid). Smooth velocity (i.e. slow ocular drifts) tracked target motion,
albeit at a low gain, and there were plenty of catch-up saccades. Thus, the slow eye movements here
appeared similar to fixational ocular drifts in terms of velocity (e.g. inset in Fig. 3A). Moreover, catch-up
saccades were quite frequent, and resulted in the eye "hovering" around target position on average, as with
fixational microsaccades. (**B**) At high frequencies, slow movements also tracked the target motion (at low
gain). However, this time, catch-up saccades were less frequent, and they were large, often deviating the
eye substantially away from the target, and for substantial periods of time (>500 ms).

It is also interesting to note that despite the large difference in smooth pursuit gain

between horizontal and vertical tracking in Experiment 1 (Fig. 2), catch-up saccade

frequency and amplitude were not at all different from each other across tracking

directions (Fig. 4). This might suggest that there is a larger tolerance for oculomotor

errors along the vertical dimension, perhaps because of potential asymmetries in

oculomotor circuits (Hafed and Chen 2016), although this remains to be just a hypothesis

at the moment.

Over-tracking of oblique small-amplitude slow motion trajectories by monkeys

We also sought to compare the effects of temporal frequency that we observed above to

those of movement amplitude and direction for a given frequency. We therefore

conducted Experiment 2 (Materials and Methods) in which temporal frequency was

pegged at 0.5 Hz but movement amplitude varied between ~15 min arc (0.25 deg) and ~2

deg (Materials and Methods). Movement direction also included oblique tracking

(Materials and Methods). Overall, we found expected results in terms of smooth pursuit

gain as a function of target motion trajectory amplitude. For example, for both cardinal

(horizontal and vertical) and oblique pursuit, smooth pursuit gain increased with

increasing target motion amplitude (Fig. 6). This might explain why our results from

Experiment 1 above showed band-pass behavior (Fig. 2) when the target amplitude was

small; this band-pass effect was primarily due to the very small motion trajectory

amplitudes (and, correspondingly, velocities) used when compared to classic studies, and

increasing amplitude in the current experiment has alleviated this.

Figure 6 Higher pursuit gain for tracking of small-amplitude oblique motion trajectories. (**A**) In Experiment 2, we varied target sinusoidal position amplitude, and also introduced oblique pursuit trajectories. For cardinal directions, we obtained expected results based on what we found in Experiment 1. Pursuit gain was higher for horizontal than vertical pursuit, and pursuit gain progressively increased with increasing target amplitude. Note, however, that gain seemed to asymptote near a value of 1. (**B**) For oblique pursuit directions, there was still an increase in pursuit gain with target amplitude. However, this increase eventually resulted in an overshoot of target speeds (gain going substantially >1). Even at 0.5 deg amplitudes, oblique pursuit showed higher gain than cardinal pursuit (compare the 0.5 deg point in this panel to the corresponding point in panel **A**). Thus, oblique pursuit was associated with overshooting of target velocity in both animals. All error bars denote 95% confidence intervals.

The most striking result from Experiment 2, which is also evident in Fig. 6, was that

smooth pursuit gain was significantly >1 for oblique tracking directions (Fig. 6B). This is

different from pursuit of large-amplitude motions, in which oblique pursuit might be

expected to be worse in gain than pursuit along cardinal directions, perhaps due to

oblique effects in motion perception (Krukowski and Stone 2005). When we analyzed the

horizontal and vertical components of oblique pursuit independently, we found that the

"overshoot" was primarily occurring in the horizontal dimension (Fig. 7). These results

are reminiscent of overshoot in visually-guided saccade amplitudes in humans for very

small retinal eccentricities (Kalesnykas and Hallett 1994), and they demonstrate an

additional difference in the control capabilities of the slow movement oculomotor sub-

system between small and large motion amplitudes. For example, in large-amplitude

pursuit, horizontal gain can be higher than vertical gain during oblique tracking (Rottach

et al. 1996), but overall gain still always stays below or very close to 1. We found marked

increases above a gain value of 1.

**Figure 7 Over-tracking in the horizontal dimension during smooth pursuit of small-amplitude
oblique motion trajectories.** We investigated the overshoot observed in Fig. 6B by plotting the horizontal
and vertical components of eye velocity separately for an example oblique pursuit condition. In both
animals, pursuit gain was higher in the horizontal component of eye velocity than the vertical component.
Thus, the over-tracking observed in Fig. 6B was primarily driven by over-tracking in the horizontal
dimension. All error bars denote 95% confidence intervals.

The over-tracking of oblique motion trajectories was also accompanied by larger catch-

up saccades for these trajectories. In Fig. 8, we plotted catch-up saccade frequency (Fig.

8A) and amplitude (Fig. 8B), as we had done earlier for Experiment 1. Consistent with

Experiment 1, increased smooth pursuit gain was associated with an increase in catch-up

saccade frequency (Fig. 8A), suggesting synergistic interactions between smooth pursuit

eye movements and saccades to optimize eye position on the target (de Brouwer et al.

2002); this is similar to us seeing the most catch-up saccades in Experiment 1 for the

temporal frequencies in which smooth velocity gain was also at a maximum. In terms of

catch-up saccade amplitude, it also increased with increasing target motion amplitude; in

addition, while an increase in saccade amplitude was expected with increasing target

position trajectory amplitude, the increase was stronger for oblique directions (Fig. 8B).

For example, in monkey A, both oblique directions had higher saccade amplitudes than

horizontal or vertical trajectories, and in monkey M, one of the oblique directions did.

This result might reflect the slightly higher peak velocities associated with oblique

tracking (for example, oblique tracking with a horizontal and vertical amplitude of 2 deg

each meant an overall radial amplitude of 2.82 deg; Materials and Methods).

Figure 8 Catch-up saccade frequency and amplitude in Experiment 2. (**A**) Consistent with Experiment
1, when smooth velocity gain was high (Figs. 6-7), catch-up saccade frequency was high. (**B**) Catch-up
saccade amplitude also expectedly increased with larger target motion excursions. All error bars denote
95% confidence intervals.

Dependence of monkey first catch-up saccade latency and smooth pursuit initiation on

target motion

We also analyzed the properties of the very first catch-up saccade during smooth pursuit

initiation, as well as the smooth component of initial eye acceleration itself. For all

saccades (including fixational microsaccades) occurring in the interval 0-300 ms after

target motion onset, we plotted these movements' amplitudes as a function of their

occurrence time. We also plotted either baseline eye velocity (in the 50 ms interval

starting at -100 ms from target motion onset) or smooth pursuit initiation eye velocity (in

the 50 ms starting 100 ms after target motion onset; ensuring no saccades within each

interval). We observed expected relationships between initial catch-up saccades and

initial smooth pursuit eye velocity. For example, in Experiment 2, with 0.5 Hz target

trajectory variation, eye position error of the target (relative to initial fixation location if

the eye did not start tracking) monotonically increased in the first 300 ms of any trial (and

up to 500 ms). Therefore, if a saccade were to occur during initiation (and there was no

associated smooth acceleration after motion onset), then saccade amplitude was expected

to increase with increasing time after motion onset. However, this was not the case for

most target amplitudes that we tested (Fig. 9A). We think that this is so because of a

concomitant increase in smooth eye velocity to track the target (Fig. 9A, B; only

horizontal tracking data is shown for clarity). In other words, after motion onset, the eye

often started to accelerate smoothly, therefore already reducing eye position error. Such

reduction may have alleviated the need to increase first catch-up saccade amplitude. Only

when target position amplitude was large enough (2 deg) did there arise a need for

increasing initial catch-up saccade amplitude. For such a larger position amplitude of the

motion trajectory, even the initial component of smooth pursuit acceleration was not

sufficient to reduce eye position error sufficiently; a larger saccade was therefore

necessary. This idea is illustrated in Fig. 9C showing raw pursuit velocities with saccades

excised from the averages as per Materials and Methods (shown from horizontal tracking

only for clarity). Pursuit initiation velocity increased for trajectories with amplitudes of 2

deg compared to, say, 0.5 deg. However, the increase in velocity did not necessarily

allow for completely eliminating eye position error, resulting in the need for an initial

catch-up saccade whose amplitude gradually increased with increasing time after motion

onset (Fig. 9A; 2 deg motion position amplitude). Therefore, there was a synergistic

interaction between smooth pursuit initiation and initial catch-up saccade execution,

which also likely occurs between slow ocular drifts and microsaccades (Chen and Hafed

2013). Note that in Fig. 9A, saccades occurring <50-60 ms after target motion onset were

fixational microsaccades and not really target-driven catch-up saccades because they

occurred too early to reflect the new visual error signal introduced by target motion onset.

This is why these movements were also small in amplitude even for 2 deg motion

amplitudes.

Figure 9 Interactions between saccades and smooth pursuit eye movements during initiation of tracking. (**A**) We found evidence of synergistic interactions between smooth pursuit and catch-up saccades during initiation. The amplitude of the first catch-up saccade after target motion onset did not increase convincingly as a function of time (even though eye position error was monotonically increasing), except for 2 deg amplitude trajectories (dark purple). (**B**) This might be because saccade-free smooth pursuit initiation may have already acted to reduce eye position error. For example, initial eye velocity during saccade-free initiation (see pink measurement intervals in the left panel of **C**) increased with increasing target amplitude. Thus, saccades did not necessarily need to increase in amplitude in **A** for small target amplitudes. For 2 deg trajectories, the increase in smooth eye velocity may have not been sufficient to completely obviate the need for a catch-up saccade, and that is why catch-up saccade amplitudes in this particular condition increased with time from motion onset when compared to all of the other conditions that we tested. Note that in **A**, saccades with latencies less than ~50-60 ms were likely not initial catch-up saccades but instead fixational microsaccades. That is why their amplitudes were low for all conditions. (**C**) Example average saccade-free smooth velocity traces from horizontal tracking (like **B**) showing the measurement intervals for **B** and also the idea that smooth velocity effects increased with increasing target amplitude. All error bars denote 95% confidence intervals.

Finally, we checked whether the first catch-up saccade latency itself depended on target

amplitude. In Experiment 2, we plotted histograms of first catch-up saccade latency in the

different amplitude conditions (Fig. 10). For the smallest amplitude trajectories, catch-up

saccade latencies were long and variable. As target amplitude increased, latency became

more precise, as well as shorter. Since Fig. 9 showed that it was more likely for catch-up

saccade amplitudes to increase with increasing target position amplitude, these results

constitute a monkey replication of human studies, showing that saccade latencies are

substantially longer for very small-amplitude visually-guided saccades compared to

larger ones (Kalesnykas and Hallett 1994).

Figure 10 Dependence of first catch-up saccade latency on target amplitude. We plotted histograms of
first catch-up saccade latency for different target amplitudes in Experiment 2. We show results only from
horizontal tracking for clarity; other directions showed similar effects. Since larger target amplitudes
resulted in larger catch-up saccades (e.g. Fig. 9), the present histograms show monkey replication of human
observations that very small visually-guided saccades are associated with longer reaction times than larger
ones (Kalesnykas and Hallett 1994). For example, for each monkey, the red dashed line marks the latency
bin with most observations at 2 deg target motion amplitude (bottom panel), and the magenta dashed line
marks the latency bin with the most observations at 0.25 deg target motion amplitude (top panel). As can be
seen, there was a substantial differential in saccade times for the different target motion amplitudes. Note
that, like in Fig. 9, the small distribution of saccades occurring with latencies <100 ms in this figure are
likely not genuine first catch-up saccades, but instead fixational microsaccades. This is further supported by
the noticeable dip in the histograms at ~100 ms, which is similar to the phenomenon of microsaccadic
inhibition reported in the literature for both humans and monkeys (Buonocore et al. 2017; Hafed and
Ignashchenkova 2013; Rolfs et al. 2008).

Discussion

We attempted to characterize monkey oculomotor behavior with small-amplitude slow motion trajectories. We catalogued both smooth velocity effects as well as catch-up saccade effects. For smooth velocity effects, we found that pursuit gain was low for both low and high temporal frequencies, only reaching a peak in mid-frequencies near 1 Hz. This is in contrast to previous human experiments with large-amplitude (Collewijn and Tamminga 1984; Fabisch et al. 2009; Rottach et al. 1996) or small-amplitude (Martins et al. 1985) sinusoidal motions, in which low-pass behavior was observed. In terms of catch-up saccades, we found that they (perhaps paradoxically) increased in frequency when smooth velocity gain was high. Moreover, we found that catch-up saccades during presentation of low temporal frequencies acted more like fixational microsaccades, whereas catch-up saccades during presentation of high temporal frequencies were large and infrequent.

Our results provide a necessary foundation for exploring the neural mechanisms subserving fixational ocular drifts in awake monkeys. This complements early characterizations of awake monkey smooth pursuit eye movements with higher speeds (Lisberger and Westbrook 1985). These early characterizations were themselves a major boon for a wide range of significant and seminal subsequent discoveries about the neural mechanisms for oculomotor control in general, and about the neural mechanisms for smooth pursuit in particular (Krauzlis 2004). Our next goal is to extend on our current results by uncovering neural substrates in the same animals.

To that end, we were careful to avoid unnecessarily penalizing the monkeys for making

saccades during tracking. Specifically, we aimed to minimize over-training on one

particular movement modality. For example, early human studies with small-amplitude

motions barely had any saccades in the experiments, to focus almost solely on slow

control effects (Martins et al. 1985). However, we wanted the animals to engage in as

naturalistic a behavior as possible, such that we could understand important interactions

between slow control and micro/saccadic control. This has allowed us to make the

interesting observation that smooth pursuit gain in our monkeys exhibited band-pass

behavior, unlike in (Martins et al. 1985). This has also allowed us to make the interesting

observation that there were actually more saccades when pursuit gain was high (i.e. when

retinal position and motion errors were presumably low) than when it was low.

Such an observation of a concomitant increase in catch-up saccade frequency along with

an increased velocity gain might suggest that catch-up saccades normally behave like

fixational microsaccades. The latter eye movements continuously re-align gaze with a

foveal target under a variety of stimulus conditions (Guerrasio et al. 2010; Ko et al.

2010), and even when competing peripheral stimuli are presented (Tian et al. 2018;

2016). Their frequency of occurrence might therefore increase when the target is sharp,

providing a clear spatial reference frame for re-aligning gaze. During pursuit, similar re-

alignment of gaze may be necessary. This is consistent with observations that foveal

targets increase catch-up saccade frequency in humans (Heinen et al. 2018; Heinen et al.

2016); presumably, foveal targets not only support good smooth velocity gain, but they

also provide the oculomotor system with a punctate spatial reference point to which gaze

can be re-directed. We also found in monkeys that there was a tendency for higher catch-

up saccade frequencies for smaller foveal pursuit targets than for bigger and fuzzier ones

(Hafed et al. 2008). It would be interesting to analyze the relationship between eye

position error and catch-up saccade likelihood with a foveal target during smooth pursuit,

such that one can uncover an almost-deterministic estimate of whether a catch-up saccade

can occur at any one moment of time or not (de Brouwer et al. 2002). This kind of

approach was recently made for microsaccades (Tian et al. 2018), and it is very intriguing

because predicting whether and when a microsaccade might take place can then, at least

in principle, be used to estimate the occurrence of distinct cognitive performance effects

associated with such movements (Bellet et al. 2017; Chen et al. 2015; Hafed 2013).

Our interest in relying on more naturalistic tracking (i.e. with combined smooth and

saccadic eye movements) may also explain why we observed band-pass smooth pursuit

gain effects in our monkeys, even though a similar human experiment with low

frequencies and small-amplitude trajectories found very good gain (Martins et al. 1985).

As stated earlier, in that study, the human subjects tested were highly trained, and they

were instructed to minimize saccade generation. As a result, substantial systematic eye

position drifts occurred in their experiments, whereas we did not observe such systematic

drifts. In our case, we relied more on the natural behavior of the monkeys in being

intrinsically interested to foveate the small white spot that was presented on the display.

Yes, the monkeys became highly trained in the lab after multiple sessions, but their

behavior was not shaped by us, say, by aborting trials whenever a saccade occurred.

Instead, we rewarded them for tracking the target to within a reasonable radius, which

may not be too different from natural variability in human fixation among untrained

individuals (Cherici et al. 2012). As a result, our monkeys tracked the target with both

smooth and saccadic eye movements.

We were also intrigued by our oblique tracking effects in Experiment 2. Both smooth

velocity gain and catch-up saccade amplitudes increased for oblique tracking, with

smooth velocity gain surpassing a value of 1 by quite a margin. While the catch-up

saccade amplitude increase may have reflected the slightly faster trajectories associated

with oblique target motions in our stimulus design relative to cardinal target motions

(Materials and Methods), the smooth velocity gain measure is a normalized measure that

should not depend on absolute target velocity. Instead, it does reflect a real observation

that pursuit in these oblique conditions substantially over-tracked the small-amplitude

motions that we tested. This is intriguing to us because small visually-guided saccades in

humans are known to overshoot their targets (Kalesnykas and Hallett 1994). We may

have thus observed a similar phenomenon for smooth pursuit, adding to evidence that

pursuit and saccades share neural resources (Krauzlis and Dill 2002; Krauzlis 2004;

Krauzlis et al. 1997; Krauzlis et al. 2017). Similarly, even our catch-up saccade effects of

Fig. 10 demonstrate a monkey correlate of human observations that visually-guided

saccade latency increases for small target eccentricities. It would be interesting to extend

these effects for other types of saccades, like delayed visually- or memory-guided

saccades.

In all, our results add to an extensive cataloguing in the literature of rhesus macaque

sensory, perceptual, cognitive, and motor capabilities, testifying to the tremendous value

of such an animal model for systems neuroscience research.

Acknowledgments

We were funded by the Werner Reichardt Centre for Integrative Neuroscience (CIN) at

the Eberhard Karls University of Tübingen. The CIN is an Excellence Cluster funded by

the Deutsche Forschungsgemeinschaft (DFG) within the framework of the Excellence

Initiative (EXC 307). We were also funded by the DFG through a Research Unit (FOR

1847). ZMH was additionally funded by the Hertie Institute for Clinical Brain Research

at the Eberhard Karls University of Tübingen.

References

Arnstein D, Junker M, Smilgin A, Dicke PW, and Thier P. Microsaccade control
signals in the cerebellum. *J Neurosci* 35: 3403-3411, 2015.

Bahill AT, and McDonald JD. Smooth pursuit eye movements in response to
predictable target motions. *Vision Res* 23: 1573-1583, 1983.

Barlow HB. Eye movements during fixation. *J Physiol* 116: 290-306, 1952.

Bellet J, Chen CY, and Hafed ZM. Sequential hemifield gating of alpha and beta
behavioral performance oscillations after microsaccades. *J Neurophysiol* jn 00253 02017,
2017.

Bellet ME, Bellet J, Nienborg H, Hafed ZM, and Berens P. Human-level saccade
detection performance using deep neural networks. *bioRxiv* 2018.

Brainard DH. The Psychophysics Toolbox. *Spatial vision* 10: 433-436, 1997.

Buonocore A, Chen CY, Tian X, Idrees S, Muench T, and Hafed ZM. Alteration of
the microsaccadic velocity-amplitude main sequence relationship after visual transients:
implications for models of saccade control. *J Neurophysiol* jn 00811 02016, 2017.

Chen CY, and Hafed ZM. Postmicrosaccadic enhancement of slow eye movements. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience 33: 5375-
5386, 2013.

Chen CY, Ignashchenkova A, Thier P, and Hafed ZM. Neuronal Response Gain
Enhancement prior to Microsaccades. *Curr Biol* 25: 2065-2074, 2015.

Cherici C, Kuang X, Poletti M, and Rucci M. Precision of sustained fixation in trained
and untrained observers. *Journal of vision* 12: 2012.

Collewijn H, and Tamminga EP. Human smooth and saccadic eye movements during
voluntary pursuit of different target motions on different backgrounds. *J Physiol* 351:
217-250, 1984.

Cunitz RJ. Relationship Between Slow Drift and Smooth Pursuit Eye Movements (Ph.D.
Dissertation). University of Maryland, 1970.

de Bie J, and van den Brink G. A model for the slow control system during monocular
fixation. *Vision Res* 26: 1129-1142, 1986.

de Brouwer S, Yuksel D, Blohm G, Missal M, and Lefevre P. What triggers catch-up
saccades during visual tracking? *Journal of neurophysiology* 87: 1646-1650, 2002.

Fabisch K, Fitz W, Fabisch H, Haas-Krammer A, Klug G, Zapotoczky S, and
Kapfhammer HP. Sinusoidal smooth pursuit eye tracking at different stimulus
frequencies: position error and velocity error before catch-up saccades in schizophrenia
and in major depressive disorder. *Aust N Z J Psychiatry* 43: 855-865, 2009.

Fuchs AF, and Robinson DA. A method for measuring horizontal and vertical eye
movement chronically in the monkey. *J Appl Physiol* 21: 1068-1070, 1966.

Guerrasio L, Quinet J, Buttner U, and Goffart L. Fastigial oculomotor region and the
control of foveation during fixation. *J Neurophysiol* 103: 1988-2001, 2010.

Hafed ZM. Alteration of visual perception prior to microsaccades. *Neuron* 77: 775-786,
2013.

Hafed ZM. Mechanisms for generating and compensating for the smallest possible
saccades. *Eur J Neurosci* 33: 2101-2113, 2011.

Hafed ZM, and Chen C-Y. Sharper, stronger, faster upper visual field representation in
primate superior colliculus. *Current Biology*
http://dx.doi.org/10.1016/j.cub.2016.04.059: 2016.

Hafed ZM, Chen C-Y, and Tian X. Vision, perception, and attention through the lens of
microsaccades: mechanisms and implications. *Frontiers in systems neuroscience* 9: 167,
2015.

Hafed ZM, Goffart L, and Krauzlis RJ. A neural mechanism for microsaccade
generation in the primate superior colliculus. *Science* 323: 940-943, 2009.

Hafed ZM, Goffart L, and Krauzlis RJ. Superior colliculus inactivation causes stable
offsets in eye position during tracking. *J Neurosci* 28: 8124-8137, 2008.

Hafed ZM, and Ignashchenkova A. On the dissociation between microsaccade rate and
direction after peripheral cues: microsaccadic inhibition revisited. *J Neurosci* 33: 16220-
16235, 2013.

Hafed ZM, and Krauzlis RJ. Goal representations dominate superior colliculus activity
during extrafoveal tracking. *J Neurosci* 28: 9426-9439, 2008.

Hafed ZM, and Krauzlis RJ. Similarity of superior colliculus involvement in
microsaccade and saccade generation. *J Neurophysiol* 107: 1904-1916, 2012.

Heinen SJ, Badler JB, and Watamaniuk SNJ. Choosing a foveal goal recruits the
saccadic system during smooth pursuit. *J Neurophysiol* 2018.

Heinen SJ, Potapchuk E, and Watamaniuk SN. A foveal target increases catch-up
saccade frequency during smooth pursuit. *J Neurophysiol* 115: 1220-1227, 2016.

Judge SJ, Richmond BJ, and Chu FC. Implantation of magnetic search coils for
measurement of eye position: an improved method. *Vision Res* 20: 535-538, 1980.

Kalesnykas RP, and Hallett PE. Retinal eccentricity and the latency of eye saccades.
Vision Res 34: 517-531, 1994.

Kleiner M, Brainard D, and Pelli DG. What's new in Psychtoolbox-3? (Abstract).
Perception 36: 2007.

Ko HK, Poletti M, and Rucci M. Microsaccades precisely relocate gaze in a high visual
acuity task. *Nat Neurosci* 13: 1549-1553, 2010.

Krauzlis R, and Dill N. Neural correlates of target choice for pursuit and saccades in the
primate superior colliculus. *Neuron* 35: 355-363, 2002.

Krauzlis RJ. Recasting the smooth pursuit eye movement system. *J Neurophysiol* 91:
591-603, 2004.

Krauzlis RJ, Basso MA, and Wurtz RH. Shared motor error for multiple eye
movements. *Science* 276: 1693-1695, 1997.

Krauzlis RJ, Goffart L, and Hafed ZM. Neuronal control of fixation and fixational eye
movements. *Philos Trans R Soc Lond B Biol Sci* 372: 2017.

Krauzlis RJ, and Miles FA. Release of fixation for pursuit and saccades in humans:
evidence for shared inputs acting on different neural substrates. *J Neurophysiol* 76: 2822-
2833, 1996.

Krukowski AE, and Stone LS. Expansion of direction space around the cardinal axes
revealed by smooth pursuit eye movements. *Neuron* 45: 315-323, 2005.

Lisberger SG, and Westbrook LE. Properties of visual inputs that initiate horizontal
smooth pursuit eye movements in monkeys. *J Neurosci* 5: 1662-1673, 1985.

Martins AJ, Kowler E, and Palmer C. Smooth pursuit of small-amplitude sinusoidal
motion. *J Opt Soc Am A* 2: 234-242, 1985.

McFarland JM, Cumming BG, and Butts DA. Variability and Correlations in Primary
Visual Cortical Neurons Driven by Fixational Eye Movements. *J Neurosci* 36: 6225-
6241, 2016.

Murphy BJ, Kowler E, and Steinman RM. Slow oculomotor control in the presence of
moving backgrounds. *Vision research* 15: 1263-1268, 1975.

Nachmias J. Determiners of the drift of the eye during monocular fixation. *J Opt Soc Am*
51: 761-766, 1961.

Ohashi N, and Mizukoshi K. Analysis of smooth pursuit by the fast Fourier transform.
Acta Otolaryngol Suppl 481: 348-351, 1991.

Peel TR, Hafed ZM, Dash S, Lomber SG, and Corneil BD. A Causal Role for the
Cortical Frontal Eye Fields in Microsaccade Deployment. *PLoS Biol* 14: e1002531, 2016.

Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers
into movies. *Spatial vision* 10: 437-442, 1997.

Rolfs M. Microsaccades: small steps on a long way. *Vision Res* 49: 2415-2441, 2009.

Rolfs M, Kliegl R, and Engbert R. Toward a model of microsaccade generation: the
case of microsaccadic inhibition. *J Vis* 8: 5 1-23, 2008.

Rottach KG, Zivotofsky AZ, Das VE, Averbuch-Heller L, Discenna AO,
Poonyathalang A, and Leigh RJ. Comparison of horizontal, vertical and diagonal
smooth pursuit eye movements in normal human subjects. *Vision Res* 36: 2189-2195,
1996.

Rucci M, and Victor JD. The unsteady eye: an information-processing stage, not a bug.
Trends Neurosci 2015.

Sun Z, Junker M, Dicke PW, and Thier P. Individual neurons in the caudal fastigial
oculomotor region convey information on both macro- and microsaccades. *Eur J*
Neurosci 44: 2531-2542, 2016.

Tian X, Yoshida M, and Hafed ZM. Dynamics of fixational eye position and
microsaccades during spatial cueing: the case of express microsaccades. *J Neurophysiol*
119: 1962-1980, 2018.

Tian X, Yoshida M, and Hafed ZM. A Microsaccadic Account of Attentional Capture
and Inhibition of Return in Posner Cueing. *Frontiers in systems neuroscience* 10: 23,
2016.

Van Horn MR, and Cullen KE. Coding of microsaccades in three-dimensional space by
premotor saccadic neurons. *J Neurosci* 32: 1974-1980, 2012.

Veale R, Hafed ZM, and Yoshida M. How is visual salience computed in the brain?
Insights from behaviour, neurobiology and modelling. *Philos Trans R Soc Lond B Biol*
Sci 372: 2017.

Wyatt HJ, and Pola J. Slow eye movements to eccentric targets. *Invest Ophthalmol Vis*
Sci 21: 477-483, 1981.
917
918