1

A total synthetic approach to CRISPR/Cas9 genome editing and homology

directed repair

Sara E. DiNapoli[1,2] [*], Raul Martinez-McFaline[1,2] [*], Caitlin K. Gribbin[1,2*], Courtney A.

Balgobin[1,2], Isabel Nelson[1,2], Abigail Leonard[1,2], Darya Mailhiot[4], Clara Kao[5], Sean

McDonnell[5], Jill L.O. de Jong[5], Yariv Houvras[1,2,3] [**]

1. Department of Surgery, Weill Cornell Medical College, New York Presbyterian

Hospital, New York, NY, USA

2. Meyer Cancer Center, Weill Cornell Medical College, New York Presbyterian

Hospital, New York, NY, USA

3. Department of Medicine, Weill Cornell Medical College, New York Presbyterian

Hospital, New York, NY, USA

4. Department of Surgery, Animal Resources Center, University of Chicago,

Chicago, IL, USA

5. Department of Pediatrics, University of Chicago, Chicago, IL, USA

* These authors contributed equally to this work.

** Correspondence and requests for materials should be addressed to Y.H. (email:

yah9014@med.cornell.edu).

ABSTRACT

CRISPR/Cas9 has become a powerful tool for genome editing in zebrafish that permits the rapid generation of loss of function mutations and the knock-in of specific alleles using DNA templates and homology directed repair (HDR). We directly compared synthetic, chemically modified sgRNAs to *in vitro* transcribed sgRNAs and demonstrate the increased activity of synthetic sgRNAs. We harnessed the efficacy of synthetic sgRNAs to generate large chromosomal deletions over 100kb in size. By developing an *in vivo* genetic assay to measure HDR efficiency, we systematically examined the characteristics of synthetic DNA templates to promote HDR. Utilizing these principles, we successfully performed precise knock-in of fluorophores at multiple genomic loci. Collectively, our studies demonstrate the utility of enhanced synthetic sgRNA activity for a wide range of applications *in vivo* including precise homology directed repair and genome editing.

INTRODUCTION

CRISPR/Cas9 has been utilized for a wide range of experimental applications, and zebrafish has been a key model organism to test and validate strategies for genome editing[1,2]. Repair of CRISPR-generated double-stranded breaks (DSBs) by non-homologous end joining (NHEJ) leads to insertions and deletions (indels) which may result in loss of function of the targeted gene product. By supplying an exogenous DNA template, DSBs are repaired through homology directed repair (HDR), allowing for precision genome editing, including base pair changes and insertion of protein tags. Although HDR has been reported in zebrafish, the efficiency varies widely and has required the construction of vectors to serve as a DNA template[3–7]. While diverse approaches and molecular tools have been reported to improve HDR efficiency in mammalian systems, a comprehensive assessment of strategies for genome editing in zebrafish has not been undertaken.

Recent reports have described an increased efficiency of CRISPR/Cas9 targeting using

chemically modified synthetic sgRNAs in human cells[8,9]. We systematically evaluated

synthetic sgRNAs and find they dramatically outperform conventional *in vitro* transcribed

sgRNAs. Synthetic sgRNAs lead to increased targeting, an expanded range of indels,

and allow the generation of large scale deletions in the genome. We developed an

assay for HDR in zebrafish utilizing an *mitfa* mutant, b692. This assay allowed us to

quantitatively compare multiple strategies for improving HDR efficiency and correlate

phenotypic and molecular efficiency. We demonstrate that synthetic reagents, including

linear dsDNA templates, can effectively be used to perform precise genomic editing

including generation of in-frame gene fusions with fluorophores.

RESULTS

Synthetic sgRNAs outperform laboratory synthesized sgRNAs

In order to directly compare editing efficiencies generated by synthetic sgRNA versus *in*

vitro transcribed sgRNA (IVT sgRNA), we targeted tyrosinase (*tyr*), a gene required for

melanin synthesis. Wild type zebrafish embryos were microinjected at the one-cell stage

with recombinant Cas9 protein (rCas9) complexed with either synthetic sgRNA (IDT,

Alt-R®) or with laboratory synthesized IVT sgRNA. Embryos were scored for

melanocyte number at 48 hours post-fertilization (hpf) and divided into phenotypic

categories representing the degree of gene editing (Figure 1b). We find that the

synthetic sgRNA leads to a significantly increased fraction of embryos with fewer

melanocytes as compared with IVT sgRNA (Figure 1a). An increasing dose of synthetic

sgRNA results in an increase in the fraction of phenotypically edited embryos. Only

synthetic sgRNAs were capable of producing large numbers of embryos with high

phenotype gene editing. We found comparable differences between synthetic and IVT

sgRNAs at a second locus, *slc24a5* (Supplementary Figure 1). In several cases

injection of synthetic sgRNAs leads to highly penetrant phenotypes in F_0 larvae that

phenocopy established germline mutants[10] (Supplementary Figure 2). In order to

characterize the molecular alterations arising in synthetic sgRNA injected embryos, we

performed a clonal analysis. We find that 10/10 clones from a synthetic sgRNA injected

high-phenotype embryo contain indels, as compared with 7/10 clones from an IVT

sgRNA injected high-phenotype embryo. An analysis of indel spectrum and distribution

reveals larger deletions in synthetic sgRNA injected embryos (Figure 1c). Taken

together, these data indicate that synthetic sgRNAs lead to penetrant phenotypes that

are associated with a broad spectrum of gene-specific edits.

To assess the extent of germline transmission of edited alleles, we applied CRISPR-

STAT fluorescence-based genotyping to F_0 founders and progeny[11]. Sperm samples

were collected from F_0 founders injected with rCas9 and synthetic sgRNA targeting

slc6a15 at the one-cell stage. Results reveal mutant alleles containing indels as the

major peaks, with additional minor peaks corresponding to wildtype and mutant alleles,

indicating a high efficiency of CRISPR-induced mutagenesis in the germline (Figure 2a).

Five distinct alleles were transmitted to F_1 progeny (Figures 2b-f). The frequencies of

observed CRISPR-induced indels in F_1 progeny were similar to those found in the

sperm of the F_0 founder (Supplementary Figure 3). These findings are consistent with

high levels of somatic mosaicism and confirm that F_0 sperm samples accurately predict

the specific mutations transmitted to F_1 progeny. These data demonstrate that CRISPR-

induced indels created with synthetic sgRNAs are found in the germline of founders and

are efficiently transmitted to F_1 progeny.

Synthetic sgRNA pairs produce large chromosomal deletions

We assessed whether the enhanced efficiency of synthetic sgRNAs would enable the

generation of large chromosomal deletions. We sought to examine the efficiency of

synthetic sgRNAs to produce deletions in a region of noncoding DNA (Figure 3a). We

designed two sgRNAs situated 453bp apart and co-injected with rCas9. We assayed

injected embryos using deletion-specific PCR. We observed the presence of a deleted

allele in 5/10 (50%) embryos (Supplementary Figure 4a). Sequencing of an amplicon

revealed a deletion of 427 nucleotides (Figure 3B). We co-injected rCas9 with sgRNAs

situated 4,286bp apart. By deletion-specific PCR, we identified the presence of a

deletion in 7/24 (29%) embryos (Supplementary Figure 4c). Deletions between 4297

and 4451 nucleotides were confirmed by Sanger sequencing (Figure 3d and

Supplementary Figure 4d). We tested our ability to delete larger regions by co-injecting

rCas9 with additional sgRNAs. We observed deletions of 52kb in 3/8 (38%) embryos by

PCR and confirmed a deletion of 52, 262 nucleotides by Sanger sequencing

(Supplementary Figure 4e,f and Figures 3e). Using sgRNAs spanning the entire non-

coding region we generated a deletion of 124KB in 2/23 (9%), which was confirmed by

Sanger sequencing (Supplementary Figure 4g,h and Figure 3f). We find that pairs of

synthetic sgRNAs are highly efficient in causing large scale deletions in zebrafish, and

that the efficiency varies with size (Figure 3a).

A genetic assay for homology-directed repair

To develop a reliable assay for detecting HDR in zebrafish embryos, we sought to target

a locus that could provide phenotypic and molecular evidence of DNA template

integration. We initially examined the widely used *mitfa*(w2) mutant, which lacks

melanocytes[12]. Unexpectedly, injection of rCas9 complexed with a synthetic sgRNA

targeting the mutation site in w2 resulted in restoration of pigmented melanocytes

(Supplementary Figure 5). We presume that indels caused by targeting the *mitfa*(w2)

mutation site (Q113Stop) restore protein function since this region of the protein is

relatively unstructured. We next examined the *mitfa*(b692) which harbors a mutation

leading to a Ile215Ser substitution and lacks melanocytes[13]. We find that a synthetic

sgRNA targeting the *mitfa*(b692) mutation site does not restore pigmented melanocytes

(Supplementary Figure 5). Therefore, we utilized *mitfa*(b692) to examine a broad range

of synthetic reagents and approaches to achieve precise HDR.

We performed an *in vivo* screen to compare the efficiency of HDR across multiple

conditions. *mitfa*(b692) embryos were microinjected with rCas9, synthetic sgRNA, and a

matrix of synthetic templates designed to restore the wildtype *mitfa* sequence and gene

function. Microinjected embryos were scored at 48hpf for phenotypic evidence of

melanocytes, indicative of template integration and mitfa function (Figure 4B).

Melanocytes were not identified in uninjected embryos or in embryos injected with non-

targeting sgRNA (Figure 4b). HDR templates were designed to feature multiple

barcoded nucleotides and mutated PAM recognition sequence, enabling unambiguous

detection by sequencing and preventing re-cleavage by rCas9 (Figure 4a). Melanocyte

positive embryos were subject to allele-specific PCR and Sanger sequencing to confirm

precise HDR (Figures 4c and d).

We found that a 951bp, double-stranded linear DNA template with an asymmetric

sgRNA site located at the 3' end leads to a reproducible 9% rate of phenotypic rescue in

mitfa(b692) embryos (Figure 4a and e). We examined DNA templates that differ in

length, single vs. double stranded DNA, linear vs. circular templates, and the symmetry

of the sgRNA position within the template (Figure 4e). We find that both longer (2kb)

and shorter (318bp, 76bp) DNA templates led to a decrease in HDR efficiency.

Asymmetric positioning of sgRNA sites led to higher efficiency than symmetric

positioning. Our data indicates that HDR efficiency is maximal with dsDNA templates

and is sensitive to template length and symmetry of the sgRNA site.

Using the b692 assay we examined multiple reagents and conditions reported to

improve HDR efficiency in other model systems (Figure 4e and Supplementary Figure

6). The addition of a second sgRNA site to the optimal 951bp template reduced HDR

efficiency to < 1%. We cloned the 951bp template into a TopoTA plasmid vector and

microinjected a circular plasmid into zebrafish. This plasmid-based template also failed

to lead to measurable HDR in b692 embryos. In light of prior studies reporting increased

HDR efficiency with the use of catalytically-mutated Cas9 (nickase Cas9)[14], we altered

the 951bp DNA template to allow us to test this approach. Microinjection of paired

sgRNAs and recombinant nickase Cas9 failed to lead to measurable HDR in b692

embryos. Microinjection of ssDNA templates and small molecule poly (ADP-ribose)

polymerase (PARP) inhibitors also failed to lead to measurable HDR efficiency. A

complete summary of the tested reagents and phenotypic editing efficiency is

summarized (Supplementary Figure 6).

Melanocyte restoration in b692 assay correlates with molecular efficiency

Using the b692 assay we sought to measure the molecular efficiency of genome editing

and correlate DNA template integration with melanocyte number. To this end, we

performed single-cell injections in a large cohort of b692 embryos using the optimized

reagents described: rCas9, synthetic b692 sgRNA, and 951bp linear dsDNA template.

Embryos were scored for the presence of melanocytes at 48hpf and binned by

phenotype into four categories: high, medium, low, and no-editing. Uninjected embryos

were included as a negative control. Genomic DNA was extracted from pools of five

embryos in each category and exon 7 was amplified by PCR. Amplicons from each

group were subjected to next-generation sequencing. We find that that the efficiency of

genome editing as determined by next-generation sequencing correlates with

phenotype in b692 edited embryos. In high rescued embryos we find that 17% of reads

(106428 edited/641281 total) harbor an edited codon restoring Ser215 to Ile, consistent

with template integration. Analysis of aligned reads at exon 7 reveals the uniform

presence of barcoded nucleotides from the HDR template across the sgRNA target site.

Phenotypic rescue in medium and low categories revealed molecular efficiency of

approximately 3% (14810/575344 and 17566/622068 reads, respectively). Injected

embryos with no phenotypic evidence of HDR had molecular efficiency of <0.3%.

Alignment of reads demonstrates a mutually exclusive pattern of either HDR and

template integration or indels at the sgRNA site (Figure 5a), consistent with a highly

efficient dsDNA cleavage. We analyzed the molecular efficiency across all phenotypes

and plotted the fraction of reads harboring barcoded nucleotides from the HDR template

(Figure 5b-e). These data confirm the sensitivity of the b692 assay to read out the

molecular efficiency of HDR *in vivo*.

Genome editing using synthetic reagents leads to efficient fluorophore knock-in

To determine the feasibility of using synthetic reagents to knock-in a fluorophore with HDR, we targeted *tyrp1b*, a melanocyte-specific gene. Using albino embryos (pigmentless due to a null mutation in *slc45a2*) we performed microinjection of rCas9 with a synthetic sgRNA and a synthetic linear dsDNA template designed to knock EGFP in frame with the terminal *tyrp1b* exon (Figure 6a). We identified 7/153 (5%) embryos with EGFP-positive melanocytes at 48hpf (Figure 6b). Confocal imaging revealed the presence of an EGFP-positive melanocyte with characteristic stellate morphology (Figure 6d). Allele-specific PCR was performed from injected embryos with EGFP-positive melanocytes (Figure 6e). Sanger sequencing revealed the presence of multiple barcoded nucleotides encoded by the HDR template, consistent with precise integration (Figures 6c).

To determine the generalizability of this approach across multiple loci, we targeted *h3f3a*, one of several genes encoding histone H3.3. We designed a linear dsDNA template to knock EGFP in frame with *h3f3a*, creating a C-terminal fusion (Figure 6f). We microinjected rCas9 with a synthetic sgRNA targeting *h3f3a* and synthetic dsDNA template into wild type embryos. We identified 12/152 (8%) embryos with EGFP-positive cells at 48hpf (Figure 4g). Confocal imaging revealed the presence of nuclear EGFP-positive cells characteristic of chromatin (Figure 6i). Allele-specific PCR and Sanger sequencing performed on EGFP positive embryos revealed precise template integration (Figure 6h and j). Taken together these data demonstrate the ability of synthetic sgRNAs and dsDNA templates to knock-in allele-specific fluorophores at several genomic loci.

DISCUSSION

Here we describe the development and application of genome editing in zebrafish using synthetic reagents. We show that synthetic sgRNAs combined with recombinant Cas9

protein leads to highly efficient indels as compared to conventional laboratory

synthesized sgRNAs. We speculate that improved efficiency of commercial sgRNAs

results from both the chemical modifications that protect sgRNAs from degradation and

the increased fidelity of solid phase chemical synthesis. Our studies demonstrate

germline transmission of synthetic sgRNA induced mutations to F_1 animals following

detection of mutant alleles in the sperm of F_0 animals. While previous reports have

generated smaller deletions in zebrafish[15], we demonstrate that synthetic sgRNAs can

produce deletions up to 124kb. Our approach to knock-in synthetic linear dsDNA

templates has broad applications for studying specific cell populations in development,

for engineering disease causing mutations, and for following clones of cells in mosaic F_0

animals.

In order to understand the parameters critical for homology-directed repair we

developed a genetic assay using *mitfa*(b692) mutant zebrafish. The b692 assay is

highly sensitive and allowed us to systematically examine multiple variables associated

with HDR efficiency. We find that template length, strandedness, and sgRNA site

symmetry appear to be the most critical variables in our model. A key aspect of our

study is the empirical test of multiple conditions reported to improve HDR in other

contexts, using tools such as single-stranded short DNA templates, PARP inhibitors,

multiple sgRNA sites, catalytically mutated rCas9[14,16,17]. In our hands these approaches

do not translate to efficient HDR in zebrafish, suggesting that requirements for HDR

may be species specific.

A total synthetic approach has the unique advantage that reagents can be designed *in*

silico and commercially manufactured in a short time frame. Design and precision

editing with a synthetic dsDNA template can generally be accomplished within 2 weeks.

The use of synthetic reagents creates unique possibilities for future studies. The use of

sgRNAs produced by commercial solid phase synthesis raises the possibility of

incorporating degeneracy in the sgRNA sequence to permit a single sgRNA to target

multiple related genes. The synthesis of linear dsDNA templates may allow

investigators to rapidly test sequence requirements for HDR. As new recombinant Cas9

proteins become available the ability to perform base editing and other enzyme

modifications may be possible. Finally, the b692 assay may be used as a screening

platform to identify chemical and genetic modifiers of HDR efficiency.

The use of CRISPR/Cas9 to generate loss of function mutations in zebrafish and other

model organisms makes it a uniquely valuable resource for forward genetics. To our

knowledge, this is the first head-to-head study investigating the effects of synthetic

versus *in vitro* transcribed guide RNAs and to perform a comprehensive optimization of

HDR efficiency in zebrafish. Our studies offer insight into the relative efficiency of HDR

and provide investigators with a highly efficient workflow for generating edited alleles,

and precise gene knock-ins. We find a single synthetic sgRNA is highly efficient in

generating on-target loss of function indels in a majority of tested genes. Our approach

to genome editing should allow investigators to pursue a broad range of *in vivo*

applications including high efficiency mutant generation, deletion of large genomic

regions to study regulatory DNA, and fluorophore and epitope tag knock-in for functional

studies. Future improvements in HDR efficiency may occur with new variations in Cas9

protein and chemical or genetic tools to further enhance the efficiency of homology

directed repair and integration of donor DNA templates.

MATERIALS AND METHODS

sgRNA sequence design

sgRNAs were designed using the design tool sgRNA Scorer 1.0 and 2.0[18,19]

GuideScan[20], or CRISPRz[21]. Using GuideScan we selected sgRNAs with zero

computed off-targets with a maximum of 3 base mismatches.

Preparation of *in vitro* transcribed sgRNAs

DNA templates for IVT were synthesized as double-stranded gBlock Gene Fragments

(Integrated DNA Technologies) containing a T7 binding site with additional 3' G (5'-

TAATACGACTCACTATAGG), 20bp target site without PAM site, and 80bp tracrRNA

site (5'-

GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAA

AGTGGCACCGAGTCGGTGCTTTT-3').

50ng of sgRNA gBlock (10ng/uL) was *in vitro* transcribed using a SureGuide gRNA

Synthesis Kits (Agilent, 5190-7718) in a 25uL reaction and incubated at 37°C for 3-4

hours. Following DNAse digestion sgRNAs were column-purified using standard kit

instructions and eluted in H_2O. Concentration was determined using a Nanodrop

spectrophotometer and 200ng of the sgRNA was run on an agarose gel for quality

control.

Synthetic sgRNA preparation

sgRNA sequences were ordered from Integrated DNA Technologies (IDT) as Alt-R®

CRISPR-Cas9 crRNAs. The common tracrRNA was purchased from IDT. The crRNA

and tracrRNA were resuspended to 100uM in nuclease-free duplex buffer (IDT)

according to manufacturer specifications. Equimolar amounts of crRNA and tracrRNA

were heteroduplexed by heating solution to 95°C for 5 min and then cooled to room

temperature for 5-10 min. Dilution of sgRNA complex to a working concentration was

done using Nuclease-free Duplex Buffer and stored in -80°C as single-use aliquots.

HDR template design

HDR templates were ordered as gBlock Gene Fragments (IDT). Synthesized gBlocks

were resuspended in nuclease-free water to a concentration of 50ng/uL and stored at -

20°C.

Microinjection

For global editing with a single sgRNA, 250pg sgRNA and 500pg rCas9 were injected

into the yolk of one-cell stage embryos. To generate deletions, 250pg sgRNA 1, 250pg

sgRNA 2 and 500pg rCas9 were injected into the yolk of one-cell stage embryos. For

HDR injections, 250pg sgRNA, 500pg rCas9 and 37.5pg HDR template were injected

into the cell one-cell stage embryos.

gDNA isolation

Individual embryos (24-48hpf) were incubated in 150 μl of DirectPCR Lysis Reagent

(Viagen, 102-T) containing 20 μg/ml Proteinase K (Qiagen, 158920) at 55°C for 60 min

followed by 45 min at 85°C. gDNA was stored at 4°C.

CRISPR-STAT protocol

Sperm samples from potential founders were collected with 10 uL capillary tubes as

previously described, and then followed the HOTShot method with 25 uL alkaline lysis

buffer to prepare DNA templates[22-23]. Tail clips were prepared similarly. The CRISPR-

STAT protocol was used to perform fluorescence-based genotyping[11]. Primers

sequences used for the CRISPR-STAT assay were *slc6a15* FOR,

tgtaaaacgacggccagtGGCCACGACCTACTACTGGTAT, which includes (lowercase)

M13 tag for binding to a fluorescent-labeled M13 primer; and *slc6a15* REV,

gtgtcttTATAGATGCTGCGTCACGTTTC, which include the (lowercase) 'PIG tail' tag

that helps ensure uniform product size, via Taq adenylation[24]. Relative amounts of each

uniquely-sized PCR product (>100 bp, and >100 peak height) were calculated with area

under the curve measurements using Applied Biosystems Peak Scanner software.

Assessment of CRISPR/Cas9 efficiency

1uL of genomic DNA was PCR-amplified using specified primers (Supplementary Figure

7). 8 uL of PCR product was mixed with 1.6uL NEB Buffer 2 and 6.4uL of water and

was incubated at 95°C for 5 minutes followed by cooling from 95°C-85°C at -

2°C/second and 85-25°C at -0.1°C/second. 0.4uL NEB Buffer 2, 3.4uL water and 2U of

T7EI endonuclease (New England BioLabs, M0302) were added to hybridized products

and were incubated for 1 hour at 37°C. The 20uL reaction was run on a 2% agarose

gel.

For clonal analysis, the PCR product was cloned into pCRII-TOPO (Invitrogen,

K460001) as per the manufacturer's instructions and Sanger sequenced (Genewiz).

Sequences were compared to reference genome and non-targeting controls to identify

indels using CrispRVariants[25] and MacVector (Version 15.5).

Assessment of deletions

gDNA was PCR amplified using primers that flanked the deletion sgRNAs. PCR

products were purified (Qiagen, 28104) and Sanger sequenced. Sequences were

compared to the reference genome using MacVector.

Imaging

Micrographs of whole embryos and larval animals were taken with a Zeiss Discovery V8

stereomicroscope (Zeiss, Oberkochen, Germany) equipped with epifluorescence and

appropriate filters. Live imaging of fluorescent larvae was acquired with a Zeiss LSM800

laser scanning confocal microscope (Zeiss). For melanocyte counting, 48hpf embryos

were visualized under the stereoscope and binned into 4 categories (WT, 100+, 51-100,

0-50 melanocytes). Scientists were blinded to the conditions when binning embryos.

Next-generation sequencing

50ng of PCR amplicon DNA was converted into blunt ends using T4 DNA polymerase

and *E. coli* DNA polymerase I Klenow fragment. Libraries were prepared using the

Kapa_LTP_Library Preparation Kit (Roche, Basel, Switzerland). Multiple indexing

adapters are ligated to the ends of the DNA fragments. Ligation reaction products were

purified by AMPure XP beads (NEB) to remove all unligated adapters and quantified

using Qubit (Thermo Fisher Scientific) and Bioanalyzer DNA chip (Agilent). Indexed

sample libraries were normalized and pooled and were sequenced using the Illumina

HiSeq4000 sequencer at 2x50 cycles single reads. Reads were aligned to the zebrafish

genome (GRCz10) using the Ensembl transcriptome[26]. Results were viewed using

Integrative Genome Browser (IGV)[27,28].

Fish lines and maintenance

Zebrafish strains were maintained according to established guidelines[29].Wild-type

animals were from the AB background. Experiments were performed in strict

accordance with the recommendations in the Guide for the Care and Use of Laboratory

Animals of the National Institutes of Health. All of the animals were handled according

to approved institutional animal care and use committee (IACUC) protocols of Weill

Cornell Medical College.

Acknowledgements

We are grateful to members of the Houvras laboratory for critical discussion and

manuscript review. We thank the Weill Cornell Genomics Resources Core Facility for

next generation sequencing. This work was supported by the Department of Surgery,

Weill Cornell Medical College (YH); National Institutes of Health/National Cancer

Institute pre-doctoral fellowship F31CA192813 (SED); National Institutes of

Health/National Cancer Institute pre-doctoral fellowship F31CA213997 (RM); the

Medical Scientist Training Program of General Medical Sciences of the NIH

(T32GM007739) to the Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-

PhD Program (RM); the Clinical and Translational Science Center at Weill Cornell

Medical College TL1-TR0002386-01 (CKG).

Author Contributions

Y.H. conceived and led this work. S.D., R.M., and C.G. designed the genetic assays,

performed experiments and conducted data analyses related to genome editing. I.N.,

C.B., and A.L. provided technical assistance and performed data analysis. D.M., C.K.,

S.M., and J.J. implemented and interpreted the entirety of the CRISPR-STAT

protocol. S.D., C.G., and Y.H. wrote the paper.

REFERENCES
1.   Jao, L.-E., Wente, S. R. & Chen, W. Efficient multiplex biallelic zebrafish genome

editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. U. S. A.* **110,**
13904–9 (2013).

2.   Shah, A. N., Davey, C. F., Whitebirch, A. C., Miller, A. C. & Moens, C. B. Rapid
reverse genetic screening using CRISPR in zebrafish. *Nat. Methods* **12,** 535–540
(2015).

3.   Auer, T. O., Duroure, K., Cian, A. De, Concordet, J. & Bene, F. Del. Highly
efficient CRISPR / Cas9-mediated knock-in in zebrafish by homology-independent
DNA repair. *Genome Res.* **24,** 142–153 (2014).

4.   Irion, U., Krauss, J. & Nusslein-Volhard, C. Precise and efficient genome editing
in zebrafish using the CRISPR/Cas9 system. *Development* 4827–4830 (2014).
doi:10.1242/dev.115584

5.   Hoshijima, K., Jurynec, M. J. & Grunwald, D. J. Precise Editing of the Zebrafish
Genome Made Simple and Efficient. *Dev. Cell* **36,** 654–667 (2016).

6.   Hwang, W. Y. *et al.* Heritable and Precise Zebrafish Genome Editing Using a
CRISPR-Cas System. *PLoS One* **8,** 1–9 (2013).

7.   Kimura, Y., Hisano, Y., Kawahara, A. & Higashijima, S. I. Efficient generation of
knock-in transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-
mediated genome engineering. *Sci. Rep.* **4,** 1–7 (2014).

8.   Hendel, A. *et al.* Chemically modified guide RNAs enhance CRISPR-Cas genome
editing in human primary cells. *Nat. Biotechnol.* **33,** 985–989 (2015).

9.   Rahdar, M. *et al.* Synthetic CRISPR RNA-Cas9-guided genome editing in human
cells. *Proc. Natl. Acad. Sci.* **112,** E7110-7117 (2015).

10.   Sheets, L., Ransom, D. G., Mellgren, E. M., Johnson, S. L. & Schnapp, B. J.
Zebrafish Melanophilin Facilitates Melanosome Dispersion by Regulating Dynein.
Curr. Biol. **17,** 1721–1734 (2007).

11.   Carrington, B., Varshney, G. K., Burgess, S. M. & Sood, R. CRISPR-STAT: An
easy and reliable PCR-based method to evaluate target-specific sgRNA activity.
Nucleic Acids Res. **43,** 1–8 (2015).

12.   Lister, J., Robertson, C., Lepage, T., Johnson, S. & Raible, D. Nacre Encodes a
Zebrafish Microphthalmia-Related Protein That Regulates Neural-Crest-Derived
Pigment Cell Fate. *Development* **126,** 3757–3767 (1999).

13.   Lister, J. A., Close, J. & Raible, D. W. Duplicate mitf genes in zebrafish:
Complementary expression and conservation of melanogenic potential. *Dev. Biol.*
237, 333–344 (2001).

14.   Richardson, C. D., Ray, G. J., DeWitt, M. A., Curie, G. L. & Corn, J. E. Enhancing
homology-directed genome editing by catalytically active and inactive CRISPR-
Cas9 using asymmetric donor DNA. *Nat. Biotechnol.* **34,** 339–344 (2016).

15.   Varshney, G. K. *et al.* High-throughput gene targeting and phenotyping in
zebrafish using CRISPR / Cas9. *Genome Res.* **25,** 1–13 (2015).

16.   Quadros, R. M. *et al.* Easi-CRISPR: A robust method for one-step generation of
mice carrying conditional and insertion alleles using long ssDNA donors and
CRISPR ribonucleoproteins. *Genome Biol.* **18,** 1–15 (2017).

17.   Anantha, R. W. *et al.* Functional and mutational landscapes of BRCA1 for
homology-directed repair and therapy resistance. *Elife* **6,** 1–21 (2017).

18. Chari, R., Mali, P., Moosburner, M. & Church, G. M. Unraveling CRISPR-Cas9
genome engineering parameters via a library-on-library approach. *Nat. Methods*
12, 1–7 (2015).
19. Chari, R., Yeo, N. C., Chavez, A. & Church, G. M. SgRNA Scorer 2.0: A Species-
Independent Model to Predict CRISPR/Cas9 Activity. *ACS Synth. Biol.* **6,** 902–
904 (2017).
20. Perez, A. R. *et al.* GuideScan software for improved single and paired CRISPR
guide RNA design. *Nat. Biotechnol.* **35,** 347–349 (2017).
21. Varshney, G. K. *et al.* CRISPRz: A database of zebrafish validated sgRNAs.
Nucleic Acids Res. **44,** D822–D826 (2016).
22. Draper, B. W. & Moens, C. B. A High-Throughput Method For Zebrafish Sperm
Cryopreservation and In Vitro Fertilization. *J. Vis. Exp.* 1–5
(2009). doi:10.3791/1395
23. Meeker, N. D., Hutchinson, S. A., Ho, L. & Trede, N. S. Method for isolation of
PCR-ready genomic DNA from zebrafish tissues. *Biotechniques* **43,** 610–614
(2007).
24. Brownstein, M. J., Carpten, J. D. & Smith, J. R. Modulation of non-templated
nucleotide addition by Taq DNA polymerase: Primer modifications that facilitate
genotyping. *Biotechniques* **20,** 1004–1010 (1996).
25. Lindsay, H. *et al.* CrispRVariants charts the mutation spectrum of genome
engineering experiments. *Nat. Biotechnol.* **34,** 701–702 (2016).
26. Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to
the human genome. *Nature* **496,** 498–503 (2013).
27. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer
(IGV): High-performance genomics data visualization and exploration. *Brief.*
Bioinform. **14,** 178–192 (2013).
28. Robinson, J. T. *et al.* Integrative Genomics Viewer. *Nat. Biotechnol.* **29,** 24–26
(2011).
29. Westerfield M. 2007. The Zebrafish Book a Guide for the Laboratory Use of
Zebrafish (Danio Rerio). 5th Edition. Eugene: University of Oregon Press.

Figure 1 Synthetic sgRNAs outperform laboratory synthesized *in vitro* transcribed
sgRNAs. a, Dose-response comparison of editing in embryos injected with rCas9 and
either an *in vitro* synthesized or synthetic sgRNA targeting *tyr*. Embryos were scored for
melanocyte dropout at 48hpf and binned into 4 categories. The percentage of embryos
in each condition is plotted. At the 1000pg sgRNA/2000pg rCas9 dose, there is a
statistically significant increase in the editing observed with the synthetic sgRNA
(p=<0.001, Chi-square). **b**, Light microscopic images of representative embryos in each
category. **c**, Distribution of wild type clones, small indels (<5bp), and large indels (>5bp)
in embryos injected with synthetic or *in vitro* transcribed sgRNAs. **d**, CrisprVariants
(Lindsay et al., 2016) plot of indels in clones isolated from embryos injected with either
synthetic or *in vitro* transcribed sgRNAs. 10/10 clones isolated from individual embryos
injected with synthetic from sgRNAs contained indels and 7/10 clones isolated from
embryos injected with IVT sgRNAs contained indels.
527
Figure 2 Germline transmission of edited alleles is achieved using synthetic
sgRNAs. a, Sperm samples were collected from a CRISPR-injected potential F_0
founder and assayed using CRISPR-STAT fluorescence-based genotyping. **b-f**,
CRISPR-STAT results for tail clips from representative F_1 progeny, each carrying one of
the distinct alleles with specific insertions and deletions (del3, del4, del7, del11), as well
as the wild type (WT) allele, transmitted from the F0 founder. The number of F1
offspring identified with each genotype is identified in the bottom right of each panel.
Relative fluorescence levels (y-axis) are shown against fragment size in base pairs (x-
axis) for each genotyping assay. PCR products consistent with predicted wild type
amplicon size are indicated by red asterisks.
538
Figure 3 Synthetic sgRNA pairs produce large chromosomal deletions. a,
Schematic of a 126kb non-coding region targeted for genomic deletions. Guide
locations are marked in purple, and primers used for PCR are displayed as orange
arrows. Guide pairs were co-injected into 1 cell stage embryos. **b**, Deletion-specific
PCR was performed to confirm the presence of a deletion in individual F_0 injected
embryos. The predicted deletion size for each pair and frequency of deletions is shown.
c-f, PCR amplicons from individual embryos were subjected to Sanger sequencing and
analysis of deletions. The observed deletion size is displayed.
547
Figure 4 A genetic assay for optimizing homology-directed repair using synthetic
reagents. a, Schematic depicting *mitfa* gene structure and location of the mutation in
exon 7 in b692 mutants. Guide locations are marked in purple, and primers used for
PCR are displayed as orange arrows. A linear 951bp dsDNA template encodes a wild-
type Ile codon at position 215 and several other nucleotide changes for template

identification after sequencing. **b**, *mitfa*(b692) embryos uninjected (top), injected with
rCas9, 951bp DNA template, and non-targeting sgRNA (middle) or *mitfa* sgRNA
(bottom). All embryos shown at 48hpf. **c**, Allele-specific PCR was performed on rescued
embryos after microinjection with 951bp template compared with uninjected embryos. **d**,
Sanger sequencing results of allele-specific PCR products from a single phenotypically
rescued embryo. **e**, Chart depicting attributes of DNA template (left), a description of the
template (middle) and observed rate of rescue (right).
560
Figure 5 Quantitative assessment of genome editing efficiency by next-generation
sequencing. a, Alignment of next generation sequencing reads from b692 genome
edited embryos from the high rescue phenotype. Exon 7 of *mitfa* is displayed. The
sgRNA sequence is highlighted in yellow. b692 mutants have a T>G mutation leading to
an isoleucine to serine substitution at codon 215(*). The 951bp dsDNA template
restores the wild-type isoleucine codon (ATC) and encodes nine additional barcoded
nucleotides as indicated. Indels in sequencing reads are represented as gapped regions
in black. **b-e**, The fraction of reads at each barcoded nucleotide in the HDR template is
plotted for each phenotype category.
570
571
Figure 6 Genome editing leads to precise fluorophore knock-in. a, Schematic
depicting *tyrp1b* locus. A linear, dsDNA template designed to lead to an in-frame EGFP
knock-in is shown. **b**, albino(b4) embryo injected with rCas9, *tyrp1b* synthetic sgRNA
and dsDNA template is photographed with light (top panel) or fluorescence (bottom
panel) microscopy at 48hpf. **c**, Sanger sequencing from allele-specific PCR to detect
template integration. The sequence was aligned to reference (top). Barcoded
nucleotides at the sgRNA binding site consistent with template integration were
identified. A representative chromatogram is shown. **d**, Confocal imaging reveals the
presence of stellate EGFP-positive melanocytes. **e**, Allele-specific PCR was performed
on uninjected or HDR edited/GFP+ embryos at 48hpf. Only HDR edited/EGFP positive
embryos had a PCR amplicon. **f**, Schematic depicting *h3f3a* locus. A linear, dsDNA
template designed to lead to an in-frame EGFP knock-in is shown. **g**, Wild type embryo
injected with rCas9, *h3f3a* sgRNA and dsDNA template is photographed with light (top
panel) or fluorescence (bottom panel) microscopy (48hpf). **h**, Sanger sequencing was
performed from allele-specific PCR to detect template integration. The sequence was
aligned to reference (top). Barcoded nucleotides at the sgRNA binding site, consistent
with template integration were identified. A representative chromatogram is shown. **i**,
Confocal imaging reveals the presence of a EGFP-positive nuclei. **j**, Allele-specific PCR
was performed on uninjected or HDR edited/GFP+ embryos at 48hpf. Only HDR
edited/EGFP positive embryos had a PCR amplicon.
592
Supplementary Figure 1 Synthetic sgRNAs outperform *in vitro* synthesized
sgRNA in targeting *slc25a2*. a, Comparison of Alt-R® synthetic sgRNA preparation
and *in vitro* synthesis of sgRNAs. Left, Alt-R® synthetic sgRNAs are prepared by
duplexing a gene specific crRNA with a common tracrRNA. Right, IVT sgRNAs are

transcribed from a DNA template using RNA polymerase and purified. Both sgRNAs are
used with rCas9 and microinjected into embryos at the one-cell stage. **b,** Light
microscopic images of representative embryos in each category. **c,** Dose-response
comparison of editing in embryos injected with rCas9 and either an *in vitro* synthesized
or synthetic sgRNA targeting *slc25a2*. Embryos were scored for melanocyte dropout at
48hpf and binned into 4 categories. The percentage of embryos in each condition is
plotted.
604
Supplementary Figure 2 Synthetic sgRNAs produce highly penetrant phenotypes
in F_0 injected animals. Wild-type (AB) embryos were injected with rCas9 and synthetic
sgRNA targeting *mlpha*. Uninjected larval animals (top) have normal stripe pattern.
Juvenille zebrafish (7 weeks post fertilization) injected with rCas9 and a synthetic
sgRNA targeting *mlpha* show reduced pigmentation and phenocopies an established
mutant.
611
612 **Supplementary Figure 3 Indels in F_0 sperm predict alleles transmitted to the**
613 **germline.** Percentages shown are calculated as area under the curve for each peak in
614 the indicated tissue from the F_0 founder. The frequencies of each mutation in the F_1
615 progeny were calculated from a total of 17 siblings from a cross between the F_0 male
616 parent and a WT female.
617
618 **Supplementary Figure 4 Detection and verification of chromosomal deletions.**
619 **a,c,e,g,** Deletion-specific PCR was performed to detect deletions from individual
620 embryos injected with rCas9 and the indicated sgRNAs. In **(a)**, the top band is the wild-
621 type allele and the lower band is indicative of a deletion. **b,d,f,h,** Sanger sequencing of
622 PCR products reveals the specific deletions resulting from injection of sgRNA pairs.
623
624 **Supplementary Figure 5 Comparison of melanocyte phenotypes in mitfa mutants**
625 **for after targeting mitfa using CRISPR/Cas9. a,** *mitfa*(w2) embryos were injected with
rCas9 and a specific sgRNA near the mutation site. Ten representative embryos at
24hpf are shown. Melanocyte rescue occurs in a significant fraction of embryos. **b,**
CrispRVariants plot from *mitfa*(w2) embryos injected with rCas9 and nacre gRNA. A
majority of embryos harbor indels. **c,** mitfa(b692) embryos were injected with rCas9 and
a synthetic sgRNA near the b692 mutation site (b692 sgRNA). Uninjected wild-type
(AB) embryos (top, left) have pigmented melanocytes at 48hpf. Wild-type embryos
injected with rCas9 and b692 sgRNA (bottom, left) have a highly penetrant loss of
melanocytes. Uninjected *mitfa*(b692) embryos (top, right) lack pigmentation because of
the b692 mutation. *mitfa*(b692) embryos injected with rCas9 and the b692 sgRNA
(bottom, right) have no evidence of melanocyte rescue.
635
Supplementary Figure 6 Strategies employed for optimizing homology-directed
repair at *mitfa*. Schematic depicts several different DNA templates employed for HDR
in vivo. Mutated sgRNA binding sites are indicated by vertical purple lines and
overlapping blue lines indicate shared genomic sequence. Descriptions of each
template along with any special considerations are noted in the middle column.

Efficiencies of HDR, as measured phenotypically by presence or absence of
melanocytes in injected *mitfa*(b692) zebrafish, are noted in the right column.
643
644   **Supplementary Figure 7 sgRNA and primer sequences**. Sequences of sgRNA
645   sequences are noted with the PAM sequence in parentheses. Primer sequences are
646   listed for each locus.

Figure 1

Figure 2

a

chr5:34397610-34600530

foxd1

20kb

fch02

b

sgRNAs	Predicted deletion size	Observed deletion frequency
sgRNA1+2	0.45kb	5/10 (50%)
sgRNA3+4	4.3kb	7/24 (29%)
sgRNA2+4	52kb	3/8 (38%)
sgRNA1+5	124kb	2/23 (9%)

c

rCas9 + sgRNA1 + sgRNA2

5' sequence	3' sequence

T T G T T G C C A T C C G A A A G T T T T A T T T G T G C T C T T T G A A G G C C G T T T C C C A C A T T T A T C T C G C T A A T A G C T A C T C A G C G A C A G C

427bp deletion

d

rCas9 + sgRNA3 + sgRNA4

5' sequence	3' sequence

T T A A G A G A T C T G G G A T G G G A T C A T A A G T G C T G A A C A C T T T G C A G G G C C A C G C T G C A C C T T A A C C T A C A A T A A T A A A G C A A

4309bp deletion

e

rCas9 + sgRNA2 + sgRNA4

5' sequence	3' sequence

C A G A C A A C A G A T G C A A T G A T G T T G G A T C T T C A T C A G G G C C A C G C T G C A C C T T A A G C T A C A A T A A T A A A G C A A G C T G C C G T T

52,262bp deletion

f

rCas9 + sgRNA1 + sgRNA5

5' sequence		3' sequence

T G T T G C C A T C C G A A A G T T T T A T T T G T G C T C T T T G A A G G C A A C G A C T T A A A T A T A A A T C T T T A A G T C G T T A C T T C A C C A A A A A A G T G T

123,847bp deletion + 9bp insertion

Figure 3

a

chr6: 43429063-43435741

mitfa

mitfa sgRNA

b692 mutation: A**G**C (215 Ser)

951nt HDR template   7

Wild-type: A**T**C (215 Ile)

b

WT

Non-targeting

mitfa HDR

c

uninjected   rescued

191bp

d

barcoded nucleotides

TCAAGAAGTAGATCAAATTTGAATTTTAATAGAGTGGTT**TTT**GAAACTTACGGATCAT
.....................................**GCAC**...............
.....................................**GCAC**...............
ГAGATCAAATTTGAATTTTAATAGAGTGGTTG**CAC**AAACTTACGGAT

e

DNA template design	Strategy employed	HDR efficiency
	Asymmetric ds 951bp template	9%
	Asymmetric ds 2000bp template	6%
	Asymmetric ds 318bp template	1%
	Asymmetric ds 951bp template with two guide sites	<1%
	5' asymmetric ds 951bp template	0%
	Symmetric ds 951bp template	0%
	Asymmetric ds 951bp template in a vector	0%

Figure 4

Figure 5

Figure 6

a

Strategies for sgRNA synthesis

IDT Alt-R sgRNA preparation | In vitro sgRNA preparation

crRNA   tracrRNA | DNA synthesis

RNA synthesis

duplex

RNA purification

+ rCas9 protein

Micro injection

b

Wildtype

Low

Intermediate

High

c

Percentage

125pg gRNA 250pg rCas9 (n=54)
250pg gRNA 500pg rCas9 (n=53)
500pg gRNA 1000pg rCas9 (n=81)
1000pg gRNA 2000pg rCas9 (n=70)

IVT sgRNA

125pg gRNA 250pg rCas9 (n=52)
250pg gRNA 500pg rCas9 (n=47)
500pg gRNA 1000pg rCas9 (n=87)
1000pg gRNA 2000pg rCas9 (n=77)

synthetic sgRNA

■ None (wild-type)
■ Low (100+ melanocytes)
■ Intermediate (51-100 melanocytes)
■ High (0-50 melanocytes)

Supplementary Figure 1

Uninjected AB

mlpha sgRNA + rCas9

Supplementary Figure 2

	F0 Sperm	F1 Progeny
del11	7.4%	5.9%
del7	58.9%	41.2%
del4	<5%	5.9%
del3	33.7%	23.5%
WT	<5%	23.5%
Total	100.0%	100.0%

Supplementary Figure 3

a rCas9 + sgRNA 1 + sgRNA 2

b
| 427bp deletion + 0bp insertion |

c rCas9 + sgRNA 3 + sgRNA 4

d
| 4297bp deletion + 21bp insertion |
| 4298bp deletion + 15bp insertion |
| 4302bp deletion + 34bp insertion |
| 4316bp deletion + 0bp insertion |
| 4451bp deletion + 40bp insertion |

e rCas9 + sgRNA 2 + sgRNA 3

f
| 52262 bp deletion + 0bp insertion |
| 52266 bp deletion + 12bp insertion |

g rCas9 + sgRNA 1 + sgRNA 5

h
| 123847bp deletion + 9bp insertion |

Supplementary Figure 4

a

b

c

Supplementary Figure 5

DNA template design	Strategy employed	HDR efficiency
	Symmetric ds 76bp template	0%
	Symmetric ds 200bp template	0%
	Asymmetric ds 951bp template with nickase-compatible guide sites	0%
	Asymmetric ds 951bp template with nonhomologous flanking guide sites	0%
	Symmetric ds 2000bp template	0%
	Symmetric ds 1214bp template with homologous flanking guide sites	0%
	Symmetric ds 896bp template with two guide sites	0%
	Asymmetric ds 951bp template with flanking IScel restriction enzyme sites	0%
	Symmetric ss 896nt template with two guide sites	0%
	Symmetric ss 200bp template	0%

Supplementary Figure 6

gRNA name	gRNA sequence
tyr	TGAAAGTTACAACCTCCGCG(AGG)
slc24a5	GGTCTCTCGCAGGATGTTGC(TGG)
mitfa(w2)	AGGACTTCAAATGACAAACA(CGG)
b692	ATGACAGAATTAAGGAGCTG(GGG)
tyrp1b	CACTATCGCTATAGTTGCAG(TGG)
mlpha	AATGACCTCCCACACATGCT(TGG)
h3f3a	CAAGGATATCCAGCTTGCA(AGG)
non-targeting	TGCTGATGTCTTGGTTGAAT(TGG)
deletion_1	TGTGCTCTTTGAAGGCCATG(AGG)
deletion_2	TGTGGGAAACGGCATATATG(AGG)
deletion_3	AATCACCGAGAAAATTGGAG(AGG)
deletion_4	CAGGCAGGATCAAAGAAGGC(AGG)
deletion_5	AATAGCTCATAAAGACGGTG(AGG)
slc6a15	GACTCCATCTCAGAGGGTGG(AGG)

Primer Name	Primer sequences
tyr_guide_validation (F)	AATGTCGTTCACTCTGCTGTTG
tyr_guide_validation (R)	GTGACCGTGCTGTTGTACCTT
mitfa_w2_guide_validation (F)	GTGGATTGAGGTTCCCTTCA
mitfa_w2_guide_validation (R)	CGGATAATTCCCTTTTGACG
mitfa_HDR_detection (F)	GCTACAGTGATGACATTCTTGG
mitfa_HDR_detection (R)	GCATGATTGCTGTACATATCAAGCAAA
mitfa_next_gen (F)	GCAGTTATTAATGATAATGTCTCG
mitfa_next_gen (F)	GGACTTAAGTAAATCAAGAAGTAGATC
downstream_foxd1_deletion_1 (F)	TCACTGAGGCCTCTGACAGA
downstream_foxd1_deletion_2 (R)	TACTCCTTTGGACAGGGCCA
downstream_foxd1_deletion_3 (F)	CTCTGTGGTCTATTTCTCTCC
downstream_foxd1_deletion_4 (R)	CAGAAGGTGAAGCAGCCTCA
downstream_foxd1_deletion_5 (R)	ACCACGTCTTCAAGGGCTTC

Supplementary Figure 7