Nuclear and mitochondrial genomic resources for the meltwater stonefly, *Lednia tumana*

(Plecoptera: Nemouridae)

Scott Hotaling[1,2], Joanna L. Kelley[1], and David W. Weisrock[2]

Affiliations:

[1] School of Biological Sciences, Washington State University, Pullman, WA, USA

[2] Department of Biology, University of Kentucky, Lexington, KY, USA

Correspondence: Scott Hotaling, School of Biological Sciences, Washington State University,

Pullman, WA, 99164, USA; Email: scott.hotaling@wsu.edu; Phone: (828) 507-9950

Abstract:

With more than 3,500 described species, stoneflies (Order Plecoptera) are an important

component of global aquatic biodiversity. The meltwater stonefly *Lednia tumana* (Family

Nemouridae) is endemic to alpine streams of Glacier National Park and has been petitioned for

listing under the U.S. Endangered Species Act (ESA) due to climate change-induced loss of

alpine glaciers and snowfields. Here, we present a *de novo* assembly of the nuclear genome

(~305 million bp) as well as the nearly complete mitochondrial genome (13,752 bp) for *L.*

tumana. To our knowledge, this represents the first high-coverage (>10x) nuclear genome for the

stonefly order (Plecoptera) and second mitochondrial genome for the family Nemouridae. Both

assemblies provide important foundational resources for future stonefly research.

Keywords: nuclear genome, Glacier National Park, mitochondrial genome, conservation

genetics, global change biology

Running head: Nuclear and mitochondrial genomes for *Lednia tumana*

Introduction:

Stoneflies are a diverse, globally distributed group of hemimetabolous insects that

diverged from their closest relatives (e.g., Orthoptera, Dermaptera, Zoraptera) ~275 million years

ago in the early Permian (Misof et al., 2014). With more than 3,500 described species, stoneflies

account for a substantial portion of freshwater biodiversity (Fochetti and De Figueroa, 2008).

The meltwater stonefly, *Lednia tumana* (Plecoptera: Nemouridae), resides in alpine streams of

Glacier National Park (GNP), USA, where it is iconic of habitat loss due to climate change in the

region (Giersch, Hotaling, Kovach, Jones and Muhlfeld, 2017). *Lednia tumana* is one of four

species in the genus *Lednia* which all have alpine, cold-water distributions in western North

America (Baumann and Kondratieff, 2010). The majority of *L. tumana*'s habitat is supported by

seasonal melting of permanent ice and snow, a habitat type that is under considerable threat of

near-term loss as the global cryosphere recedes (Hotaling, Finn, Joseph Giersch, Weisrock and

Jacobsen, 2017). The recent evolutionary history of *L. tumana* is also closely tied to glacier

dynamics with present-day genetic clusters arising in parallel with ice sheet recession at the end

of the Pleistocene (~20,000 years ago, Hotaling et al., 2018). Genetic evidence has also

highlighted a possible loss of mitochondrial genetic diversity on even more recent, decadal

timescales for the species (Jordan et al., 2016). With such a narrow habitat niche in a small,

mountainous region of the northern Rocky Mountains, *L. tumana* has been recommended for

listing under the U.S. Endangered Species Act (US Fish & Wildlife Service, 2016). In this study,

we present an assembly of the nuclear genome for *L. tumana*, the first high-coverage (>10x)

nuclear genome for the order Plecoptera. This resource also represents one of only three high-

coverage genomes for any EPT taxon (Ephemeroptera, Plecoptera, and Trichoptera), a globally

important group of aquatic organisms commonly used for biological monitoring (e.g., Tronstad,

Hotaling and Bish, 2016). We also present a mitochondrial genome assembly (mitogenome) for

L. tumana, the second for the stonefly family Nemouridae after *Nemoura nankinensis* (Chen and

Du, 2017).

Materials and Methods:

Genomic DNA was extracted using a Qiagen DNeasy Blood & Tissue Kit from a single

L. tumana nymph collected in 2013 from Lunch Creek in GNP. Prior to extraction, both the head

and as much of the digestive tract as possible were removed. A whole-genome shotgun

sequencing library targeting a 250-bp fragment size was constructed and sequenced by the

Florida State University Center for Genomics. The library was sequenced twice on 50% of an

Illumina HiSeq2500 flow cell each time with paired-end, 150-bp chemistry, resulting in

242,208,840 total reads. The size of the nuclear genome for *L. tumana* was estimated to be 536.7

mega basepairs (Mb) using a kmer-based approach in sga preQC (Simpson, 2014). Read quality

was assessed with fastQC (Andrews, 2010) and low-quality reads were either trimmed or

removed entirely using TrimGalore (Krueger, 2015) with the flags: --stringency 3 --quality 20 --

length 40. We assembled the nuclear genome using SPAdes v3.11.1 with default settings

(Bankevich et al., 2012) and generated summary statistics with the Assemblathon2 perl script

(assemblathon_stats.pl, Bradnam et al., 2013). The completeness of our nuclear assembly as well

as two other publicly available stonefly genomes (Macdonald, Cunha and Bruford, 2016;

Macdonald, Ormerod and Bruford, 2017) was assessed by calculating the number of conserved

single copy orthologs (BUSCOs) in the assemblies using BUSCO v3 and the 1,658

"insecta_ob9" set of reference genes (Simão, Waterhouse, Ioannidis, Kriventseva and Zdobnov,

2015). These BUSCO analyses provided a proxy for how complete genic regions are in both our

own and the two existing stonefly genomes.

We assembled the *L. tumana* mitogenome with NOVOPlasty v2.6.7 (Dierckxsens,

Mardulyn and Smits, 2016) using an 872-bp segment of the *L. tumana cytb* gene (GenBank

KX212756.1) as the "seed" sequence. The mitogenome was annotated through a combination of

the MITOS web server with default settings (Bernt et al., 2013) and comparison to the *Nemoura*

nankinensis mitogenome (Plecoptera: Nemouridae; Chen and Du, 2017).

Results and Discussion:

The *L. tumana* nuclear genome assembly comprises just over 304.5 Mb with 50% of the

assembly in contigs equal to or larger than 4.69 kilobases (kb; Table 1). The size of the genome

is in line with two other publicly available stonefly genomes, *Isoperla grammatica* (509.5 Mb;

Family Perlodidae) and *Amphinemura sulcicollis* (271.9 Mb; Family Nemouridae) which are

both deposited under GenBank BioProject PRJNA315680 (Macdonald et al., 2016; Macdonald

et al., 2017). All associated data for the resources detailed in this study, including both raw reads

and assemblies, are available as part of GenBank BioProject PRJNA472568 (mitogenome:

MH374046, nuclear genome: SAMN09295077, raw reads: SRP148706).

The meltwater stonefly's nuclear genome is similarly A/T-rich (58.4%) to the other

stoneflies (58.2-59.9%; Table 1), ants (55-67%; Gadau et al., 2012), *Drosophila melanogaster*

(58%), *Anopheles gambiae* (56%), and the honeybee, *Apis mellifera* (67%, The Honeybee

Genome Sequencing Consortium, 2006). Our assembly is more intact in terms of genic regions
than both existing stonefly assemblies with 92.8% of BUSCOs either complete (70.6%) or
fragmented (22.2%) versus 80.8% for *A. sulcicollis* (50.5% complete, 31.3% fragmented) and
just 50.1% for *I. grammatica* (13.3% complete, 36.8% fragmented; Table 1). However, our
assembly is still highly fragmented, making it difficult to draw more nuanced conclusions about
its structure and evolution.

The mitogenome is also nearly complete, covering 13,752 bp, including all 13 protein-
coding genes, 21 tRNA genes, the 12S rRNA gene, and only missing the 16S rRNA gene and
control region (Figure 1). The organization of the *L. tumana* mitogenome is similar to that of *N.*
nankinensis, the only other mitogenome available for the family Nemouridae. In *N. nankinensis*,
the regions missing from the *L. tumana* mitogenome assembly are ~3 kb, indicating that the
complete *L. tumana* mitogenome is likely around 16.7 kb, which falls in line with other stonefly
mitogenomes (Chen and Du, 2017). Our inability to resolve the control region is perhaps
unsurprising as the *N. nankinensis* control region contains a large, ~1 kb repeat region which is
difficult to resolve without targeted long-range PCR re-sequencing or long-read high-throughput
sequencing.

Future efforts to refine both assemblies, including the incorporation of longer reads (e.g.,
generated using Pacific Biosciences sequencing technology, Utturkar et al., 2014) will yield
greater insight into the genome biology of *L. tumana* and related stoneflies. Still, the resources
included here provide an important step towards empowering modern stonefly research, a
globally relevant group of aquatic insects that has been largely overlooked in the genomic age.
115

Acknowledgements:
We thank Alan and Emily Lemmon for advice and assistance with sequencing. This research was
supported by the University of Kentucky (UK) and National Science Foundation (DEB-
0949532). Computational resources were provided by the UK Center for Computational Sciences
and the Lipscomb High Performance Computing Cluster, as well as the Washington State
University Center for Institutional Research Computing.
122

Declaration of Interest:
The authors declare no financial interest or benefit stemming from this research.

References:

Andrews, S. (2010), 'FastQC: a quality control tool for high throughput sequence data'.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
Nikolenko, S.I., Pham, S. and Prjibelski, A.D. (2012), 'SPAdes: a new genome assembly
algorithm and its applications to single-cell sequencing', *Journal of computational biology*,
19, 455-477.

Baumann, R.W. and Kondratieff, B.C. (2010), 'The stonefly genus Lednia in North America
(Plecoptera: Nemouridae)', *Illiesia*, 6, 315-327.

Bernt, M., Donath, A., Jühling, F., Externbrink, F., Florentz, C., Fritzsch, G., Pütz, J.,
Middendorf, M. and Stadler, P.F. (2013), 'MITOS: improved de novo metazoan
mitochondrial genome annotation', *Molecular phylogenetics and evolution*, 69, 313-319.

Bradnam, K.R., Fass, J.N., Alexandrov, A., Baranay, P., Bechner, M., Birol, I., Boisvert, S.,
Chapman, J.A., Chapuis, G. and Chikhi, R. (2013), 'Assemblathon 2: evaluating de novo
methods of genome assembly in three vertebrate species', *GigaScience*, 2, 10.

Chen, Z.-T. and Du, Y.-Z. (2017), 'First mitochondrial genome from Nemouridae (Plecoptera)
reveals novel features of the elongated control region and phylogenetic implications',
International journal of molecular sciences, 18, 996.

Dierckxsens, N., Mardulyn, P. and Smits, G. (2016), 'NOVOPlasty: de novo assembly of
organelle genomes from whole genome data', *Nucleic Acids Res*, 45, e18-e18.

Fochetti, R. and De Figueroa, J.M.T. (2008), 'Global diversity of stoneflies (Plecoptera; Insecta)
in freshwater', *Hydrobiologia*, 595, 365-377.

Gadau, J., Helmkampf, M., Nygaard, S., Roux, J., Simola, D.F., Smith, C.R., Suen, G., Wurm,
Y. and Smith, C.D. (2012) 'The genomic impact of 100 million years of social evolution in
seven ant species', *Trends in Genetics*, 28, 14-21.

Giersch, J.J., Hotaling, S., Kovach, R.P., Jones, L.A. and Muhlfeld, C.C. (2017), 'Climate-
induced glacier and snow loss imperils alpine stream insects', *Global Change Biology*, 23,
2577-2589.

Honeybee Genome Sequencing Consortium. (2006), 'Insights into social insects from the
genome of the honeybee Apis mellifera', *Nature*, 443, 931.

Hotaling, S., Finn, D.S., Joseph Giersch, J., Weisrock, D.W. and Jacobsen, D. (2017), 'Climate
change and alpine stream biology: progress, challenges, and opportunities for the future',
Biological Reviews, 92, 2024-2045.

Hotaling, S., Muhlfeld, C.C., Giersch, J.J., Ali, O.A., Jordan, S., Miller, M.R., Luikart, G. and
Weisrock, D.W. (2018), 'Demographic modelling reveals a history of divergence with gene
flow for a glacially tied stonefly in a changing post-Pleistocene landscape', *Journal of
Biogeography*, 45, 304-317.

Jordan, S., Giersch, J.J., Muhlfeld, C.C., Hotaling, S., Fanning, L., Tappenbeck, T.H. and
Luikart, G. (2016), 'Loss of genetic diversity and increased subdivision in an endemic
alpine stonefly threatened by climate change', *PLoS One*, 11, e0157386.

Krueger, F. (2015), 'Trim Galore!: A wrapper tool around Cutadapt and FastQC to consistently
apply quality and adapter trimming to FastQ files'.

Macdonald, H.C., Cunha, L. and Bruford, M.W. (2016), 'Development of genomic resources for
four potential environmental bioindicator species: *Isoperla grammatica, Amphinemura*
sulcicollis, Oniscus asellus and *Baetis rhodan*i', *bioRxiv*, 046227.

Macdonald, H.C., Ormerod, S.J. and Bruford, M.W. (2017), 'Enhancing capacity for freshwater
conservation at the genetic level: a demonstration using three stream
macroinvertebrates', *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27, 452-
461.

Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen, P.B., Ware, J.,
Flouri, T., Beutel, R.G., Niehuis, O., Petersen, M., Izquierdo-Carrasco, F., Wappler, T.,
Rust, J., Aberer, A.J., Aspock, U., Aspock, H., Bartel, D., Blanke, A., Berger, S., Bohm,
A., Buckley, T.R., Calcott, B., Chen, J., Friedrich, F., Fukui, M., Fujita, M., Greve, C.,
Grobe, P., Gu, S., Huang, Y., Jermiin, L.S., Kawahara, A.Y., Krogmann, L., Kubiak, M.,
Lanfear, R., Letsch, H., Li, Y., Li, Z., Li, J., Lu, H., Machida, R., Mashimo, Y., Kapli, P.,
McKenna, D.D., Meng, G., Nakagaki, Y., Navarrete-Heredia, J.L., Ott, M., Ou, Y., Pass,
G., Podsiadlowski, L., Pohl, H., von Reumont, B.M., Schutte, K., Sekiya, K., Shimizu, S.,
Slipinski, A., Stamatakis, A., Song, W., Su, X., Szucsich, N.U., Tan, M., Tan, X., Tang,
M., Tang, J., Timelthaler, G., Tomizuka, S., Trautwein, M., Tong, X., Uchifune, T., Walzl,
M.G., Wiegmann, B.M., Wilbrandt, J., Wipfler, B., Wong, T.K., Wu, Q., Wu, G., Xie, Y.,
Yang, S., Yang, Q., Yeates, D.K., Yoshizawa, K., Zhang, Q., Zhang, R., Zhang, W., Zhang,
Y., Zhao, J., Zhou, C., Zhou, L., Ziesmann, T., Zou, S., Li, Y., Xu, X., Zhang, Y., Yang,
H., Wang, J., Wang, J., Kjer, K.M. and Zhou, X. (2014), 'Phylogenomics resolves the
timing and pattern of insect evolution', *Science*, 346, 763-767.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V. and Zdobnov, E.M. (2015),
'BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs', *Bioinformatics*, 31, 3210-3212.

Simpson, J.T. (2014), 'Exploring genome characteristics and sequence quality without a
reference', *Bioinformatics*, 30, 1228-1235.

Tronstad, L.M., Hotaling, S. and Bish, J.C. (2016), 'Longitudinal changes in stream invertebrate
assemblages of Grand Teton National Park, Wyoming', *Insect Conservation and Diversity*,
9, 320-331.

US Fish & Wildlife Service. (2016), 'Endangered and threatened wildlife and plants; 12-month
finding on a petition to list the western glacier stonefly as an endangered or threatened
species; proposed threatened species status for Meltwater Lednian Stonefly and Western
Glacier Stonefly', *Federal Register*, 81, 68379-68397.

Utturkar, S.M., Klingeman, D.M., Land, M.L., Schadt, C.W., Doktycz, M.J., Pelletier, D.A. and
Brown, S.D. (2014), 'Evaluation and validation of de novo and hybrid assembly techniques
to derive high-quality genome sequences', *Bioinformatics*, 30, 2709-2716.

Tables:

Table 1. Assembly statistics for the nuclear genome of *Lednia tumana* (Plecoptera: Nemouridae)

and two other stonefly species, *Amphinemura sulcicollis* and *Isoperla grammatica* (Macdonald et

al., 2016; Macdonald et al., 2017). For *L*. tumana, only contigs larger than 500-bp were included.

"BUSCOs" are single-copy, orthologous genes known to be highly conserved among insects. A

total of 1,658 BUSCOs were searched.

	L. tumana	*A. sulcicollis*	*I. grammatica*
Estimated genome size	536,700,000	n/a	n/a
Assembly size	304,502,267	271,924,966	509,522,935
Coverage	~60x	~1.4x	~0.7x
Contigs >1 kb	74,445	51,555	53,204
Contigs >10 kb	4,608	849	4
Contig N50	4.69 kb	0.85 kb	0.46 kb
%A/T	58.4	58.2	59.9
%G/C	41.5	42.8	40.1
%N	0.1	0	0
Complete BUSCOs	1172 (70.6%)	837 (50.5%)	221 (13.3%)
Fragmented BUSCOs	368 (22.2%)	519 (31.3%)	610 (36.8%)
Missing BUSCOs	118 (7.2%)	302 (18.2%)	827 (49.9%)

210   **Figures:**

Figure 1. (a) The nearly complete mitogenome of *Lednia tumana* (Plecoptera: Nemouridae). (b)

Location of tRNAs in the *L. tumana* mitogenome. (c) Location of protein coding genes (PCGs)

and rRNA genes in the *L. tumana* mitogenome.