Diversity of acquired adaptation to fluconazole is influenced by genetic background and ancestral fitness in *Candida albicans*

Aleeza C. Gerstein[1,2] and Judith Berman[1]

[1] Department of Molecular Microbiology and Biotechnology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel
[2] Current Address: Departments of Microbiology & Statistics, The University of Manitoba, Winnipeg, Canada

Correspondence
Aleeza C. Gerstein: acgerstein@gmail.com
Judith Berman: jberman@post.tau.ac.il

Abstract

The role of genetic background in determining the evolutionary potential of a strain to acquire drug resistance is not well understood. We evolved replicates of twenty different *Candida albicans* strains for 100 generations in the common antifungal drug fluconazole. The majority of replicates rapidly increased in fitness in the evolutionary environment, and the degree of improvement was inversely correlated with ancestral strain fitness under those conditions. Unexpectedly, very few replicates in any genetic background acquired *bona fide* drug resistance. Rather, changes in drug tolerance (slow subpopulation growth at high concentrations of drug) and karyotype (aneuploidy and ploidy) were extremely prevalent. Importantly, poor ancestral strain growth in drug was correlated with high levels of evolved variation among replicates in growth, tolerance, and karyotype. This provides a link between strain background, mediated in part through ancestral growth ability, and the degree of acquired genotypic and phenotypic diversity during adaptation.

Introduction

The importance of genetic background in the link between genotype and phenotype for *de novo* acquired mutations is increasingly appreciated in both laboratory and natural settings (Chandler et al. 2013; Fournier and Schacherer 2017; Wong 2017, and references within). For example, the phenotype of deleted or repressed genes can vary significantly in different backgrounds (Paaby et al. 2015; Vu et al. 2015) and even the classification of genes as essential for growth under a specific condition can differ between closely related strains (Dowell et al. 2010). The same is true of mutations classified as beneficial in one strain background (including different mutations in the same gene, Peng et al. 2018), which can be neutral or deleterious in other backgrounds (e.g., MacLean et al. 2010; Wang et al. 2013, 2016, Hou et al. 2016). Similarly, when human disorders are caused by variation in a single gene (monogenic traits), background-specific modifiers can lead to significant clinical phenotypic heterogeneity (i.e., variable expressivity, Cooper et al. 2013, and references within).

The acquisition of drug resistance is an inherently evolutionary process, and genetic background may directly determine the nature or scope of available mutations and their evolutionary potential. In principle, a small, finite number of cellular mechanisms can lead to drug resistance, defined as the ability of all cells in a population to grow at drug concentrations above a specific, clinically-established breakpoint, measured as the minimal inhibitory concentration (MIC) of the drug. However, even for well-studied pathogen/drug pairs, the factors that influence the rate at which resistance can be acquired have not been well elucidated. Thus, understanding the processes that lead to the appearance and spread of reduced drug susceptibility can help inform clinical decisions.

Drug tolerance, a property distinct from drug resistance, is broadly defined as the ability of cells to grow slowly at drug concentrations above the minimum inhibitory concentration (MIC) (Fridman et al. 2014; Delarze and Sanglard 2015), and correlates well with the size of the subpopulation of cells that can form small colonies in drug as well as the rate at which they do so (Rosenberg et al. 2018). Importantly, strains that exhibit tolerance are more likely to cause clinically persistent infections (i.e., those not resolved by a single course of antifungal therapy) compared to strains with similar MIC levels that are cleared by a single course of the drug (Rosenberg et al. 2018). Tolerance can be quantified using either disk diffusion assays or broth microdilution measurements (Gerstein et al. 2016;

Rosenberg et al. 2018). Of the handful of experimental evolution experiments that have been conducted with *C. albicans* (Cowen et al. 2000, 2001; Selmecki et al. 2009; Anderson et al. 2015), none have distinguished between changes in drug tolerance and drug resistance during sustained exposure to drug.

The acquisition of drug resistance differs fundamentally between fungi and bacteria (Levin and Bergstrom 2000). Fungi do not acquire drug resistance via the lateral gene transfer mechanisms commonly used by many bacterial pathogens. Rather, they evolve primarily via vertical transmission of *de novo* mutations, as well as via copy number variations. Karyotypic variation, whether via changes in ploidy (the number of homologous chromosome sets) or aneuploidy (chromosome gain or loss), is frequently observed in laboratory strains subjected to stress (Wellington and Rustchenko 2005; Bouchonville et al. 2009), in strains passaged through mice (Forche et al. 2009, 2018), as well as in clinical isolates of *C. albicans, C. glabrata* and *S. cerevisiae* (Ford et al. 2015; Zhu et al. 2016). Karyotype changes appear much more frequently than point mutations (Zhu et al. 2014), and are especially prevent in strains exposed to azole drug stress (Perepnikhatka et al. 1999; Gerstein et al. 2015; Hickman et al. 2015; Anderson et al. 2017; Hirakawa et al. 2017).

A mechanistic link between karyotypic variation and drug resistance has been determined in some pathogenic fungi. In *C. albicans*, the most prevalent human fungal pathogen, isolates that carry an isochromosome 5L (i[5L]) were detected frequently among isolates resistant to the antifungal drug fluconazole [(Selmecki et al. 2006)](#) and resistance is conferred by the extra copies of *ERG11*, the target of fluconazole, and *TAC1*, a positive regulator of drug efflux (Selmecki et al. 2008). Similarly, *Cryptococcus neoformans* isolates aneuploid for chromosome 1, which carries both the *ERG11* ortholog and *AFR1,* encoding a drug efflux pump, are transiently fluconazole-resistant (Sionov et al. 2010; Ngamskulrungroj et al. 2012). Additional aneuploidies that increase the ability to grow in fluconazole without a concrete link to overexpression of a particular gene have also been noted in both species (e.g., chromosome 4 in *C. neoformans,* Sionov et al. 2010; Ngamskulrungroj et al. 2012*)*, chromosomes 3, 4, and 6 in *C. albicans* (Perepnikhatka et al. 1999; Anderson et al. 2017; Hirakawa et al. 2017). Intriguingly, certain aneuploidies appear to be beneficial in some strain backgrounds but not in others (e.g., Anderson et al. 2017; Hirakawa et al. 2017). Thus, karyotype changes have the potential to provide a rapid, potentially beneficial and reversible, response to drug stress. Furthermore, exposure to fluconazole and other azole drugs also can drive karyotypic change by inducing unconventional cell cycle events (Harrison et al. 2014; Altamirano et al. 2017). Hence, fluconazole can both cause, and select for,

karyotypic variation, and distinguishing between cause and effect is necessary for understanding the mechanistic routes to acquire drug resistance.

Here, we set out to explore the interplay between genetic background, resistance, tolerance, and karyotypic variation by following the evolutionary trajectories of 20 different *C. albicans* strains, 12 replicates per strain, for 100 generations in a low concentration (1µg/ml) of fluconazole. The majority of replicates rapidly acquired the ability to grow more in the evolutionary level of drug, with the degree of improvement inversely correlated with ancestral strain growth in the low drug environment. While very few replicates from any background acquired *bona fide* drug resistance, changes in tolerance and karyotype were prevalent, especially in strains with low ancestral growth. Thus, tolerance to drug is an evolvable phenotype, distinct from the acquisition of drug resistance and it changes much more frequently than resistance. Importantly, ancestral strain fitness was inversely correlated with evolved variation in growth, tolerance, and karyotype, providing a link between strain background and the degree of parallelism among the adaptive trajectories of replicate isolates.

Results

Twenty different clinical strains of *C. albicans* spanning the phylogenetic diversity of the species (Ropars et al. 2018), and varying in mating type zygosity, geographic origin and site of isolation (Table 1) were evolved in fluconazole (YPD + 1 µg/mL FLC) for ~100 generations (20 strains x 12 replicates per strain = 240 replicates total). Growth was measured as optical density (OD at A_{600}) at 24 and 72 h for all ancestral and evolved replicates. OD correlates with cell size and density, and reflects the ability to convert nutrients into cellular biomass. OD at 24 h is also consistent with clinical assessment of drug resistance, and correlates with growth rate (Gerstein et al. 2012). Drug resistance (MIC_{50}) was measured using standard liquid broth microdilution methods (Clinical and Laboratory Standards Institute, CLSI, http://clsi.org/) at 24 h. Tolerance was measured as average growth above the MIC_{50} normalized to total growth without drug in a liquid assay (SMG: supra-MIC_{50} growth, Rosenberg et al. 2018) and was assessed at both 24 and 72 h.

Table 1. Strain used in this study. Ancestral minimum inhibitory concentration of drug (MIC_{50}) was measured in YPD at 30 °C.

Strain #	Strain Name	Clade	*MTL* genotype, patient status (if known), site of isolation, country of origin	Aneuploidies	Ancestral MIC	Original Reference
A1	L26	1	a/a, vaginitis, vagina, USA	trisomic chr7	4	Wu 2007
A2	P87	4	a/a, HIV, oral, South Africa	-	1	Wu 2007
A3	GC75	4	α/α, healthy, oral, South Africa	-	0.0125	Wu 2007
A4	P78048	1	α/α, bloodstream, Canada	-	0.5	Wu 2007
A5	P57072	2	α/α, bloodstream, USA	-	4	Wu 2007
A6	P34048	3	a/α, bloodstream, Turkey	-	1	Wu 2007
A7	P37037	1	a/α, healthy, oral, USA	-	1	Wu 2007
A8	P75016	4	a/α, bloodstream, Israel	-	0.5	Wu 2007
A9	P75063	4	a/α, bloodstream, France	-	0.0125	Wu 2007
A10	P76055	2	a/α, bloodstream, USA	-	0.0125	Wu 2007
A11	P78042	3	a/α, bloodstream, USA	trisomic chr4	0.0125	Wu 2007
A12	T101	3	a/a, oropharynx, Canada	-	32	Odds 2007
A13	OKP90	2	a/a, healthy, oral, South Africa	-	0.0125	Lockhard 2002
A14	AM2003.089	2	a/α, oropharynx, UK	-	0.0125	Odds 2007
A15	AM2003.0165	2	α/α, bloodstream, UK	-	0.0125	Odds 2007
A16	AM2003.0069	3	a/α, vagina, UK	-	0.0125	Odds 2007
A17	SC5314	1	a/α, bloodstream, USA	-	0.0125	Fonzi 1993
A18	FH1	3	a/α, marrow transplant, rectal, USA	-	4	Marr 1997
A19	DSY294	11	a/α, HIV, oral, France	-	0.0125	Sanglard 1995, MacCallum 2010
A20	T118	1	a/α, HIV, oral, Canada	-	4	Cowen 2000

Adaptation is influenced by strain background

The majority of evolved replicates acquired significantly higher growth ability than their respective ancestral strains in the evolutionary drug environment (Figure 1; detailed statistical results in Table S1). The major exception was A12, which had the highest ancestral MIC_{50} of 32 µg/mL, well above the 1 µg/mL drug concentration evolutionary environment, and evolved significantly lower 24 h growth, with no change in average growth at 72 h. Four strains with initial MIC ≤ 1 (A2, A9, A10 and A14) also did not acquire improved average growth among replicates at 24 h, though on average all had significantly increased growth at 72 h.

Figure 1. The majority of evolved replicates improved growth in the evolutionary environment after ~100 generations of evolution. Fitness was measured as optical density in YPD + 1 µg/mL fluconazole, the evolutionary environment after 24 h (top) and 72 h (bottom). Strains are ordered by ancestral growth in the evolutionary environment at 24 h. Ancestral growth ability (the median growth among 12 ancestral replicates) is indicated for each strain by a grey bar. Each coloured point represents one of twelve evolved replicates while the coloured bars indicate median evolved growth for visual comparison to ancestral growth.

Ancestral strain growth ability strongly influenced the mean degree of growth improvement among replicates, while neither mating type nor clade had a significant effect (Figure 2, ANOVA tests; 24 h—ancestral growth: $F_{1, 13} = 7.99$, $p = 0.014$, clade: $F_{4, 13} = 0.64$, $p = 0.64$, *MAT* zygosity: $F_{1, 13} = 0.27$, $p = 0.61$; 72 h—ancestral growth: $F_{1, 13} = 158.73$, $p < 0.0001$, clade: $F_{4, 13} = 1.97$, $p = 0.16$, MAT zygosity: $F_{1, 13} = 0.25$, $p = 0.63$). The change in growth (improvement in growth between evolved and ancestral replicates) and the ancestral growth were very strongly correlated at both time points (Pearson's correlation; 24 h: $t18 = -3.01$, $p = 0.007$, cor = -0.58. 72 h : $t_{18} = -11.63$, $p < 0.00001$, cor = -0.94). The influence of strain background is evident in those strains that deviated considerably from the correlation line of best fit, particularly at 24 h (Figure 2).

The variance among evolved replicates from each strain reflects stochasticity in the evolutionary process. The variance in evolved growth and the degree of growth improvement were inversely correlated: evolved replicates that were least fit initially and that increased in growth the most also had the most variability among replicates from the same strain background (Spearman's rank correlation test; 24 h: S = 694, $p = 0.035$, rho = 0.48. 72 h: S = 476, $p = 0.003$, rho = 0.64). Overall, strain background influenced both the degree of growth improvement and the growth variability among replicates, both mediated in part by the ancestral growth ability of the strain in the evolutionary environment.

Figure 2. Significant negative correlation between ancestral growth and change in growth. Growth was measured as optical density in YPD+1 µg/mL FLC, the low drug evolutionary environment after 24 h (top) and 72 h (bottom). The colour matches Figure 1, based on ancestral growth (low ancestral growth strains are coloured red, with colours proceeding as the colours of the rainbow to the highest initial 24 h growth strain in violet).

Minimal tradeoff to permissive environment

A tradeoff is often seen between the cost and benefit of mutations that enable microbes to grow well in antimicrobial drugs, although several drug classes and pathogen species do not impart a major cost (Melnyk et al. 2015, and references within). A similar pattern of species/drug specificity on tradeoffs appears for mutations that provide an advantage to fungi exposed to antifungal drugs. For example, the ability to grow in nystatin incurs a significant cost in *S. cerevisiae* (Gerstein et al. 2012), yet mutations that confer an advantage to high concentrations of fluconazole in *C. albicans* incur no appreciable cost (Anderson et al. 2003; Selmecki et al. 2009). The degree to which there is a cost-benefit tradeoff for growth in lower concentrations of fluconazole is not known.

Strain background had a significant impact on the tradeoff between growth in 1 µg/ml fluconazole and growth in the absence of the drug. After 100 generations of evolution in 1µg/ml fluconazole (Figure 3), 80% of evolved replicates grew more slowly in the drug-free environment at 24 h, with an average growth reduction of 21% compared to the ancestral replicates. Exceptions were strain A5, where the majority of evolved replicates showed a significant growth improvement (6% on average), and strains A8, A10 and A11, where evolved replicates had no significant change on average (Table S2). Evolved growth measured at 72 h depended on strain background (Table S2; average growth improvement of 1%: A1, A4, A5, A10, A11, A12, A20; average growth reduction of 5%: A3, A13, A14, A15, A17). Overall, the majority of the evolved replicates exhibited slower initial growth in a drug-free environment, yet reached the same or slightly higher growth by 72 h. The primary trade-off for the drug-free environment was thus in the initial phase of growth, suggesting an increased amount of time spent adjusting to the growth conditions.

Mean growth in the drug and drug-free environments at 24 h was significantly correlated among strain backgrounds (Spearman's rank correlation test; S = 396, *p* = 0.0008, rho = 0.702, Figure S1; there was not enough variation in the drug free environment at 72 h for correlative purposes). A significant positive relationship was also recovered when comparing among evolved replicates of the same strain background for 12 of 16 strain backgrounds with variable replicates (the four backgrounds with the highest ancestral growth had extremely low variation in growth in both drug-free and low drug environments; Figure S2). Thus, the overall pattern among strain backgrounds and different replicates of the same strain was similar: strains (and replicates) that evolved higher growth in low drug also had an advantage in the no drug environment. Hence, mutations acquired in response to a

low concentration of fluconazole had an initial cost at 24 h in the no-drug environment, though this was mitigated when growth was allowed to proceed for 72 h.

Figure 3. Growth in a drug-free environment. Growth was measured as optical density in YPD, the standard lab rich medium, after 24 h (top) and 72 h (bottom). Strains are ordered by ancestral growth ability in the low drug evolutionary environment. Ancestral growth in YPD is indicated by a grey bar. Each coloured bar represents the median of twelve replicates evolved from each ancestral strain.

Increases in *bona fide* drug resistance were rare and changes in drug tolerance were common

Drug susceptibility/resistance is a concentration-dependent property of individual strains that is measured as the MIC_{50} using broth microdilution assays. An increase in MIC_{50} beyond the evolutionary level appeared in only 18 of the 240 replicates, which hailed from seven of the 20 strain backgrounds (Figure 4), with only slight

increases in MIC_{50} in the majority of them (to 4 µg/mL; five replicates from strain A3, four replicates from A8, and one replicate from A4 and A7). Clinical resistance ($MIC_{50} \geq 8$ µg/mL at 24 h, Fothergill et al. 2014) was acquired in replicates from only three backgrounds: six from strain A5 (initial MIC_{50} of 4 µg/mL) and one each from strains A17 and A19, (initial MIC_{50} of 0.0125 µg/mL). Interestingly, MIC_{50} decreased to 1µg/mL in some replicates from the five strains with initial MIC_{50} >1 µg/mL (two replicates from A5, eight replicates from A12, all 12 replicates from A20, Figure 4). This suggests a fitness cost of initially high MIC_{50} during prolonged growth in low concentrations of drug, albeit in some, but not all, strain backgrounds.

Figure 4. The majority of evolved replicates did not acquire resistance. Drug resistance was measured through broth microdilution assays. The clinical cutoff for resistance at 8 µg/mL fluconazole is indicated by the dashed line. Strains are arranged on the x-axis (and coloured) by initial growth in the evolutionary environment. Each point represents an individually-evolved replicate line.

Increases and decreases in tolerance were prevalent among evolved replicates (Figure 5, Figure S3), unlike the infrequent evolution of increased resistance. Approximately one third of the evolved replicates had increased tolerance (24h: 72 replicates from 13 different strain backgrounds; 72h: 73 replicates from 13 strain backgrounds), while a similar number of replicates had decreased tolerance at both time points (24 h: 79 replicates from 17 strain backgrounds; 72 h: 91 replicates from 17 strain backgrounds). The magnitude of tolerance change increased sharply

between 24 and 48 h of drug exposure (Figure S3), consistent with a previous examination of SMG (Rosenberg et al. 2018). To match the timing of our transfers and growth measurements, we primarily considered SMG at 72 h as a measure of tolerance in evolved replicates (SMG at 48 h and 72 h are very similar, Figure S3). The number of replicates that increased or decreased in tolerance varied considerably among strain backgrounds (Figure 5, Figure S4). Notably, in some strain backgrounds nearly all replicates changed in the same direction, while in others the replicates were split between those that increased and those that decreased evolved tolerance. Change in tolerance was not significantly correlated with initial growth, clade, or mating type (*MAT* locus*)* status in the evolutionary environment (linear mixed-effects model with change in tolerance as the response variable and initial growth, zygosity and clade as predictor variables and strain as a random effect; initial growth: $F_{1,217} = 0.51$, $p = 0.47$, clade: $F_{4,14} = 1.79$, $p = 0.19$, MAT zygosity: $F_{1,14} = 2.53$, $p = 0.13$). Variation in evolved growth and the variance in tolerance were positively correlated (Spearman's rank correlation: growth measured at 24 h: S = 588, p-value = 0.012, rho = 0.56; this relationship was not statistically significant for variability in growth at 72 h though rho was still relatively high: S = 762, p = 0.06, rho = 0.43). Thus, genetic background, mediated in part by ancestral growth, influenced the evolved variation among replicates in the evolution of tolerance as well as resistance.

Genome size changes are pervasive

Given that exposure to fluconazole induces the formation of tetraploid, and subsequently aneuploid, cells (Harrison et al. 2014), and that specific aneuploidies can be selectively advantageous (Selmecki et al. 2006; Yang et al. 2013; Anderson et al. 2017), we asked if evolved replicates exhibited altered ploidy and, if so, whether the degree to which ploidy was changed was a strain background-dependent characteristic.

Evolved replicates underwent a significant increase in median genome size across the majority of strain backgrounds (Figure 6A, Table 2). The three exceptions were A12, the resistant strain; and A4 and A9, initial MIC_{50} values < 1 µg/mL. The coefficient of variance for ploidy level (a measure of variability among replicates) also increased for the majority of strains (Figure 6B). Thus, genome size variation among replicates increased over time in drug, as expected if evolved replicates acquired different final numbers of chromosomes.

Table 2. Welch two sample t-test comparing ancestral and evolved genome size.

Strain	Statistics
A1	$t_{20.6} = 11.70$, $p < 0.0001$
A2	$t_{11.2} = 5.34$, $p = 0.002$
A3	$t_{11.1} = 5.69$, $p = 0.0001$
A4	$t_{11.4} = 1.91$, $p = 0.08$
A5	$t_{20.14} = 18.07$, $p < 0.0001$
A6	$t_{11.1} = 3.82$, $p = 0.003$
A7	$t_{11.1} = 2.80$, $p = 0.017$
A8	$t_{11.1} = 2.80$, $p = 0.017$
A9	$t_{11.2} = 0.88$, $p = 0.40$
A10	$t_{11.4} = 3.31$, $p = 0.007$
A11	$t_{11.5} = 5.24$, $p = 0.0002$
A12	$t_{18.8} = 0.71$, $p = 0.49$
A13	$t_{11.1} = 5.46$, $p = 0.0002$
A14	$t_{11.1} = 4.10$, $p = 0.002$
A15	$t_{11.1} = 2.79$, $p = 0.018$
A16	$t_{11.3} = 4.17$, $p = 0.0015$
A17	$t_{11.1} = 2.94$, $p = 0.013$
A18	$t_{18.4} = 2.98$, $p = 0.008$
A19	$t_{11.4} = 2.93$, $p = 0.013$
A20	$t_{11.5} = 5.91$, $p < 0.0001$

Importantly, the least ploidy variation in evolved replicates was seen in replicates evolved from the four parental strain backgrounds that had the highest initial growth levels in low drug and MIC_{50} above 1 µg/mL (Figure 6C, bottom four panels). Median evolved genome size was neither correlated with ancestral low drug growth (Pearson's correlation, 24h: $t14 = -0.34$, $p = 0.74$; 72 h: $t14 = -0.32$, $p = 0.75$), nor with the change in low drug growth (Spearman's rank correlation, 24 h: $S = 541.19$, $p = 0.45$; 72 h: $S = 953.4$, $p = 0.12$), when these four strains were removed from the analysis, indicating that a different factor influences karyotypic changes among strain backgrounds.

Figure 6. Genome size and variation in genome size increased after evolution to low fluconazole. a) Median genome size, and b) coefficient of variation (CV; i.e., variability among replicates) in each strain background. A dashed line indicates a non-significant change between ancestral and evolved replicates. c) Flow cytometry traces of each replicate evolved line, arranged by ancestral strain growth in low fluconazole from lowest (A13) to highest (A18).

These median numbers, however, obscure the tremendous variation in ploidy level and evolved growth observed among independently evolved replicates. Comparison of evolved genome size and evolved growth for replicates from the 16 strain backgrounds that evolved variable genome size revealed that evolved growth at 24 h and evolved genome size correlated significantly in five strains: negatively in A6, A8, A10 and A16 (replicates that increased in genome size tended to have less growth at 24 h), and positively in A13 (replicates that increased in genome size tended to have more growth at 24 h; Figure 7, Table S3). Evolved growth in strain A13 at 72 h remained positively correlated with genome size. By contrast, two different strains (A2 and A4) had weak, but significant negative correlations between growth at 72 and evolved genome size (Figure S4, Table S3). Across strain

backgrounds, no clear association between alterations in genome size and increases in MIC_{50} (Figures 7, filled points), or evolved changes in tolerance (Figure S5) were evident. Importantly, once again, the variance in evolved genome size correlated with ancestral growth (Pearson's correlation, $t_{18} = -3.22$, $p = 0.0047$, cor = -0.60) and the variance in evolved growth (Pearson's correlation, $t_{18} = 5.65$, $p = 0.00002$, cor = 0.80). Hence, there was no direct link between evolved genome size and evolved growth in most strain backgrounds, yet the variability among replicates is consistently larger in some strain backgrounds than others, mediated in part by ancestral strain growth in the evolutionary environment.

Figure 7. Evolved growth and evolved genome size are not consistently correlated. Growth was measured as optical density after 24 hours in low fluconazole. Evolved genome size indicates the most prominent G1 peak; when multiple G1 peaks were present this is indicated with a triangle symbol. Solid lines indicate a significant correlation. Each point represents an independently evolved replicate line. Filled-points are those lines that increased in MIC beyond the level of drug stress (1 μg/mL fluconazole) they were evolved in. Strains are arranged from the lowest initial growth (top left) to highest (bottom right). Replicates from the four strains that had ancestrally high MIC levels above the evolutionary environment did not evolve variation in genome size and are not presented here.

Discussion

The acquisition of *bona fide* drug resistance by microbial pathogens is inherently an evolutionary process that is dictated in part by the specifics of the population and the evolutionary environment. Yet few experiments have directly quantified the influence of genetic background on the nature and rate of adaptation. Here, we evolved 12 replicates from 20 *C. albicans* strains to a relatively low concentration of fluconazole (1 µg/ml) that is nevertheless above the ancestral MIC_{50} of the majority of strains. The short period of laboratory evolution (100 generations) was chosen to capture the first one or several beneficial mutations in each replicate, based on the rationale that mutations that increase MIC_{50} in a clinical setting would typically be the first beneficial mutations that confer the ability to grow in the presence of the drug stress. Importantly, the most prevalent changes were increases in growth in the drug environment and changes in drug tolerance while changes in MIC_{50} were rare.

The majority of replicates rapidly increased in growth ability in the evolutionary environment in a manner inversely correlated with ancestral strain growth in the drug. Similar results from experimental evolution studies were reported for eight divergent bacteriophage genotypes (Rokyta et al. 2009) and for five different *E. coli* strain backgrounds (Moore and Woods 2006). This pattern of diminishing-returns for fitness is also consistent with a suite of related microbial experiments: a negative correlation between ancestral fitness and rate of adaptation in experimental evolution studies that were initiated with closely related strains of the same ancestral background that differed by only one or a few beneficial (Kryazhimskiy et al. 2014; Wünsche et al. 2017) or deleterious mutations (Perfeito et al. 2014); the effect size of a beneficial mutation tends to be smaller in a background that already contains a different beneficial mutation (Chou et al. 2011; Khan et al. 2011; Ono et al. 2017); the effect size of beneficial mutations transferred onto a diversity of strain backgrounds is negatively correlated with ancestral fitness (Wang et al. 2016); and compensatory adaptation occurs more rapidly for severely deleterious mutations compared to weakly deleterious mutations (Moore et al. 2000; Sanjuán et al. 2005; Barrick et al. 2010). Couce and Tenaillon (2015) found a robust match between these observed patterns and Fisher's abstract geometric model (Fisher 1930, Tenaillon 2014) through simulation analyses, providing a theoretical underpinning for the negative relationship between the distance from the fitness optimum (ancestral fitness) and the number and/or effect size of beneficial mutations (rate of adaptation). We thus predict to uncover a negative correlation

between ancestral growth ability and the effect size of the mutations that underpin the observed fitness improvements, a hypothesis we will test in future work.

Of note, in our study the correlation between ancestral growth and change in growth was stronger when growth was measured at 72 hours (adjusted r-squared from the log-log regression = 0.96) compared to 24 hours, the time when MIC is clinically determined (adjusted r-squared = 0.72). Thus, although the pattern of a declining rate of adaptability with increased ancestral growth holds across species (e.g., Couce and Tenaillon 2015), it is worth considering whether some aspects of the strain background influence deviations from this pattern, particularly during adaptation to a fungistatic drug such as fluconazole, which inhibits cell growth. Six strains had a notably lower growth improvement than expected at 24 h: A2, A3, A9, A10, A12, A14. Nothing that we know about these strains distinguishes them from the others—they are from different clades, include both homozygous and heterozygous mating loci, and were isolated from diverse geographic locations and clinical niches. Nevertheless, as we conduct additional experiments with these strains, it will be interesting to see whether the same strains repeatedly behave as outliers, and to further probe how the role of tolerance vs resistance affects growth measurements and the interpretations of the role of ancestral fitness.

Very few (<5%) of replicates with initial MIC_{50} at or below 1 µg/mL fluconazole increased their MIC_{50} to > 1 µg/mL, in contrast to the rapid growth improvements observed in the evolutionary environment and widespread changes in tolerance (SMG). Thus, there was only a small window of MIC_{50} improvement that did not extend beyond the concentration of drug in which the replicates were evolved. The influence of subinhibitory drug concentrations on MIC_{50} remains under-studied in fungi. In *Sclerotinia sclerotiorum,* a plant pathogenic fungus, no consistent relationship was found between increased resistance after repeated exposure to sublethal concentrations of different antifungal drugs. In three different drugs, approximately half of the strains increased in MIC_{50}, while in two additional drugs almost no strains acquired increased MIC_{50} (Amaradasa and Everhart 2016)**. By contrast,** many studies in bacteria have, however, found that exposure to subinhibitory concentrations of antibiotics can select for *de novo* mutations that confer resistance (Wu et al. 1999; Henderson-Begg et al. 2006; Gullberg et al. 2011; Chow et al. 2015; Torres-Barceló et al. 2015; Wistrand-Yuen et al. 2018) or augment existing resistance mutations (e.g., Gullberg et al. 2011; Liu et al. 2011; McVicker et al. 2014; Mira et al. 2015; Lundström et al. 2016). This highlights the important influence of different drugs as well as the potential differences between different organisms on the breadth and molecular influence of drug resistance mutations.

Strikingly, one-third of evolved replicates with ancestral MIC_{50} above 1μg/mL fluconazole (22 of 60) evolved a reduced MIC_{50} (i.e., higher of susceptibility to fluconazole). The four strains that initially had an MIC_{50} of 4 μg/mL exhibited patterns consistent with an effect of strain background on the propensity for MIC_{50} reduction: all replicates from two backgrounds (A1, A18) maintained an MIC_{50} of 4 μg/mL, while the MIC_{50} decreased to 1 μg/mL in all 12 replicates from strain A20 (fisher-exact test, $p < 0.00001$). Replicates from the fourth strain (A5) were extremely variable, as two replicates decreased to MIC_{50} 1 μg/mL, while seven increased to > 4 μg/mL, and the remainder did not change. Replicates from a fifth strain, A12, which was initially clinically resistant at MIC_{50} 32 μg/mL, also evolved to lower MIC_{50}. That few ancestrally low-MIC_{50} replicates increased in MIC_{50}, while many ancestrally-high MIC_{50} replicates decreased in $MIC_{50,}$ suggests that a cost-benefit trade-off exists between alleles that provide high fitness in low drug and those that provide high fitness in high drug, in at least some strain backgrounds. Furthermore, it demonstrates that *C. albicans* MIC_{50} can rapidly decrease upon exposure to low concentrations of fluconazole, a result that future experiments in an *in vivo* context (e.g., *in vivo* evolution in mice) will seek to replicate.

We demonstrate for the first time that drug tolerance, the ability of a subpopulation to grow slowly at high drug concentration, often evolves during exposure to drug. Over half of the replicates evolved an altered tolerance level, with both increased and decreased detected. This implies that the ability of a subpopulation to grow in concentrations of drug beyond the MIC_{50} evolves at a higher frequency than does the acquisition of *bona fide* resistance. There were significant differences among strains in the propensity for tolerance levels to change. For example, 12/12 replicates from strain A3 increased in tolerance, while 9/12 replicates from strains A14 and A16 decreased in tolerance (with no change in the other three replicates). Other strains exhibited variability among replicates (e.g., strain A9—four replicates decreased while six replicates increased tolerance). The lack of correlation of evolved tolerance with ancestral growth levels, MIC_{50}, or known strain characteristics (i.e., *MAT* genotype or clade), suggest that the evolution of tolerance is not easily predicted based on resistance or fitness and may well depend on the genetic background of a given strain.

The appearance of karyotypic variability among evolved replicates also was strain-dependent. The four strains with the highest initial growth in low drug (all with ancestral $MIC_{50} > 1$ μg/mL) had stable genomes with no change in their

variability after evolution. By contrast, karyotypic variability increased significantly for all other strain backgrounds. There also was diversity in the variability among replicates: replicates from some backgrounds were more variable than others, and the degree of variability in karyotype among replicates correlated significantly with variability both in evolved growth and in evolved tolerance. There was not, however, a consistent correlation between a specific evolved ploidy level and evolved growth, or a specific evolved ploidy level and tolerance. This implies that there are different ploidy states that can confer altered levels of tolerance. Thus, some strain backgrounds were more likely to yield evolved populations with variable phenotypes than others, regardless of the phenotype being measured (i.e., growth ability, tolerance, genome size), yet the actual phenotype parameter values were not necessarily correlated with each other or even in the same direction.

Aneuploidy was both caused and selected for in earlier studies using higher concentrations of fluconazole (10 μg/ml) (Selmecki et al. 2006; Harrison et al. 2014). Here, a 10-fold lower concentration of fluconazole also rapidly resulted in widespread aneuploidy within some strain populations that were under drug stress, presumably via similar mechanisms. The variation that remains to be sampled seems to be largely neutral. This study highlights the importance of the relative relationship between the drug concentration used and the ability of the initial strain to grow at that drug concentration: the same drug concentration can be 'low stress' in one strain (e.g., with an MIC_{50} well above that concentration), and likely yields a 'high stress' response in a different strain (with an MIC_{50} below that concentration). The low evolved ploidy variability observed for strains with initial MIC above the drug evolutionary environment is indicative of this difference. Nonetheless, the fact that very few replicates acquired *bona fide* drug resistance, while nearly all improved in growth in the evolutionary environment and many acquired altered tolerance levels, indicates that the responses involved in growth and tolerance to 1μg/mL fluconazole are distinct from those that yielded much higher MIC_{50} levels. Furthermore, it suggests that many more genes and genetic pathways likely influence overall growth and tolerance than the small number of genes that directly affect *bona fide* fluconazole resistance. In other words, the target size for genes affecting tolerance is likely much larger than that for *bona fide* resistance.

Conclusion

Experimental evolution studies provide an extremely useful framework to isolate important factors that influence adaptation. Here, we evolved replicates from 20 different strain backgrounds to probe the importance of strain background on *Candida albicans* evolution to 1 µg/mL fluconazole. We found that genetic background had a significant influence on the rate of adaptation, mediated in part through ancestral fitness relative to the selective conditions used, in accordance with theoretical predictions. Fitness improvements were largely restricted to the drug concentration used in the evolution experiment, indicating that adaptation to low fluconazole levels often does not select for mutants resistant to higher levels of drug. Indeed, a number of replicates from strains with initially high MIC evolved an increase in drug susceptibility. Genome size changes were prevalent among replicates evolved from strains that had ancestral MIC below the drug concentration used in the evolutionary environment, yet was largely absent from ancestral strains with MIC levels above it. This implies that drug stress is context-dependent: the level of stress conferred by exposure to a high concentration of drug to one strain may be relatively benign to another. Importantly, variability among replicates was correlated for multiple evolved phenotypes, and strains with similar ancestral fitness did not always yield the same degree of variability. Thus, strain background is an important factor that needs to be taken into consideration explicitly when making generalizations about stress responses.

Materials and Methods

Strains

Twenty diploid clinical strains of *Candida albicans* were selected to represent the phylogenetic diversity of the species. The strain set includes at least four strains from each of the four major clades that encapsulate ~70% of the typed *C. albicans* strains and nearly span the known phylogenetic diversity of the species (Odds et al. 2007; Ropars et al. 2018) as well as four commonly-studied laboratory strains (SC5314, FH1, DSY294, T188). For each clade, strains with both heterozygous (*MATa*/*MATα*) and homozygous (*MATa/MATa, MATα*/*MATα*) mating loci were chosen. The strains were chosen blind with respect to the initial minimum inhibitory concentration of drug (MIC). Full strain information including clade designation, country of origin, and infection niche were obtained from the original manuscripts (Table 1) and Tavanti et al. (2005). Mating type genotype was confirmed by PCR with *MATa* and *MATα* specific primers (*MATa* F-TTGAAGCGTGAGAGGCAGGAG, *MATa* R-GTTTGGGTTCCTTCTTTCTCATTC, *MATα* F-TTCGAGTACATTCTGGTCGCG, *MATα* R-TGTAAACATCCTCAATTGTACCCG). All strains were initially streaked onto YPD and grown for 48h at 30°C. A single colony was frozen down in 15% glycerol and stored at –80 °C. Thus, minimal genetic variation should be present in our initial freezer stock, which we refer to throughout as the "ancestral strains".

Evolution Experiment

Strains were evolved in 1 µg/mL fluconazole, the epidemiological cut off value that denotes the upper limit of drug susceptibility (MIC_{50}) in the wild type *C. albicans* population (Delarze and Sanglard 2015). This concentration of drug was equivalent to the ancestral MIC_{50} of three strains, above the MIC_{50} for 11 strains and below the MIC_{50} for one highly resistant strain (MIC_{50} = 32) and four additional strains (MIC_{50} = 4) (Table 1).

To initiate the evolution experiment we generated twelve independent replicates from each ancestral strain. Cultures were struck from frozen stocks onto YPD (1% yeast extract, 1% peptone, 2% dextrose; the standard lab rich medium) plates and incubated at 30°C overnight. For each strain, twelve colonies were randomly chosen by spotting each plate and picking the closest colony to each dot. Colonies were separately inoculated into 1 mL YPD in a deep well (3 mL) 96-culture tube box and grown shaking overnight at 30 °C.

From the overnight cultures, we froze 100 μL from each replicate in duplicate in 50% glycerol as the ancestral replicats. Overnight cultures were diluted 1:1000 into YPD+fluconazole in two steps: first, 10 μL of overnight culture was transferred into 990 μL YPD (a 1:100 dilution), followed by a transfer of 20uL diluted culture into 180 μL of YPD + 1.11 μg/mL FLC in round bottom microtiter plates. To minimize the likelihood of contamination and keep environmental conditions similar, culture from replicates from one strain was inoculated into row A while culture from replicates from a second strain was inoculated into row H. Plates were sealed with breathe-easy sealing membranes (Sigma Z380058) and incubated at 30 °C in small sealed rubbermaid containers with wet paper towels inside to minimize evaporation.

After 72 hours, wells were mixed by pipetting and another two-step 1:1000 transfer was conducted into fresh YPD + fluconazole medium. In total 10 transfers were conducted, yielding 100 generations of evolution (9.97 generations between transfers, $\log_2(1000) = 9.97 * 10$ transfers = 99.7 generations).

50μl of the evolved replicate cultures were frozen in duplicate in 50% glycerol after the third, fifth, eighth and tenth (final) transfer and maintained at -80 °C.

Resistance and tolerance

The initial susceptibility and tolerance of all strains was tested using broth microdilution liquid assays to measure minimal inhibitory concentration (MIC_{50}) and supraMIC growth (SMG), respectively. The liquid assay experiments followed the initial cell dilution regulations from the clinical CLSI M27-A guidelines, using YPD incubated at 30 °C as the base medium and optical density at A_{600} (OD), instead of a McFarland 0.5 standard, to determine the initial density of cells. OD readings were taken at 24 hours after inoculation to calculate MIC_{50} (the point at which optical density was reduced by 50% compared to optical density in YPD without drug).

Four liquid broth microdilution assays were conducted on both the ancestral and evolved replicates, and an additional two assays conducted on the ancestral replicates. We were not able to assay all drug concentrations in each assay due to capacity (2 time points x 20 strain backgrounds x 12 replicates). A single measurement was taken for each replicate at each concentration of drug measured in a given experiment (see Table S4). The median OD among experiments was

determined at each concentration of drug for each replicate. The MIC_{50} was then calculated as the highest concentration of drug with an OD greater than 50% of the measured OD in YPD (i.e., optical density in medium without drug).

Tolerance was measured from the liquid assay results as in Rosenberg et al. (2018). The sum of growth (measured as optical density, A_{600}) in the concentrations of drug that were measured beyond MIC_{50} was divided by the number of measured drug concentrations to obtain the average supra-MIC_{50} growth for each ancestral and evolved replicate. Tolerance was assessed at 24, 48 and 72 h. We consider evolved replicates to have increased or decreased in tolerance when their measured tolerance was respectively above or below the range of ancestral tolerance levels for replicates from that strain.

We used "growth" as a proxy measurement of fitness in the evolutionary (low drug) environment, measured as optical density (A_{600}) at both 24 and 72 h. Here, "growth" reflects the ability to convert nutrients from the environment into cellular biomass and is consistent with the clinical determination of drug resistance from standardized liquid assays.

Ploidy variation

Flow cytometry was performed on a BD Biosciences BD LSR II. In all cases, all replicates from the same time point were fixed, stained and measured in parallel. Ancestral replicates were measured twice independently from the freezer stocks maintained at -80 °C. Freezer cultures were thawed, mixed, and 10 µL was added to 500 µL, YPD in deep 96-well boxes, covered with a breathe-easy membrane, and shaken at 200 RPM, 30 °C overnight. After ~ 16 hours of growth, 20 µL of culture was washed in 180 µL of TE, in a round bottom microtiter plate, pelleted, and resuspended in 20 µL TE and fixed by adding 180 µL 95% cold ethanol.

Samples from evolved replicates were fixed in ethanol at the end of the last growth cycle; 50 µL of 72 h culture from t10 was washed in 150 µL TE in a round bottom microtiter plate, pelleted and resuspended in 20 µL TE and 180 µL 95% cold ethanol. Ethanol-fixed cultures were stored at -20 °C for up to 4 weeks. The remainder of the protocol was identical for both time points, following Gerstein et al. 2017. As in Gerstein et al. (2017) we used the cell cycle analysis in Flow-Jo (Treestar) to determine the mean G1 peak for each replicate; when more than one peak was evident we recorded both the major and minor G1 peaks.

Although we always used the same machine settings, subtle but significant variation is always observed in flow cytometry data. To better compare between t0 and t10 data we performed a day-correction based on the median G1 intensity of the A12 ancestral and evolved replicates, which always measured cleanly as diploids.

Statistical methods

All analysis was conducted in the R Programming Language (R Core Team 2017). To maximize statistical power when testing the influence of mating type, we examined the effect of a heterozygous mating type vs. homozygous mating type, i.e., we combined $MATa/a$ and $MAT\alpha/\alpha$ strains.

MIC_{50} values were log transformed prior to statistical analysis. When parametric tests were used, all assumptions were tested and met. When data transformations were insufficient to meet the test assumptions, nonparametric tests were used. In all cases the specific test is indicated inline.

Acknowledgements

We thank R Urbitas, D Abbey, MA Hickman, M McClellan, E Shtifman Segal, D Tank and the University of Minnesota Flow Cytometry core facility for technical support, A Selmecki for thoughtful discussions and J Hill for helpful comments on the manuscript. JB was funded by an European Research Council Advanced Award 340087 (RAPLODAPT). ACG is grateful to the Azrieli Foundation for the award of an Azrieli Fellowship, and for support from a National Sciences and Engineering Research Council of Canada Postdoctoral Fellowship and a Banting Fellowship from the Canadian Institutes of Health Research.

References

Altamirano, S., D. Fang, C. Simmons, S. Sridhar, P. Wu, K. Sanyal, and L. Kozubowski. 2017. Fluconazole-induced ploidy change in *Cryptococcus neoformans* results from the uncoupling of cell growth and nuclear division. mSphere 2.

Amaradasa, B. S., and S. E. Everhart. 2016. Effects of sublethal fungicides on mutation rates and genomic variation in fungal plant pathogen, *Sclerotinia sclerotiorum*. PLoS One 11:e0168079.

Anderson, J. B., C. Sirjusingh, A. B. Parsons, C. Boone, C. Wickens, L. E. Cowen, and L. M. Kohn. 2003. Mode of selection and experimental evolution of antifungal drug resistance in *Saccharomyces cerevisiae*. Genetics 163:1287–1298.

Anderson, M. Z., A. Saha, A. Haseeb, and R. J. Bennett. 2017. A chromosome 4 trisomy contributes to increased fluconazole resistance in a clinical isolate of *Candida albicans*. Microbiology 163:856–865.

Anderson, M. Z., L. J. Wigen, L. S. Burrack, and J. Berman. 2015. Real-time evolution of a subtelomeric gene family in *Candida albicans*. Genetics 200:907–919.

Barrick, J. E., M. R. Kauth, C. C. Strelioff, and R. E. Lenski. 2010. *Escherichia coli* rpoB mutants have increased evolvability in proportion to their fitness defects. Mol. Biol. Evol. 27:1338–1347.

Bouchonville, K., A. Forche, K. E. S. Tang, A. Selmecki, and J. Berman. 2009. Aneuploid chromosomes are highly unstable during DNA transformation of *Candida albicans*. Eukaryot. Cell 8:1554–1566. Am Soc Microbiol.

Chandler, C. H., S. Chari, and I. Dworkin. 2013. Does your gene need a background check? How genetic background impacts the analysis of mutations, genes, and evolution. Trends Genet. 29:358–366.

Chou, H.-H., H.-C. Chiu, N. F. Delaney, D. Segrè, and C. J. Marx. 2011. Diminishing returns epistasis among beneficial mutations decelerates adaptation. Science 332:1190–1192.

Chow, L., L. Waldron, and M. R. Gillings. 2015. Potential impacts of aquatic pollutants: sub-clinical antibiotic concentrations induce genome changes and promote antibiotic resistance. Front. Microbiol. 6:803.

Cooper, D. N., M. Krawczak, C. Polychronakos, C. Tyler-Smith, and H. Kehrer-Sawatzki. 2013. Where genotype is not predictive of phenotype: towards an understanding of the molecular basis of reduced penetrance in human inherited disease. Hum. Genet. 132:1077–1130.

Couce, A., and O. A. Tenaillon. 2015. The rule of declining adaptability in microbial

evolution experiments. Front. Genet. 6:99.

Cowen, L. E., D Sanglard, D. Calabrese, C. Sirjusingh, J. Anderson, and L. M. Kohn. 2000. Evolution of drug resistance in experimental populations of *Candida albicans*. J. Bacteriol. 182:1515–1522.

Cowen, L. E., L. M. Kohn, and J. B. Anderson. 2001. Divergence in fitness and evolution of drug resistance in experimental populations of *Candida albicans*. J. Bacteriol.

Delarze, E., and D. Sanglard. 2015. Defining the frontiers between antifungal resistance, tolerance and the concept of persistence. Drug Resist. Updat. 23:12–19.

Dowell, R. D., O. Ryan, A. Jansen, D. Cheung, S. Agarwala, T. Danford, D. A. Bernstein, P. A. Rolfe, L. E. Heisler, and B. Chin. 2010. Genotype to phenotype: a complex problem. Science 328:469–469.

Forche, A., G. Cromie, A. C. Gerstein, N. V. Solis, T. Pisithkul, W. Srifa, E. Jeffery, D. Abbey, S. G. Filler, A. M. Dudley, and J. Berman. 2018. Rapid phenotypic and genotypic diversification after exposure to the oral host niche in *Candida albicans*. Genetics, doi: 10.1534/genetics.118.301019.

Forche, A., P. T. Magee, A. Selmecki, J. Berman, and G. May. 2009. Evolution in *Candida albicans* populations during a single passage through a mouse host. Genetics 182:799–811. Genetics Soc America.

Ford, C. B., J. M. Funt, D. Abbey, L. Issi, C. Guiducci, D. A. Martinez, T. Delorey, B. yu Li, T. C. White, C. Cuomo, R. P. Rao, J. Berman, D. A. Thompson, and A. Regev. 2015. The evolution of drug resistance in clinical isolates of *Candida albicans*. Elife 4:e00662.

Fothergill, A. W., D. A. Sutton, D. I. McCarthy, and N. P. Wiederhold. 2014. The impact of new antifungal breakpoints on antifungal resistance in Candida species. J. Clin. Microbiol., doi: 10.1128/JCM.03044-13.

Fournier, T., and J. Schacherer. 2017. Genetic backgrounds and hidden trait complexity in natural populations. Curr. Opin. Genet. Dev. 47:48–53.

Fridman, O., A. Goldberg, I. Ronin, N. Shoresh, and N. Q. Balaban. 2014. Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. Nature 418–421. Nature Publishing Group.

Gerstein, A. C., M. S. Fu, L. Mukaremera, Z. Li, K. L. Ormerod, J. A. Fraser, J. Berman, and K. Nielsen. 2015. Polyploid titan cells produce haploid and aneuploid progeny to promote stress adaptation. MBio 6:e01340–15.

Gerstein, A. C., D. S. Lo, and S. P. Otto. 2012. Parallel genetic changes and nonparallel gene–environment interactions characterize the evolution of drug resistance in yeast. Genetics 192:241–252. Genetics.

Gerstein, A. C., A. Rosenberg, I. Hecht, and J. Berman. 2016. diskImageR:

quantification of resistance and tolerance to antimicrobial drugs using disk diffusion assays. Microbiology 162:1059–1068.

-G. Moore, F. B., and R. Woods. 2006. Tempo and constraint of adaptive evolution in Escherichia coli (Enterobacteriaceae, Enterobacteriales). Biol. J. Linn. Soc. Lond. 88:403–411.

Gullberg, E., S. Cao, O. G. Berg, C. Ilbäck, L. Sandegren, D. Hughes, and D. I. Andersson. 2011. Selection of resistant bacteria at very low antibiotic concentrations. PLoS Pathog. 7:e1002158.

Harrison, B. D., J. Hashemi, M. Bibi, R. Pulver, D. Bavli, Y. Nahmias, M. Wellington, G. Sapiro, and J. Berman. 2014. A tetraploid intermediate precedes aneuploid formation in yeasts exposed to fluconazole. PLoS Biol. 12:e1001815.

Henderson-Begg, S. K., D. M. Livermore, and L. M. C. Hall. 2006. Effect of subinhibitory concentrations of antibiotics on mutation frequency in *Streptococcus pneumoniae*. J. Antimicrob. Chemother. 57:849–854.

Hickman, M. A., C. Paulson, A. M. Dudley, and J. Berman. 2015. Parasexual ploidy reduction drives population heterogeneity through random and transient aneuploidy in *Candida albicans*. Genetics. Genetics Soc America.

Hirakawa, M. P., D. E. Chyou, D. Huang, A. R. Slan, and R. J. Bennett. 2017. Parasex generates phenotypic diversity de novo and impacts drug resistance and virulence in *Candida albicans*. Genetics 207:1195–1211.

Hou, J., T. Fournier, and J. Schacherer. 2016. Species-wide survey reveals the various flavors of intraspecific reproductive isolation in yeast. FEMS Yeast Res. 16.

Khan, A. I., D. M. Dinh, D. Schneider, R. E. Lenski, and T. F. Cooper. 2011. Negative epistasis between beneficial mutations in an evolving bacterial population. Science 332:1193–1196.

Kryazhimskiy, S., D. P. Rice, E. R. Jerison, and M. M. Desai. 2014. Microbial evolution. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science 344:1519–1522.

Levin, B. R., and C. T. Bergstrom. 2000. Bacteria are different: observations, interpretations, speculations, and opinions about the mechanisms of adaptive evolution in prokaryotes. Proc. Natl. Acad. Sci. U. S. A. 97:6981–6985.

Liu, A., A. Fong, E. Becket, J. Yuan, C. Tamae, L. Medrano, M. Maiz, C. Wahba, C. Lee, K. Lee, K. P. Tran, H. Yang, R. M. Hoffman, A. Salih, and J. H. Miller. 2011. Selective advantage of resistant strains at trace levels of antibiotics: a simple and ultrasensitive color test for detection of antibiotics and genotoxic agents. Antimicrob. Agents Chemother. 55:1204–1210.

Lundström, S. V., M. Östman, J. Bengtsson-Palme, C. Rutgersson, M. Thoudal, T. Sircar, H. Blanck, K. M. Eriksson, M. Tysklind, C.-F. Flach, and D. G. J.

Larsson. 2016. Minimal selective concentrations of tetracycline in complex aquatic bacterial biofilms. Sci. Total Environ. 553:587–595.

MacLean, R. C., G. G. Perron, and A. Gardner. 2010. Diminishing returns from beneficial mutations and pervasive epistasis shape the fitness landscape for rifampicin resistance in Pseudomonas aeruginosa. Genetics 186:1345–1354.

McVicker, G., T. K. Prajsnar, A. Williams, N. L. Wagner, M. Boots, S. A. Renshaw, and S. J. Foster. 2014. Clonal expansion during Staphylococcus aureus infection dynamics reveals the effect of antibiotic intervention. PLoS Pathog. 10:e1003959.

Melnyk, A. H., A. Wong, and R. Kassen. 2015. The fitness costs of antibiotic resistance mutations. Evol. Appl. 8:273–283.

Mira, P. M., J. C. Meza, A. Nandipati, and M. Barlow. 2015. Adaptive landscapes of resistance genes change as antibiotic concentrations change. Mol. Biol. Evol. 32:2707–2715.

Moore, F. B., D. E. Rozen, and R. E. Lenski. 2000. Pervasive compensatory adaptation in Escherichia coli. Proc. Biol. Sci. 267:515–522.

Ngamskulrungroj, P., Y. Chang, B. Hansen, C. Bugge, E. Fischer, and K. J. Kwon-Chung. 2012. Characterization of the chromosome 4 genes that affect fluconazole-induced disomy formation in *Cryptococcus neoformans*. PLoS One 7:e33022.

Odds, F. C., M. E. Bougnoux, D. J. Shaw, J. M. Bain, A. D. Davidson, D. Diogo, M. D. Jacobsen, M. Lecomte, S. Y. Li, A. Tavanti, M. C. J. Maiden, N. A. R. Gow, and C. d'Enfert. 2007. Molecular phylogenetics of *Candida albicans*. Eukaryot. Cell 6:1041–1052. Am Soc Microbiol.

Ono, J., A. C. Gerstein, and S. P. Otto. 2017. Widespread genetic incompatibilities between first-step mutations during parallel adaptation of *Saccharomyces cerevisiae* to a common environment. PLoS Biol. 15:e1002591.

Paaby, A. B., A. G. White, D. D. Riccardi, and K. C. Gunsalus. 2015. Wild worm embryogenesis harbors ubiquitous polygenic modifier variation. Elife.

Peng, F., S. Widmann, A. Wünsche, K. Duan, K. A. Donovan, R. C. J. Dobson, R. E. Lenski, and T. F. Cooper. 2018. Effects of beneficial mutations in pykf gene vary over time and across replicate populations in a long-term experiment with bacteria. Mol. Biol. Evol. 35:202–210.

Perepnikhatka, V., F. J. Fischer, M. Niimi, R. A. Baker, R. D. Cannon, Y. K. Wang, F. Sherman, and E. Rustchenko. 1999. Specific chromosome alterations in fluconazole-resistant mutants of *Candida albicans*. J. Bacteriol. 181:4041–4049.

Perfeito, L., A. Sousa, T. Bataillon, and I. Gordo. 2014. Rates of fitness decline and rebound suggest pervasive epistasis. Evolution 68:150–162.

R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rokyta, D. R., Z. Abdo, and H. A. Wichman. 2009. The genetics of adaptation for eight microvirid bacteriophages. J. Mol. Evol. 69:229–239.

Ropars, J., C. Maufrais, D. Diogo, M. Marcet-Houben, A. Perin, N. Sertour, K. Mosca, E. Permal, G. Laval, C. Bouchier, L. Ma, K. Schwartz, K. Voelz, R. C. May, J. Poulain, C. Battail, P. Wincker, A. M. Borman, A. Chowdhary, S. Fan, S. H. Kim, P. Le Pape, O. Romeo, J. H. Shin, T. Gabaldon, G. Sherlock, M.-E. Bougnoux, and C. d'Enfert. 2018. Gene flow contributes to diversification of the major fungal pathogen *Candida albicans*. Nat. Commun. 9:2253.

Rosenberg, A., I. V. Ene, M. Bibi, S. Zakin, E. S. Segal, N. Ziv, A. M. Dahan, A. L. Colombo, R. J. Bennett, and J. Berman. 2018. Antifungal tolerance is a subpopulation effect distinct from resistance and is associated with persistent candidemia. Nat. Commun. 9:2470.

Sanjuán, R., J. M. Cuevas, A. Moya, and S. F. Elena. 2005. Epistasis and the adaptability of an RNA virus. Genetics 170:1001–1008.

Selmecki, A., A. Forche, and J. Berman. 2006. Aneuploidy and isochromosome formation in drug-resistant *Candida albicans*. Science 313:367–370.

Selmecki, A., M. Gerami-Nejad, C. Paulson, A. Forche, and J. Berman. 2008. An isochromosome confers drug resistance in vivo by amplification of two genes, ERG11 and TAC1. Mol. Microbiol. 68:624–641.

Selmecki, A. M., K. Dulmage, L. E. Cowen, J. B. Anderson, and J. Berman. 2009. Acquisition of aneuploidy provides increased fitness during the evolution of antifungal drug resistance. PLoS Genet. 5:e1000705.

Sionov, E., H. Lee, Y. C. Chang, and K. J. Kwon-Chung. 2010. *Cryptococcus neoformans* Overcomes stress of azole drugs by formation of disomy in specific multiple chromosomes. PLoS Pathog. 6:e1000848. Public Library of Science.

Tavanti, A., A. D. Davidson, M. J. Fordyce, N. A. R. Gow, M. C. J. Maiden, and F. C. Odds. 2005. Population structure and properties of *Candida albicans*, as determined by multilocus sequence typing. J. Clin. Microbiol. 43:5601–5613. Am Soc Microbiol.

Tenaillon, O. 2014. The utility of fisher's geometric model in evolutionary genetics. Annu. Rev. Ecol. Evol. Syst. 45:179–201.

Torres-Barceló, C., M. Kojadinovic, R. Moxon, and R. C. MacLean. 2015. The SOS response increases bacterial fitness, but not evolvability, under a sublethal dose of antibiotic. Proc. Biol. Sci. 282:20150885.

Vu, V., A. J. Verster, M. Schertzberg, T. Chuluunbaatar, M. Spensley, D. Pajkic, G. T. Hart, J. Moffat, and A. G. Fraser. 2015. Natural variation in gene expression modulates the severity of mutant phenotypes. Cell 162:391–402. Elsevier Inc.

Wang, Y., C. D. Arenas, D. M. Stoebel, and T. F. Cooper. 2013. Genetic background affects epistatic interactions between two beneficial mutations. Biol. Lett. 9:20120328.

Wang, Y., C. Diaz Arenas, D. M. Stoebel, K. Flynn, E. Knapp, M. M. Dillon, A. Wünsche, P. J. Hatcher, F. B.-G. Moore, V. S. Cooper, and T. F. Cooper. 2016. Benefit of transferred mutations is better predicted by the fitness of recipients than by their ecological or genetic relatedness. Proc. Natl. Acad. Sci. U. S. A. 113:5047–5052.

Wellington, M., and E. Rustchenko. 2005. 5-Fluoro-orotic acid induces chromosome alterations in *Candida albicans*. Yeast 22:57–70.

Wistrand-Yuen, E., M. Knopp, K. Hjort, S. Koskiniemi, O. G. Berg, and D. I. Andersson. 2018. Evolution of high-level resistance during low-level antibiotic exposure. Nat. Commun. 9:1599.

Wong, A. 2017. Epistasis and the evolution of antimicrobial resistance. Front. Microbiol. 8:246.

Wünsche, A., D. M. Dinh, R. S. Satterwhite, C. D. Arenas, D. M. Stoebel, and T. F. Cooper. 2017. Diminishing-returns epistasis decreases adaptability along an evolutionary trajectory. Nat Ecol Evol 1:61.

Wu, Y. L., E. M. Scott, A. L. Po, and V. N. Tariq. 1999. Development of resistance and cross-resistance in *Pseudomonas aeruginosa* exposed to subinhibitory antibiotic concentrations. APMIS 107:585–592.

Yang, F., A. Kravets, G. Bethlendy, S. Welle, and E. Rustchenko. 2013. Chromosome 5 monosomy of candida albicans controls susceptibility to various toxic agents, including major antifungals. Antimicrob. Agents Chemother. 57:5026–5036.

Zhu, Y. O., G. Sherlock, and D. A. Petrov. 2016. Whole genome analysis of 132 clinical *Saccharomyces cerevisiae* strains reveals extensive ploidy variation. G3 6:2421–2434.

Zhu, Y. O., M. L. Siegal, D. W. Hall, and D. A. Petrov. 2014. Precise estimates of mutation rate and spectrum in yeast. Proc. Natl. Acad. Sci. U. S. A. 111:E2310–E2318.

Supporting Tables and Figures

Table S1. The majority of strains evolved an increased growth ability in the evolutionary environment. The test column indicates the test that was run: a t-test when both ancestral and evolved replicate groups were normally-distributed with equal variance, a t-test with welch approximation for degrees of freedom when variances were unequal, or the Wilcoxon Rank Sum test when the data from at least one group was not normally distributed. Equal variance was assessed with an F test, normality with the Shapiro-Wilk test (assumption test results not shown).

Line	Growth ability at 24 h					Growth ability at 72 h				
	test	t-value	df	p	evol-anc	test	t-value	df	p	evol-anc
A1	t-test	-6.85	22	<0.0001	0.09	t-test	-12.49	22	<0.0001	0.07
A2	wilcox	89	-	0.35	-0.02	wilcox	0	-	<0.0001	0.45
A3	wilcox	0	-	<0.0001	0.42	t-test	-12.96	22	<0.0001	0.64
A4	welch	-9.02	11.47	<0.0001	0.7	wilcox	0	-	<0.0001	0.84
A5	welch	-5.58	13.31	0.0001	0.21	t-test	-8.49	22	<0.0001	0.05
A6	t-test	-4.29	22	0.0003	0.27	wilcox	0	-	<0.0001	0.34
A7	t-test	-4.17	22	0.0004	0.31	t-test	-7.95	22	<0.0001	0.26
A8	welch	-6.67	11.79	<0.0001	0.67	welch	-19.22	15.01	<0.0001	0.57
A9	wilcox	96	-	0.18	0.03	welch	-11.17	12.63	<0.0001	0.69
A10	welch	1.03	13.14	0.32	-0.08	wilcox	7	-	<0.0001	0.14
A11	welch	-3.58	11.5	0.0041	0.29	welch	-8.09	12.45	<0.0001	0.74
A12	t-test	7.26	22	<0.0001	-0.18	wilcox	36	-	0.0387	0.01
A13	welch	-7.45	11.16	<0.0001	0.74	welch	-10.56	15.23	<0.0001	0.4
A14	wilcox	94	-	0.22	0.06	wilcox	21	-	0.0023	0.36
A15	welch	-7.35	12.41	<0.0001	0.72	t-test	-9.91	22	<0.0001	0.27
A16	welch	-8	14.74	<0.0001	0.58	welch	-21.39	12.27	<0.0001	0.36
A17	welch	-4.77	11.07	0.0006	0.64	welch	-18.68	14.55	<0.0001	0.99
A18	t-test	-4.92	22	0.0001	0.05	t-test	-6.34	22	<0.0001	0.03
A19	welch	-3.9	11.83	0.0022	0.42	t-test	-12.61	22	<0.0001	0.41
A20	t-test	-9.59	22	<0.0001	0.41	t-test	-11.5	22	<0.0001	0.08

Table S2. Replicates from the majority of strain backgrounds evolved a reduced growth ability at 24 hours in a drug-free environment in the evolutionary environments. The test column indicates the test that was run: a t-test when both ancestral and evolved replicate groups were normally-distributed with equal variance, a t-test with welch approximation for degrees of freedom when variances were unequal, or the Wilcoxon Rank Sum test when the data from at least one group was not normally distributed. Equal variance was assessed with an F test, normality with the Shapiro-Wilk test (assumption test results not shown).

Line	Growth ability at 24 h					Growth ability at 72 h				
	test	t-value	df	p	evol-anc	test	t-value	df	p	evol-anc
A1	welch	2.38	14.79	0.0311	-0.08	t-test	-3.61	22	0.0016	0.02
A2	welch	7.54	12.91	<0.0001	-0.55	t-test	0.93	22	0.36	-0.01
A3	welch	4.75	12.75	0.0004	-0.31	t-test	2.98	22	0.0069	-0.02
A4	welch	2.56	12.68	0.0241	-0.21	t-test	-3.63	22	0.0015	0.03
A5	welch	-2.78	16.69	0.0129	0.09	t-test	-3.59	22	0.0016	0.02
A6	welch	4.06	12.35	0.0015	-0.21	welch	1.11	15.03	0.28	-0.01
A7	wilcox	134	-	0.0001	-0.3	wilcox	76.5	-	0.82	-0.01
A8	wilcox	104	-	0.068	-0.25	t-test	-0.85	22	0.4	0.01
A9	wilcox	142	-	<0.0001	-0.46	wilcox	77.5	-	0.77	-0.03
A10	wilcox	99	-	0.13	-0.22	t-test	-2.47	22	0.0219	0.01
A11	welch	0.35	12.66	0.73	-0.02	t-test	-3.03	22	0.0062	0.02
A12	t-test	2.95	22	0.0074	-0.07	wilcox	33	-	0.0242	0.01
A13	welch	6.6	11.07	<0.0001	-0.62	welch	6.61	11.63	<0.0001	-0.13
A14	welch	5.59	11.55	0.0001	-0.53	welch	3.12	14.1	0.0075	-0.03
A15	welch	4.73	11.69	0.0005	-0.2	welch	9.55	11.41	<0.0001	-0.15
A16	welch	2.98	13.84	0.0101	-0.15	t-test	2.05	22	0.053	-0.01
A17	welch	6.19	11.03	0.0001	-0.66	welch	3.86	11.27	0.0025	-0.13
A18	welch	4.11	12.17	0.0014	-0.07	t-test	-0.13	22	0.89	0
A19	welch	10.51	12.86	<0.0001	-0.47	wilcox	58	-	0.44	-0.06
A20	t-test	4.59	22	0.0001	-0.13	t-test	-2.33	22	0.0291	0.01

Table S3. Minimum inhibitory concentration experiments

Exper	Time	Fluconazole level (µg)							
		0	0.5	1	4	8	32	128	512
1	Anc&Evol	x		x		x	x		
2	Anc	x		x					
3	Anc&Evol	x	x	x	x				
4	Anc	x		x	x	x	x		
5	Anc&Evol	x				x	x	x	
6	Anc&Evol	x					x	x	x

Figure S1. Evolved growth in a drug-free environment correlates with evolved growth in the low drug environment at 24 h. Each point represents the mean of 12 replicates evolved from each strain background.

Figure S2. Evolved growth in a drug-free environment (YPD) is correlated with evolved growth in the evolutionary (low drug) environment in many strain backgrounds. Growth in both environments was measured as optical density at 24 h. Each point represents an independently evolved replicate from the same strain background (indicated by the label in the top left corner of each figure panel). The solid red lines indicate a significant Spearman correlation test, dotted lines are for insignificant correlations. The replicate lines from the four strains plotted on the bottom (A1, A5, A12 and A18) were not variable enough to test for a correlation.

Figure S3. Tolerance of evolved strains measured at a) 24 h, b) 48 h, and c) 72 h.
Tolerance was measured as the growth observed in supra-MIC levels of fluconazole (as appropriate to each replicate) normalized to the growth in the absence of drug. Strains are arranged on the x-axis (and coloured) by initial growth in the evolutionary environment. Each point represents an individually-evolved replicate line. The black line indicates the median ancestral level and the two grey lines indicate the upper and lower bounds of twelve measured ancestral replicates.

Figure S4. Evolved growth at 72 h and evolved genome size are not consistently correlated. Growth was measured as optical density after 72 hours in low drug. Evolved genome size indicates the most prominent G1 peak; when multiple G1 peaks were present this is indicated with a triangle symbol. Solid lines indicate a significant Spearman correlation test. Each point represents an independently evolved replicate line.

Figure S5. Evolved tolerance at 24 h and evolved genome size are only correlated in a single strain background. Each point represents an independently evolved replicate line. Tolerance was measured as SMG24. Evolved genome size represents the most prominent G1 peak; when multiple G1 peaks were present this is indicated with a triangle symbol. A solid line indicates a significant Spearman correlation test.

Figure S6. Evolved tolerance at 72 h and evolved genome size are only correlated in a single strain background. Each point represents an independently evolved replicate line. Tolerance was measured as SMG72. Evolved genome size represents the most prominent G1 peak; when multiple G1 peaks were present this is indicated with a triangle symbol. A solid line indicates a significant Spearman correlation test.