3   Systems biology reveals NR2F6 and TGFB1 as key regulators of feed efficiency in beef

4   cattle

6   Pâmela A. Alexandre[1,2]; Marina Naval-Sanchez[2]; Laercio R. Porto-Neto[2]; José Bento S.

7   Ferraz[1]; Antonio Reverter[2] and Heidge Fukumasu[1§]

[1]Department of Veterinary Medicine, College of Animal Sciences and Food Engineering,

University of São Paulo, Av. Duque de Caxias Norte 225, Pirassununga, 13635-900, Brazil

[2]Commonwealth Scientific and Industrial Research Organisation (CSIRO), Agriculture &

Food, 306 Carmody Road, St. Lucia, 4067, QLD, Australia

[§]corresponding author

Heidge Fukumasu, DVM, PhD, Associate Professor

Phone: 55-19-35656864

Fax: 55-19-3565

e-mail: fukumasu@usp.br

26   **Abstract**

Systems biology approaches are used as strategy to uncover tissue-specific perturbations and

regulatory genes related to complex phenotypes. We applied this approach to study feed

efficiency (FE) in beef cattle, an important trait both economically and environmentally.

Poly-A selected RNA of five tissues (adrenal gland, hypothalamus, liver, skeletal muscle and

pituitary) of eighteen young bulls, selected for high and low FE, were sequenced (100bp,

pared-end). From the 17,354 expressed genes, 1,317 were prioritized by five selection

categories (differentially expressed, harbouring SNPs associated with FE, tissue-specific,

secreted in plasma and key regulators) and used for network construction. NR2F6 and TGFB

were identified and validated by motif discovery as key regulators of hepatic inflammatory

response and muscle tissue development, respectively, two biological processes demonstrated

to be associated to FE. Moreover, we indicated potential biomarkers of FE which are related

to hormonal control of metabolism and sexual maturity. By using robust methodologies and

validation strategies, we confirmed main biological processes related to FE in *Bos indicus*

and indicated candidate genes as regulators or biomarkers of superior animals.

Keywords: *Bos indicus*/co-expression/motif discovery/inflammation/residual feed intake

Introduction

Since the domestication of the first species, animal selection aims to meet human needs and their changes over time. The current main selection goals in livestock production are increase of productivity, reduction of the environmental impact and reduction of competition for grains with human nutrition (Hayes *et al*, 2013). Thus, feed efficiency (FE) has become a relevant trait of study, as animals considered of high feed efficiency are those presenting reduced feed intake and lower production of methane and manure without compromising animal's weight gain (Gerber *et al*, 2013). However, the incorporation of FE as selection criteria in animal breeding programs is costly and time consuming. Daily feed intake and weight gain for a large number of animals need to be recorded for at least 70 days to obtain accurate estimates of FE (Archer *et al*, 1997).

In the past years, several studies have been carried out with the aim to identify molecular markers associated with FE to enable the faster and cost-effectively identification of superior animals (de Oliveira *et al*, 2014; Rolf *et al*, 2011; Santana *et al*, 2014; Seabury *et al*, 2017). However, for each population, different biological processes seem to be identified (de Oliveira *et al*, 2014; Rolf *et al*, 2011; Santana *et al*, 2014; Seabury *et al*, 2017). Probably, that is because FE is a multifactorial trait and many different biological mechanisms seems to be involved in its regulation (Herd *et al*, 2004; Herd & Arthur, 2009). It has been indicated that high FE animals present increased mitochondrial function (Lancaster *et al*, 2014; Connor *et al*, 2010), less oxygen consumption (Gonano *et al*, 2014) and delayed puberty (Randel & Welsh, 2013; Shaffer *et al*, 2011; Fontoura *et al*, 2016). On the other hand, low FE animals have increased physical activity, ingestion frequency and stress (Francisco *et al*, 2015; Cafe *et al*, 2011; Kelly *et al*, 2010; Chen *et al*, 2014), increased leptin and cholesterol levels (Alexandre *et al*, 2015; Foote *et al*, 2016; Nkrumah *et al*, 2007; Mota *et al*, 2017), higher subcutaneous and visceral fat (Santana *et al*, 2012; Gomes *et al*, 2012; Mader *et al*, 2009),

higher energy wastage as heat (Montanholi *et al*, 2010, 2009; Archer *et al*, 1999) and more

hepatic lesions associated with inflammatory response (Alexandre *et al*, 2015; Paradis *et al*,

2015).

In the context of such a complex trait, we perform a multiple-tissue transcriptomic

analyses of high and low FE Nellore cattle across tissues related to endocrine control of

hanger/satiety, hydric and energy homeostasis, stress and immune response, physical and

sexual activity, as is the case of hypothalamus-pituitary-adrenal axis and organs as liver and

skeletal muscle. Based on gene co-expression across tissues and conditions we derived a

regulatory network revealing NR2F6 and TGFB signalling as key regulators of hepatic

inflammatory response and muscle tissue development, respectively. Next, we apply

advanced motif discovery methods which i) validate that co-expressed genes are enriched for

NR2F6 and TGFB signalling effector molecule SMAD3 binding sites in their 10KB upstream

regions and ii) predict direct transcription factor (TF) – Target gene (TG) interactions at the

sequence level. These binding interactions were experimentally validated with public TF

ChIP-seq from ENCODE. Regulatory activity in the tissues of interest was also confirmed by

performing an enrichment analysis on open chromatin tracks and histone chromatin marks

across cell types and tissues in the human and cow genome. Moreover, we propose a

hormonal control of differences in metabolism and sexual maturity between high and low FE

animals, indicating potential biomarkers for further validation such as adrenomedullin, FSH,

oxytocin, somatostatin and TSH.

Results

Multi-tissue transcriptomic data reveal differences between high and low feed efficient

animals

Feed efficiency is a complex trait characterized by multiple distinct biological

processes including metabolism, ingestion, digestion, physical activity and thermoregulation

(Herd *et al*, 2004; Herd & Arthur, 2009). To study FE at transcriptional level we performed

RNAseq of five tissues (i.e. adrenal gland, hypothalamus, liver, muscle and pituitary) from

nine male bovines of high feed efficiency (HFE, characterized by low residual feed intake

(RFI) (Koch *et al*, 1963)) and nine of low FE (LFE, characterized by high RFI). In total, we

analysed 18 samples of liver, hypothalamus and pituitary; 17 of muscle and 15 of adrenal

gland, yielding 13 million reads per sample on average (S1 Supporting Information). Gene

expression was estimated for 24,616 genes present in the reference genome (UMD 3.1) and

after quality control (refer to methods), 17,354 genes were identified as being expressed in at

least one of the five tissues analysed.

Differential expression (DE) analysis between high and low FE animals resulted in

471 DE genes across tissues (P<0.001, S2 Supporting Information), namely, 111 in adrenal

gland, 125 in hypothalamus, 91 in liver, 104 in muscle and 98 in pituitary (S3A-E Supporting

Information). Although no significant functional enrichment was found for the 281 genes up-

regulated in high feed efficiency, the 248 genes down-regulated presented a significant

enrichment of GO terms such as response to hormone (Padj=5.43 x 10^{-6}), regulation of

hormone levels (Padj=3.48 x 10^{-6}), cell communication (Padj=3.18 x 10^{-4}), regulation of

signaling receptor activity (Padj=3.20 x 10^{-4}), hormone metabolic process (Padj=5.86 x 10^{-4}),

response to corticosteroid (Padj=6.28 x 10^{-4}), regulation of secretion (Padj=7.2 x 10^{-4}),

response to lipopolysaccharide (Padj=7.9 x 10^{-4}) and regulation of cell proliferation

(Padj=1.86 x 10^{-3}). Refer to S4 Supporting Information to see all enriched terms.

Overlap between gene selection criteria prioritizes genes associated with feed efficiency

The genetic architecture behind complex traits involves a large variety of genes with

coordinated expression pattern, which can be represented by gene regulatory networks as a

blueprint to study their relationships and to identify central regulatory genes (Swami, 2009).

Therefore, it is important to select relevant genes and gene families according to the

phenotype of interest to be used for network analysis. We defined five categories of genes

(see methods for further information) for inclusion in co-expression analysis: 1 -

differentially expressed (DE), 2 - genes harbouring SNPs previously associated with FE

(harbouring SNP), 3 - tissue specific (TS), 4 - genes coding proteins secreted in plasma by

any of the five tissues analysed (secreted) and 5 - key regulators.

As reported before, we have identified 471 DE genes between high and low FE

animals (Figure 1A, S5A Supporting Information). In addition, 267 genes were selected for

harbouring SNPs previously associated with FE, as not only differences in expression levels

can influence the phenotype but also polymorphism in the DNA sequence that can alter the

translated protein behaviour (S5B Supporting Information). Moreover, 396 were selected for

being tissue specific (refer to methods for definition); 22 in adrenal gland, 32 in

hypothalamus, 215 in liver, 218 in muscle and 9 in pituitary (S5C Supporting Information). A

total of 244 genes coding proteins secreted in plasma were selected because of their potential

as biomarkers of FE (S5D Supporting Information). From those, 135 had liver as the tissue of

maximum expression and were functionally enriched for GO terms such as complement

activation (Padj=1.82×10^{-19}), regulation of acute inflammatory response (Padj=1.89×10^{-14}),

innate immune response (Padj=9.71×10^{-12}), negative regulation of endopeptidase activity

(Padj=2.35×10^{-10}), platelet degranulation (Padj=1.08×10^{-10}), regulation of coagulation

(Padj=3.39×10^{-9}), triglyceride homeostasis (Padj=1.23×10^{-6}), cholesterol efflux

(Padj=$1.03E^{-5}$) (S6 Supporting Information). Finally, from 1570 potential regulators in public

available TFdb, 78 were identified as key regulators of the genes selected by all the other

categories, i.e. 78 genes presented a coordinated expression level with many of the genes in

the network reflecting a tight control of expression pattern across tissues (S5E Supporting

Information).

Considering all the inclusion criteria, 1,317 genes were selected to be included in co-

expression network analysis (Figure 1B, S7 Supporting Information), some of them selected

by more than one category (Figure 1C). Regarding DE genes, six of them were also reported

before as harbouring SNPs associated with the phenotype (*LUZP2, MAOB, SFRS5,*

SLC24A2, SOCS3 and *WIF1*) and 13 of them were key regulators (*HOPX, PITX1, CRYM,*

PLCD1, ND6, cytb, ND1, MT-ND4L, ND5, ATP8, ND4, ENSBTAG00000046711 and

ENSBTAG00000048135). Many of the genes that are both DE and regulators are involved in

respiratory chain (*ND6, cytb, ND1, MT-ND4L, ND5, ATP8* and *ND4*) and were all up-

regulated in high FE group.

insert Figure 1

Considering both DE and secreted genes, 18 were identified (*NOV, SPP1, CTGF,*

OXT, PTX3, VGF, CCL21, COL1A2, PGF, SOD3, SERPINE1, PRL, PON1, SST, JCHAIN,

PCOLCE, IGFBP6 and *SCG2*). In addition, four genes were DE, secreted and tissue specific,

two from liver (*CXCL3* and *IGFBP1*) and two from pituitary (*NPY* and *CYP17A1*). Genes

RARRES2 and *PENK* (proenkephalin) were DE, secreted and had been previously reported as

harbouring SNP associated with FE [30, AnimalQTLdb]. Other DE genes worthy to

highlight, due to their well-known role in metabolic processes, are *AMH* (anti-mullerian

hormone), *TSHB* (thyroid stimulating hormone beta), *FGF21* (Fibroblast growth factor 21)

and *FST* (follistatin), up-regulated in high FE group, and *PMCH* (pro-melanin concentrating

hormone), *ADM* (adrenomedullin) and *FSHB* (follicle stimulating hormone beta), up-

regulated in low FE group.

Co-expression network reveals regulatory genes and biological processes related to feed

efficiency

The co-expression network (Figure 2) was composed by 1,317 genes and 91,932

connections, with a mean of 70 connections per gene. Most of the connections (51%)

involved a DE gene and 23% of those were between two DE genes. Tissue specific (TS) gene

were involved in 49% of the connections with 119 connections per gene in average, which

was higher than the overall network mean and reflects the close relationship between genes

involved in tissue specific functions. Key regulators was the least represented category in the

network (only 78 genes) but accounted for 11% of the connections in the network with the

highest value of mean connections per gene, 131 connections, which is in accordance with

their regulatory role. Regarding the connections within tissues, when we ranked all the genes

in the network by the number of connections and looked at the top 50 genes, 29 were from

liver, 15 were from muscle and 3, 2 and 1 were from pituitary, adrenal gland and

hypothalamus, respectively. This result indicates a very well-coordinated expression pattern

in liver and muscle that could be a reflex of the number of TS genes in those tissues and the

presence of central regulatory genes coordinating the expression of many other genes.

In the network (Figure 2), genes grouped together by tissue which was mostly driven

by TS genes. As mentioned before, most of the secreted proteins coding genes were locate in

the liver. Most of the key regulators were located peripherally in relation to the clusters which

could be reflecting their regulatory nature independent of tissue specificity. Despite that,

some regulators draw attention because of their high number of connections.

insert Figure 2

The top five most connected regulators were *EPC1, NR2F6, MED21,*
ENSBTAG00000031687 and *CTBP1*, varying from 317 to 284 connections. They were all
first neighbours of each other and were connected mainly to genes with higher expression in
liver and essentially enriched for acute inflammatory response (Padj=4.5 x 10^{-13}, S8
Supporting Information). The next most connected regulator is *TGFB1* with 217 connections.
It is mainly connected to genes from muscle that are primarily enriched for muscle organ
development (Padj=6.87 x 10^{-5}) and striated muscle contraction (Padj=1.39 x 10^{-5}, S9
Supporting Information). Besides indicating main regulator genes, network approach can be
useful to access the role of specific genes. For instance, gene *FGF21*, a hormone up regulated
in liver of high FE animals, is directly connected to genes enriched for plasma lipoprotein
particle remodelling, regulation of lipoprotein oxidation and cholesterol efflux (Padj=5.64E-
3, S10 Supporting Information). Indeed, according to the literature, this gene is associated to
decrease in body weight, blood triglycerides and LDL-cholesterol (Cheung & Deng, 2014).

Motif discovery confirms NR2F6 as a key regulator of liver transcriptional changes
between high and low feed efficiency

By means of the power-law theory, co-expression networks present many nodules
with few connections and few central nodules with many connections (de la Fuente, 2010),
being the last ones indicated as central regulatory genes responsible for the transcriptional
changes between the divergent phenotypes analysed. In our study, the most connected
regulators were indicated, together with their target genes, i.e. their first neighbours in the
network. Those genes are a mixture of direct and indirect regulator targets. In order to
validate the regulatory role of the most connected regulators in the network and identify their

core direct targets we performed motif discovery in their co-expressed target genes. It is noteworthy motif discovery should confirm the presence of DNA motifs of a TF in the regulatory regions of co-expressed genes. From the top five most connected regulators from our previous co-expression analysis, only NR2F6 has the ability to bind DNA. In contrast, the other four regulators act mainly as cofactors (corepressor, i.e. CTBP1; coactivator, i.e. MED21; or histones modifier, i.e. EPC1), that is co-binding through protein-protein interactions.

The analysis of 313 co-expressed genes with NR2F6 yield the Nuclear Factor motif HNF4-NR2F2 (transfac_pro-M01031) as the second motif most enriched out of 9732 PWMs with a Normalized Enrichment Score (NES) of 7.99 (Figure 3B). In addition, a total of 19 motifs associated with HNF4-NR2F2 were enriched in the dataset, associating HNF4-NR2F2 to 168 direct target genes (Figure 3C). Due to motif redundancy or highly similarity between a plethora of TFs, these motifs can be associated with multiple TFs from HNF4 (direct) to several nuclear factors such as NR2F6 (motif similarity score FDR 1.414E-5). However, our co-expression analysis strongly indicates NR2F6 is the key TF, since it was the TF with the highest number of nodes in the co-expression network (Figure 3C) and neither HNF4 nor NR2F2 were prioritized by any selection category to be included in the network.

Each of the NR2F6 inferred direct target genes contain one or more predicted enhancers, i.e. regions with high-scoring motif binding sites for NR2F6 or TFs with highly similar motifs. To validate the binding of these genomic regions by NR2F6 or TFs with highly similar motifs to NR2F6 we performed a region enrichment analysis of our predicted NR2F6 binding sequences against public TF ChiP-seq bound regions in human cell lines from ENCODE the ENCODE consortium (1394 TF binding site tracks). This analysis, confirms the experimental binding of TFs with similar binding as NR2F6 in HepG2 cells, HNF4A (NES=8.57), HNF4G (NES=7.83), RXRA (NES=6.85), and NR2F2 (NES=4.45) as

the most enriched tracks (S11 Supporting Information). Recent NR2F6 ChIP-seq data in

HepG also confirms an enrichment for NR2F6 (Figure 3D), indicating predicted NR2F6

binding regions are experimentally bound by NR2F6 in hepatocyte cell lines (Figure 3D).

Next, to validate that the NR2F6 binding in those regions is functional in liver we

performed an enrichment analysis for open-chromatin (tracks=655) and histone modifications

(tracks =2450) related to active regulatory elements (S12 Supporting Information). This

analysis yielded FAIRE-seq on HepG2 cell lines and H3K9ac and H3K4me3 in adult liver

(E066 Roadmap Epigenomics Track) as the most enriched tracks respectively, strongly

indicating not only predicted target enhancers are bound by NR2F6 in Hepatocyte cell lines

but these regulatory regions are functionally active in hepatocytes and human liver (Figure

3D).

Regarding the cow genome, a recent open-chromatin study (Villar *et al*, 2015) has

delineated the map active promoters and enhancers by H3K4me3 and H3K27ac ChIP-seq in

cow liver resulting in 13796 promoter and 45786 enhancers (S13 Supporting Information).

We performed an enrichment analysis of predicted NR2F6 enhancers converted to cow

coordinates (n=779) resulting in 446 regions being identified as functional regulatory regions

in cow liver. This number is significantly higher compared to only 43 regions are expected to

overlap by random (1000 permutation tests) (Figure 4E).

Finally, in addition to NR2F6 motif, HNF1A motif was found as a potential co-

regulator in liver, in particular swissregulon-HNF1A.p2 with a NES =10.17 and in total 20

enriched motifs and 170 direct targets were associated to HNF1A (Figure 3B). HNF1 is a

master regulator of liver gene expression (Tronche & Yaniv, 1992), thus making its finding

justified.

insert Figure 3

Motif discovery validates TGF-beta signalling through Smad3/MyoD1 binding as
drivers of transcriptional differences in muscle of divergent feed efficient cattle

The analysis of the 217 genes co-expressed with *TGFB1* (Figure 4A) showed most

target genes motifs were enriched for master regulators of muscle differentiation, namely,

MEF2 (NES=10.42) a MADS box Transcription factor with 148 target genes, and *MYOD1*

(NES=8.12), a bHLH transcription factor (CANNTG) with 135 direct target genes (Figure

4B, S14 Supporting Information). To evaluate the precision of our predicted *MYOD1* (bHLH)

target genes we assessed how many of these TF-TG relationships had been previously

experimentally reported. Based on *MYOD1* ChIP-seq binding in mouse myotubules, 86 genes

had already been associated with *MYOD1* resulting in 63% success rate (hypergeometric test

1.72E-22). *SMAD3*, the effector molecule of *TGFB1* signalling is known to recruit *MYOD1* to

drive transcriptional changes during muscle differentiation (Mullen *et al*, 2011). Thus, we

evaluated whether predicted *MYOD1* target genes were enriched for known *SMAD3* target

genes resulting in 21 out of 135 *MYOD1* predicted target genes presented *SMAD3* ChIP-seq

binding in myotubes. Thus indicating there is a statistically significant association between

MYOD1 target genes and *SMAD3* target genes in myotubes (hypergeometric test 1.98 E-6)

(Figure 4 C) (Mullen *et al*, 2011) In contrast, no significant association was found between

predicted *MYOD1* target genes in this study and *SMAD3* target genes in other cell lines, such

as pro-B and ES cell (hypergeometric test 0.056 and 0.076, respectively) (Mullen *et al*,

2011). That is in agreement that the effect of TGFB signalling driven by SMAD3 DNA

binding is tissue-specific (Liu *et al*, 2001). Our analysis predicted 621 potential *MYOD1*

binding sites, of which 114 (18%) and 153 (24.5%) present a *MYOD1* ChIP-seq signal in

mouse C2C12 myotubes cells (Mullen *et al*, 2011) and in primary myotubes (Cao *et al*,

2009), respectively.

Finally, we evaluate whether predicted MYOD1 binding regions were regulatory

regions active in muscle cells across different species, namely human (S15 Supporting

Information), mouse (S16 and S17 Supporting Information) and cow (S18 Supporting

Information). To tackle this issue we performed an enrichment analysis across 2113 open-

chromatin ENCODE tracks. This analysis resulted in a clear enrichment of our predicted

MYOD1 binding regions with H3K27ac (NES=15.98) and H3K9ac (NES=8.78) regions in

skeletal muscle (Figure 4D). Both chromatin marks are associated with active transcription,

H3K27ac related to active enhancers and H3K9ac related to active gene transcription (Shin *et*

al, 2012) thus, validating most of our enhancer predictions in human are active in skeletal

muscle. Once in cow, we assess the overlap of predicted MYOD1 enhancers and promoter

regions in cow muscle experimentally detected with H3K4me3 (Cao *et al*, 2009). This

resulted in 282 regions out of 671 (42 %) overlap when only 11 regions are expected to

overlap by random 1000 permutation test) (Figure 4E).

insert Figure 4

Differential co-expression

Although the general co-expression network give us important insights about

regulatory genes and their behaviour, by creating specific networks for high and low FE and

comparing the connectivity of the genes in each one, we can identify genes that change their

behaviour depending on the situation, moving from highly connected to lowly connected and

vice-versa. We were able to identify 87 differentially connected genes between high and low

FE (P<0.05); 63 mainly expressed in liver, 19 in muscle and 3, 1 and 1 in hypothalamus,

adrenal gland and pituitary, respectively (S19 Supporting Information). Those genes were

enriched for terms such as regulation of blood coagulation (Padj=3.14 x 10^{-10}), fibrinolysis

(Padj=7.71 x10^{-7}), platelet degranulation (Padj=7.49 x10^{-6}), regulation of peptidase activity

(Padj=6.16 x 10^{-4}), antimicrobial humoral response (Padj=2.49 x10^{-3}), acute inflammatory

response (Padj=2.18 x10^{-4}) and induction of bacterial agglutination (Padj=3.58 x 10^{-2}) (S20

Supporting Information). It is important to highlight 20 of the differentially connected genes

were also differentially expressed (Table 1) and three of them, i.e. *SST, JCHAIN* and

IGFBP1, were secreted in plasma as well, which make them very promising potential

biomarkers.

insert Table 1

Comparing the two networks there was no large difference regarding the number of

genes and connections. While high FE network contained 1,074 genes in total and 28,018

connections, low FE network was composed by 1,098 genes and 30,705 connections. For all

tissues, low FE networks showed more connections but the difference is slight, being the

bigger difference of 44 versus 40 connections per gene in liver.

Discussion

Feed efficiency is a complex trait, regulated by several biological processes. Thus, the

indication of genomic regions associated with this phenotype, as well as regulators genes and

biomarkers to select superior animals and to direct management decisions is still a great

challenge. In this work, multi-tissue transcriptomic data of high and low feed efficient

Nellore bulls were analysed through robust co-expression network methodologies in order to

uncover some of the biology that governs this traits and put forward candidate genes to be

focus of further research. In this sense, the validation of target genes of main transcription

factors (key regulators) in the network by motif search proves the efficacy of the

methodology for network construction and prioritizes some transcription factors as central

regulators (Aerts *et al*, 2010; Naval-Sańchez *et al*, 2013; Potier *et al*, 2014). Moreover, the

addition of a category of genes coding proteins secreted in plasma in the co-expression

analysis highlight genes with potential to be explored as biomarkers of feed efficiency. We

were able to identify genes related to main biological process associated with feed efficiency

and indicate key regulators, as can be seen on the following lines.

Firstly, it is worthy to mention the 98 animals used to select the high and low FE

groups in this study have been previously analysed regarding several phenotypic and

molecular measures (Alexandre *et al*, 2015; Mota *et al*, 2017; Novais *et al*, 2018). It was

observed high and low FE groups had similar body weight gain, carcass yield and loin eye

area but low FE animals had higher feed intake, greater fat deposition, higher serum

cholesterol levels, as well as hepatic inflammatory response, indicated by transcriptome

analysis of liver biopsy and proved by the higher number of periportal mononuclear infiltrate

(histopathology) and increased serum gamma-glutamyl-transferase (GGT, a biomarker of

liver injury) in this group (Alexandre *et al*, 2015). In the present study, the simultaneous

analysis of five distinct tissues revealed a prominence of the hepatic tissue. Liver presented

the most connected genes in the network, the higher number of differentially connected genes

and higher number of secreted genes, which although can be explained by its biological

function, are enriched mostly for terms related to lipid homeostasis and inflammatory

response. Moreover, the top five most connected regulators in the network are co-expressed

mainly with genes highly expressed in liver and also enriched for inflammatory response.

The relationship between FE and genes or pathways related to immune response and

lipid metabolism is becoming more evident, as recent studies also reported it in beef cattle

(Karisa *et al*, 2014; Weber *et al*, 2016; Paradis *et al*, 2015; Zarek *et al*, 2017; Mukiibi *et al*,

2018) and pigs (Gondret *et al*, 2017; Ramayo-Caldas *et al*, 2018). In our previous work

(Alexandre *et al*, 2015), we proposed increased liver lesions associated with higher

inflammatory response in liver of low FE animals could be due to increased lipogenesis

and/or higher bacterial infection in the liver. While further evidence is needed to test these

hypotheses, the enrichment of terms such as induction of bacterial agglutination and response

to lipopolysaccharide makes bacterial infection a strong possibility. Indeed, pigs with low FE

were reported to have higher intestinal inflammation, neutrophil infiltration biomarkers and

increased serum endotoxin (lipopolysaccharide and other bacterial products) which could be

related to increased bacterial infection or to decreased capacity to neutralize endotoxins

(Mani *et al*, 2013). The authors hypothesized differences in bacterial population could

partially explain the increase in circulating endotoxins, which could also be true for cattle

given that differences in intestinal and ruminal bacterial population between high and low FE

animals have already been reported (Myer *et al*, 2015, 2016). Furthermore, the literature

reports lipopolysaccharides (LPS) may cause up-regulation of adrenomedullin (*ADM*)

hormone (Shindo *et al*, 1998), an up-regulated gene in low FE individuals as showed here. It

was also demonstrated in rats that intravenous infusion of LPS caused up-regulation of *ADM*

in ileum, liver, lung, aorta, skeletal muscle and blood vessels (Shoji *et al*, 1995) whereas in

our study, *ADM* presented differential expression in muscle, but not in liver.

Against pathogen invasion, a tightly regulated adaptive immune response must be

triggered in order to allow T lymphocytes to produce cytokines or chemokines and B cells to

differentiate and produce antibodies (Hermann-Kleiter & Baier, 2014). This regulation is

known to be strongly influenced by the expression level and transcriptional activity of several

nuclear receptors, including the NR2F-family, which consists of three orphan receptors:

NR2F1, NR2F2 and NR2F6 (Hermann-Kleiter & Baier, 2014). Those receptors present

highly conserved DNA and ligand binding domains among each other and across species

(Pereira *et al*, 2000), and all three are expressed in adaptive and immune cells (Hermann-

Kleiter & Baier, 2014). In our study, NR2F6 appeared as the second most connected

regulator gene in the network while the other family members, although present in our

expression data, were not selected by any of our inclusion criteria, thus indicating they might

not be so relevant in our conditions. Indeed, NR2F6 appears to be a critical regulatory factor

in the adaptive immune system by directly repressing the transcription of key cytokine genes

in T effector cells (Hermann-Kleiter *et al*, 2008; Klepsch *et al*, 2016). The role of NR2F6 as a

key regulator of inflammatory response in our network was validated at gene level by the

identification of the binding motif HNF4-NR2F2 (transfac_pro-M01031) as one of the most

enriched in NR2F6 target genes, due to the high similarity between NR2F2 and NR2F6

biding sites. Furthermore, using open chromatin data public available, we provided

experimental evidence of the binding of TFs with highly similar binding motifs as NR2F6 in

hepatocyte cells in humans and in cattle, thus, indicating predicted target enhancers are

functional in this tissue.

Another regulator prioritized in our analysis is *TGFB1*, the sixth most connected gene

in the co-expression network, and a potential driver of transcriptional changes between high

and low FE cattle in muscle. This gene has been previously pointed as a master regulator of

FE in beef cattle, using genomics and metabolomics data (Widmann *et al*, 2015). Moreover,

our motif discovery analysis showed *TGFB1* co-expressed genes are mostly enriched for

binding site of master regulators of muscle differentiation as *MEF2* and *MYOD*. Indeed,

public available data show many of *TGFB1* target genes were associated with *MYOD*

(Mullen *et al*, 2011). It is known signalling pathways are an effective mechanism for cells to

respond to environmental cues by regulation gene expression. *TGFB1* signalling triggers the

phosphorylation of *SMAD2/3* transcription factors, which co-bind with cell-type master

regulators at the nuclear level allowing/triggering/leading to cell-type specific transcriptional

changes (Schmierer & Hill, 2007; Mullen *et al*, 2011). In skeletal muscle cells, myoblasts and

myotubes, *SMAD3* co-binds with *MYOD1* (Mullen *et al*, 2011). The overlap between

MYOD1 and *SMAD3* target genes demonstrate the significant association between both genes

in skeletal muscle, in agreement with the tissue-specificity *TGFB1* signalling response

(Mullen *et al*, 2011). The overlap percentage between our predicted binding sites and

MYOD1 Chip-seq data (18 and 24.5%) confirms previous analysis in mice where they

reported only 20% of experimental validated distal enhancers in mouse myotubes with a

bHLH (MyoD1) binding were actually bound by *MYOD* ChIP-seq data (Blum et al. 2012).

Thus, suggesting additional transcription-factors and/or histone modification have a key role

in *MYOD1* binding. The *SMAD3/MYOD1* co-bound regions for known target genes are also

captured, such as the promoter regions of *ACTA1* and *ANKRD1*, both genes involved in

skeletal muscle differentiation (Figure 4C). We also demonstrated predicted *MYOD1* binding

regions are enriched for muscle regulatory regions across species (human, mouse and cow).

Altogether, we showed co-expressed genes with *TGFB1* are enriched for

SMAD3/MYOD1 binding sites, which we validate at the gene and enhancer level by proving

not only *MYOD1* and *SMAD3* binding, but also their accessibility, in human, mouse and cow.

In pigs, it has been indicated increased feed efficiency is associated with stimulation of

muscle growth through TGF-β signalling pathway (Jing *et al*, 2015). Finally, although not

directly co-expressed with *TGFB1*, oxytocin (*OXT*) was DE in muscle and despite the lack of

knowledge on its role in this tissue, previous work in cattle have shown a massive increase of

OXT expression in muscle of bovines chronically exposed to anabolic steroids (Jager *et al*,

2011). It is not known yet if oxytocin alone have an anabolic activity, but in a context where

muscle growth seems to be associated with high FE animals, this is a hormone that worth

further investigation.

From the 13 regulator genes that are DE between groups, six are involved in

respiratory chain and are up-regulated in high FE group. Genes *ND1, ND4, ND4L, ND5, ND6*

and *ND2,* which is DE but not identified as key regulator, are core subunits of the

mitochondrial membrane respiratory chain Complex I (CI) which functions in the transfer of

electrons from NADH to the respiratory chain, while *ATP8* is part of Complex V and

produces ATP from ADP in the presence of the proton gradient across the membrane.

Interestingly, greater quantity of mitochondrial CI protein were associated with high FE cattle

by Ramos and Kerley (2013) whereas Davis et al. (2016) found higher CI-CII and CI-CIII

concentration ratios for the same group. Other studies demonstrated high FE animals

consume less oxygen (Chaves *et al*, 2015) and present lower plasma CO_2 concentrations,

which suggests a decreased oxidation process (Gonano *et al*, 2014). In general, the literature

suggests mitochondrial ADP has greater control of oxidative phosphorylation in high FE

individuals (Lancaster *et al*, 2014) and their increased mitochondrial function may contribute

to feed efficiency (Connor *et al*, 2010). In pigs, differences in mitochondrial function were

reported when analysing muscle (Vincent *et al*, 2015), blood (Liu *et al*, 2016) and adipose

tissue transcriptomes (Louveau *et al*, 2016). Differences in metabolic rate associated with FE

has long been discussed (Herd & Arthur, 2009) and here is corroborated by the up-regulation

of *TSHB* in high FE animals, which stimulates production of T3 and T4 in thyroid thus

increasing metabolism. It is inhibited by *SST*, a down-regulated hormone in this group which

was also found to be differentially connected between high and low FE.

Looking at the DE genes, many hormones can be identified. Hormones are signalling

proteins that are transported by the circulatory system to target distant organs in order to

regulate physiology. Regarding the relationship between FE and other production traits of

economic importance, *FSHB*, responsible for spermatozoa production by activating Sertoli

cells in the testicles (Walker & Cheng, 2005), is up-regulated in low FE group and is

inhibited by follistatin (*FST*), a gene found to be down-regulated in the same group.

Moreover, in rats, it was already demonstrated *FSH* secretion is stimulated by somatostatin

expression, which is up-regulated in low FE animals (Kitaoka *et al*, 1989). In this scenario,

one could argue that selection for high FE delay reproduction traits, something that could be

related to the lower fat deposition in this group, as previously observed (Alexandre *et al*,

2015; Santana *et al*, 2012; Gomes *et al*, 2012). Indeed, differences in body composition and

in intermediary metabolism can impact on reproductive traits (Shaffer *et al*, 2011) and it has

been observed before that feed efficient bulls present features of delayed sexual maturity, i.e.

decreased progressive motility of the sperm and higher abundance of tail abnormalities

(Montanholi *et al*, 2016; Fontoura *et al*, 2016). Moreover, high FE heifers presented less fat

deposition and later sexual maturity which results in calving later in the calving season than

their low FE counterparts (Randel & Welsh, 2013; Shaffer *et al*, 2011). It is important to

point that low FE animals also present down-regulation of *AMH* and the fall of this hormone

in serum was pointed as an excellent marker of Sertoli cells pubertal development (Rey *et al*,

1993).

Concerning the differences in lipid metabolism in divergent FE phenotypes, *FGF21*, a

hormone up-regulated in liver of high FE animals, is associated in humans to decrease in

body weight, blood triglycerides and LDL-cholesterol, with improvement in insulin

sensitivity (Cheung & Deng, 2014). It is an hepatokine released to the bloodstream and an

important regulator of lipid and glucose metabolism (Giralt *et al*, 2015). When we select its

first neighbours in the network and perform an enrichment analysis we indeed found terms

related to plasma lipoprotein particle remodelling, regulation of lipoprotein oxidation and

cholesterol efflux mostly due to *FGF21* co-expression with the apolipoproteins *APOA4*,

APOC3 and *APOM*. In the same context, pro-melanin-concentrating hormone (*PMCH*)

encodes three neuropeptides: neuropeptide-glycine-glutamic acid, neuropeptide-glutamic

acid-isoleucine and melanin-concentrating hormone (*MCH*) being the last one the most

extensively studied (Helgeson & Schmutz, 2008). *MCH* up-regulation has been related to

obesity and insulin resistance, as well as increased appetite and reduced metabolism in

murine models (Ludwig *et al*, 2001; Ito *et al*, 2003). *PMCH* gene is up-regulated in low FE

animals and harbour SNPs found to be associated with higher carcass fat levels and marbling

score (Walter *et al*, 2014; Helgeson & Schmutz, 2008).

In this work, we were able to identify several biological processes known to be related

to feed efficiency, which together with the validation of the main transcription factors of the

network, demonstrate the quality of the data and the robustness of the analyses, giving us the

confidence to indicate candidate genes to be regulators or biomarkers of superior animals for

this trait. The transcription factors NR2F6 and TGFB1 play central roles in liver and muscle,

respectively, by regulating genes related to inflammatory response and muscle development

and growth, two main biological mechanisms associated to feed efficiency. Likewise,

hormones and other proteins secreted in plasma as oxytocin, adrenomedulin, TSH,

somatostatin, follistatin and AMH are interesting molecules to be explored as potential

biomarkers of feed efficiency.

Material and methods

Phenotypic data and biological sample collection

All animal protocols were approved by the Institutional Animal Care and Use

Committee of Faculty of Food Engineering and Animal Sciences, University of São Paulo

(FZEA-USP – protocol number 14.1.636.74.1). All procedures to collect phenotypes and

biological samples were carried out at FZEA-USP, Pirassununga, State of São Paulo, Brazil.

Ninety eight Nellore bulls (16 to 20 months old and 376 ± 29 kg BW) were evaluated in a

feeding trial comprised of 21 days of adaptation to feedlot diet and place and a 70-day period

of data collection. Total mixed ration was offered *ad libitum* and daily dry matter intake

(DMI) was individually measured. Animals were weighted at the beginning, at the end and

every 2 weeks during the experimental period. Feed efficiency was estimated by residual feed

intake (RFI) which is the residual of the linear regression that estimates DMI based on

average daily gain and mid-test metabolic body weight (Koch *et al*, 1963). Forty animals

selected either as high feed efficiency (HFE) or low feed efficiency (LFE) groups were

slaughtered on two days with a 6-day interval. Adrenal gland, hypothalamus, liver, muscle

and pituitary samples were collected from each animal, rapidly frozen in liquid nitrogen and

stored at -80 °C. Further information about management and phenotypic measures of the

animals used in this study can be found in Alexandre et al. (2015).

527   RNAseq data generation

Samples of nine animals from each feed efficiency group (high and low) were

selected for RNAseq using RFI measure. For hypothalamus and pituitary, the nitrogen frozen

tissue was macerated with crucible and pistil and stored in aliquots at -80 °C. Then, RNA was

extracted using AllPrep DNA/RNA/Protein Mini kit (QIAGEN, Crawley, UK). For liver,

muscle and adrenal gland, a cut was made in the frozen tissue and the RNA was extracted

using RNeasy Mini Kit (QIAGEN, Crawley, UK). RNA quality and quantity were assessed

using automated capillary gel electrophoresis on a Bioanalyzer 2100 with RNA 6000 Nano

Labchips according to the manufacturer's instructions (Agilent Technologies Ireland, Dublin,

Ireland). Samples that presented an RNA integrity number (RIN) less than 8.0 were

discarded.

RNA libraries were constructed using the TruSeq™ Stranded mRNA LT Sample Prep

Protocol and sequenced on Illumina HiSeq 2500 equipment in a HiSeq Flow Cell v4 using

HiSeq SBS Kit v4 (2x100pb). Liver, pituitary and hypothalamus were sequenced on the same

run, each one in a different lane. Muscle and adrenal gland were sequenced in a second run,

in different lanes.

Gene expression estimation

The quality of the sequencing was evaluated using the software FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Sequence alignment against the

bovine reference genome (UMD3.1) was performed using STAR (Dobin *et al*, 2013),

according to the standard parameters and including the annotation file (Ensembl release 89)

and secondary alignments, duplicated reads and reads failing vendor quality checks were

removed using Samtools (Li *et al*, 2009). Then, HTseq (Anders *et al*, 2014) was used to

generate gene read counts and expression values were estimated by reads per kilobase of gene

per million mapped reads (RPKM). Genes with average value lower than 0.2 FPKM across

all samples and tissues were discarded.

Gene expression normalization was performed using the following mixed effect

model (Reverter *et al*, 2005):

$$Y_{ijkl} = \mu + L_i + G_j + GT_{jk} + GP_{jl} + e_{ijkl}$$

where the log2-transformed FPKM value for i-th library (86 levels), j-th gene (17,354 levels),

k-th tissue (5 levels), l-th RFI phenotype (2 levels), corresponding to Y_{ijkl}, was modelled as a

function of the fixed effect of library (L_i) and the random effects of gene (G_j), gene by tissue

(GT_{jk}) and gene by RFI phenotype (GP_{jl}). Random residual (e_{ijkl}) was assumed to be

independent and identically distributed. Variance component estimates and solutions to the

model were obtained using VCE6 (Eildert Groeneveld, Milena Kovac and Norbert Mielenz,

ftp://ftp.tzv.fal.de/pub/vce6/doc/vce6-manual-3.1-A4.pdf). Normalized mean expression

(NME) values for each gene were defined as the linear combination of the solutions for

random effects.

The mixed model used to normalize the expression data explained 96% of the

variation in gene expression, of which the largest proportion (0.30) was due to tissue-

specificity. Contrariwise, differences between high and low FE represented no variation

(0.27E-11). For that reason, normalized mean expression (NME) were only used to identify

tissue specific genes and the raw FPKM values were used for differential expression and co-

expression analysis.

Gene selection for network construction

In order to select a set of relevant genes for network analysis, we defined five

categories based on the following inclusion criteria:

1. Differential expression (DE) - The mean expression value of each gene, for

each group (high and low FE) and each tissue was calculated and then the expression of low

FE group was subtracted from the expression in high FE group. Next, genes were ranked

according to their mean expression in all samples for each tissue and divided in five bins.

Genes were considered differentially expressed when the difference between the expression

in high and low FE groups were greater than 3.1 or smaller than -3.1 standard deviation from

the mean in each bin, corresponding to a t-test $P<0.001$.

2. Harbouring SNPs - Genes harbouring SNPs associated with feed efficiency,

mainly indicated by GWAS, were identified using PubMed database

(www.ncbi.nlm.nih.gov/pubmed/) and AnimalQTL database (www.animalgenome.org/cgi-

bin/QTLdb/index) and only bovine data were considered regardless of breed.

3. Tissue specific (TS) - A gene was considered as tissue specific when the

average NME in that tissue was greater than one standard deviation from the mean of all

genes AND the average NME in all the other four tissues was smaller than zero.

4. Secreted - The human secretome database

(www.proteinatlas.org/humanproteome/secretome) was used to select genes encoding

proteins secreted in plasma by any of the analysed tissues (adrenal gland, hypothalamus,

liver, muscle and pituitary).

594   5. Key regulators - In order to identify key regulatory genes to be included in the

co-expression network, a list of genes were obtained from the Animal Transcription Factor

Database (http://www.bioguo.org/AnimalTFDB/) and it was compared to a set of potential

target genes in each tissue, composed by the categories: TS, DE, harbouring SNPs and

secreted. The analysis was based on regulatory impact factor metrics (Reverter *et al*, 2010),

which comprises a set of two metrics designed to assign scores to regulator genes consistently

differentially co-expressed with target genes and to those with the most altered ability to

predict the abundance of target genes. Those scores deviating ± 1.96 standard deviation from

the mean (corresponding to $P < 0.05$) were considered significant. Genes presenting mean

expression value less than the mean of all genes expressed were not considered in this

analysis.

Some of the genes selected by the categories above were represented by more than

one ensemble ID. Those duplications were removed for further analysis, keeping only the

expression value of the most meaningful ensemble ID. Additionally, genes with mean

expression across the samples equal to zero were also removed from further analysis.

Co-expression network analysis

For gene network inference, genes selected by the five categories described

previously were used as nodes and significant connections (edges) between them were

identified using partial correlation and information theory (PCIT) algorithm (Reverter &

Chan, 2008), considering all animals and all tissues. PCIT determinates the significance of

the correlation between two nodes after accounting for all the other nodes in the network. The

output of PCIT was visualized on Cytoscape (Shannon *et al*, 2003).

Network validation through transcription factor biding motifs analysis

Using the regulatory impact factor metric (RIF) we prioritize key regulator genes from gene expression data and predict target genes based on co-expression network (PCIT). In order to assess whether those target genes were enriched for motifs associated to the top most connected regulators in the network with a DNA biding domain (transcription factors - TF), we performed motif discovery analysis in the set of co-expressed target genes (first neighbours of the TF) using i-cistarget method (Herrmann *et al*, 2012) and i-Regulon, a Cytoscape plug-in (Janky *et al*, 2014). These tools use human (hg19) as the reference species, therefore only genes with human orthology are assessed. Then, to validate the binding of the identified genomic regions by the TFs, we performed a region enrichment analysis across experimentally available TF bound regions from ChiP-seq in cell lines from the ENCODE consortium (1,394 TF binding site tracks). Finally, we converted identified enhancer regions to cow coordinates and searched for regions of open-chromatin using data from a public available studies in cow tissues.

Differential connectivity

In order to explore differentially connected genes between high and low FE, two networks were created, one for each condition, using the same methodology described before. Then, the number of connections of each gene in each condition was computed and scaled so that connectivity varied from 0 to 1, making possible to compare the same gene in the two networks. The connectivity in high RFI group was subtracted from the connectivity in low RFI group and results deviating ± 1.96 standard deviation from the mean were considered significant (P<0.05).

Functional Enrichment

Functional enrichment analysis was performed on the online platform GOrilla (Gene Ontology enRIchment anaLysis and visuaLizAtion tool, http://cbl-gorilla.cs.technion.ac.il/), using all genes that passed FPKM filter as background, hypergeometric test and multiple test correction (FDR - false discovery rate). The human database was used to take advantage of a more comprehensive knowledge regarding gene functions. GO terms were considered significant when $Padj<0.05$. For genes in co-expression networks, visualized using Cytoscape (Shannon *et al*, 2003), the functional enrichment was performed with BiNGO plug-in (Maere *et al*, 2005) using the same background genes and statistical test.

Acknowledgments

The authors thank Mrs. Elisângela C. M. Oliveira for all technical support. This study and PAA scholarships were funded by São Paulo Research Foundation (FAPESP) - Proc. 2014/07566-2, 2015/22276-3 and 2017/14707-0. M.N.S was funded by the CSIRO Science Excellence Research Office.

Conflict of interest

The authors declare that they have no conflict of interest.

Author contributions

HF was the overall project leader who conceived this study and supervised PAA in all data generation. TR performed the network analysis and supervised PAA in bioinformatics and data interpretation. MNS performed the motif discovery analysis. LRPN and JBSF provided informatics and statistical support. All authors contributed to and approved the final version of this manuscript.

Availability of supporting data

Datasets supporting the results of this article is public available in the European Nucleotide

Archive (ENA) as part of FAANG consortium under de study ID PRJEB27337 and can be

accessed following the link https://www.ebi.ac.uk/ena/data/view/PRJEB27337.

References

Aerts S, Quan XJ, Claeys A, Sanchez MN, Tate P, Yan J & Hassan BA (2010) Robust target gene discovery through transcriptome perturbations and genome-wide enhancer predictions in drosophila uncovers a regulatory basis for sensory specification. *PLoS Biol.* **8:**

Alexandre P a., Kogelman LJ a., Santana MH a., Passarelli D, Pulz LH, Fantinato-Neto P, Silva PL, Leme PR, Strefezzi RF, Coutinho LL, Ferraz JBS, Eler JP, Kadarmideen HN & Fukumasu H (2015) Liver transcriptomic networks reveal main biological processes associated with feed efficiency in beef cattle. *BMC Genomics* **16:** 1073 Available at: http://www.biomedcentral.com/1471-2164/16/1073

Anders S, Pyl PT & Huber W (2014) HTSeq - A Python framework to work with high-throughput sequencing data. *Bioinformatics* **31:** 166–9 Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4287950&tool=pmcentrez&rendertype=abstract [Accessed September 29, 2014]

Archer JA, Arthur PF, Herd RM, Parnell PF & Pitchford WS (1997) Optimum postweaning test for measurement of growth rate, feed intake, and feed efficiency in British breed cattle. *J. Anim. Sci.* **75:** 2024–32 Available at: http://www.ncbi.nlm.nih.gov/pubmed/9263047 [Accessed September 5, 2014]

Archer JA, Richardson EC, Herd RM & Arthur PF (1999) Potential for selection to improve efficiency of feed use in beef cattle: a review. *Aust. J. Agric. Res.* **50:** 147–61 Available at: http://www.publish.csiro.au/paper/A98075.htm [Accessed February 29, 2016]

Bolormaa S, Hayes BJ, Savin K, Hawken R, Barendse W, Arthur PF, Herd RM & Goddard ME (2011) Genome-wide association studies for feedlot and growth traits in cattle. *J. Anim. Sci.* **89:** 1684–97 Available at: http://www.ncbi.nlm.nih.gov/pubmed/21239664 [Accessed March 21, 2012]

Cafe LM, Robinson DL, Ferguson DM, Geesink GH & Greenwood PL (2011) Temperament and hypothalamic-pituitary-adrenal axis function are related and combine to affect growth, efficiency, carcass, and meat quality traits in Brahman steers. *Domest. Anim. Endocrinol.* **40:** 230–40 Available at: http://www.sciencedirect.com/science/article/pii/S0739724011000063 [Accessed February 21, 2016]

Cao P, Hanai J-I, Tanksale P, Imamura S, Sukhatme VP & Lecker SH (2009) Statin-induced muscle damage and atrogin-1 induction is the result of a geranylgeranylation defect. *FASEB J.* **23:** 2844–2854 Available at: http://www.ncbi.nlm.nih.gov/pubmed/19406843%5Cnhttp://www.fasebj.org/content/23/9/2844.full.pdf

Chaves AS, Nascimento ML, Tullio RR, Rosa AN, Alencar MM & Lanna DP (2015)

Relationship of efficiency indices with performance, heart rate, oxygen consumption, blood parameters, and estimated heat production in Nellore steers. *J. Anim. Sci.* **93:** 5036–46 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26523596 [Accessed February 28, 2016]

Chen L, Mao F, Crews DH, Vinsky M & Li C (2014) Phenotypic and genetic relationships of feeding behavior with feed intake, growth performance, feed efficiency, and carcass merit traits in Angus and Charolais steers. *J. Anim. Sci.* **92:** 974–83 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24492561 [Accessed February 21, 2016]

Cheung BM & Deng HB (2014) Fibroblast growth factor 21: a promising therapeutic target in obesity-related diseases. *Expert Rev Cardiovasc Ther* **12:** 659–666

Connor EE, Kahl S, Elsasser TH, Parker JS, Li RW, Van Tassell CP, Baldwin RL & Barao SM (2010) Enhanced mitochondrial complex gene function and reduced liver size may mediate improved feed efficiency of beef cattle during compensatory growth. *Funct. Integr. Genomics* **10:** 39–51 Available at: http://www.ncbi.nlm.nih.gov/pubmed/19777276 [Accessed February 28, 2016]

Davis MP, Brooks MA & Kerley MS (2016) Relationship between residual feed intake and lymphocyte mitochondrial complex protein concentration and ratio in crossbred steers. *J. Anim. Sci.* **94:** 1587–1591

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M & Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29:** 15–21 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23104886 [Accessed February 17, 2017]

Fontoura ABP, Montanholi YR, Diel de Amorim M, Foster RA, Chenier T & Miller SP (2016) Associations between feed efficiency, sexual maturity and fertility-related measures in young beef bulls. *Animal* **10:** 96–105 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26351012 [Accessed March 2, 2016]

Foote AP, Tait RG, Keisler DH, Hales KE & Freetly HC (2016) Leptin concentrations in finishing beef steers and heifers and their association with dry matter intake, average daily gain, feed efficiency, and body composition. *Domest. Anim. Endocrinol.* **55:** 136–41 Available at: http://www.sciencedirect.com/science/article/pii/S0739724016000035 [Accessed April 15, 2016]

Francisco CL, Resende FD, Benatti JMB, Castilhos AM, Cooke RF & Jorge AM (2015) Impacts of temperament on Nellore cattle: physiological responses, feedlot performance, and carcass characteristics. *J. Anim. Sci.* **93:** 5419 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26641061 [Accessed February 21, 2016]

Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Falcucci A & Tempio G (2013) Tackling Climate Change Through Livestock - A Global Assessment of Emissions and Mitigation Opportunities Rome Available at: http://www.fao.org/docrep/018/i3437e/i3437e.pdf

Giralt M, Gavaldà-Navarro A & Villarroya F (2015) Fibroblast growth factor-21, energy balance and obesity. *Mol. Cell. Endocrinol.* **418:** 66–73

Gomes RC, Sainz RD, Silva SL, César MC, Bonin MN & Leme PR (2012) Feedlot performance, feed efficiency reranking, carcass traits, body composition, energy requirements, meat quality and calpain system activity in Nellore steers with low and high residual feed intake. *Livest. Sci.* **150:** 265–273 Available at: http://www.sciencedirect.com/science/article/pii/S1871141312003587 [Accessed February 19, 2014]

Gonano C, Montanholi Y, Schenkel F, Smith B, Cant J & Miller S (2014) The relationship between feed efficiency and the circadian profile of blood plasma analytes measured in beef heifers at different physiological stages. *Animal* **8:** 1684–1698 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/24923431 [Accessed February 28, 2016]

Gondret F, Vincent A, Houée-Bigot M, Siegel A, Lagarrigue S, Causeur D, Gilbert H &
Louveau I (2017) A transcriptome multi-tissue analysis identifies biological pathways
and genes associated with variations in feed efficiency of growing pigs. *BMC Genomics*
18: 1–17

Hayes BJ, Lewin H a & Goddard ME (2013) The future of livestock breeding: genomic
selection for efficiency, reduced emissions intensity, and adaptation. *Trends Genet.* **29:**
206–14 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23261029 [Accessed
October 15, 2014]

Helgeson SC & Schmutz SM (2008) Genetic variation in the pro-melanin-concentrating
hormone gene affects carcass traits in Bos taurus cattle. *Anim. Genet.* **39:** 310–315

Herd RM & Arthur PF (2009) Physiological basis for residual feed intake. *J. Anim. Sci.* **87:**
E64-71 Available at: http://jas.fass.org/cgi/content/abstract/87/14_suppl/E64 [Accessed
March 28, 2012]

Herd RM, Oddy VH & Richardson EC (2004) Biological basis for variation in residual feed
intake in beef cattle. 1. Review of potential mechanisms. *Aust. J. Exp. Agric.* **44:** 423
Available at:
http://www.publish.csiro.au/view/journals/dsp_journal_fulltext.cfm?nid=72&f=EA0222
0 [Accessed April 18, 2012]

Hermann-Kleiter N & Baier G (2014) Orphan nuclear receptor NR2F6 acts as an essential
gatekeeper of Th17 CD4+ T cell effector functions. *Cell Commun. Signal.* **12:** 1–12

Hermann-Kleiter N, Gruber T, Lutz-Nicoladoni C, Thuille N, Fresser F, Labi V,
Schiefermeier N, Warnecke M, Huber L, Villunger A, Eichele G, Kaminski S & Baier G
(2008) The Nuclear Orphan Receptor NR2F6 Suppresses Lymphocyte Activation and T
Helper 17-Dependent Autoimmunity. *Immunity* **29:** 205–216

Herrmann C, Van De Sande B, Potier D & Aerts S (2012) i-cisTarget: An integrative
genomics method for the prediction of regulatory features and cis-regulatory modules.
Nucleic Acids Res. **40:**

Ito M, Gomori A, Ishihara A, Oda Z, Mashiko S, Matsushita H, Yumoto M, Ito M, Sano H,
Tokita S, Moriya M, Iwaasa H & Kanatani A (2003) Characterization of MCH-mediated
obesity in mice. *Am. J. Physiol. - Endocrinol. Metab.* **284:** E940–E945

Jager N De, Hudson NJ, Reverter A, Wang Y, Nagaraj SH, Cafe LM, Greenwood PL,
Barnard RT, Kongsuwan KP, Dalrymple BP, Pl G, Rt B, Kp K & Bp D (2011) Chronic
exposure to anabolic steroids induces the muscle expression of oxytocin and a more than
fiftyfold increase in circulating oxytocin in cattle. **:** 467–478

Janky R, Verfaillie A, Imrichová H, van de Sande B, Standaert L, Christiaens V, Hulselmans
G, Herten K, Naval Sanchez M, Potier D, Svetlichnyy D, Kalender Atak Z, Fiers M,
Marine JC & Aerts S (2014) iRegulon: From a Gene List to a Gene Regulatory Network
Using Large Motif and Track Collections. *PLoS Comput. Biol.* **10:**

Jing L, Hou Y, Wu H, Miao Y, Li X, Cao J, Brameld JM, Parr T & Zhao S (2015)
Transcriptome analysis of mRNA and miRNA in skeletal muscle indicates an important
network for differential Residual Feed Intake in pigs. *Sci. Rep.* **5:** 11953 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4493709&tool=pmcentrez&
rendertype=abstract [Accessed April 6, 2016]

Karisa B, Moore S & Plastow G (2014) Analysis of biological networks and biological
pathways associated with residual feed intake in beef cattle. *Anim. Sci. J.* **85:** 374–87
Available at: http://www.ncbi.nlm.nih.gov/pubmed/24373146 [Accessed February 1,
2015]

Kelly AK, McGee M, Crews DH, Fahey AG, Wylie AR & Kenny DA (2010) Effect of
divergence in residual feed intake on feeding behavior, blood metabolic variables, and

body composition traits in growing beef heifers. *J. Anim. Sci.* **88:** 109–23 Available at: http://www.ncbi.nlm.nih.gov/pubmed/19820067 [Accessed February 21, 2016]

Kitaoka M, Takano K, Kojima I & Ogata E (1989) A stimulatory effect of somatostatin: Enhancement of activin A-mediated FSH secretion in rat pituitary cells. *Biochem. Biophys. Res. Commun.* **162:** 958–962

Klepsch V, Hermann-Kleiter N & Baier G (2016) Beyond CTLA-4 and PD-1: Orphan nuclear receptor NR2F6 as T cell signaling switch and emerging target in cancer immunotherapy. *Immunol. Lett.* **178:** 31–36

Koch RM, Swiger LA, Chambers D & Gregory KE (1963) Efficiency of Feed Use in Beef Cattle. *J. Anim. Sci.* **22:** 486–494

de la Fuente A (2010) From 'differential expression' to 'differential networking' - identification of dysfunctional regulatory networks in diseases. *Trends Genet.* **26:** 326–33 Available at: http://www.ncbi.nlm.nih.gov/pubmed/20570387 [Accessed July 18, 2014]

Lancaster PA, Carstens GE, Michal JJ, Brennan KM, Johnson KA & Davis ME (2014) Relationships between residual feed intake and hepatic mitochondrial function in growing beef cattle. *J. Anim. Sci.* **92:** 3134–41 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24894006 [Accessed February 29, 2016]

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G & Durbin R (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25:** 2078–9 Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2723002&tool=pmcentrez&rendertype=abstract [Accessed July 9, 2014]

Liu D, Black BL & Derynck R (2001) TGF-beta inhibits muscle differentiation through functional repression of myogenic transcription factors by Smad3. *Genes Dev.* **15:** 2950–2966

Liu H, Nguyen YT, Nettleton D, Dekkers JCM & Tuggle CK (2016) Post-weaning blood transcriptomic differences between Yorkshire pigs divergently selected for residual feed intake. *BMC Genomics* **17:** 73 Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4724083&tool=pmcentrez&rendertype=abstract [Accessed February 1, 2016]

Louveau I, Vincent A, Tacher S, Gilbert H & Gondret F (2016) Increased expressions of genes and proteins involved in mitochondrial oxidation and antioxidant pathway in adipose tissue of pigs selected for a low residual feed intake. *J. Anim. Sci.* **94:** 5042–5054

Ludwig DS, Tritos NA, Mastaitis JW, Kulkarni R, Kokkotou E, Elmquist J, Lowell B, Flier JS & Maratos-Flier E (2001) Melanin-concentrating hormone overexpression in transgenic mice leads to obesity and insulin resistance. *J. Clin. Invest.* **107:** 379–386

Mader CJ, Montanholi YR, Wang YJ, Miller SP, Mandell IB, McBride BW & Swanson KC (2009) Relationships among measures of growth performance and efficiency with carcass traits, visceral organ mass, and pancreatic digestive enzymes in feedlot cattle. *J. Anim. Sci.* **87:** 1548–57 Available at: http://www.ncbi.nlm.nih.gov/pubmed/18952722 [Accessed January 30, 2015]

Maere S, Heymans K & Kuiper M (2005) BiNGO: A Cytoscape plugin to assess overrepresentation of Gene Ontology categories in Biological Networks. *Bioinformatics* **21:** 3448–3449

Mani V, Harris AJ, Keating AF, Weber TE, Dekkers JCM & Gabler NK (2013) Intestinal integrity, endotoxin transport and detoxification in pigs divergently selected for residual feed intake. *J. Anim. Sci.* **91:** 2141–50 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23463550 [Accessed March 1, 2016]

Montanholi YR, Fontoura ABP, Diel de Amorim M, Foster RA, Chenier T & Miller SP (2016) Seminal plasma protein concentrations vary with feed efficiency and fertility-related measures in young beef bulls. *Reprod. Biol.* **16:** 147–156

Montanholi YR, Swanson KC, Palme R, Schenkel FS, McBride BW, Lu D & Miller SP (2010) Assessing feed efficiency in beef steers through feeding behavior, infrared thermography and glucocorticoids. *Animal* **4:** 692–701 Available at: http://www.ncbi.nlm.nih.gov/pubmed/22444121 [Accessed February 29, 2016]

Montanholi YR, Swanson KC, Schenkel FS, McBride BW, Caldwell TR & Miller SP (2009) On the determination of residual feed intake and associations of infrared thermography with efficiency and ultrasound traits in beef bulls. *Livest. Sci.* **125:** 22–30 Available at: http://www.sciencedirect.com/science/article/pii/S1871141309000766 [Accessed February 29, 2016]

Mota LFM, Bonafé CM, Alexandre PA, Santana MH, Novais FJ, Toriyama E, Pires AV, da Luz Silva S, Leme PR, Ferraz JBS & Fukumasu H (2017) Circulating leptin and its muscle gene expression in Nellore cattle with divergent feed efficiency. *J. Anim. Sci. Biotechnol.* **8:**

Mukiibi R, Vinsky M, Keogh KA, Fitzsimmons C, Stothard P, Waters SM & Li C (2018) Transcriptome analyses reveal reduced hepatic lipid synthesis and accumulation in more feed efficient beef cattle. **:** 1–12

Mullen AC, Orlando DA, Newman JJ, Lovén J, Kumar RM, Bilodeau S, Reddy J, Guenther MG, Dekoter RP & Young RA (2011) Master transcription factors determine cell-type-specific responses to TGF-β signaling. *Cell* **147:** 565–576

Myer PR, Smith TPL, Wells JE, Kuehn LA & Freetly HC (2015) Rumen microbiome from steers differing in feed efficiency. *PLoS One* **10:** e0129174 Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4451142&tool=pmcentrez&rendertype=abstract [Accessed March 1, 2016]

Myer PR, Wells JE, Smith TPL, Kuehn LA & Freetly HC (2016) Microbial community profiles of the jejunum from steers differing in feed efficiency. *J. Anim. Sci.* **94:** 327 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26812338 [Accessed March 27, 2017]

Naval-Sańchez M, Potier D, Haagen L, Sánchez M, Munck S, Van De Sande B, Casares F, Christiaens V & Aerts S (2013) Comparative motif discovery combined with comparative transcriptomics yields accurate targetome and enhancer predictions. *Genome Res.* **23:** 74–88

Nkrumah JD, Keisler DH, Crews DH, Basarab JA, Wang Z, Li C, Price MA, Okine EK & Moore SS (2007) Genetic and phenotypic relationships of serum leptin concentration with performance, efficiency of gain, and carcass merit of feedlot cattle. *J. Anim. Sci.* **85:** 2147–55 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17468416 [Accessed February 23, 2016]

Novais FJ, Pires PRL, Alexandre PA, Dromms RA, Iglesias AH, Ferraz JBS, Styczynski MP & Fukumasu H (2018) Serum metabolomics predicts biological processes associated to feed efficiency in beef cattle [under review]. *Sci. Rep.*

de Oliveira PSN, Cesar ASM, do Nascimento ML, Chaves AS, Tizioto PC, Tullio RR, Lanna DPD, Rosa AN, Sonstegard TS, Mourao GB, Reecy JM, Garrick DJ, Mudadu MA, Coutinho LL & Regitano LCA (2014) Identification of genomic regions associated with feed efficiency in Nelore cattle. *BMC Genet.* **15:** 100 Available at: http://www.biomedcentral.com/1471-2156/15/100 [Accessed December 27, 2014]

Paradis F, Yue S, Grant JR, Stothard P, Basarab JA & Fitzsimmons C (2015) Transcriptomic analysis by RNA sequencing reveals that hepatic interferon-induced genes may be associated with feed efficiency in beef heifers. *J. Anim. Sci.* **93:** 3331 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/26440002 [Accessed November 8, 2015]

Pereira FA, Tsai MJ & Tsai SY (2000) COUP-TF orphan nuclear receptors in development and differentiation. *Cell Mol Life Sci* **57:** 1388–1398

Potier D, Davie K, Hulselmans G, NavalSanchez M, Haagen L, Huynh-Thu VA, Koldere D, Celik A, Geurts P, Christiaens V & Aerts S (2014) Mapping Gene Regulatory Networks in Drosophila Eye Development by Large-Scale Transcriptome Perturbations and Motif Inference. *Cell Rep.* **9:** 2290–2303

Ramayo-Caldas Y, Ballester M, Sánchez JP, González-Rodríguez O, Revilla M, Reyer H, Wimmers K, Torrallardona D & Quintanilla R (2018) Integrative approach using liver and duodenum RNA-Seq data identifies candidate genes and pathways associated with feed efficiency in pigs. *Sci. Rep.* **8:**

Ramos MH & Kerley MS (2013) Mitochondrial complex I protein differs among residual feed intake phenotype in beef cattle. *J. Anim. Sci.* **91:** 3299–304 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23798519 [Accessed February 29, 2016]

Randel RD & Welsh TH (2013) Joint Alpharma-Beef Species Symposium: interactions of feed efficiency with beef heifer reproductive development. *J. Anim. Sci.* **91:** 1323–8 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23048157 [Accessed March 2, 2016]

Reverter A, Barris W, McWilliam S, Byrne KA, Wang YH, Tan SH, Hudson N & Dalrymple BP (2005) Validation of alternative methods of data normalization in gene co-expression studies. *Bioinformatics* **21:** 1112–1120

Reverter A & Chan EKF (2008) Combining partial correlation and an information theory approach to the reversed engineering of gene co-expression networks. *Bioinformatics* **24:** 2491–2497 Available at: http://www.ncbi.nlm.nih.gov/pubmed/18784117 [Accessed June 11, 2017]

Reverter A, Hudson NJ, Nagaraj SH, Pérez-Enciso M & Dalrymple BP (2010) Regulatory impact factors: Unraveling the transcriptional regulation of complex traits from expression data. *Bioinformatics* **26:** 896–904

Rey R, Lordereau-Richard I, Carel JC, Barbet P, Cate RL, Roger M, Chaussain JL & Josso N (1993) Anti-müllerian hormone and testosterone serum levels are inversely during normal and precocious pubertal development. *J. Clin. Endocrinol. Metab.* **77:** 1220–1226

Rolf MM, Taylor JF, Schnabel RD, McKay SD, McClure MC, Northcutt SL, Kerley MS & Weaber RL (2011) Genome-wide association analysis for feed efficiency in Angus cattle. *Anim. Genet.* Available at: http://www.ncbi.nlm.nih.gov/pubmed/22497295 [Accessed April 24, 2012]

Santana MHA, Rossi P, Almeida R & Cucco DC (2012) Feed efficiency and its correlations with carcass traits measured by ultrasound in Nellore bulls. *Livest. Sci.* **145:** 252–257 Available at: http://linkinghub.elsevier.com/retrieve/pii/S1871141312000686 [Accessed January 13, 2015]

Santana MH, Utsunomiya YT, Neves HH, Gomes RC, Garcia JF, Fukumasu H, Silva SL, Oliveira Junior G a, Alexandre P a, Leme PR, Brassaloti R a, Coutinho LL, Lopes TG, Meirelles F V, Eler JP & Ferraz JB (2014) Genome-wide association analysis of feed intake and residual feed intake in Nellore cattle. *BMC Genet.* **15:** 21 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24517472 [Accessed February 19, 2014]

Schmierer B & Hill CS (2007) TGFβ-SMAD signal transduction: Molecular specificity and functional flexibility. *Nat. Rev. Mol. Cell Biol.* **8:** 970–982

Seabury CM, Oldeschulte DL, Saatchi M, Beever JE, Decker JE, Halley YA, Bhattarai EK, Molaei M, Freetly HC, Hansen SL, Yampara-Iquise H, Johnson KA, Kerley MS, Kim JW, Loy DD, Marques E, Neibergs HL, Schnabel RD, Shike DW, Spangler ML, et al (2017) Genome-wide association study for feed efficiency and growth traits in U.S. beef

cattle. *BMC Genomics* **18:**

Shaffer KS, Turk P, Wagner WR & Felton EED (2011) Residual feed intake, body composition, and fertility in yearling beef heifers. *J. Anim. Sci.* **89:** 1028–34 Available at: http://www.ncbi.nlm.nih.gov/pubmed/21112981 [Accessed March 2, 2016]

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B & Ideker T (2003) Cytoscape: A software Environment for integrated models of biomolecular interaction networks. *Genome Res.* **13:** 2498–2504

Shin JH, Li RW, Gao Y, Baldwin VI R & Li CJ (2012) Genome-wide ChIP-seq mapping and analysis reveal butyrate-induced acetylation of H3K9 and H3K27 correlated with transcription activity in bovine cells. *Funct. Integr. Genomics* **12:** 119–130

Shindo T, Kurihara H, Kurihara Y, Morita H & Yazaki Y (1998) Upregulation of endothelin-1 and adrenomedullin gene expression in the mouse endotoxin shock model. *J Cardiovasc Pharmacol* **31 Suppl 1:** S541-4

Shoji H, Minamino N, Kangawa K & Matsuo H (1995) Endotoxin markedly elevates plasma concentration and gene transcription of adrenomedullin in rat. *Biochem. Biophys. Res. Commun.* **215:** 531–537

Swami M (2009) Networking complex traits. *Nat. Rev. Genet.* **10:** 2566

Tronche F & Yaniv M (1992) HNF1, a homeoprotein member of the hepatic transcription regulatory network. *BioEssays* **14:** 579–587

Villar D, Berthelot C, Aldridge S, Rayner TF, Lukk M, Pignatelli M, Park TJ, Deaville R, Erichsen JT, Jasinska AJ, Turner JMA, Bertelsen MF, Murchison EP, Flicek P & Odom DT (2015) Enhancer evolution across 20 mammalian species. *Cell* **160:** 554–566

Vincent A, Louveau I, Gondret F, Tréfeu C, Gilbert H & Lefaucheur L (2015) Divergent selection for residual feed intake affects the transcriptomic and proteomic profiles of pig skeletal muscle. *J. Anim. Sci.* **93:** 2745–58 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26115262 [Accessed February 29, 2016]

Walker WH & Cheng J (2005) FSH and testosterone signaling in Sertoli cells. *Reproduction* **130:** 15–28

Walter LJ, Gasch CA, Mcevers TJ, Hutcheson JP, Defoor P, Marquess FLS & Lawrence TE (2014) Association of pro-melanin concentrating hormone genotype with beef carcass quality and yield. *J. Anim. Sci.* **92:** 325–331

Weber KL, Welly BT, Van Eenennaam AL, Young AE, Port-Neto LR, Reverter A & Rincon G (2016) Identification of Gene networks for residual feed intake in Angus cattle using genomic prediction and RNA-seq. *PLoS One* **11:** 1–19

Widmann P, Reverter A, Weikard R, Suhre K, Hammon HM, Albrecht E & Kuehn C (2015) Systems biology analysis merging phenotype, metabolomic and genomic data identifies Non-SMC Condensin I Complex, Subunit G (NCAPG) and cellular maintenance processes as major contributors to genetic variability in Bovine feed efficiency. *PLoS One* **10:** 1–22

Zarek CM, Lindholm-Perry AK, Kuehn LA & Freetly HC (2017) Differential expression of genes related to gain and intake in the liver of beef cattle. *BMC Res. Notes* **10:** 1 Available at: http://www.ncbi.nlm.nih.gov/pubmed/28057050 [Accessed March 27, 2017]

Figure 1. Genes selected for co-expression network construction. A) Heatmap of
normalized mean expression (NME) of 471 differentially expressed (DE) genes between high
(HFE) and low (LFE) feed efficient animals in adrenal gland (ADR), hypothalamus (HYP),
liver (LIV), muscle (MUS) and pituitary (PIT). Genes (rows) and samples (columns) are
organized by hierarchical clustering based on Euclidean distances. **B)** NME heatmap of all
1,317 genes selected for network construction. Genes (columns) and samples (rows) are
organized by hierarchical clustering based on Euclidean distances. **C)** Venn diagram of 1,317
genes selected for network construction. The inclusion criteria for selecting genes were
divided in five categories: differentially expressed genes (DE), tissue specific genes (TS),
genes harbouring SNPs reported by literature as being associated with feed efficiency in beef
cattle (SNP), genes encoding proteins secreted by at least one of the tissues in plasma (SEC)
and key regulators (REG). Numbers between brackets indicate the total number of genes in
each category.
1022

Figure 2. Gene co-expression network constructed using PCIT algorithm on 1,317
selected genes (see methods). Nodes with diamond shape correspond to secreted proteins
coding genes and triangles correspond to key regulators; all the other genes are represented
by ellipses. Nodes with black borders are differentially expressed between high and low feed
efficiency. Colours are relative to the tissue of maximum expression: blue represent liver, red
represent muscle, yellow represent pituitary, green represent hypothalamus and orange
represent adrenal gland. The size of the nodules is relative to the normalized mean expression
values in all samples. Only correlations above 0.9 and bellow -0.9 and its respective genes are
shown in this figure.
1032

Figure 3. Mapping of NR2F6 direct targets. A) Heatmap of the 313 genes coexpressed
with NR2F6 across all samples (derived from the co-expression analysis), B) i-cisTarget
motif discovery results on the genes shown in (A), C) Predicted NR2F6 targetome. A red
node indicates genes known to be targeted my NR2F6 in human Hepatocytes. D) Example of
predicted NR2F6 target regions for SERPINA1 gene. The predicted enhancer overlaps the
exact position for NR2F6 and NR2F2 binding in HepG sites from ENCODE dataset as well
as histone chromatin marks related with active regulatory regions, namely H3K27ac, and
promoters, H3K4me3 in human primary tissue from RoadMap Epigenetics E) The enhancer
prediction in cow coordinates (bosTau6) overlaps a region marked with H3K4me3 in cow
liver (Villar *et al*, 2015).
1043

Figure 4. Mapping the downstream network of TGFB signaling through
SMAD3/MyoD1 DNA binding. A) Heatmap of the 217 genes coexpressed with TGFB1
(derived from the co-expression analysis). B) i-cisTarget motif discovery results on the genes
shown in (A), C) Predicted MyoD targetome. A red node indicate genes know to be targeted
my MyoD1 in murine myotubes (Mullen *et al*, 2011). Blue nodes indicate genes to be
targeted by SMAD3, the effector DNA binding molecular of TGFB signalling, in murine
myotubes (Mullen *et al*, 2011). D) Example of predicted MyoD1 target regions for Acta1
gene. The predicted enhancer overlaps the exact position for SMAD3 and MyoD1 ChIP-seq
binding in murine myotubes (Mullen *et al*, 2011). E) The enhancer prediction in cow
coordinates (bosTau6) overlaps a promoter region marked with H3K4me3 in muscle tissue in
cow (Cao *et al*, 2009).
1055
1056

Table 1. Differentially connected and differentially expressed genes between high and low feed efficiency.

Gene name	Number of connections		Category*	Tissue of maximum expression	Tissue of differential expression
	Low feed efficiency	High feed effciency			
SST	0	45	DE, SEC	Hypothalamus	Hypothalamus
SNORA73	41	108	DE	Liver	Liver
ENSBTAG00000047700	56	111	DE	Liver	Liver
ENSBTAG00000047121	62	111	DE	Liver	Liver
ENSBTAG00000047816	53	96	DE	Liver	Liver
ENSBTAG00000039928	50	89	DE	Liver	Liver
ANXA13	115	63	DE	Liver	Liver
FST	113	56	DE	Liver	Liver
PBLD	115	55	DE	Liver	Liver
ENSBTAG00000021368	95	0	DE	Liver	Liver
JCHAIN	52	113	DE, SEC	Liver	Liver
IGFBP1	55	0	DE, TS, SEC	Liver	Liver
SBK2	0	70	DE	Muscle	Muscle
ACTC1	54	0	DE	Muscle	Muscle
MYH1	0	47	DE, TS	Muscle	Muscle
HR	119	50	DE	Pituitary	Muscle
TAGLN	83	31	DE	Adrenal	Muscle, Pituitary
SFRP2	41	91	DE	Hypothalamus	Pituitary
FN1	119	69	DE	Liver	Pituitary
CAV1	98	50	DE	Muscle	Pituitary

*Differentially expresses genes between high and low feed efficiency (DE), tissue specific genes (TS) and genes encoding proteins secreted in plasma (SEC).

S1 Supporting Information. Information of reads mapping to bovine genome (UMD3.1) per sample.

S2 Supporting Information. Scatter plots showing differentially expressed genes between high and low feed efficiency (FE) from adrenal gland, hypothalamus, liver, muscle and pituitary. Dots represent the mean expression for a gene in high FE subtracted from the mean expression of the same gene in low FE (M) by the average expression value in both groups (A). Pink dots represent significant genes (P<0.001).

S3 Supporting Information. Differentially expressed genes between high and low feed efficiency in adrenal gland (A), hypothalamus (B), liver (C), muscle (D) and pituitary (E).

S4 Supporting Information. Functional enrichment of the 248 genes down-regulated in high feed efficiency considering the five tissues (adrenal gland, hypothalamus, liver, muscle and pituitary). Colour intensity increase with the significance of the term; white represents $P>10^{-3}$ and the darkest orange represents $P<10^{-9}$.

S5 Supporting Information. Genes selected for network construction for being differentially expressed between high and low feed efficiency (A), harbouring SNPs previously associated with feed efficiency (B), tissue specific (C), coding proteins secreted in plasma (D) and key regulators (E).

S6 Supporting Information. Functional enrichment for the 135 genes coding proteins secreted in plasma which the tissue of maximum expression is liver. Colour intensity increase with the significance of the term; white represents $P>10^{-3}$ and the darkest orange represents $P<10^{-9}$.

S7 Supporting Information. Genes included in co-expression analysis.

S8 Supporting Information. Top five key regulator genes network and enrichment. A)
Network of genes *EPC1, NR2F6, MED21, ENSBTAG00000031687* and *CTBP1* and their
first neighbours. Nodes with diamond shape correspond to secreted proteins coding genes
and triangles correspond to key regulators; all the other genes are represented by ellipses.
Nodes with black borders are differentially expressed between high and low feed efficiency.
Colours are relative to the tissue of maximum expression: blue represent liver, red represent
muscle, yellow represent pituitary, green represent hypothalamus and orange represent
adrenal gland. The size of the nodules is relative to the normalized mean expression values in
all samples. Only correlations above 0.9 and bellow -0.9 and its respective genes are shown
in this figure. **B) Functional enrichment of the 345 genes in the network (A).** Colour
intensity increase with the significance of the term; white represents $P>5x10^{-3}$.

1104

S9 Supporting Information. *TGFB1* network and enrichment. A) Network of *TGFB1*
gene and its first neighbours. Nodes with diamond shape correspond to secreted proteins
coding genes and triangles correspond to key regulators; all the other genes are represented
by ellipses. Nodes with black borders are differentially expressed between high and low feed
efficiency. Colours are relative to the tissue of maximum expression: blue represent liver, red
represent muscle, yellow represent pituitary, green represent hypothalamus and orange
represent adrenal gland. The size of the nodules is relative to the normalized mean expression
values in all samples. Only correlations above 0.9 and bellow -0.9 and its respective genes are
shown in this figure. **B) Functional enrichment of the 157 genes in the network (A).**
Colour intensity increase with the significance of the term; white represents $P>5x10^{-3}$.

1115

S10 Supporting Information. *FGF21* network and enrichment. A) Network of *FGF21*
gene and its first neighbours. Nodes with diamond shape correspond to secreted proteins
coding genes and triangles correspond to key regulators; all the other genes are represented
by ellipses. Nodes with black borders are differentially expressed between high and low feed
efficiency. Colours are relative to the tissue of maximum expression: blue represent liver, red
represent muscle, yellow represent pituitary, green represent hypothalamus and orange
represent adrenal gland. The size of the nodules is relative to the normalized mean expression
values in all samples. Only correlations above 0.9 and bellow -0.9 and its respective genes are
shown in this figure. **B) Functional enrichment of the 98 genes in the network (A).** Colour
intensity increase with the significance of the term; white represents $P>5x10^{-3}$.

1126
1127 **S11 Supporting Information. NR2F6 i-cis Target results.**
1128
1129 **S12 Supporting Information. NR2F6 predicted regions binding hg19.**
1130
1131 **S13 Supporting Information. NR2F6 predcited regions bindg bosTau.**
1132
1133 **S14 Supporting Information. TGBF1 i-cis Target results.**
1134
1135 **S15 Supporting Information. MYOD predicted transcription factors binding sites in**
1136 **hg19.**
1137
1138 **S16 Supporting Information. MYOD predicted transcription factors binding sites in**
1139 **mm8.**
1140
1141 **S17 Supporting Information. MYOD predicted transcription factors binding sites in**
1142 **mm9.**

1143
1144 **S18 Supporting Information. MYOD predicted transcription factors binding sites in**
1145 **bosTau6.**
1146
1147 **S19 Supporting Information. Differentially connected genes between high and low feed**
1148 **efficiency.**
1149
1150 **S20 Supporting Information. Functional enrichment for the 87 differentially co-**
1151 **expressed genes between high and low feed efficiency.** Colour intensity increase with the
1152 significance of the term; white represents $P>10^{-3}$ and the darkest orange represents $P<10^{-9}$.
1153
1154

A

B

Cluster rank (PWM rank)	Associated TF	NES	# Targets	Logo
1 (1)	HNF1A/B	10.17	170	
2 (4)	NR2F6	7.98	168	
3 (16)	HNF4A/G	5.09	168	
4 (27)	CEBPA	4.25	68	

C

D

chr14: 94906801 - 9492122

SERPINA11

E

chr21: 59646366 - 59666543

SERPINA11

B

Cluster rank (PWM rank)	Associated TF	NES	# Targets	Logo
1 (1)	MEF2A/B/C/D	10.42	148	
2 (3)	SRF	8.12	128	
3 (34)	MyoD1	5.09	118	
4 (43)	TBP/POLRA	4.31	67	
5 (54)	Myog	4.07	119	

MyoD1 ChiP-seq in C2C12 murine cells Mullen *et al.* 2011

Smad3 ChiP-seq in C2C12 murine cells Mullen *et al.* 2011

D chr8: 126775857 - 126782286

Acta1

E chr28: 425010 - 431766

Acta1

Table 1. Differentially connected and differentially expressed genes between high and low feed efficiency.

Gene name	Number of connections		Category*	Tissue of maximum expression	Tissue of differential expression
	Low feed efficiency	High feed effciency			
SST	0	45	DE, SEC	Hypothalamus	Hypothalamus
SNORA73	41	108	DE	Liver	Liver
ENSBTAG00000047700	56	111	DE	Liver	Liver
ENSBTAG00000047121	62	111	DE	Liver	Liver
ENSBTAG00000047816	53	96	DE	Liver	Liver
ENSBTAG00000039928	50	89	DE	Liver	Liver
ANXA13	115	63	DE	Liver	Liver
FST	113	56	DE	Liver	Liver
PBLD	115	55	DE	Liver	Liver
ENSBTAG00000021368	95	0	DE	Liver	Liver
JCHAIN	52	113	DE, SEC	Liver	Liver
IGFBP1	55	0	DE, TS, SEC	Liver	Liver
SBK2	0	70	DE	Muscle	Muscle
ACTC1	54	0	DE	Muscle	Muscle
MYH1	0	47	DE, TS	Muscle	Muscle
HR	119	50	DE	Pituitary	Muscle
TAGLN	83	31	DE	Adrenal	Muscle, Pituitary
SFRP2	41	91	DE	Hypothalamus	Pituitary
FN1	119	69	DE	Liver	Pituitary
CAV1	98	50	DE	Muscle	Pituitary

*Differentially expresses genes between high and low feed efficiency (DE), tissue specific genes (TS) and genes encoding proteins secreted in plasma (SEC).