Genetic mapping with Allele Dosage Information in Tetraploid *Urochloa decumbens* (Stapf) R.D. Webster Reveals Insights into Spittlebug (*Notozulia entreriana* Berg) Resistance

Rebecca Caroline Ulbricht Ferreira[1], Letícia Aparecida de Castro Lara[2], Lucimara Chiari[3],

Sanzio Carvalho Lima Barrios[3], Cacilda Borges do Valle[3], José Raul Valério[3,] Fabrícia

Zimermann Vilela Torres[3], Antonio Augusto Franco Garcia[2], Anete Pereira de Souza[4*]

[1]Center for Molecular Biology and Genetic Engineering (CBMEG), University of Campinas

(UNICAMP), Campinas, SP, Brazil

[2]Genetics Department, ESALQ, University of São Paulo (USP), Piracicaba, SP, Brazil

[3]Embrapa Beef Cattle, Brazilian Agricultural Research Corporation, Campo Grande, MS, Brazil

[4]Plant Biology Department, Biology Institute, UNICAMP, Campinas, SP, Brazil

[¶]Both authors contributed equally to this work.

* Correspondence:

Anete Pereira de Souza

E-mail: anete@unicamp.br

Abstract

Urochloa decumbens (Stapf) R.D. Webster is one of the most important African forage grasses in

Brazilian beef production. Currently available genetic-genomic resources for this species are

restricted mainly due to polyploidy and apomixis. Therefore, crucial genomic-molecular studies such

as the construction of genetic maps and the mapping of quantitative trait loci (QTLs) are very

challenging and consequently affect the advancement of molecular breeding. The objectives of this

work were to (i) construct an integrated *U. decumbens* genetic map for a full-sibling progeny using

GBS-based markers with allele dosage information, (ii) detect QTLs for spittlebug (*Notozulia*

entreriana) resistance, and (iii) seek putative candidate genes involved in resistance/defense against

pathogens. We used the *Setaria viridis* genome as reference to align GBS reads and selected 4,240

high-quality SNPs markers with allele dosage information. Of these markers, 1,000 were distributed

throughout nine homologous groups with a cumulative map length of 1,335.09 cM and an average

marker density of 1.33 cM. We detected QTLs for resistance to spittlebug, an important pasture

insect pest, that explained between 4.66% and 6.24% of the phenotypic variation. These QTLs are in

regions containing putative candidate genes related to resistance/defense against pathogens. Because

this is the first genetic map with SNP autotetraploid dosage data and QTL detection in *U. decumbens*,

it will be useful for future evolutionary studies, genome assembly, and other QTL analyses in

Urochloa spp. Moreover, the results might facilitate the isolation of spittlebug-related candidate

genes and help clarify the mechanism of spittlebug resistance. These approaches will improve

selection efficiency and accuracy in *U. decumbens* molecular breeding and shorten the breeding

cycle.

Keywords: allele dosage, *Brachiaria*, linkage map, polyploidy, signalgrass, SNP, quantitative traits

36   1   Introduction

Brazilian cultivated pastures are the basis for the production of beef cattle, and they cover extensive
areas, most of which are populated by grasses of the genus *Urochloa* (syn. *Brachiaria*) (Jank et al.,
2011; Associação Brasileira das Indústrias Exportadoras de Carne (ABIEC), 2016). This genus
belongs to the Poaceae family and comprises species of plants distributed in tropical and subtropical
regions, mainly in the African continent (Renvoize et al., 1996; Valle et al., 2009).

One of the most widely cultivated species of genus is *Urochloa decumbens* (Stapf) R.D. Webster,
also known as signalgrass. This forage grass has relevant agronomic attributes, including exceptional
adaptation to poor and acidic soils that are typical of the tropics, leading to good animal performance
(Valle et al., 2010). However, the species is susceptible to several types of spittlebug, including
Notozulia entreriana Berg (Hemiptera: Cercopidae)*,* which is the most damaging pest for tropical
pastures (Gusmão et al., 2016).

Spittlebugs reduce biomass production, decrease palatability, and reduce the carrying capacity of
pastures (Valério and Nakano, 1988). Therefore, breeding programs aim to develop new cultivars
with reasonable or high resistance to spittlebugs to reduce forage loss. To achieve this advance, it is
essential to understand the genetic mechanisms involved in the spittlebug resistance response in
forage grass and to identify valuable markers for selection in breeding schemes. Currently, spittlebug
resistance in *Urochloa* grasses is thought to be under relatively simple genetic control and is highly
heritable (Miles et al., 1995). Therefore, despite being a trait with little genetic information,
spittlebug resistance can probably be easily manipulated in a plant breeding program.

Genetic breeding of *U. decumbens* is recent and has proven challenging because this grass is
predominantly tetraploid ($2n = 4x = 36$) and apomictic (Naumova et al., 1999; Valle et al., 2008). In
this scenario, the number of currently available molecular genetic resources is limited. Previous
molecular studies have increased knowledge of *U. decumbens* genetics, including the development of
sets of microsatellite markers (Ferreira et al., 2016; Souza et al., 2018) and the first transcriptome
(Salgado et al., 2017) of the species as well as linkage maps from the interspecific progeny of *U.
decumbens* x *U. ruziziensis* (Worthington et al., 2016). Other studies with molecular markers have
analyzed the genetic relationships of this species to other species in the *Urochloa* genus (Almeida et
al., 2011; Pessoa-Filho et al., 2017; Triviño et al., 2017).

Until recently, it was not possible to perform intraspecific crossings due to the ploidy barriers
between apomictic (polyploid) and sexual (diploid) accessions, but this restriction changed when
diploid accessions were artificially tetraploidized using colchicine (Simioni and Valle, 2009). This
advance allowed the generation of a base population and the exploration of a genetic variability
previously conserved by apomixis, favoring new molecular studies, including building specific
genetic maps to increase the genetic knowledge of the species.

Genetic maps are essential tools for genetics and genomics research, and they are becoming
increasingly common in polyploids species, mainly due to the development of next-generation-
sequencing (NGS) technologies and advances in genetic and statistical methods (Garcia et al., 2013).
In this study, we utilized the genotyping-by-sequencing (GBS) technique, which enables the
discovery of thousands of SNPs at low cost (Elshire et al., 2011; Poland et al., 2012). In addition,
robust software has been developed for polyploid organisms, including programs that can estimate
the allele dosage of SNPs and provide more genetic information to generate linkage maps with high

marker densities (Serang et al., 2012; Garcia et al., 2013; Mollinari and Serang, 2015; Bourke et al.,
2016, 2018).

The mapping process is more complex in polyploid species, mainly because of the larger number of
possible genotypes (Ripol et al., 1999; Vigna et al., 2016). Most polyploid genetic maps have been
limited to the use of single-dose markers (SDMs). In this case, either each marker is considered as a
single-allele copy from only one of the parents of the cross, with a 1:1 segregation ratio, or the SDMs
in both parents segregate in a 3:1 ratio (Wu et al., 1992). This method has been developed and
successfully adopted for various forage grass species, such as *Panicum maximum* (Ebina et al., 2005),
Paspalum notatum (Stein et al., 2007), *Panicum virgatum* (Okada et al., 2010) and *Urochloa*
humidicola (Vigna et al., 2016)*.* However, the use of only SDMs represents a loss of genetic
information, because in autotetraploid species, for example, five possible dosage classes exist:
nulliplex (aaaa), simplex (Aaaa), duplex (AAaa), triplex (AAAa), and quadruplex (AAAA). Thus,
many other marker segregations can be considered. Knowledge of the dosage of an SNP is essential
for genetic studies in polyploids species and can significantly increase the information imparted by
each locus (Voorrips et al., 2011; Garcia et al., 2013; Bourke et al., 2016, 2018).

High-resolution genetic linkage mapping for polyploid species can identify beneficial trait loci and
allow genomics-based breeding programs (Shirasawa et al., 2017). Molecular markers have been
widely used to locate quantitative trait loci (QTL) associated with quantitative resistance to insects in
many crop plants; however, genomic loci related to resistance to spittlebugs have not been
determined in any plant species. Particularly in *U. decumbens*, the identification of loci involved in
this trait is a promising tool for characterizing the genetic architecture and can assist in the design of
strategies to be introduced into breeding programs to increase the efficiency of the selection
processes and accelerate the release of new cultivars (Valle et al., 2009).

This study reports the development and application of GBS for mapping studies in an intraspecific
progeny of *U. decumbens*. To the best of our knowledge, there have been no reports of QTL mapping
in signalgrass. Our goals were to (i) build a GBS-based integrated genetic map using autotetraploid
allele dosage information in a bi-parental progeny, (ii) identify QTLs related to spittlebug resistance
on the integrated genetic map, and (iii) search for putative candidate genes that may be involved in
spittlebug resistance.

2   Materials and Methods

2.1   Plant Material and DNA Extraction

At the Brazilian Agricultural Research Corporation (Embrapa) Beef Cattle (EBC), Campo
Grande/MS, an intraspecific cross was made between *U. decumbens* D24/27 (sexual accession
tetraploidized by colchicine) and *U. decumbens* cv. Basilisk (tetraploid, apomictic cultivar used as
the pollen donor); therefore, both parents were tetraploid. The full-sib progeny of 239 F_1 individuals
was analyzed, from which 217 hybrids were identified using SSRs markers obtained from Ferreira et
al. (2016).

Leaf samples from each hybrid and both parents were collected, and genomic DNA was extracted
using the DNeasy 96 Plant Kit (Qiagen GMbH, Germany). DNA concentrations were determined
using a Qubit fluorometer (Invitrogen, Carlsbad, CA).

2.2   GBS Library Construction and Sequencing

The GBS library was prepared at the University of Campinas following the protocol described by
Elshire et al. (2011). Samples from both parents of the progeny were replicated five times for
sequencing. Each individual within a library was part of a 96-plex reaction. Each DNA sample (300
ng in a volume of 10 µL) was digested with the restriction enzyme NsiI (NEB) to reduce genomic
complexity and then ligated to a unique barcoded adapter plus a common adapter. The 96-plex
libraries were checked for quality using an Agilent DNA 1000 Kit on an Agilent 2100 Bioanalyzer
(Agilent Technologies, Santa Clara, CA, United States). Libraries sequencing was performed as 150-
bp single-end reads on the Illumina NextSeq 500 platform. The quality of the resulting sequence data
was then evaluated with the NGS QC Toolkit v2.3.3 (Patel and Jain, 2012).

2.3   SNP Calling

SNP discovery and genotype calling were performed using Tassel-GBS pipeline (Glaubitz et al.,
2014), which was modified to obtain the original count of the number of reads for each SNP allele
(Pereira et al., submitted). Because this pipeline requires a reference genome, and because the *U.*
decumbens genome has not yet been sequenced, we aligned the GBS tags using the *Setaria viridis*
genome (v 1.0; ~394.9 Mb arranged in 9 chromosomes and 724 scaffolds; diploid forage) as an
alternative pseudo-genome. This genome is available from the Phytozome website
(http://www.phytozome.net/) (Goodstein et al., 2012), and it sequence data were produced by the US
Department of Energy Joint Genome Institute. The Bowtie2 algorithm version 2.1 (Langmead and
Salzberg, 2012) was used to align tags against the pseudo-reference with -D and -R parameters
defined as 20 and 4, respectively, and with the very-sensitive-local argument.

2.4   Allele Dosage Estimation and Data Filtering

SuperMASSA software (Serang et al., 2012; Pereira et al., submitted) was used to estimate the allele
dosage (number of copies of each allele) of each individual. The minimum overall depth considered
was 25 reads, and the model used was the F_1 Population Model. Markers were fitted and filtered to
ploidy 4.

Additionally, monomorphic SNPs were removed, and markers with a maximum of 25% missing data
were selected manually using R software.

2.5   Linkage Map

An integrated linkage map was built using the TetraploidSNPMap (TPM, BioSS) software (Hackett
et al., 2017), following the methodology described in Hackett et al. (2013, 2014). Homologous
groups were constructed separately using the SNPs identified on each relative *S. viridis* chromosome.
Markers with significance of the χ^2 goodness-of-fit statistic less than 0.001 for simplex and 0.01 for
higher segregations were considered distorted and were removed. A total of 2,725 distorted SNPs
were excluded from the cluster analyses. The remaining 1,515 non-distorted SNPs were clustered
into 9 homologous groups using a Chi-square test for independent segregation (Hackett et al., 2013).

Considering the ordination of markers, a 2-point analysis was used to calculate the recombination
frequency and LOD score for each pair of SNPs in each possible phase by an expectation-
maximization (EM) algorithm to maximize the likelihood. Duplicate and near-duplicate SNPs were
removed in this step. The recombination frequencies were converted to map distances using
Haldane's mapping function, and a multi-dimensional scaling analysis (MDS) was then performed to

calculate the best order for the SNPs in the homology group. After this step, we excluded some SNPs
that showed low LOD scores or that were distant from the rest of the group (Hackett et al., 2017).

Some phases of the ordered SNPs were inferred by the automated phase analysis in
TetraploidSNPMap, and others were completed manually based on the most likely phase for previous
informative pairs prior to carrying out QTL analysis.

The genetic map was drawn using MapChart 2.32 (Voorrips, 2002), in which SNPs configurations
were identified with different colors.

2.6   Phenotypic Evaluation

The resistance of *U. decumbens* to spittlebugs (*N. entreriana*) was evaluated in greenhouse
experiments at Embrapa Beef Cattle, Brazil, according to the methodology described by Lapointe et
al. (1992) and Valério et al. (1997). From 2011 to 2013, a total of 12 experiments were conducted;
each experiment had a randomized complete block design (RCBD) with different numbers of blocks
and treatments. Common treatments were repeated in all experiments and were used as checks in the
statistical analysis. A total of 349 individuals were evaluated for spittlebug resistance, of which 157
were hybrids of the mapping population.

During 2011 and 2012, five RCBD experiments were conducted, with ten blocks and 114 individuals
evaluated. The best hybrids were selected with respect to spittlebug resistance and included in the
experiments conducted in 2012 and 2013, when seven RCBD experiments were conducted, with four
blocks and 259 individuals. The common checks for all experiments were *U. decumbens* cv. Basilisk,
U. brizantha cv. Marandu, *Urochloa* spp. cv. BRS Ipyporã, and *Urochloa* spp. MulatoII. The
variable evaluated was the percentage of nymphal survival (number of nymphs that reached the adult
stage, as a percentage). The following statistical linear mixed model was used for the analysis:

$$y_{ijlk} = \mu + b_{j(l)} + p_l + a_k + t_{ik} + \varepsilon_{ijlk}$$

where y_{ijlk} is the phenotype of the *i*-th individual, at *j*-th block, *l*-th experiment, and *k*-th year; μ is
the overall mean; $b_{j(l)}$ is the random effect of the *j*-th block within the *l*-th experiment
$(b_{j(l)} \sim N(0, \sigma_B^2))$; p_l is the random effect of the *l*-th experiment $(p_l \sim N(0, \sigma_P^2))$; a_k is the fixed
effect of the *k*-th year; t_{ik} is the random effect of the *i*-th individual in the *k*-th year $(t_{ik} \sim N(0, \boldsymbol{G}))$,
in which $\boldsymbol{G}$ is the variance-covariance (VCOV) matrix for genetic effects; and ε_{ijlk} is the residual
error $(\varepsilon_{ijlk} \sim N(0, \ \hat{\sigma}^2))$. The $\boldsymbol{G}$ matrix is indexed by two factors (genotype and year) written as the
Kronecker product of matrices, $\boldsymbol{G} = \boldsymbol{I_n} \otimes \boldsymbol{G_a}$, in which $\boldsymbol{I_n}$ is an identity matrix relative to genotype
effects and $\boldsymbol{G_a}$ is the VCOV matrix relative to year effects. The $\boldsymbol{G_a}$ matrix was evaluated considering
six different VCOV structures: independent (ID), diagonal (DIAG), compound symmetry (CS),
compound symmetry heterogeneous (CS-Het), first-order factor analytic (FA1), and unstructured
(US). The selection model was performed considering the Akaike Information Criteria (AIC)
(Akaike, 1974) and Schwarz Information Criteria (SIC) (Schwarz, 1978).

Phenotypic analysis was performed using the packages ASReml-R v3 (Butler et al., 2009; Gilmour et
al., 2009) and ASRemlPlus (Brien, 2016). Heritability between genotype means was estimated using
the index as presented by Cullis et al. (2006) and Piepho and Möhring (2007), where:

198
$$\hat{H}_C^2 = 1 - \frac{PEV}{2\sigma_G^2}$$

199   in which PEV is the prediction error variance, and σ_G^2 is the genetic variance.

2.7   QTL Mapping

QTL mapping for spittlebug resistance was performed by applying an Interval Mapping (IM) model
based on autotetraploid allele dosage information with TetraploidSNPMap software (Hackett et al.,
2017). This analysis was performed separately for each homology group using the SNP data for each
homology group, the integrated genetic map with phase information and the phenotypic trait data.
The interval mapping was fit to a model of additive effects using a weighted regression approach. A
significant QTL was declared if the LOD score was above the 95% threshold obtained from a
permutation test with 1,000 permutations. From a significant QTL, simple models such as a simplex,
duplex or double-simplex QTL models were then tested using the SIC (Schwarz, 1978) to determine
if the estimated genotype corresponded to the most likely QTL location. The SIC is calculated in
TetraploidSNPMap as follows:

SIC= -2 log L + p log m_o,

where L is the likelihood for the simple model, p is the number of parameters in the simple model,
and m_o is the number of observations (the 36 genotype means). Models with the lowest value for the
SIC were considered the best models (Hackett et al., 2014).

If two closely linked significant QTLs were identified on the same chromosome, TetraploidSNPMap
could not estimate the individual QTL LOD and parameter values, yielding data only pertaining to
the largest effect.

2.8   Search for Candidate Genes in Detected QTL Regions

We investigated the functional annotation of the candidate genes located close to the QTL regions
associated with spittlebug resistance. Using the chromosomal locations of the markers adjacent to
QTLs in the *S. viridis* genome as a reference, we aligned the sequences found in these regions (100
kb approximately) with a BLASTX search (e-value cutoff of 1e-0.5) against plants databases using
the JBrowse tool (http://www.phytozome.org/). Finally, from this alignment, we selected only genes
reported in the literature as associated with resistance/defense against pathogens that were present in
the transcriptome of *U. decumbens* (Salgado et al., 2017), our species of study.

3   Results

3.1   SNP Calling

The GBS library generated a total of 1,183,089,925 raw reads (94% Q20 bases), with an average of
4,151,193 reads per individual. Using the Tassel-GBS pipeline modified to polyploid, alignment with
the *S. viridis* genome enabled the identification of 58,370 SNPs markers. The output of the Tassel-
GBS pipeline was used as the input for the SuperMASSA software, and 9,345 SNPs markers with
allele dosage were selected with a minimum overall depth of 25 reads. Afterward, markers with up to
25% missing data and monomorphic markers were removed, resulting in 4,240 SNPs markers that
were used in the subsequent steps.

Using three different SNPs markers, Figure 1 shows how the SuperMASSA software uses the ratio of
allele count to classify individuals according to their genotypes using a probabilistic graphical model
(Serang et al., 2012). The markers were named with the allele dosage segregation, the homology
group number, and the chromosomal position of the SNP in *S. viridis* genome.

GBS sequence data has been submitted to the NCBI Sequence Read Archive (SRA) under accession
number SRP092493.

3.2   Genetic Map

An integrated genetic map was built using 217 full sibs obtained from a cross between *U. decumbens*
cv. Basilisk (apomictic cultivar) and *U. decumbens* D24/27 (sexual accession). Of the 4,240 markers
used for linkage analysis, 1,515 showed significance of the χ^2 goodness-of-fit statistic, and 1,000
were placed in the linkage map (Table 1).

The markers were distributed throughout nine homology groups (HGs), with a cumulative map
length of 1,335.09 cM and an average marker density of 1.33 cM. The length of each group ranged
from 129.70 cM (HG7) to 166.57 cM (HG9), with an average of 148 cM (Table 1). Homology group
9 showed the highest average density and the highest number of mapped markers (Table 1; Figure 3).

The distances between adjacent markers on the genetic map were plotted as a histogram, which
revealed that the majority of markers were within 1 cM of each other (Figure 2). The two largest
distances between adjacent markers were located in HG8 and had lengths of 11.2 cM and 12.4 cM.

In Figure 3, the possible genotype configuration of each marker is indicated by the marker name
prefix and the different colors. For example, markers with the prefix S and the color black represent
the simplex configuration (Table 2). Four homologous groups showed almost all configurations
(HG1, HG3, HG5 and HG9), and simplex markers (S) were the most frequent. The X-double-simplex
(XSS) configuration was less frequent and appeared in only HG3 and HG5 (Table 2; Figure 3).

3.3   Phenotypic Analysis and QTL Mapping

The selected VCOV matrix for $\boldsymbol{G_a}$ based on AIC was CS-Het and/or US, in which both account for
genetic correlation and heterogeneous variance across years. Considering the SIC, the selected
VCOV matrix for $\boldsymbol{G_a}$ was CS, in which the genetic correlation and homogeneous variance across
years are considered, in addition to requiring estimation of a lower number of parameters than the
previous VCOV matrices. As the difference in SIC values between CS and CS-Het/US (18880.37 −
18877.39 = 2.98) was greater than the respective AIC difference (18848.03 − 18845.13 = 2.9), the CS
matrix was used for $\boldsymbol{G_a}$ (Table 3).

Predicted genotypic values for spittlebug (*Notozulia entreriana*) resistance in hybrids of the mapping
population, genetic and residual variance components and trait heritability are shown in Table 4. The
assumption of normality of the residuals can be visualized in Supplementary Figure 1.

Best linear unbiased prediction values were used to perform QTL mapping for spittlebug resistance
by applying an IM model to the integrated genetic map. Using this method, three significant QTLs
were identified for spittlebug resistance (Table 5) that were located in three homologous groups:
HG1, HG2 and HG6. The obtained threshold values for LOD scores using 1,000 permutations ranged
from 4.39 to 5.49 and were used to declare the presence of these QTLs. The percentages of
phenotypic variation (R^2) explained by the QTLs ranged from 4.66% to 6.24%. The male progenitor

of the mapping population, *U. decumbens* D62 (cv. Basilisk), contributed with the resistance alleles
for these three QTLs

In homology group 1, the QTL had a LOD score of 4.68 and explained 4.80% of the phenotypic
variation for spittlebug resistance. This LOD score was above the upper 95% LOD permutations
threshold of 3.11. This first QTL was located at position 95 cM (Figure 3). Analyses of different
simple genetic models were performed with TetraploidSNPMap to determine the best simple fitting
model for the studied trait. For this QTL, the best model was a duplex genotype (AAAA x BBAA),
with the allele B present on parent 2 associated with spittlebug resistance. This model had the lowest
SIC (SIC = 85.34) compared with the full model (SIC = 102.99) (Table 5).

In homology group 2, the maximum LOD score was 4.39, and this QTL explained 4.66% of the
phenotypic variation for spittlebug resistance. The LOD peak was located at position 60 cM (Figure
3), and its score was above the upper 95% LOD permutation threshold of 3.29 (Supplementary
Figure 2). Again, the analysis with the simpler models estimated a duplex genotype (AAAA x
BBAA) for this QTL, with the resistance allele present in parent 2. This model had the lowest SIC
(SIC = 93.03) in comparison with the full model (SIC = 89.55) (Table 5).

The last detected QTL, in homology group 6, had the highest LOD score, 5.49, explained 6.24% of
the phenotypic variation and was above the 95% LOD permutation upper threshold of 2.79
(Supplementary Figure 2). The QTL peak was located at 23 cM (Figure 3), and an analysis of
different genetic models indicated that an allele explains the phenotypic variation. This QTL is linked
to a simplex SNP (AAAA x ABAA), with the B allele present on parent 2 associated with spittlebug
resistance. This model had the lowest SIC of 99.07 compared to the full model with SIC of 115.47
(Table 5).

3.4   Search for Candidate Resistance Genes

Sequence similarity was detected in the regions of the markers adjacent to the mapped QTLs, with
homologies for *Arabidopsis thaliana* and *Oryza sativa.* Functional annotation of these sequences
showed possible candidate genes involved in the spittlebug resistance response (Supplementary
Table 1) that are also present in the *U. decumbens* transcriptome (Salgado et al., 2017).

The first identified QTL, located in HG1, had two adjacent markers, D.1_40431816 and
S.1_21570712, identified from the *S. viridis* genome reference. The region of this QTL showed
similarity to pathogenesis-related gene 1 (*PR1*), leucine-rich repeat (LRR) family protein and the
WRKY11 gene, which are involved in pathogen defense (Supplementary Table 1).

In HG2, the region of the GBS-based adjacent markers (DS.2_39902899 and S.2_399022936) to
QTL2 showed similarity to NBS-LRR and NB-ARC protein domains and with the pentatricopeptide
repeat (PPR) superfamily proteins (Supplementary Table 1).

The region of the GBS-based markers (D.6_25761630 and S.6_35357677) adjacent to QTL3 in the
HG6 showed similarity to a WRKY DNA-binding protein, a kinase superfamily protein, an F-box
family protein, NB-ARC protein domains and kinase superfamily proteins. This QTL region also
showed similarity to the thaumatin-like protein superfamily (TLP), which is classified within the
pathogenesis-related (PR) protein family, to LRR protein domains and to F-box family proteins
(Supplementary Table 1).

4   Discussion

The development of GBS technology enables the identification of thousands of SNP markers across
the genome at relatively low cost. This advance represents an evolution in genetic studies of
polyploids species, since a larger part of the genome can be represented, favoring approaches such as
the construction of genetic maps. The map reported here was developed using an intraspecific
progeny of *U. decumbens* of Embrapa Beef Cattle and includes GBS-based markers with
autotetraploid allele dosages, an approach never previously employed in a forage grass species.
Although not completely elucidated, some evidence suggests that *U. decumbens* is a segmental
allopolyploid (Mendes-Bonato et al., 2001; Worthington et al., 2016), considered to have arisen from
hybridization between very closely related species (Stebbins, 1947).

Our molecular markers fit a tetrasomic segregation model (Supplementary Figure 3), probably
because autotetraploids as well as segmental allotetraploids, do not behave as diploids with respect to
the mode of inheritance (Bourke et al., 2018). Another explanation is the fact that one of the parents
of the mapping population was tetraploidized artificially and therefore this parent behaves as
autotetraploid. We choose to use all possible SNP configurations to add even more genetic
information to genetic map. Our map provides important genomic information, such as the detection
of QTLs involved in spittlebug (*N. entreriana* Berg) resistance, a relevant trait for breeding programs
in the *Urochloa* genus.

We used the genome of *S. viridis*, a member of the Poaceae, or grass, family, to align the tags and
discover GBS-based markers. This alignment was sufficient to identify a relevant number (4,240) of
high-quality SNPs, with allele dosage estimated. In previous analyses, we tested other reference
genomes for tag alignment, but the *S. viridis* genome alignment produced the best results in terms of
the number of markers identified (Supplementary Table 2). The sequences obtained herein by the
GBS libraries are dynamic, allowing the raw data to be reanalyzed as bioinformatics techniques
evolve (Poland and Rife, 2012), as well as when some tetraploid *Urochloa* spp. genome is
sequenced. Therefore, our sequences may be reanalyzed in the future to obtain a higher alignment
rate, as well as to identify a larger number of markers.

Worthington et al. (2016) detected 3,912 SNPs markers after filtering using the UNEAK (Universal
Network-Enabled Analysis Kit) pipeline (Lu et al., 2013), which is similar to the number of markers
that we detected using the *S. viridis* genome as reference. In a previous analysis, we tested the
UNEAK pipeline for SNP identification in our intraspecific progeny, but this method detected fewer
markers (1,210) compared with alignment to the genome reference (4,240). Most likely, the cross
between two different species (*U. decumbens* x *U. ruziziensis*) favored the discovery of more SNPs
through the UNEAK pipeline by Worthington et al. (2016), in addition to the different methodology
used for GBS library construction.

Previous polyploid genetic studies have been developed with markers scored as presence/absence of
alleles, but this method does not make use of all the information available in the SNP at multiple
doses. The GBS data obtained here were sufficient in terms of read depth to call allele dosage in
9,345 markers from the *S. viridis* genome using SuperMASSA software (Serang et al., 2012; Pereira
et al., submitted) (Supplementary Table 2). This methodology provided the distribution of alleles in
the progeny and the relative intensities of each allele, increasing the amount of genetic information
obtained (Serang et al., 2012; Garcia et al., 2013; Pereira et al., submitted).

After analysis using SuperMASSA software (Serang et al., 2012) and subsequent filtering steps, were
obtained 4,240 SNPs that represented all possible configurations for an autotetraploid species.
Approximately 55% (2,345) of the SNPs markers identified from the *S. viridis* genome alignment

361 followed the parental genotype configurations of simplex (AAAA x AAAB/ABBB x BBBB) and
362 double-simplex (AAAB x AAAB/ABBB x ABBB). Slightly less than 1% of the identified markers
363 followed the nulliplex (AAAA x BBBB) and simplex-duplex (AAAB x AABB) configurations
364 (Table 2). These results were satisfactory and relevant for genetic studies with a tetraploid species,
365 but a GBS library construction methodology with more than one restriction enzyme might provide a
366 larger read depth and, accordingly, increase the accuracy of allele dosage estimation and the number
367 of markers detected.

368 The 4,240 high-quality SNPs markers were inserted in TetraploidSNPMap software (Hackett et al.,
369 2017), but only 1,515 showed significance in the χ^2 goodness-of-fit statistic, and 1,000 were placed
370 in the linkage map (Table 1). These 1,000 SNPs markers were distributed across nine homology
371 groups, with a total length of 1,335.09 cM and an average map density of 1.33 cM (Table 1; Figure
372 3), reflecting the molecular source and chosen mapping methodology. Comparatively, Thaikua et al.
373 (2016) and Vigna et al. (2016) presented the largest maps, with 1,702.82 cM and 1,423.2 cM of total
374 size, respectively. Although small, this length difference may be due to the smaller number of non-
375 NGS markers used (89 and 64, respectively) and the exclusive usage of SDMs for mapping. In
376 addition to having a smaller length, our genetic map has a higher marker density than these others,
377 reflecting the ability of the GBS methodology to allow the simultaneous detection of thousands of
378 markers.

379 In the Worthington et al. (2016) study, genetic maps for an interspecific progeny resulting from a
380 cross between *U. decumbens* x *U. ruziziensis* was built with GBS-based markers, and 1,916 SDA
381 (single-dose allele) markers were distributed across the parental linkage maps. The different mapping
382 method used in our study may have influenced the number of markers linked in the map. However,
383 although our map has a lower marker density, it contain SNP markers with different allele dosage,
384 adding more genetic information to the map and enabling more robust genetic analyses (Garcia et al.,
385 2013).

386 In our study, the density of markers per HG ranged from 63 (HG6) to 187 (HG9) (Table 1). The
387 greater density of markers for certain HGs may correspond to a greater recombination frequency. On
388 the other hand, less saturated groups may have fewer SNP in these regions and/or correspond to
389 highly homozygous regions that have lower recombination frequencies (Bai et al., 2016).

390 The largest distance between adjacent markers, 12.4 cM, was observed in homology group eight
391 (Figure 3), one of the groups with the fewest mapped markers. Compared with other linkage map
392 studies, our map has low average marker intervals in all HGs (Figures 2 and 3). Nevertheless,
393 substantial intervals are common and even expected using the GBS technique, mainly due to
394 centromeric regions, which are not reached with this methodology (Conson et al., 2018).
395 Additionally, it is possible that the number of hybrids in the mapping population was not sufficient to
396 observe recombination in these intervals (Ma et al., 2014) or that these intervals represent regions
397 that were not captured with the methodology used. One possible method to fill these distances is to
398 use different combinations of enzymes to enlarge the sequencing pools and thereby enable the
399 capture of markers in other genomic regions. Furthermore, we mapped only markers identified by
400 aligning with the *S. viridis* genome. Therefore, these observed intervals may contain markers that are
401 exclusive to the *U. decumbens* genome and thus cannot be represented with the methodology used.
402 Therefore, advances in the assembly of complex polyploid genomes are expected to enable the use of
403 the full signalgrass genome as a reference for future studies (Margarido and Heckerman, 2015).

Of the 1000 mapped SNPs, 59% (591) followed the simplex parental genotype configuration (Table
2; Figure 3). This configuration was also the most common in other genetic mapping studies (Hackett
et al., 2013; da Silva et al., 2017), probably because it is the most common in plant genomes and is
easily scored. The least common configuration was X-double-simplex (XSS), followed by double-
duplex (DD) (Table 2), which is the most complete configuration because it allows the representation
of all types of segregation for autotetraploid species. Overall, the map displays most of the possible
SNP configurations, and these data significantly increase the information about each locus and
provide several advantages for genetic analysis (Garcia et al., 2013). When building genetic maps for
polyploid species, the use of markers with allele dosage information allows the development of
robust genetic maps that allow the detection of QTLs (Hackett et al., 2013; da Silva et al., 2017).

Until recently, no intraspecific progeny have been available for *U. decumbens*; therefore, no QTL
mapping studies of this species have been performed to date. The identification of genes/loci
underlying spittlebug (*N. entreriana*) resistance is critical for forage grass molecular breeding
programs because these pests severely damage pastures (Valle et al., 2009). In our study, QTL
mapping for spittlebug resistance was performed by applying the statistical model IM for the
tetraploid progeny (Hackett et al., 2017). Although more accurate models are available, these models
have not been extended to polyploidy and therefore do not use molecular markers with allele dosage
information.

For the multi-allelic QTL analysis, we performed a permutation test to obtain the threshold for
declaring significant QTLs. This analysis allowed the identification of three significant QTLs which
explained between 4.66% and 6.24% of the phenotypic variation (Table 5). The identification of a
small number of QTLs with small effects reflects the median heritability found for spittlebug
resistance ($H_C^2 = 0.37$). Spittlebug resistance in forage grasses is probably not genetically complex
but likely involves more than a single major resistance gene (Miles et al., 1995). Therefore, values of
phenotypic variation and the number of QTLs detected in our analysis may be mainly due the low
genetic variability for spittlebug resistance in the analyzed progeny, since the parents do not differ
greatly for this trait. Moreover, the methodology used for QTL detection and/or map saturation may
have influenced these results.

Although *U. decumbens* is considered a susceptible species to the pasture spittlebug, our results
showed that the segregation of different alleles in each tetraploid parent contributed to the observed
polymorphisms in the progeny. Therefore, our results can be useful for breeders to identify the alleles
from the male parent that contribute to increased values of this phenotypic trait, consequently
allowing genotypes that are more resistant to the development and survival of spittlebugs to be found
more rapidly.

QTLs with greater effects on spittlebug resistance are necessary and more efficient for use in
breeding programs, but our results represent a great gain as the only genetic architecture study using
an *U. decumbens* intraspecific progeny. Therefore, the detection of these QTLs is the first step in the
advancement of genomic studies involving spittlebug resistance. The annotation of sequences that
originated markers with mapped QTLs is important to identify candidate genes involved in spittlebug
resistance in *U. decumbens*. Thus, the regions that gave rise to these QTLs should be more
thoroughly evaluated to identify more genetic information for future applications in the marker-
assisted selection in signalgrass breeding programs.

N. entreriana spittlebug nymphs constantly suck the sap of roots, causing yellowing of the plant, and
the saliva of the adults induces phytotoxemia that causes plant death (Valério and Nakano, 1992;

448 Gusmão et al., 2016). Thus, to respond mechanical damage due to these insect attacks, plants possess
449 multiple molecular defense mechanisms (War et al., 2012). Based on an investigative and
450 comparative analysis, the QTL regions of this study showed similarity to genes containing protein
451 domains reported to be involved in disease resistance and defense responses to pathogens that attack
452 plants (Supplementary Table 1).

453 For spittlebug resistance trait, we can highlight the similarity of the region of the QTL1 with
454 pathogenesis-related gene 1 (*PR1*). PR proteins are induced in response to attack by various types of
455 pathogens, being that the *PR1* gene is involved in biological processes related to defense responses
456 and systemic acquired resistance (Metzler et al., 1991; Uknes et al., 1992; Van Loon and Van Strien,
457 1999; Van Loon et al., 2006; Ali et al., 2018). This region also showed similarity to the *WRKY11*
458 gene, which acts as a negative regulator of basal resistance to bacterial pathogens (Journot-Catalino
459 et al., 2006) and plays a crucial role in cellular defense responses and defense-related gene
460 transcriptional regulation (Jiang et al., 2016) (Supplementary Table 1).

461 For QTL3, we can highlight the similarity of the region to the adjacent markers with WRKY DNA
462 binding domains. These transcription factors are involved in plant defense responses (Du and Chen,
463 2000). Moreover, F-box and WRKY DNA binding domains have orthologs in *A. thaliana*, with genes
464 that develop a function in plant defense. This QTL region also showed similarity to TLPs, which play
465 a variety of roles including plant protection against pathogen attacks (Liu et al., 2010; Mukherjee et
466 al., 2010; Misra et al., 2016).

467 The regions of the markers adjacent to the mapped QTLs 2 and 3 showed similarity to NBS-LRR
468 protein domains, which are widely distributed in plants and thought to respond to pathogen attacks,
469 including viruses, bacteria, fungi, and even insects (Song et al., 2017). NBS-LRRs are the most
470 prevalent class of resistance proteins (Gupta et al., 2012). F-box proteins form one of the largest
471 multigene superfamilies and control many important biological functions such as pathogen resistance
472 (Lechner et al., 2006), and the PPR superfamily proteins (Barkan and Small, 2014) have been
473 implicated in plant defense mechanisms. Moreover, the NB-ARC protein domain is a functional
474 ATPase domain containing a nucleotide-binding site that is proposed to regulate plant disease
475 resistance (Van Ooijen et al., 2008).

476 Although we used the *S. viridis* genome as a reference to investigate the marker regions adjacent to
477 the identified QTLs, we selected only genes found in the *U. decumbens* transcriptome (Salgado et al.,
478 2017) to lend greater reliability to our results. Therefore, our results provide a starting point,
479 indicating possible candidate genes involved in spittlebug resistance in signalgrass. However further
480 studies should be conducted to validate these genomic regions as well as the role of the candidate
481 genes in signalgrass and their effects on phenotypic expression.

482 We can conclude that this recent opportunity to analyze a *U. decumbens* full-sib progeny has opened
483 new genetic molecular perspectives for this and related species. The genetic map presented herein
484 includes important approaches such as the estimation of tetraploid dosage of each molecular SNP and
485 greatly increases genetic information and represents an important evolution for polyploid studies
486 (Bourke et al., 2016). In addition, the map allowed the identification of genomic regions related to
487 spittlebug (*N. entreriana*) resistance, the main insect that attacks forage grasses, providing new
488 insights about this trait. Furthermore, the molecular data developed here are dynamic and can be
489 applied in future studies, such as genome assembly and other QTL analyses in *U. decumbens*.
490 Therefore, our results are the first step towards possible marker-assisted selection (MAS) and

491 genomic selection (GS), a subject of great interest in genetic improvement programs for
492 economically important plant species.

Conflict of Interest

495 The authors declare that the research was conducted in the absence of any commercial or financial
496 relationships that could be construed as a potential conflict of interest.

Author Contributions

498 LC, SB, CV, JV, FZ and AS conceived and designed the experiments. RF, LC, SB, CV, JV and FZ
499 conducted the experiments and collected data. RF, LL and AG analyzed the data. RF, LL and AS
500 wrote the manuscript. All authors read and approved the manuscript.

Funding

This work was supported by grants from the Fundação de Amparo à Pesquisa de do Estado de São
Paulo (FAPESP 08/52197-4) and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
(CAPES - Computational Biology Programme). RCUF received PhD fellowships from the CAPES-
Embrapa Programme and the CAPES Computational Biology Programme, and LACL received a PD
fellowship from the CAPES Computational Biology Programme. AAFG and APS were recipients of
a research fellowship from Conselho Nacional de Desenvolvimento Científico e Tecnológico
(CNPq).

Acknowledgments

510 We would like to acknowledge the Brazilian Agricultural Research Corporation (Embrapa Beef
511 Cattle) for providing the *Urochloa decumbens* progeny used in this study. We thank the Aline da
512 Costa Lima Moraes for the assistance in the GBS library construction and performing DNA
513 sequencing.

References

Akaike, H. (1974). A new look at the statistical model identification. IEEE Trans. Automat. Contr.
19, 716-723.

Ali, S., Mir, Z.A., Bhat, J.A., Tyagi, A., Chandrashekar, N., Yadav, P., et al. (2018). Isolation and
characterization of systemic acquired resistance marker gene PR1 and its promoter from
Brassica juncea. 3 Biotech 8, 10. doi: 10.1007/s13205-017-1027-8

Almeida, M.C.D.C., Chiari, L., Jank, L., and Valle, C.B.D. (2011). Diversidade genética molecular
entre cultivares e híbridos de *Brachiaria* spp. e *Panicum maximum*. Ciênc. Rural 41, 1998-
2003.

Associação Brasileira das Indústrias Exportadoras de Carne (ABIEC) (2016). *Perfil da Pecuária do*
Brasil: Relatório Anual. Available at: http://abiec.siteoficial.ws/images/upload/sumario-pt-
010217.pdf

Bai, Z.-Y., Han, X.-K., Liu, X.-J., Li, Q.-Q., and Li, J.-L. (2016). Construction of a high-density
genetic map and QTL mapping for pearl quality-related traits in *Hyriopsis cumingii*. Sci. Rep.
6, 32608. doi: 10.1038/srep32608

Barkan, A., and Small, I. (2014). Pentatricopeptide repeat proteins in plants. Annu. Rev. Plant Biol.
65, 415-442. doi: 10.1146/annurev-arplant-050213-040159

Bourke, P.M., Voorrips, R.E., Kranenburg, T., Jansen, J., Visser, R.G.F., and Maliepaard, C. (2016).
Integrating haplotype-specific linkage maps in tetraploid species using SNP markers. Theor.
Appl. Genet. 129, 2211-2226. doi: 10.1007/s00122-016-2768-1

Bourke, P.M., Voorrips, R.E., Visser, R.G.F., and Maliepaard, C. (2018). Tools for genetic studies in
experimental populations of polyploids. Front. Plant Sci. 9, 513. doi:10.3389/fpls.2018.00513

Brien, C. (2016). ASRemlPlus: augments the use of 'ASReml' in fitting mixed models. R package
version 2.0–9.

Butler, D.G., Cullis, B.R., Gilmour, A.R., and Gogel, B.J. (2009). ASReml-R reference manual,
release 3. Technical report. Brisbane, QLD: Queensland Department of Primary Industries.

Conson, A.R.O., Taniguti, C.H., Amadeu, R.R., Andreotti, I.A.A., de Souza, L.M., Dos Santos,
L.H.B., et al. (2018). High-resolution genetic map and QTL analysis of growth-related traits
of *Hevea brasiliensis* cultivated under suboptimal temperature and humidity conditions.
Front. Plant Sci. 9, 1255. doi: 10.3389/fpls.2018.00513

Cullis, B.R., Smith, A.B., and Coombes, N.E. (2006). On the design of early generation variety trials
with correlated data. J. Agric. Biol. Environ. Stat. 11, 381-393. doi:
10.1198/108571106X154443

Da Silva, W.L., Ingram, J., Hackett, C.A., Coombs, J.J., Douches, D., Bryan, G.J., et al. (2017).
Mapping loci that control tuber and foliar symptoms caused by PVY in Autotetraploid potato
(*Solanum tuberosum* L.). G3 7, 3587-3595. doi: 10.1534/g3.117.300264

Du, L., and Chen, Z. (2000). Identification of genes encoding receptor-like protein kinases as
possible targets of pathogen- and salicylic acid-induced WRKY DNA-binding proteins in
Arabidopsis. Plant J. 24, 837-847. doi: 10.1046/j.1365-313x.2000.00923.x
553
554 Ebina, M., Nakagawa, H., Yamamoto, T., Araya, H., Tsuruta, S.-i., Takahara, M., et al. (2005). Co-
555   segregation of AFLP and RAPD markers to apospory in Guineagrass (*Panicum maximum*
556   Jacq.). Grassl. Sci. 51, 71-78. doi: 10.1111/j.1744-697X.2005.00011.x
557
558 Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., et al. (2011). A
559   robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS
560   One 6, e19379. doi: 10.1371/journal.pone.0019379
561
Ferreira, R.C.U., Cançado, L.J., Do Valle, C.B., Chiari, L., and De Souza, A.P. (2016). Microsatellite
loci for *Urochloa decumbens* (Stapf) R.D. Webster and cross-amplification in other *Urochloa*
species. BMC Res. Notes 9, 152. doi: 10.1186/s13104-016-1967-9

Garcia, A.A.F., Mollinari, M., Marconi, T.G., Serang, O.R., Silva, R.R., Vieira, M.L.C., et al. (2013).
SNP genotyping allows an in-depth characterisation of the genome of sugarcane and other
complex autopolyploids. Sci. Rep. 3, 3399. doi: 10.1038/srep03399

Gilmour, A.R., Gogel, B.J., Cullis, B.R., and Thompson, R. (2009). ASReml User Guide Release 3.0.
Hempstead, NY: VSN Int. Ltd.

Glaubitz, J.C., Casstevens, T.M., Lu, F., Harriman, J., Elshire, R.J., Sun, Q., et al. (2014). TASSEL-
GBS: a high capacity genotyping by sequencing analysis pipeline. PLoS One 9, e90346. doi:
10.1371/journal.pone.0090346

Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., et al. (2012). Phytozome:
a comparative platform for green plant genomics. Nucl. Acids Res. 40, D1178-D1186. doi:
10.1093/nar/gkr944

Gupta, S.K., Rai, A.K., Kanwar, S.S., and Sharma, T.R. (2012). Comparative analysis of zinc finger
proteins involved in plant disease resistance. PLoS One 7, e42578. doi:
10.1371/journal.pone.0042578

Gusmão, M.R., Valério, J.R., Matta, F.P., Souza, F.H.D., Vigna, B.B.Z., Fávero, A.P., et al. (2016).
Warm-season (C4) turfgrass genotypes resistant to spittlebugs (Hemiptera: Cercopidae). J.
Econ. Entomol. 109, 1914-1921. doi: 10.1093/jee/tow135

Hackett, C.A., Boskamp, B., Vogogias, A., Preedy, K.F., and Milne, I. (2017). TetraploidSNPMap:
software for linkage analysis and QTL mapping in autotetraploid populations using SNP
dosage data. J. Hered. 108, 438-442. doi: 10.1093/jhered/esx022

Hackett, C.A., Bradshaw, J.E., and Bryan, G.J. (2014). QTL mapping in autotetraploids using SNP
dosage information. Theor. Appl. Genet. 127, 1885-1904. doi: 10.1007/s00122-014-2347-2

Hackett, C.A., McLean, K., and Bryan, G.J. (2013). Linkage analysis and QTL mapping using SNP
dosage data in a tetraploid potato mapping population. PLoS One 8, e63939. doi:
10.1371/journal.pone.0063939

Heffelfinger, C., Fragoso, C.A., Moreno, M.A., Overton, J.D., Mottinger, J.P., Zhao, H., et al. (2014).
Flexible and scalable genotyping-by-sequencing strategies for population studies. BMC
Genom. 15, 979. doi: 10.1186/1471-2164-15-979

Jank, L., Valle, C.B., and Resende, R.M.S. (2011). Breeding tropical forages. Crop Breed. Appl.
Biotechnol. 11, 27-34. doi: 10.1590/S1984-70332011000500005

Jiang, C.-H., Huang, Z.-Y., Xie, P., Gu, C., Li, K., Wang, D.-C., et al. (2016). Transcription factors
WRKY70 and WRKY11 served as regulators in rhizobacterium *Bacillus cereus* AR156-
induced systemic resistance to *Pseudomonas syringae* pv.tomato DC3000 in *Arabidopsis*. J.
Exp. Bot. 67, 157-174. doi: 10.1093/jxb/erv445

Journot-Catalino, N., Somssich, I.E., Roby, D., and Kroj, T. (2006). The transcription factors
WRKY11 and WRKY17 act as negative regulators of basal resistance in *Arabidopsis*
thaliana. Plant Cell Online 18, 3289-3302. doi: 10.1105/tpc.106.044149

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat. Methods
9, 357-359. doi: 10.1038/nmeth.1923

Lapointe, S.L., Serrano, M.S., Arango, G.L., Sotelo, G., and Cordoba, F. (1992). Antibiosis to
spittlebugs (Homoptera: Cercopidae) in accessions of *Brachiaria* spp. J. Econ. Entomol. 85,
1485-1490. doi: 10.1093/jee/85.4.1485

Lechner, E., Achard, P., Vansiri, A., Potuschak, T., and Genschik, P. (2006). F-box proteins
everywhere. Curr. Opin. Plant Biol. 9, 631-638. doi: 10.1016/j.pbi.2006.09.003

Liu, J.-J., Sturrock, R., and Ekramoddoullah, A.K.M. (2010). The superfamily of thaumatin-like
proteins: its origin, evolution, and expression towards biological function. Plant Cell Rep. 29,
419-436. doi: 10.1007/s00299-010-0826-8

Lu, F., Lipka, A.E., Glaubitz, J., Elshire, R., Cherney, J.H., Casler, M.D., et al. (2013). Switchgrass
genomic diversity, ploidy, and evolution: novel insights from a network-based SNP discovery
protocol. PLoS Genet. 9, e1003215. doi: 10.1371/journal.pgen.1003215

Ma, J.-Q., Yao, M.-Z., Ma, C.-L., Wang, X.-C., Jin, J.-Q., Wang, X.-M., et al. (2014). Construction
of a SSR-based genetic map and identification of QTLs for catechins content in tea plant
(*Camellia sinensis*). PLoS One 9, e93131. doi: 10.1371/journal.pone.0093131

Margarido, G.R.A., and Heckerman, D. (2015). ConPADE: genome assembly ploidy estimation from
next-generation sequencing data. PLoS Comput. Biol. 11, e1004229. doi:
10.1371/journal.pcbi.1004229

Mendes-Bonato, A.B., Pagliarini, M.S., Da Silva, N., and Do Valle, C.B. (2001). Meiotic instability
in invader plants of signal grass *Brachiaria decumbens* Stapf (Gramineae). Acta Sci. 23, 619-
625.

Metzler, M.C., Cutt, J.R., and Klessig, D.F. (1991). Isolation and characterization of a gene encoding
a PR-1-like protein from *Arabidopsis thaliana*. Plant Physiol. 96, 346-348.

Miles, J.W., Lapointe, S.L., Escandén, M.L., and Sotelo, G. (1995). Inheritance of resistance to
spittle bug (Homoptera: Cercopidae) in inter specific *Brachiaria* spp. hybrids. J. Econ.
Entomol. 88, 1477-1481. doi: 10.1093/jee/88.5.1477

Misra, R.C., Sandeep, Kamthan, M., Kumar, S., and Ghosh, S. (2016). A thaumatin-like protein of
Ocimum basilicum confers tolerance to fungal pathogen and abiotic stress in transgenic
Arabidopsis. Sci. Rep. 6, 25340. doi: 10.1038/srep25340

Mollinari, M., and Serang, O. (2015). Quantitative SNP genotyping of polyploids with MassARRAY
and other platforms. Methods Mol. Biol. 1245, 215-241. doi: 10.1007/978-1-4939-1966-6_17

Mukherjee, A.K., Carp, M.-J., Zuchman, R., Ziv, T., Horwitz, B.A., and Gepstein, S. (2010).
Proteomics of the response of *Arabidopsis thaliana* to infection with *Alternaria brassicicola*.
J. Proteom. 73, 709-720. doi: 10.1016/j.jprot.2009.10.005

Naumova, T.N., Hayward, M.D., and Wagenvoort, M. (1999). Apomixis and sexuality in diploid and
tetraploid accessions of *Brachiaria decumbens*. Sex Plant Reprod. 12, 43-52.

Okada, M., Lanzatella, C., Saha, M.C., Bouton, J., Wu, R., and Tobias, C.M. (2010). Complete
switchgrass genetic maps reveal subgenome collinearity, preferential pairing and multilocus
interactions. Genetics 185, 745-760. 10.1534/genetics.110.113910

Patel, R.K., and Jain, M. (2012). NGS QC toolkit: a toolkit for quality control of next generation
sequencing data. PLoS One 7, e30619. doi: 10.1371/journal.pone.0030619

Pereira, G.S., Garcia, A.A.F., and Margarido, G.R.A. A fully automated pipeline for quantitative
genotype calling from next generation sequencing data in polyploids. Submitted.

Pessoa-Filho, M., Martins, A.M., and Ferreira, M.E. (2017). Molecular dating of phylogenetic
divergence between *Urochloa* species based on complete chloroplast genomes. BMC Genom.
18, 516. doi: 10.1186/s12864-017-3904-2

Piepho, H. P., and Möhring, J. (2007). Computing heritability and selection response from
unbalanced plant breeding trials. *Genetics* 177, 1881–1888. doi: 10.1534/genetics.107.074229

Poland, J.A., Brown, P.J., Sorrells, M.E., and Jannink, J.-L. (2012). Development of high-density
genetic maps for barley and wheat using a novel two-enzyme genotyping-by-sequencing
approach. PLoS One 7, e32253. doi: 10.1371/journal.pone.0032253

Poland, J.A., and Rife, T.W. (2012). Genotyping-by-sequencing for plant breeding and genetics.
Plant Genome J. 5, 92-102. doi:10.3835/plantgenome2012.05.0005

Rajput, S.G., Santra, D.K., and Schnable, J. (2016). Mapping QTLs for morpho-agronomic traits in
proso millet (*Panicum miliaceum* L.). Mol. Breed. 36, 37. doi: 10.1007/s11032-016-0460-4

Renvoize, S.A., Clayton, W.D., and Kabuye, C.H.S. (1996). "Morphology, taxonomy, and natural
distribution of *Brachiaria* (Trin.). Griseb," in Brachiaria: Biology, Agronomy, and
Improvement, eds. J.W. Miles, B.L. Maass and C.B. Valle. (Cali, Colombia: Centro
Internacional de Agricultura Tropical (CIAT), Empresa Brasileira de Pesquisa Agropecuaria
(EMBRAPA)), 1-15.

Ripol, M.I., Churchill, G.A., da Silva, J.A.G., and Sorrells, M. (1999). Statistical aspects of genetic
mapping in autopolyploids. Gene 235, 31-41.

Salgado, L.R., Lima, R., Santos, B.F.d., Shirakawa, K.T., Vilela, M.D.A., Almeida, N.F., et al.
(2017). De novo RNA sequencing and analysis of the transcriptome of signalgrass (*Urochloa*
decumbens) roots exposed to aluminum. Plant Growth Regul. 83, 157-170. doi:
10.1007/s10725-017-0291-2

Schwarz, G. (1978). Estimating the dimension of a model. Ann. Stat. 6, 461-464.

Serang, O., Mollinari, M., and Garcia, A.A.F. (2012). Efficient exact maximum a posteriori
computation for Bayesian SNP genotyping in polyploids. PLoS One 7, e30906. doi:
10.1371/journal.pone.0030906

Shirasawa, K., Tanaka, M., Takahata, Y., Ma, D., Cao, Q., Liu, Q., et al. (2017). A high-density SNP
genetic map consisting of a complete set of homologous groups in autohexaploid sweetpotato
(*Ipomoea batatas*). Sci. Rep. 7, 44207. doi: 10.1038/srep44207

Simioni, C., and Valle, C.B. (2009). Chromosome duplication in *Brachiaria* (A. Rich.) stapf allows
intraspecific crosses. Crop Breed. Appl. Biotechnol. 9, 328-334. doi: 10.12702/1984-
7033.v09n04a07

Song, H., Wang, P., Li, C., Han, S., Zhao, C., Xia, H., et al. (2017). Comparative analysis of NBS-
LRR genes and their response to *Aspergillus flavus* in *Arachis*. PLoS One 12, e0171181. doi:
10.1371/journal.pone.0171181

Souza, J.S., Chiari, L., Simeão, R.M., de Mendonça Vilela, M., and Salgado, L.R. (2018).
Development, validation and characterization of genic microsatellite markers in *Urochloa*
species. Am. J. Plant Sci. 09, 281-295. doi: 10.4236/ajps.2018.92023

Stebbins, G. L. (1947). Types of polyploids: their classification and significance. Adv. Genet.
1, 403–429.

Stein, J., Pessino, S.C., Martínez, E.J., Rodriguez, M.P., Siena, L.A., Quarin, C.L., et al. (2007). A
genetic map of tetraploid *Paspalum notatum* Flügge (bahiagrass) based on single-dose
molecular markers. Mol. Breed. 20, 153-166. doi: 10.1007/s11032-007-9083-0

Thaikua, S., Ebina, M., Yamanaka, N., Shimoda, K., Suenaga, K., and Kawamoto, Y. (2016). Tightly
clustered markers linked to an apospory-related gene region and quantitative trait loci
mapping for agronomic traits in *Brachiaria* hybrids. Grassl. Sci. 62, 69-80. doi:
10.1111/grs.12115

Triviño, N.J., Perez, J.G., Recio, M.E., Ebina, M., Yamanaka, N., Tsuruta, S.-i., et al. (2017). Genetic
diversity and population structure of species and breeding populations. Crop Sci. 57, 1-12.
doi: 10.2135/cropsci2017.01.0045

Uknes, S., Mauch-Mani, B., Moyer, M., Potter, S., Williams, S., Dincher, S., et al. (1992). Acquired
resistance in *Arabidopsis*. Plant Cell 4, 645-656. doi: 10.1105/tpc.4.6.645

Valério, J., and Nakano, O. (1988). Influência do adulto de *Zulia entreriana* (Berg, 1879)
(Homoptera: Cercopidae) na produção e qualidade de *Brachiaria decumbens*. Pesqui.
Agropecu. Bras. 23, 447-453.

Valério, J., and Nakano, O. (1992). Sintomatologia dos danos causados pelo adulto da cigarrinha
Zulia entreriana (Berg, 1879) (Homoptera: Cercopidae) em *Brachiaria decumbens* Stapf.
Ann. Soc. Entomol. Bras 21, 95-100.

Valério, J.R., Jeller, H., and Peixer, J. (1997). Seleção de introduções do gênero *Brachiaria* (Griseb)
resistentes à cigarrinha *Zulia entreriana* (Berg) (Homoptera: Cercopidae). An. Soc. Entomol.
Bras. 26, 383-387.

Valle, C.B., Jank, L., and Resende, R.M.S. (2009). O melhoramento de forrageiras tropicais no
Brasil. Rev. Ceres 56, 460-472.

Valle, C.B., Macedo, M.C.M., Euclides, V.P.B., Jank, L., and Resende, R. (2010). "Gênero
Brachiaria," in *Plantas Forrageiras,* eds. D.M.D.F. Martuscello and J. Azevedo. (Viçosa,
Brasil: Editora UFV), 30-77.

Valle, C.B., Simioni, C., Resende, R.M.S., and Jank, L. (2008). "Melhoramento genético de
Brachiaria," in Melhoramento de Forrageiras Tropicais, eds. R.M.S. Resende, C.B. Valle and
L. Jank. (Campo Grande, MS: Embrapa Gado de Corte), 13-54.

Van Loon, L.C., Rep, M., and Pieterse, C.M.J. (2006). Significance of inducible defense-related
proteins in infected plants. Annu. Rev. Phytopathol. 44, 135-162. doi: 10.1146/
annurev.phyto.44.070505.143425

Van Loon, L.C., and Van Strien, E.A. (1999). The families of pathogenesis-related proteins, their
activities, and comparative analysis of PR-1 type proteins. Physiol. Mol. Plant Pathol. 55, 85-
97.

Van Ooijen, G., Mayr, G., Kasiem, M.M.A., Albrecht, M., Cornelissen, B.J.C., and Takken, F.L.W.
(2008). Structure–function analysis of the NB-ARC domain of plant disease resistance
proteins. J. Exp. Bot. 59, 1383-1397. doi: 10.1093/jxb/ern045

Vigna, B.B.Z., Santos, J.C.S., Jungmann, L., do Valle, C.B., Mollinari, M., Pastina, M.M., et al.
(2016). Evidence of allopolyploidy in *Urochloa humidicola* based on cytological analysis and
genetic linkage mapping. PLoS One 11, e0153764. doi: 10.1371/journal.pone.0153764

Voorrips, R.E. (2002). MapChart: software for the graphical presentation of linkage maps and QTLs.
J. Hered. 93, 77-78. doi: 10.1093/jhered/93.1.77
737

Voorrips, R.E., Gort, G., and Vosman, B. (2011). Genotype calling in tetraploid species from bi-
allelic marker data using mixture models. BMC Bioinform. 12, 172. doi: 10.1186/1471-2105-
12-172

War, A.R., Paulraj, M.G., Ahmad, T., Buhroo, A.A., Hussain, B., Ignacimuthu, S., et al. (2012).
Mechanisms of plant defense against insect herbivores. Plant Signal Behav. 7, 1306-1320.
doi: 10.4161/psb.21663

Worthington, M., Heffelfinger, C., Bernal, D., Quintero, C., Zapata, Y.P., Perez, J.G., et al. (2016). A
parthenogenesis gene candidate and evidence for segmental allopolyploidy in apomictic
Brachiaria decumbens. Genetics 203, 1117-1132. doi: 10.1534/genetics.116.190314

Wu, K.K., Burnquist, W., Sorrells, M.E., Tew, T.L., Moore, P.H., and Tanksley, S.D. (1992). The
detection and estimation of linkage in polyploids using single-dose restriction fragments.
Theor. Appl. Genet. 83, 294-300. doi: 10.1007/BF00224274

Tables

Table 1. The number of GBS-based markers after significance testing, the number of mapped
markers within each homology group (HG), and the length and marker density of each HG (cM) in
the genetic map.

HG	No. GBS-based markers*	No. mapped markers	Length of HG (cM)	Marker density (cM)
1	132	100	136.74	1.37
2	181	121	164.95	1.36
3	219	138	151.56	1.10
4	169	101	139.70	1.38
5	183	123	129.70	1.05
6	100	63	161.72	2.57
7	154	99	140.08	1.41
8	82	68	144.07	2.12
9	295	187	166.57	0.89
Total	1,515	1,000	1,335.09	1.33

755 *Markers were selected if the χ^2 goodness-of-fit statistic was higher than 0.001 for simplex and 0.01
756 for higher segregations.

758   **Table 2.** Distribution of SNPs into genotype classes.

Type	Parent 1	Parent 2	P1 dosage	P2 dosage	SNP initial number	Mapped SNP
Null (N)	AAAA	BBBB	4	0	8	0
Simplex (S)	AAAA	AAAB	4	3	1,332	591
Simplex (S)	ABBB	BBBB	1	0		
Triplex (T)	AAAA	ABBB	4	1	126	0
Triplex (T)	AAAB	BBBB	3	0		
Duplex (D)	AAAA	AABB	4	2	720	172
Duplex (D)	AABB	BBBB	2	0		
Double-simplex (SS)	AAAB	AAAB	3	3	1,013	78
Double-simplex (SS)	ABBB	ABBB	1	1		
X-double-simplex (XSS)	AAAB	ABBB	3	1	65	8
Simplex-duplex (SD)	AAAB	AABB	3	2	11	0
Duplex-simplex (DS)	AABB	ABBB	2	1	773	116
Double-duplex (DD)	AABB	AABB	2	2	192	35
Total					4,240	1,000

760 **Table 3.** Values of AIC and SIC for G_a matrix, considering different VCOV structures.

G_a matrix	Df	AIC	SIC	logREML	
ID	4	18868.02	18891.51	-9430.010	
DIAG	5	18866.02	18895.41	-9428.021	
CS	5	18848.03	**18877.39**	-9419.014	Selected
CS-Het	6	**18845.13**	18880.37	-9416.565	
FA1	7	18847.13	18888.24	-9416.565	
US	6	**18845.13**	18880.37	-9416.565	

762   **Table 4.** Summary of the predicted genotypic values, genetic ($\hat{\sigma}^2_G$) and residual ($\hat{\sigma}^2_R$) variance and
763   heritability ($\hat{H}^2_C$) for spittlebug resistance.

Predicted Means			$\hat{\sigma}^2_G$	$\hat{\sigma}^2_R$	$\hat{H}^2_C$
Min.	Mean	Max.			
56.16	70.81	85.07	68.86	382.32	0.3730

765   **Table 5.** QTL information for the spittlebug resistance trait analyzed in the *U. decumbens* progeny.

Trait	HG	QTL Position (cM)	LOD	Variance Explained (%)	SIC	Configuration
Spittlebug	1	95	4.68	4.80	85.33	Duplex (D)
Spittlebug	2	60	4.39	4.66	93.03	Duplex (D)
Spittlebug	6	23	5.49	6.24	99.07	Simplex (S)

766   HG, homology group; QTL, quantitative trait loci; LOD, logarithm of the odds; SIC, Schwarz
767   Information Criterion

Figures

Figure 1. Allele dosage in the parents and progeny, and the frequency histograms. A) Marker S.5_20487325. Red squares represent the Aaaa parent and offspring, and blue circles represent the aaaa parent and offspring. B) Marker D.2_28922605. Red squares represent the AAaa parent and offspring, blue circles represent the aaaa parent and the Aaaa offspring, and green triangles represent the aaaa offspring. C) Marker XSS.1_23891474. Red squares represent AAAa parent and offspring, blue circles represent the Aaaa parent and the AAaa offspring, and green triangles represent the Aaaa offspring.

779   **Figure 2.** Histogram of the distance between adjacent markers on the *U. decumbens* genetic map.

Figure 3. Linkage map for *U. decumbens*: homologous groups from 1 to 9. The genotype
configuration of each marker is indicated by the marker name prefix and color (Table 2). The QTLs
are identified in HG1, HG2 and HG6.

Supplementary Figure and Tables Legends

Supplementary Figure 1. Residual plots.

Supplementary Figure 2. Interval mapping (IM) for spittlebug resistance from the *U. decumbens*
mapping population in chromosomes 1 (A), 2 (B) and 6 (C). Dotted lines indicate the LOD
thresholds of 90% and 95% obtained after the permutation tests.

Supplementary Figure 3. Theoretical distribution of genotypes in the mapping population and the
distribution of individuals assigned to each genotype. The genotype distributions are nearly identical
to the theoretical distribution in a F_1 population.

Supplementary Table 1. Functional description of the sequences found in the regions of the markers
adjacent to mapped *Urochloa decumbens* QTLs for spittlebug resistance based on similarity to the
Setaria viridis genome.

Supplementary Table 2. Number of markers identified in the alignment with five reference
genomes using Tassel-GBS pipeline after allele dosage estimation with SuperMASSA software and
R data filtration analysis.